


# A simple molecular identification method of the *Thrips tabaci* (Thysanoptera: Thripidae) cryptic species complex

Péter Farkas<sup>1</sup>, Zsuzsanna György<sup>2</sup>, Annamária Tóth<sup>3</sup>, Annamária Sojnóczki<sup>1</sup>  
and József Fail<sup>1</sup> 

## Research Paper

**Cite this article:** Farkas P, György Zs, Tóth A, Sojnóczki A, Fail J (2020). A simple molecular identification method of the *Thrips tabaci* (Thysanoptera: Thripidae) cryptic species complex. *Bulletin of Entomological Research* **110**, 397–405. <https://doi.org/10.1017/S0007485319000762>

Received: 20 September 2018

Revised: 13 September 2019

Accepted: 23 October 2019

First published online: 9 December 2019

### Keywords:

CAPS marker; COI; cryptic species complex; identification; mitochondrial DNA; genetic analysis; *Thrips tabaci*

### Author for correspondence:

József Fail, Email: [fail.jozsef@kertk.szie.hu](mailto:fail.jozsef@kertk.szie.hu)

<sup>1</sup>Department of Entomology, Faculty of Horticultural Science, Szent István University, Budapest, Hungary; <sup>2</sup>Department of Genetics and Plant Breeding, Faculty of Horticultural Science, Szent István University, Budapest, Hungary and <sup>3</sup>Department of Plant Pathology, Faculty of Horticultural Science, Szent István University, Budapest, Hungary

### Abstract

The onion thrips (*Thrips tabaci* Lindeman, 1889) is a key pest of a wide range of crops because of its ecological attributes such as polyphagy, high reproduction rate, ability to transmit tospoviruses and resistance to insecticides. Recent studies revealed that *T. tabaci* is a cryptic species complex and it has three lineages (leek-associated arrhenotokous L1-biotype, leek-associated thelytokous L2-biotype and tobacco-associated arrhenotokous T-biotype), however, the adults remain indistinguishable. *T. tabaci* individuals were collected from different locations of Hungary to create laboratory colonies from each biotypes. Mitochondrial COI (mtCOI) region was sequenced from morphologically identified individuals. After sequence analysis SNPs were identified and used for CAPS marker development, which were suitable for distinguishing the three *T. tabaci* lineages. Genetic analysis of the *T. tabaci* species complex based on mtCOI gene confirmed the three well-known biotypes (L1, L2, T) and a new biotype because the new molecular evidence presented in this study suggests T-biotype of *T. tabaci* forming two distinct (sub)clades (T1 and T2). This genetic finding indicates that the genetic variability of *T. tabaci* populations is still not fully mapped. We validated our developed marker on thrips individuals from our thrips colonies. The results demonstrated that the new marker effectively identifies the different *T. tabaci* biotypes. We believe that our reliable genotyping method will be useful in further studies focusing on *T. tabaci* biotypes and in pest management by scanning the composition of sympatric *T. tabaci* populations.

## Introduction

The onion thrips (*Thrips tabaci* Lindeman, 1889) has already been known as a cosmopolitan, polyphagous insect of economic importance due to causing significant damage on cultivated crops globally, mainly on alliaceous crops, cabbage and tobacco (Jenser and Szénási, 2004; Trdan *et al.*, 2006, 2007; Shelton *et al.*, 2008; Diaz-Montano *et al.*, 2011). The onion thrips was known as a single cosmopolitan species before Zawirska (1976) recognized different biological types of *T. tabaci* ('communis type' and 'tabaci type') based on some ecological traits, distinct reproductive modes, different host range, virus transmission capability (TSWV) and morphological differences on the abdominal tergites of second-stage larvae. In the last three decades nucleic acid-based methods have been used to distinguish various population of *T. tabaci*. Klein and Gafni (1996) reported intraspecific molecular variability between *T. tabaci* populations collected from onion fields by using randomly amplified polymorphic DNA-polymerase chain reaction (RAPD-PCR) technique. Kraus *et al.* (1999) found molecular genetic differences between three different populations of *T. tabaci*, however, the computed cluster analysis was not able to distinguish the populations. Jenser *et al.* (2001) used RAPD-PCR to show differences between populations of *T. tabaci* collected from onion and tobacco. Their results indicated characteristic molecular differences among specimens from onion and tobacco. These results, at the nucleic acid level indicated the existence of subspecies within onion thrips, which colonize different host plants. Later Brunner *et al.* (2004) described that onion thrips is a cryptic species complex, based on the DNA sequences of the mitochondrial COI gene and divided the species into three lineages based on host preferences: leek-associated types (L1 and L2) and tobacco-associated type (T). Since then, this genetic divergence was confirmed in many other onion thrips populations as well (Toda and Murai, 2007; Kobayashi and Hasegawa, 2012; Jacobson *et al.*, 2013; Kobayashi *et al.*, 2013; Westmore *et al.*, 2013; Fekrat *et al.*, 2014; Li *et al.*, 2015).

In the different populations of *T. tabaci* three reproductive modes have been reported: thelytoky, arrhenotoky (Lewis, 1973; Jenser and Szénási, 2004) and deuterotoky (Nault *et al.*,

2006). In thelytoky (asexual reproduction) virgin females propagate solely females, whereas in arrhenotoky (sexual reproduction) unfertilized females produce only males and fertilized females can produce both sexes (diploid females are produced from fertilized eggs and haploid males from unfertilized eggs) (Moritz, 1997; Jenser and Szénási, 2004) and through deuterotoky virgin females produce both females and males (Moritz, 1997).

Accurate identification of tiny arthropod species such as thrips is often difficult by using exclusively morphological traits, therefore, new identification methods were introduced to separate thrips species based on PCR. Closely related taxa could be identified based on several loci. It has already been demonstrated that the internal transcribed spacer (ITS) region of the nuclear ribosomal DNA (rDNA) (Moritz *et al.*, 2002; Toda and Komazaki, 2002; Rugman-Jones *et al.*, 2006; Farris *et al.*, 2010) and mtCOI gene (Brunner *et al.*, 2002; Frey and Frey, 2004; Asokan *et al.*, 2007; Sabahi *et al.*, 2017) can be used to make distinction at species level between thrips. However, other PCR-based techniques also provide opportunity for identifying thrips species (Mehle and Trdan, 2012).

Although, to make a distinction within a species is particularly difficult and require comprehensive knowledge, the leek-associated lineages (L1 and L2) of onion thrips populations can be discriminated by their reproductive mode using PCR-based techniques. Takeuchi and Toda (2011) suggested PCR-restriction fragment length polymorphism (PCR-RFLP) while Kobayashi and Hasegawa (2012) developed PCR-strain specific primers (PCR-SSP) based on sequence differences in the mtCOI gene.

Morphological identification of different lineages within a species is very difficult and problematic. There is no external morphological difference between the adults of the *T. tabaci* 'communis' and *T. tabaci* 'tabaci' according to Zawirska's (1976) experience. However, Fekrat *et al.* (2014) observed differences between tobacco-associated and onion-associated individuals using morphometric analysis between populations from tobacco and onion fields, nonetheless, no external morphological character was singled out which could provide a definite separation.

Despite all the efforts, there is still no available method to discriminate the members of the currently known subspecies in the *T. tabaci* cryptic species complex. The present study describes a CAPS marker system targeting a part of the mitochondrial COI gene, which can be applied to identify the different biotypes in each developmental stages. A better understanding of the genetic diversity of *T. tabaci* cryptic species complex could be useful for improving pest management strategies. The recognition of the different lineages of *T. tabaci* by the presented identification tool is accurate, simple (the method requires basic laboratory skills), rapid (identification without DNA sequencing) and economic (no need for DNA sequencing equipment and purification kit). We investigated the effectiveness of the proposed method by using *T. tabaci* individuals collected in Hungary by analyzing the genetic relationship of haplotypes reported in the present work with published mtCOI haplotypes obtained from NCBI GeneBank.

## Materials and methods

### Insect collection

*T. tabaci* specimens were taken from our stock laboratory cultures. The cultures were established with the following procedure in 2013 and 2014. The thelytokous *T. tabaci* (L2) samples were collected from different plants (*Filipendula vulgaris* Moench, *Santolina*

*chamaecyparissus* L., *Lonicera caprifolium* L., *Disaphora fruticosa* (L.) Rydberg, *Coriandrum sativum* L., *Stenactis annua* (L.) Persoon, *Sorbaria sorbifolia* (L.) A. Braun) from the Botanical Garden of Szent István University (47°28'N, 19°02'E, 115 m altitude), located in Budapest and from cabbage (*Brassica oleracea* L. convar. *capitata* var. *alba*) Central Hungary (47°20'N, 18°44'E, 177 m altitude). Leek-associated arrhenotokous (L1) populations were collected from onion bulbs (*Allium cepa* L.), which were obtained from a traditional onion growing area, Makó (46°14'N, 20°28'E, 76 m altitude) Southern-Hungary. Tobacco-associated arrhenotokous (T) populations were sampled on tobacco fields (*Nicotiana tabacum* L.) in Apagy (47°57'N, 21°55'E, 118 m altitude), Pócspetri (47°52'N, 21°59'E, 133 m altitude), Encsencs (47°44'N, 22°06'E, 153 m altitude) from East-Hungary (fig. 1).

### Reproductive mode identification

The collected individuals were kept isolated individually because *T. tabaci* had already been considered as a cryptic species complex with different reproductive strategies. To create pure colonies it was desirable to have thrips specimens collected from known host plants then isolated individually in order to study their reproductive mode by identifying the sex of progeny from virgin females.


To discriminate the reproductive modes, the virgin females of the next generation produced by the originally collected living females were isolated individually in 2 ml centrifuge tubes on leaf sections and allowed to oviposit through their entire lifespan. Small leaf discs (5 mm in diameter) were provided and served as a food source and oviposition substrate to the thrips samples according to the original host plant from which the given samples were collected. If virgin females produced exclusively female progenies the reared specimens were determined as thelytokous. If only males were found in the progenies then the female was considered arrhenotokous (all progenies were kept in isolation also individually and reared to adult stage to confirm the sex). Deuterotokous reproduction was not observed at all. All thrips samples were kept in a climate chamber (SANYO, MLR-352H, Panasonic Corporation, Osaka, Japan) under the identification process in 16:8 L:D cycle at 23°C temperature.

The field-collected females were preserved in 75% ethanol until the taxonomic identification. All adult females were slide-mounted using Hoyer's solution and morphologically identified at species level by using a compound light microscope (LEICA DM LB, Leica Microsystems GmbH, Wetzlar, Germany) based on the identification guide of Mound and Kibby (1998) and the key of Moritz *et al.* (2001).


### Stock laboratory colonies

The thelytokous (L2) stock colonies used in this study were established only from the unambiguously identified thelytokous progenies of *T. tabaci* females. The arrhenotokous individuals were split into two groups: those that were originally collected from tobacco were used to establish the culture of the T lineage, whereas those collected from onion were used to establish the pure culture of the L1 lineage.

The stock thelytokous colony was maintained on cabbage head leaves, leek-associated colony on leek leaf sections, and tobacco-associated colony on whole tobacco leaves (*Nicotiana tabacum* L. 'Hevesi 9') that were planted in small glass vials filled with agar medium (Agar powder for microbiology, Merck KGaA, Darmstadt, Germany) in ventilated translucent plastic containers.



**Figure 1.** Map showing sampling sites for *T. tabaci* populations in Hungary. 1, Central Hungary - Tordas; 2, Central Hungary - Budapest the source of leek-associated thelytokous (L2) colony. 3, East Hungary - Apagy; 4, East Hungary - Pócspetri; 5, East Hungary - Encs the source of tobacco-associated arrhenotokous (T) colony; 6, Southern Hungary - Makó the source of leek-associated arrhenotokous (L1) colony.




**Figure 2.** Restriction fragment patterns of the amplified mtCOI gene of *T. tabaci* digested with PstI and PvuII endonucleases. Lane M is a 100 bp DNA ladder size marker. L1: three fragments of leek-associated arrhenotokous lineage after digestion (345 bp/274 bp/161 bp), L2: two fragments of leek-associated thelytokous lineage after digestion (619 bp/161 bp), T: undigested amplicon of tobacco-associated arrhenotokous lineage (780 bp); C: Negative control with no template DNA.

A double-layered paper towel was placed on the bottom of the plastic containers (under the plant material), which served as a hiding-place for the inactive developmental stages (prepupae, pupae) and it absorbed the surplus humidity as well. The cover of the containers was cut out to provide air ventilation and was covered by thrips-proof densely woven metal mesh to prevent

the escape of thrips. All used plant materials (leek, tobacco, cabbage) were grown by ourselves and carefully examined before introduced to the culture in order to prevent unwanted thrips contamination (thrips-free plant material). The stock colonies were kept in climate chambers separately in 16 h light and 8 h dark period at 20°C.





**Figure 3.** Minimum spanning network (MSN) calculated with mtCOI sequences using PoPART (Bandelt *et al.*, 1999). Different circle sizes are proportional to the haplotype frequency in the dataset. The numbers within circles are the haplotype designations as per Table 2. Perpendicular tick marks and the numbers in the parenthesis on the lines represent the number of the nucleotide substitutions between the linked haplotypes.

transilluminator and photographed using gel documentation system (UVP BioDoc-It™ Imaging Systems LMS-20E, Upland, USA). PCR products were purified with High Pure PCR Product Purification Kit (Merck KGaA, Darmstadt, Germany) and the amplicons were sequenced directly from both directions commercially at BaseClear B.V. (Leiden, The Netherlands).

#### Identification of restriction sites and digestion by restriction enzymes

In order to develop a discrimination method sequences available in NCBI GeneBank were used by BLASTn search tool (Altschul *et al.*, 1990) was used, but we filtered the sequences and used only those where reproductive mode was confirmed by checking the sex ratio in the virgin female's progenies.

The restriction site map was generated for each *T. tabaci* biotypes and finally the biotype-specific restriction sites were identified by SnapGene® Software (<http://www.snapgene.com/>, Free Trial version, GSL Biotech LLC).

Each digestion reaction contained 2 U *PsuI* and 2 U *PsyI* *FasDigest* enzymes (ThermoFisher Scientific), 0.6 µl 10× *FastDigest* Green Buffer (ThermoFisher Scientific), 3.3 µl unpurified PCR product and 5.7 µl nuclease-free water in 10 µl total reaction volume. The solution was digested at 37°C for 10 min according to the manufacturer's protocol.


#### Sequence analysis

The raw sequence chromatograms were visually checked, assembled and edited using Chromas Version 2.6.5. (Technelysium Pty Ltd, Australia) (<https://technelysium.com.au/>

[wp/](http://www.qiagenbioinformatics.com/)) and CLC Sequences Viewer Programme Version 7.8.1. (<https://www.qiagenbioinformatics.com/>). The identity of the species was checked by BLASTn search tool (Altschul *et al.*, 1990) (<https://blast.ncbi.nlm.nih.gov/Blast.cgi>).

Haplotype sequence reconstruction was done employing the data generated in this study for each biotype (L1, L2, T), however, other sequences were also used (NCBI GeneBank Accession: MG519810-MG519825) from our previous study (Sojnowski *et al.*, 2015). Haplotype sequences were created as follows: sequence alignments for all biotypes were imported into DNA Sequence Polymorphism Software (DnaSP version 5.10) (Librado and Rozas, 2009) to reconstruct haplotypes and generate haplotype data file. The haplotype data file was used to construct a minimum spanning network (MSN) in PoPART (Population Analysis with Reticulate Trees, <http://popart.otago.ac.nz>) (Bandelt *et al.*, 1999).

Sequence alignments were done using ClustalW algorithm and Maximum Likelihood (ML) and Maximum Parsimony (MP) methods were used to analyze the genetic relationship in MEGA6 Software (Molecular Evolutionary Genetics Analysis, Version 6) (Tamura *et al.*, 2013). JModelTest 2.1.10. (Darriba *et al.*, 2012) was used to specify the substitution model that best fits the data set for the ML analysis. The HKY + G model (Hasegawa *et al.*, 1985) was selected by JModelTest2 and MP analysis was performed under the heuristic search option. *Thrips palmi* (Accession number: AF378690) and *Thrips angusticeps* (Accession number: AF378679) sequences were obtained from NCBI GenBank and served as reference species because they are closely related species (Brunner *et al.*, 2002). Branch support was assessed with 1000 bootstrap (Felsenstein, 1985) pseudoreplicates.



**Figure 4.** Dendrogram of *T. tabaci* based on 328 bp of the mtCOI gene. The tree is generated by the Maximum Likelihood method based on the HKY+G model (Hasegawa *et al.*, 1985) using MEGA6 (Tamura *et al.*, 2013). Sequence positions containing gaps and missing data were eliminated. Bootstrap values (percentage of 1000 replicates) are shown above the branches. Accession numbers shows the sequences data of *T. tabaci* obtained from DNA databases. *Thrips palmi* (AF378690) and *Thrips angusticeps* (AF378679) served as outgroups.

## Results

### Identification of restriction patterns


Restriction enzymes were selected by the following criteria: the enzymes produce easily detectable differences in fragment size and all biotypes of *T. tabaci* can be identified in one process. Among many potential restriction endonucleases based on the summarized restriction site map, *PvuI* and *PvuII* endonucleases were tested for restriction patterns preliminary in silico digestions

of the amplified fragments of mtDNA gene by using SnapGene® Software (<http://www.snapgene.com/>, Free Trial version, GSL Biotech LLC). After the *in silico* digestion, the fragment size of the two products was detected by in silico electrophoresis. This analysis indicated that *PvuI* and *PvuII* restriction endonucleases facilitate the distinction of the three biotypes of *T. tabaci*. The cleavage with *PvuI* and *PvuII* produced banding patterns that allowed indubitable identification of biotypes according to the preliminary test by using double digestion. The leek-associated arrhenotokous type (L1) has restriction sites both for *PvuI* and for *PvuII*, which results in three different sized fragments (sized 345 bp/274 bp/161 bp), the leek-associated thelytokous type (L2) has only one restriction site producing two fragments (sized 619 bp/161 bp) because *PvuI* is a noncutter enzyme for this type. The tobacco-associated type (T) does not have restriction sites for either of the two enzymes thus it remains in one single fragment (sized 780 bp) (fig. 2) (Table 2). The 116 *T. tabaci* COI sequences used in this work were digested by restriction endonucleases and 41 sequences belonged to the leek-associated thelytokous group (L2-biotype), 36 sequences to the leek-associated arrhenotokous group (L1-biotype) and 39 sequences to the tobacco-associated arrhenotokous group (T-biotype) (Table 2).

### Sequence analysis

A new thrips-specific primer pair was designed and successfully used for amplifying a fragment of COI gene. Minimum spanning network (MSN) was created by PoPART (Bandelt *et al.*, 1999) and produced two main haplotype groups: one group is the leek-associated group, which was divided to arrhenotokous (L1) and thelytokous (L2) subgroups and the other main group is the tobacco-associated arrhenotokous group, which was also divided into two (sub)groups as T1 and T2, which two subgroups were not observed earlier. 21 different haplotypes were identified among 116 *T. tabaci* sequences. The leek-arrhenotokous individuals (L1) clustered together just like the leek-thelytokous ones and these subgroups were connected closer to each other than the leek-associated group and tobacco-associated group. The MSN revealed six haplotypes within the leek-arrhenotokous subgroup, four haplotypes within the leek-thelytokous and 11 within the tobacco-associated arrhenotokous group (fig. 3). The sequences are deposited in NCBI GeneBank under accession numbers MH782433-MH782453. L2 haplotype 2 was the most common haplotype within subgroup L2 (37 sequences belonging to this haplotype), L1 haplotype 1 within subgroup L1 (30 sequences belonging to this haplotype) and T haplotype 1 within the group T (17 sequences belonging to this haplotype). Variable nucleotide positions among 21 haplotypes are shown in Table 1. The mtCOI sequences were polymorphic at 105 nucleotide positions (14.19%) and in 21 positions resulted in amino acid replacement. The amino acid replacements occurred mainly at three haplotype sequences, such as L2 haplotype 4, L1 haplotype 4, T haplotype 9 (fig. 3).

Dendrograms were prepared based on estimation using ML, MP (Supplementary fig. S1) and Bayesian analysis (Supplementary fig. S2) supported by high bootstrap values and furthermore, the principle coordinate analysis (PCoA) plot of mtDNA data of 116 *T. tabaci* sequences (fig. 5) supported the results obtained by ML analysis resulted in largely congruent topologies, therefore, we show only the ML tree (fig. 4). The genetic tree revealed four distinct subgroups (L1, L2 and T1, T2) based on the host preferences suggested by Brunner *et al.* (2004). The group



**Figure 5.** Principle coordinate analysis (PCoA) of mtDNA data (740 bp long mtCOI sequences) of 116 Hungarian *T. tabaci* samples. Pairwise distance matrix was calculated based on the model of Tamura and Nei (1993). First and second axis explains 96% and 3.6% of variability, respectively. The shape of the dots indicates the different biotypes of *T. tabaci*: circle represents T biotype, triangle represents L1 biotype and cubic represents L2 biotype. Darker colour indicates more samples from the same type.

L can be separated to leek-associated arrhenotokous (L1) and leek-associated thelytokous (L2) subgroup, while group T can be segregated tobacco-associated arrhenotokous T1 and T2 subclades (fig. 4). The six L1 haplotype sequences clustered to leek-associated arrhenotokous subgroup L1, while four L2 haplotype sequences to the leek-associated thelytokous subgroup L2 and 11 T haplotype sequences corresponded to tobacco-associated arrhenotokous group. The position of each of our constructed 21 *T. tabaci* mtCOI haplotypes in the genetic tree (fig. 4) confirmed the predicted result of the CAPS marker for all 116 thrips individuals (Table 2). Interestingly, the group T was divided into two subgroups based on the genetic analysis and the haplotype network analysis (fig. 3 and 4), but the PCoA did not show these two groups (fig. 5).

## Discussion

A CAPS marker was developed for distinguishing mtCOI sequences of *T. tabaci* lineages and employed to genotype 116 *T. tabaci* specimens from Hungary. Using the new CAPS marker the different *T. tabaci* biotypes (L1, L2 and T) can be unambiguously discriminated, which is particularly relevant for arrhenotokous biotypes. The reliability of the CAPS marker was confirmed by genetic analysis presented in this study. The genetic structure of the tobacco-associated group within *T. tabaci* species complex is limited based on our results and the few available tobacco-associated mtCOI sequences, thus further research is required to make correct consequences and reveal the whole genetic variability.

We consider that population genetic approach using a combination of mitochondrial sequences and nuclear markers (for example ITS) are practical and effective ways to detect the reproductive barriers between *T. tabaci* lineages. Use of these genes would serve directly to detect reproductive barriers between different mitochondrial lineages because mitochondrial lineages

are inherited independently from nuclear genes and they do not undergo recombination. Sogo *et al.* (2015) found DNA sequences of two ‘new arrhenotokous’ specimens in field populations in Japan, which have the same mtCOI characteristics as thelytokous haplotypes in the L2 lineage. It is assumed that these rare arrhenotokous individuals were produced by successful mating between thelytokous females and arrhenotokous males. However, due to the maternal origin of mitochondria in the progeny, it is not possible to identify paternal gene transfer with this technique. It is worth pointing out that in its current form, our method is not able to detect incidental gene flow between thelytokous and arrhenotokous individuals. Li *et al.* (2015) published evidence about interbreeding by detecting gene flow between the progeny of leek-associated arrhenotokous (L1) males and leek-associated thelytokous (L2) females. Due to the successful mating between different lineages, the male-originated nuclear gene allele (histone H3 gene) was present in the progenies, although in only very low proportions. Nevertheless Sogo *et al.* (2015) based on different gene regions (EF, 28S, mtCOI) and Jacobson *et al.* (2016) based on microsatellite and COI markers found similar evidence with regards to potential interbreeding in field-collected populations between L1 and L2 lineages. Recently, heteroplasmy was found in thelytokous (L2) lineages in India and it was also speculated to be derived from paternal leakage, which indicates hybridization events outdoors in the *T. tabaci* species complex (Gawande *et al.*, 2017). Hybridization makes the identification of the different *T. tabaci* subspecies more complicated because of the imperfect reproductive isolation among lineages, especially if they form sympatric populations in the same habitat. Interbreeding among subspecies – besides of increasing genetic diversity – provides opportunities for hybridization in thelytokous and arrhenotokous subspecies as well. In contrast, no research has been carried out about the mating behaviours of the two arrhenotokous strains. This leaves several open questions about this topic, for instance, whether

tobacco-associated arrhenotokous males can mate with leek-associated arrhenotokous females and vice versa?

Consequently, CAPS markers need to be designed in order to measure the genotypes of nuclear genes within the *T. tabaci* cryptic species complex. However, these procedures are not considered to be difficult when dealing with several closely related biotypes or subspecies. The population genetic approach in the present study is appropriate to recognize species boundaries between closely related (sub)species, however, it is very complicated to determine the boundary between subspecies and species. The population genetic approach used in this study is relevant for studying the ecological traits and evolution of *T. tabaci*, because failure to differentiate between lineages may lead to incorrect conclusions regarding ecological traits (e.g. host associations, virus transmission capabilities, the occurrence of gene flow and insecticide resistance). Therefore, recognizing lineages within a cryptic species complex using specific markers has enormous potential to improve our ecological and evolutionary understanding of the role of insects in natural ecosystems.

**Supplementary material.** The supplementary material for this article can be found at <https://doi.org/10.1017/S0007485319000762>

**Acknowledgements.** The authors were supported by the NKFI (OTKA) K 109594 project and the bilateral co-operation projects of TET\_15-1-2016-0135 0135 and 2017-2.2.4-TÉT-AT-2017-00003. Farkas P. was supported by the Ministry for Innovation and Technology within the framework of the Higher Education Institutional Excellence Program (NKFIH-1159-6/2019) in the scope of plant breeding and plant protection researches of Szent István University. Fail J. was supported by the János Bolyai Research Scholarship of the Hungarian Academy of Sciences as well as the UNKP Bolyai + fellowship of the Hungarian Ministry of Human Capacities.

The authors acknowledge the contribution of The Department of Plant Pathology at Szent István University by providing expertise and opportunity to use the infrastructure in the laboratory.

## References

- Altschul SF, Gish W, Miller W, Myers EW and Lipman DJ (1990) Basic local alignment search tool. *Journal of Molecular Biology* **215**, 403–410. <https://doi.org/10.1006/jmbi.1990.9999>.
- Asokan R, Krishna Kumar N, Kumar V and Ranganath H (2007) Molecular differences in the mitochondrial cytochrome oxidase I (mtCOI) gene and development of a species-specific marker for onion thrips, *Thrips tabaci* Lindeman, and melon thrips, *T. palmi* Karny (Thysanoptera: Thripidae), vectors of tospoviruses (Bunyaviridae). *Bulletin of Entomological Research* **97**, 461–470. <https://doi.org/10.1017/S0007485307005147>.
- Bandelt H, Forster P and Röhl A (1999) Median-joining networks for inferring intraspecific phylogenies. *Molecular Biology and Evolution* **16** (1), 37–48. <https://doi.org/10.1093/oxfordjournals.molbev.a026036>.
- Brunner PC, Fleming C and Frey JE (2002) A molecular identification key for economically important thrips species (Thysanoptera:Thripidae) using direct sequencing and a PCR-RFLP-based approach. *Agricultural and Forest Entomology* **4**, 127–136. <https://doi.org/10.1046/j.1461-9563.2002.00132.x>.
- Brunner PC, Chatzivassiliou EK, Katis NI and Frey JE (2004) Host-associated genetic differentiation in *Thrips tabaci* (Insecta; Thysanoptera), as determined from mtDNA sequence data. *Heredity* **93**, 364–370. <https://dx.doi.org/10.1038/sj.hdy.6800512>.
- Darriba D, Taboada GL, Doallo R and Posada D (2012) JModeltest2: more models, new heuristics and high-performance computing. *Nature Methods* **9**(8), 772. <https://doi.org/10.1038/nmeth.2109>.
- De Barro PJ and Driver F (1997) Use of RAPD PCR to distinguish the B biotype from other biotypes of *Bemisia tabaci* (Gennadius) (Hemiptera: Aleyrodidae). *Australian Journal of Entomology* **36**(2), 149–152. <https://doi.org/10.1111/j.1440-6055.1997.tb01447.x>.
- Diaz-Montano J, Fuchs M, Nault BA, Fail J and Shelton AM (2011) Onion thrips (Thysanoptera: Thripidae): a global pest of increasing concern in onion. *Journal of Economic Entomology* **104**, 1–13. <https://doi.org/10.1603/EC10269>.
- Farris RE, Ruiz-Arce R, Ciomperlik M, Vasquez JD and DeLeón R (2010) Development of a ribosomal DNA ITS2 marker for the identification of the thrips, *Scirtothrips dorsalis*. *Journal of Insect Science* **10**, 1–15. <https://dx.doi.org/10.1673/031.010.2601>
- Fekrat L, Manzari S and Shishehbor P (2014) Morphometric and molecular variation in *Thrips tabaci* Lindeman (Thysanoptera:Thripidae) populations on onion and tobacco in Iran. *Journal of Agricultural Science and Technology* **16**, 1505–1516. <https://doi.org/10.1111/j.1558-5646.1985.tb00420.x>.
- Felsenstein J (1985) Confidence limits on phylogenies: an approach using the bootstrap. *Evolution* **39**, 783–791. <https://doi.org/10.1111/j.1558-5646.1985.tb00420.x>.
- Frey JE and Frey B (2004) Origin of intra-individual variation in PCR amplified mitochondrial cytochrome oxidase I of *Thrips tabaci* (Thysanoptera: Thripidae): mitochondrial heteroplasmy or nuclear integration. *Heredity* **140**, 92–98. <https://doi.org/10.1111/j.1601-5223.2004.01748.x>.
- Gawande SJ, Anandhan S, Ingle AA and Jacobson A (2017) Heteroplasmy due to coexistence of mtCOI haplotypes from different lineages of the *Thrips tabaci* cryptic species group. *Bulletin of Entomological Research* **107**(4), 534–542. <https://doi.org/10.1017/S0007485317000025>.
- Hasegawa M, Kishino H and Yano T (1985) Dating the human-ape split by a molecular clock of mitochondrial DNA. *Journal of Molecular Evolution* **22**, 160–174.
- Jacobson AL, Booth W, Vargo EL and Kennedy GG (2013) *Thrips tabaci* population genetic structure and polyploidy in relation to competency as a vector of Tomato Spotted Wilt Virus. *PLoS ONE* **8**(1), e54484. <https://doi.org/10.1371/journal.pone.0054484>.
- Jacobson AL, Nault BA, Vargo EL and Kennedy GG (2016) Restricted gene flow among lineages of *Thrips tabaci* supports genetic divergence among cryptic species groups. *PLoS ONE* **11**(9), e0163882. <https://doi.org/10.1371/journal.pone.0163882>
- Jenser G and Szénási Á (2004) Review of the biology and vector capability of *Thrips tabaci* Lindeman (Thysanoptera: Thripidae). *Acta Phytopathologica et Entomologica Hungarica* **39**, 137–155.
- Jenser G, Szénási A, Törjek O, Gyulai G, Kiss E, Heszky L and Fail J (2001) Molecular polymorphism between population of *Thrips tabaci* Lindeman (Thysanoptera: Thripidae) propagating on tobacco and onion. *Acta Phytopathologica et Entomologica Hungarica* **36**, 365–368.
- Klein M and Gafni R (1996) Morphological and molecular variations in thrips populations collected on onion plants in Israel. *Folia Entomologica Hungarica* **57**, 57–59.
- Kobayashi K and Hasegawa E (2012) Discrimination of reproductive forms of *Thrips tabaci* (Thysanoptera: Thripidae) by PCR with sequence specific primers. *Journal of Economic Entomology* **105**(2), 555–559. <https://doi.org/10.1603/EC11320>
- Kobayashi K, Yoshimura J and Hasegawa E (2013) Coexistence of sexual individuals and genetically isolated asexual counterparts in a thrips. *Scientific Report* **3**, 3286. <https://doi.org/10.1038/srep03286>.
- Kraus M, Schreiter G and Moritz G (1999) Molecular genetic studies of thrips species. In Vierbergen G and Tunc I (eds), *Proceedings of the 6th International Symposium on Thysanoptera, 27 April–1 May 1998*, Akdeniz University, Antalya, Turkey 1999. 77–80. Orkun Ozan Medya, Hizmetleri A.S. Antalya, Turkey.
- Lewis T (1973) *Thrips-Their Biology, Ecology and Economic Importance*. London: New York Academic Press.
- Li X-W, Wang P, Fail J and Shelton AM (2015) Detection of gene flow from sexual to asexual lineages in *Thrips tabaci* (Thysanoptera: Thripidae). *PLoS ONE* **10**(9), e0138353. <https://doi.org/10.1371/journal.pone.0138353>.
- Librado P and Rozas J (2009) DnaSP v5: a software for comprehensive analysis of DNA polymorphism data. *Bioinformatics (Oxford, England)* **25**, 1451–1452. <https://doi.org/10.1093/bioinformatics/btp187>.
- Mehle S and Trdan S (2012) Traditional and modern methods for the identification of thrips (Thysanoptera) species. *Journal of Pest Science* **85** (2), 179–190. <https://doi.org/10.1007/s10340-012-0423-4>.



- Moritz G** (1997) Structure, growth and development. In Lewis T (ed.), *Thrips as Crop Pests*. New York, NY: CAB International, pp. 15–64.
- Moritz G, Morris D and Mound LA** (2001) *ThripsID – Pest Thrips of the World. An Interactive Identification and Information System*. CD-ROM published by ACIAR, Australia. CSIRO Publishing, Melbourne, Australia.
- Moritz G, Paulsen M, Delker C, Picl S and Kumm S** (2002) Identification of thrips using ITS-RFLP analysis. In Marullo R and Mound LA (eds), *Thrips and Tospoviruses: Proceedings of the 7th International Symposium on Thysanoptera*. Canberra: Australian National Insect Collection CSIRO, pp. 365–367.
- Mound LA and Kibby G** (1998) *Thysanoptera. An Identification Guide*, 2nd Edn. Wallingford, UK: CAB International.
- Nault BA, Shelton AM, Gangloff-Kaufmann JL, Clark ME, Werren JL, Cabrera-LA Rosa JC and Kennedy GG** (2006) Reproductive modes in onion thrips (Thysanoptera: Thripidae) populations from New York onion fields. *Environmental Entomology* **35**, 1264–1271. <https://doi.org/10.1093/ee/35.5.1264>.
- Rugman-Jones PF, Hoddle MS, Mound LA and Stouthamer R** (2006) Molecular identification key for pest species of *Scirtothrips* (thysanoptera: Thripidae). *Journal of Economic Entomology* **99**(5), 1813–1819. <https://doi.org/10.1093/jee/99.5.1813>.
- Rychlik W** (2007) OLIGO 7 primer analysis software. In Yuryev A (eds), *PCR Primer Design. Methods in Molecular Biology™*, vol. **402**. Humana Press, pp. 35–59. <https://doi.org/10.1007>.
- Sabahi S, Fekrat L and Zakiagh M** (2017) A simple and rapid molecular method for simultaneous identification of four economically important thrips species. *Journal of Agricultural Science and Technology* **19**(6), 1279–1290.
- Shelton AM, Plate J and Chen M** (2008) Advances in control of onion thrips (Thysanoptera: Thripidae) in cabbage. *Journal of Economic Entomology* **101**(2), 438–443. <https://doi.org/10.1093/jee/101.2.438>.
- Sogo K, Miura K, Aizawa M, Watanabe T and Stouthamer R** (2015) Genetic structure in relation to reproduction mode in *Thrips tabaci* (Insecta: Thysanoptera). *Applied Entomology Zoology* **50**, 73–77. doi: 10.1007/s13355-014-0306-7. <https://doi.org/10.1007/s13355-014-0306-7>.
- Sojnoczki A, Pajtli É, Reiter D, Farkas P and Fail J** (2015) Comparative study of *Thrips tabaci* (Lindeman) cytochrome-c-oxidase gene subunit I (COI) sequences data. *Die Bodenkultur* **66**(3–4), 41–45.
- Takeuchi R and Toda S** (2011) Discrimination of two reproductive forms of *Thrips tabaci* by PCR-RFLP, and distribution of arrhenotokous *T. tabaci* in Tottori prefecture. *Japanese Journal of Applied Entomology and Zoology* **55**, 254–257. <https://doi.org/10.1303/jjaez.2011.254>.
- Tamura K and Nei M** (1993) Estimation of the number of nucleotide substitutions in the control region of mitochondrial DNA in humans and chimpanzees. *Molecular Biology and Evolution* **10**, 512–526. <https://doi.org/10.1093/oxfordjournals.molbev.a040023>.
- Tamura K, Stecher G, Peterson D, Filipski A and Kumar S** (2013) MEGA6: molecular evolutionary genetics analysis version 6.0. *Molecular Biology and Evolution* **30**, 2725–2729. <https://doi.org/10.1093/molbev/mst197>.
- Toda S and Komazaki S** (2002) Identification of thrips species (Thysanoptera: Thripidae) on Japanese fruit trees by polymerase chain reaction and restriction fragment length polymorphism of the ribosomal ITS2 region. *Bulletin of Entomological Research* **92**, 359–363. <https://doi.org/10.1079/BER2002177>.
- Toda S and Murai T** (2007) Phylogenetic analysis based on mitochondrial COI gene sequences in *Thrips tabaci* Lindeman (Thysanoptera: Thripidae) in relation to reproductive forms and geographic distribution. *Applied Entomology and Zoology* **42**, 309–316. <https://doi.org/10.1303/aez.2007.309>.
- Trdan S, Žnidar D, Vali N, Rozman L and Vidrih M** (2006) Intercropping against onion thrips, *Thrips tabaci* Lindeman (Thysanoptera: Thripidae) in onion production: on the suitability of orchard grass, lacy phacelia, and buckwheat as alternatives for white clover. *Journal of Plant Diseases and Protection* **113**(1), 24–30. <https://www.jstor.org/stable/44754883>.
- Trdan S, Valič N and Žnidarčič D** (2007) Field efficacy of deltamethrin in reducing damage caused by *Thrips tabaci* Lindeman (Thysanoptera: Thripidae) on early white cabbage. *Journal of Pest Science* **80**, 217. <https://doi.org/10.1007/s10340-007-0174-9>.
- Westmore GC, Poke FS, Allen GR and Wilson CR** (2013) Genetic and host-associated differentiation within *Thrips tabaci* Lindeman (Thysanoptera: Thripidae) and its links to Tomato spotted wilt virus-vector competence. *Heredity* **111**(3), 210–215. <https://doi.org/10.1038/hdy.2013.39>.
- Zawirska I** (1976) Untersuchungen über zwei biologische Typen von *Thrips tabaci* Lind. (Thysanoptera, Thripidae) in der VR Polen. *Archives of Phytopathology & Plant Protection* **12**, 411–422. <https://doi.org/10.1080/03235407609431780>.