

# Prospective study of family adversity and maladaptive parenting in childhood and borderline personality disorder symptoms in a non-clinical population at 11 years

C. Winsper<sup>1</sup>, M. Zanarini<sup>2</sup> and D. Wolke<sup>1,3\*</sup>

<sup>1</sup> Department of Psychology, University of Warwick, Coventry, UK

<sup>2</sup> McLean Hospital, Harvard Medical School, Belmont, USA

<sup>3</sup> Division of Mental Health and Wellbeing (Warwick Medical School), University of Warwick, Coventry, UK

**Background.** Retrospective studies have consistently indicated an association between maladaptive parenting and borderline personality disorder (BPD). This requires corroboration with prospective, longitudinal designs. We investigated the association between suboptimal parenting and parent conflict in childhood and BPD symptoms in late childhood using a prospective sample.

**Method.** A community sample of 6050 mothers and their children (born between April 1991 and December 1992) were assessed. Mothers' family adversity was assessed during pregnancy and parenting behaviours such as hitting, shouting, hostility and parent conflict across childhood. Intelligence quotient (IQ) and DSM-IV Axis I diagnoses were assessed at 7–8 years. Trained psychologists interviewed children at 11 years (mean age 11.74 years) to ascertain BPD symptoms.

**Results.** After adjustment for confounders, family adversity in pregnancy predicted BPD probable 1 to 2 adversities: odds ratio (OR) = 1.34 [95% confidence interval (CI) 1.01–1.77]; >2 adversities: OR 1.99 (95% CI 1.34–2.94) and definite 1 to 2 adversities: OR 2.48 (95% CI 1.01–6.08) symptoms. Each point increase in the suboptimal parenting index predicted BPD probable: OR 1.13 (95% CI 1.05–1.23) and definite: OR 1.28 (95% CI 1.03–1.60) symptoms. Parent conflict predicted BPD probable: OR 1.19 (95% CI 1.06–1.34) and definite: OR 1.42 (95% CI 1.06–1.91) symptoms. Within the path analysis, the association between suboptimal parenting and BPD outcome was partially mediated by DSM-IV diagnoses and IQ at 7–8 years.

**Conclusions.** Children from adverse family backgrounds, who experience suboptimal parenting and more conflict between parents, have poor cognitive abilities and a DSM-IV diagnosis, are at increased risk of BPD symptoms at 11 years.

Received 18 May 2011; Revised 12 February 2012; Accepted 29 February 2012; First published online 5 April 2012

**Key words:** Avon Longitudinal Study of Parents and Children (ALSPAC), borderline personality disorder, DSM-IV Axis I diagnoses, parent conflict, suboptimal parenting.

## Introduction

Maladaptive experiences during childhood have been consistently linked with borderline personality disorder (BPD) including: abuse and neglect (Guzder *et al.* 1999; Zanarini *et al.* 2006), parent hostility and resentment (Hooley & Hoffman, 1999; Johnson *et al.* 2006) and exposure to domestic violence and parent conflict (Herman *et al.* 1989; Weaver & Clum, 1993). Most studies have been retrospective, however,

with concomitant methodological issues, such as the tendency of patients with BPD to misinterpret or misreport past experiences with family members (Bailey & Shriver, 1999). Furthermore, domestic conflict and child maltreatment usually occur in family environments characterized by multiple risk factors (Fergusson *et al.* 2006) difficult to disentangle with retrospective designs.

A series of prospective, longitudinal studies has revealed an association between abuse, neglect, parenting and BPD features (Johnson *et al.* 1999, 2000, 2001, 2006). Associations were focused on scales of personality disorder symptoms assessed in early adulthood, however, rather than on a collection of symptoms

\* Address for correspondence: D. Wolke, Ph.D., Department of Psychology, University of Warwick, Coventry CV4 7AL, UK.  
Email: DWolke@warwick.ac.uk

comparable in composition with a DSM-IV BPD diagnosis. Subsequently, large prospective longitudinal studies are now necessary to identify younger individuals with comparable symptom constellations (including subsyndromal levels of manifestation) to those identified in adult BPD. Such studies are challenging due to low base rates of BPD and protracted duration before formal diagnosis, typically during early adulthood, is made. However, BPD is unlikely to suddenly appear in early adulthood; rather, it may be considered within a developmental trajectory as the end point following the appearance of BPD symptoms during childhood or adolescence.

The importance of early identification of such symptoms, as manifest in a childhood phenotype, has been highlighted, both for the facilitation of intervention programmes (Chanen *et al.* 2008) and delimitation of aetiological factors (Geiger & Crick, 2001). Furthermore, BPD assessments for children have been developed (Crick *et al.* 2005; Rogosch & Cicchetti, 2005), and it appears that BPD-related features may be identified as early as 6 years of age, and remain relatively stable over time (Stepp *et al.* 2010). Nevertheless, it has not been ascertained whether factors associated with BPD in adulthood are also associated with BPD symptoms during late childhood.

In the current study we investigated whether exposure to family adversity and maladaptive parent behaviour, during preschool and school periods, was predictive of BPD probable and definite symptoms (five or more) in late childhood. Additionally, the developmental pathways through which this association manifests were explored by considering the mediating effects of potential markers: Axis I DSM-IV diagnoses and intelligence quotient (IQ) at age 7–8 years.

## Method

### Participants

The Avon Longitudinal Study of Parents and Children (ALSPAC) is a birth cohort study, set in the UK, examining the determinants of development, health and disease during childhood and beyond (Golding *et al.* 2001). A total of 14541 women were enrolled, provided they were resident in Avon while pregnant, and had an expected delivery date between 1 April 1991 and 31 December 1992. As shown in Fig. 1, 13971 children, alive at 12 months, formed the original cohort. From the first trimester of pregnancy parents completed postal questionnaires about themselves and the study child's health and development. Children were invited to attend annual assessment clinics, including face-to-face interviews,

and psychological and physical tests from 7 years onwards.

During the planning stage of the study, children from the Avon area were compared with 13135 children from across the UK, participating in the Child Health and Education Study, on a number of demographic variables. Results suggested that the Avon population was fairly similar to that of the whole of Great Britain (Golding *et al.* 2001). There were 11510 children living in the study area and eligible for invitation to the 11-year annual assessment clinic; 6423 attended and started the interview session, incorporating the BPD questions (Fig. 1), though 373 of these children were excluded because they did not answer at least eight of the nine BPD questions. This study is, therefore, based on 6050 children (age range 10.4–13.6 years, mean age 11.7 years).


### Ethical approval

Ethical approval was obtained from the ALSPAC Law and Ethics committee and the local research ethics committees.

### BPD features interview

Borderline features were assessed using a face-to-face semi-structured interview: the UK Childhood Interview for DSM-IV Borderline Personality Disorder (UK-CI-BPD; Zanarini *et al.* 2004), based on the borderline module of the Diagnostic Interview for DSM-IV Personality Disorders (Zanarini *et al.* 1996), which is a widely used semi-structured interview for all DSM-IV Axis II disorders. The inter-rater and test-retest reliability of the DSM-III, DSM-III-R and DSM-IV versions of this measure have all proven to be good to excellent (Zanarini *et al.* 2000; Zanarini & Frankenberg, 2001). The UK-CI-BPD was adapted from the CI-BPD (US version), with small changes in wording making it appropriate for a UK sample, e.g. 'being angry' was changed to 'being cross'. The convergent validity of the CI-BPD was investigated using 171 adolescents (boys and girls) 13–17 years of age; 111 met criteria for BPD and 60 were normal comparison subjects. A Spearman's  $\rho$  of 0.89 was obtained when comparing a dimensional score for BPD on the CI-BPD and the total score on the Revised Diagnostic Interview for Borderlines.

The UK-CI-BPD differs from the adult interview in three ways: (1) the language is simpler; (2) two forms of impulsivity are omitted (reckless driving and promiscuity) due to lack of developmental appropriateness; and (3) the childhood interview is more structured than the adult version, with the answer to each question, and not just the rating for each of the



**Fig. 1.** Flow of participants from pregnancy to 11-year assessment in the cohort study the Avon Longitudinal Study of Parents and Children (ALSPAC). <sup>a</sup> Includes multiple births (195 twins, three triplets, one quadruplet). <sup>b</sup> An additional 359 children were invited who were previously missed pregnancies, born and residing in the Avon area.

nine criteria, entered into the dataset (Zanarini *et al.* 2011).

The inter-rater reliability ( $\kappa$ ) of the UK-CI-BPD, assessed from taped interviews of 30 children, ranged from 0.36 to 1.0 (median value 0.88), and 86% of the  $\kappa$  values were within the excellent range of  $>0.75$  (Zanarini *et al.* 2011).

The UK-CI-BPD is the first semi-structured interview designed to assess DSM-IV BPD in latency-aged children. Similar to DSM-IV criteria, the interview consists of nine sections: intense inappropriate anger; affective instability; emptiness; identity disturbance; paranoid ideation; abandonment; suicidal or self-mutilating behaviours; impulsivity and intense unstable relationships. Once a trained assessor had explored each section, a judgment was made as to whether each symptom was definitely present,

probably present or absent. A symptom was classed as definitely present if it occurred daily or approximately 25% of the time, and probably present if it had occurred repeatedly, but did not meet the criterion for definitely present.

Two outcome variables were constructed for use in the logistic regression analyses: BPD symptoms probably present (symptoms present less than daily or 25% of the time) and BPD symptoms definitely present, both of which were based on the presence of five or more symptoms. Diagnosis of BPD according to the DSM-IV is based on the presence of five or more definite features; thus the probable BPD outcome represents a dimensional adjunct to the traditional categorical approach, i.e. children with five or more (categorical) subsyndromal symptoms (dimensional) are identified (Kraemer, 2007).

### *Sociodemographic and birth variables*

The mother-reported sociodemographic information during the antenatal period included marital status (married *versus* single); home ownership (home owner *versus* rented); parent social class (based on the highest of the mother's or partner's occupational social class: dichotomized into non-manual *versus* manual); and maternal education, dichotomized into below O level *versus* O level or above (O levels being the standard school-leaving qualifications at age 16 years in the UK until recently). The ethnic origin of the child (white *versus* black or minority ethnic) and birth weight were obtained from birth records. Birth weight was dichotomized into  $\leq 2499$  g (low birth weight) and  $\geq 2500$  g.

### *Exposure variables: family adversity, suboptimal parenting and parent conflict*

#### *Family adversity*

Multiple family risk factors were indicated using the Family Adversity Index (FAI; Bowen *et al.* 2005), which consists of 18 items taken from questionnaires administered throughout pregnancy (8, 12, 18 and 32 weeks gestation) (see Table 1 for more details). The FAI consists of items pertaining to young maternal age at first pregnancy (<17 years) or birth of study child (<20 years); housing (e.g. inadequacy: overcrowding or periods of homelessness); financial difficulties; problematic partner relationship; maternal affective disorder (depression, anxiety, suicidality); substance abuse (drugs or alcohol); or involvement in crime (i.e. in trouble with police or convictions). For the current analysis the item reflecting partner cruelty (emotional or physical) was removed from the FAI to prevent confounding with the domestic violence predictor variables. The remaining adversity items were summed and trichotomized into: none (no adversity); mild (one or two adversities) and severe (more than two adversities).

#### *Suboptimal parenting index*

Selection of the suboptimal parenting predictors was based on a previous study (Waylen *et al.* 2008), which factor analysed questions pertaining to maternal attitudes, behaviours and feelings within the ALSPAC cohort. Three factors were evidenced, reflecting suboptimal parenting (hostility, resentment and hitting/shouting), which were found to be predictive of a variety of health outcomes during mid-childhood. These factors have been prospectively linked to personality disorders (and BPD features) within the literature (Johnson *et al.* 2006), and thus were combined to create a suboptimal parenting index.

Scales assessing parent behaviour, as reported by the mother, were dichotomized, indicating whether the maladaptive behaviour was present or absent. Where available, variables were constructed for the preschool (birth to up to 5 years) and school (5–8 years) periods. The suboptimal parenting index was constructed by summing seven items across the preschool and school periods to create an index of increasing exposure to suboptimal parenting on a scale of 0–7. Items were: hitting (preschool, school); shouting (preschool, school); hostility (preschool, school); and resentment (preschool).

Maternal hitting and shouting were indicated by the following two items: 'When you are at home with your child how often do you slap him?' and 'When you are at home with your child how often do you shout at him?' (Waylen *et al.* 2008). For the preschool period (24 and 42 months), hitting was coded as present if it occurred daily or every week at either time point, and shouting if it occurred daily at either time point. For the school period (77 months), hitting was recorded as present if reported often or sometimes, and shouting if reported often. We used less stringent criteria for the school period to reflect the observed reduction in hitting and shouting, as the child grows older (Hyman, 1997).

Hostility and resentment were constructed from a number of items loading on two distinct factors (Waylen *et al.* 2008). Preschool hostility items included: 'mum feels that whining makes her want to hit child' (21 months); 'mum often irritated by child' (47 months); 'mum has battle of wills with child' (47 months); and 'child gets on mum's nerves' (47 months). Preschool hostility was classed as present if reported in three or all items. Preschool resentment items included: 'mum dislikes mess from child' (47 months); 'mum feels unbearable when child cries' (21 months); 'mum feels child's desires cause anger' (21 months); and 'mum feels has no time alone' (33 months).

Preschool resentment was classed as present if reported for two or more items. For the school period, only hostility items were available: 'mum often irritated by child' (85 months); 'mum has battle of wills with child' (85 months); 'child gets on mum's nerves' (85 months). School hostility was considered present if answered positively for all three items.

#### *Conflicting partnership index*

Domestic violence and conflicting partnership measures were chosen according to reported prospective associations with negative child outcomes, generally (Kitzman *et al.* 2003), and BPD, specifically, in retrospective studies (Herman *et al.* 1989; Weaver & Clum,

**Table 1.** Individual items comprising the Family Adversity Index<sup>a</sup>

Main categories and subcategories	Individual items	Number of points per subcategory
Age of mother (maximum 1 point)	First pregnancy at <17 years or birth study child at <20 years	1 point
Housing (maximum 3 points)		
Adequacy	Crowding index (< one room per person) or Became homeless (yes)	1 point
Basic living	Bath/shower (no) or Hot water (no) or Indoor toilet (no) or Kitchen (no)	1 point
Defects/infestation	Mould (yes) or Roof leaks (yes) or Rats, mice or cockroaches (yes)	1 point
Education (maximum 1 point)	Maternal (none) or Paternal (none)	1 point
Financial status (maximum 1 point)	Financial difficulties <sup>b</sup> (yes)	1 point
Critical partner relationship (maximum 4 points)		
Status	Have partner (no)	1 point
Affection	Intimate bond <sup>b</sup> (no) or Affection <sup>b</sup> (no) or Aggression <sup>b</sup> (yes)	1 point
Partner cruelty <sup>c</sup>	Physical or Emotional	1 point
Support	Emotional (no) or Practical (no) or Partner might leave (yes)	1 point
Family (maximum 2 points)	Family size (more than three children)	1 point
	Taken into care/at-risk register	1 point
Social network (maximum 2 points)	Emotional support (no)	1 point
	Practical/financial support (no)	1 point
Psychopathology of mother (maximum 1 point)	Anxiety <sup>d</sup> or depression <sup>d</sup> or attempted suicide (yes)	1 point
Substance abuse (maximum 1 point)	Drugs (yes) and/or Alcohol (yes)	1 point
Crime (maximum 2 points)	In trouble with police (yes)	1 point
	Convictions (yes)	1 point

<sup>a</sup> Maximum of 18 points in total.

<sup>b</sup> Each item was derived from a series of questions.

<sup>c</sup> These two items were removed from the index to prevent confounding with the domestic violence variable.

<sup>d</sup> Derived from the Crown Crisp Inventory.

1993). The parent conflict index was constructed across the preschool and school periods from five items, on a scale of 0–5, reflecting increasing exposure to conflict between primary caregivers. Items were: conflicting partnership (preschool, school); partner broken or thrown things (preschool); physically hurt by partner (preschool); and emotional domestic violence (preschool).

Physical and emotional domestic violence variables (Bowen *et al.* 2005) were available for the preschool period only. Two physical domestic violence variables were constructed: physically hurt by partner and partner broken or thrown things. The variable 'physically hurt by partner' was constructed from the two items 'physically hurt by partner' (8, 21, 33 and 47 months) and 'slapped or hit by partner' (21 and 33



months), which were coded as present if the mother responded yes to one or more of the six items. The variable 'partner broken or thrown things' (21 and 33 months) was considered present if answered with yes at either time point. An emotional domestic violence variable was constructed from the item 'your partner was emotionally cruel to you' (8, 21, 33 and 47 months) (Bowen *et al.* 2005). Emotional domestic violence was considered present if reported at one or more time points.

#### *Conflicting partnership*

A conflicting partnership variable was derived for the preschool (33 months, or 22 months if the 33-month response was missing) and school (73 months) periods. It was constructed from the following items: 'mum and partner argued'; 'not speaking to partner for more than 30 min'; 'one of you walking out of the house'; and 'shouting or calling partner names'. For the preschool and school periods, each of these items was dichotomized; if either the mother, her partner or both parties had engaged in the behaviour, the item was coded as present. Conflicting partnership was considered positive if reported in three or all four items.

#### *Potential confounders or markers for BPD symptoms*

Study child IQ was assessed with the Wechsler Intelligence Scale for Children III (UK version) (Wechsler *et al.* 1992) during the focus at 8 years clinic. DSM-IV psychiatric diagnoses were derived at 7–8 years using the Development and Well-Being Assessment (DAWBA; Goodman *et al.* 2000), completed by parents and teachers. Teachers were asked to complete the DAWBA for all the children in their class with a birth date between April 1991 and December 1992. The teacher completion rate was 5155/10 431 eligible children, and the mother completion rate was 8269/11 251. Mother and teacher reports were combined (where available), otherwise the mother report only was used. The diagnoses were made using a DSM-IV-TR algorithm, and reviewed by two experienced child psychiatrists (Robert Goodman and Tamsin Ford). The DAWBA has been validated for Axis I diagnoses and shown to have utility as a clinical assessment tool (Goodman *et al.* 2000) (for further information, see <http://www.dawba.com/>). A dichotomous variable, indicating the presence of any major Axis I disorder [attention-deficit hyperactivity disorder (ADHD), conduct disorder, oppositional defiant disorder, depression or anxiety] was constructed.

#### *Statistical analysis*

Initial analyses were carried out with SPSS version 17 statistical software (SPSS, Inc., USA). Selective drop-out was determined by comparing those who completed the borderline interview with those lost to follow-up (Table 2). Odds ratios (ORs) and 95% confidence intervals (CIs) were computed to test for gender differences in parenting variables and BPD probable and definite symptoms (Table 3). Crude associations between family adversity, maladaptive parenting and BPD probable and definite symptoms were computed. Associations were then adjusted for age and gender, then, additionally, DSM-IV diagnoses and IQ. ORs with 95% CIs are reported for the preschool and school periods, respectively (Tables 4 and 5). Path analysis was carried out, using Mplus version 6 (<http://www.statmodel.com/>), to elucidate the direct and indirect relationships between exposure to family adversity, suboptimal parenting and parent conflict, manifestation of DSM-IV Axis I diagnoses, IQ and the BPD outcome. A categorical ordinal BPD outcome was utilized in the path analysis, reflecting increasing severity of BPD [less than five symptoms (92.7%); five or more probable symptoms (6.4%); five or more definite symptoms (0.9%)]. Mplus version 6 software is suitable for the analysis of categorical outcomes, producing estimates in the form of probit coefficients. Probit coefficients indicate the strength of relationship between predictor variables and probability of group membership. They represent the difference that a one-unit change in the predictor variable makes in the cumulative normal probability of the outcome variable (Lee *et al.* 2007). For ordinal outcomes one coefficient per predictor is produced. This may be interpreted in the same way as a continuous dependent variable, as an ordinal dependent variable is comparable with a continuous latent response variable, which exceeds thresholds to give various outcome categories (Muthén, 1998–2004).

#### **Results**

##### *Differences between participants with and without the completed borderline interview*

The frequencies of sociodemographic factors, psychiatric diagnoses and IQ are shown for ALSPAC participants with and without borderline interviews in Table 2. Those lost to follow-up were more often boys, ethnic minority children, of low birth weight, born to single mothers of lower education level, from rented properties and with parents in manual jobs. They were more likely to have been born into family adversity, and have had psychiatric diagnoses at 7–8 years. Children who dropped out had a lower IQ at 8 years.

**Table 2.** Drop-out analysis comparing those where BPD symptoms interview was not available with those who completed the borderline interview at age 11 years

Characteristic	BPD interview not available, <i>n</i> (%)	BPD interview available, <i>n</i> (%)	BPD interview available <i>v.</i> not available, OR (95% CI) <sup>a</sup>
Gender			
Male	4328 (59.6)	2938 (40.4)	
Female	3669 (54.1)	3112 (45.9)	1.25 (1.17–1.34)
Ethnicity			
White	5967 (51.9)	5541 (48.2)	
Black and minority ethnic	395 (64.6)	216 (35.4)	0.59 (0.49–0.69)
Birth weight			
>2499 g	7370 (56.4)	5707 (43.6)	
<2500 g	517 (65.4)	273 (34.6)	0.68 (0.58–0.79)
Marital status			
Single	2206 (66.8)	1095 (33.2)	
Married	5031 (51.1)	4821 (48.9)	1.93 (1.77–2.10)
Home ownership			
Mortgage	4701 (49)	4901 (51)	
Rented	2532 (72.6)	958 (27.4)	0.36 (0.33–0.39)
Education of mother			
Below O level	2476 (66.2)	1262 (33.8)	
O level or above	4142 (47.5)	4577 (52.5)	2.17 (2.0–2.35)
Social class			
Non-manual	2729 (46.4)	3152 (53.6)	
Manual	3210 (56.9)	2430 (43.1)	0.66 (0.60–0.70)
FAI			
None	2565 (47.9)	2791 (52.1)	
Moderate; one or two adversities	3125 (56.0)	2454 (44.0)	0.72 (0.67–0.78)
Severe: more than two adversities	1577 (68.7)	717 (31.3)	0.41 (0.37–0.46)
DSM-IV Axis I diagnoses by DAWBA			
None	2791 (36.6)	4839 (63.4)	
At least one diagnosis	257 (45.5)	308 (54.5)	0.69 (0.58–0.82)
Mean IQ (s.d.) <sup>b</sup>	100.6 (17.2)	105.8 (15.8)	1.02 (1.01–1.03)

BPD, Borderline personality disorder; OR, odds ratio; CI, confidence interval; FAI, Family Adversity Index; DSM-IV, Diagnostic and Statistical Manual of Mental Disorders, 4th edn; DAWBA, Development and Well-Being Assessment; IQ, intelligence quotient.

<sup>a</sup> None of the 95% CIs includes 1.00.

<sup>b</sup> For BPD interview not available, *n* = 1669; for BPD interview available, *n* = 4787.

Thus, participants remaining in the analysis were less severely disadvantaged than those who dropped out.

### Frequency of BPD and maladaptive parenting variables

Table 3 reports the frequencies of BPD probably and definitely present and parenting variables (total and by gender). Of the ALSPAC cohort, 6.4% had five or more probable, and 0.9% had five or more definite, symptoms at 11 years. These findings are largely concordant with a previous community study, which reported that 7.8% of 9- to 19-year-olds had moderate BPD, and 3% had severe BPD (Bernstein *et al.* 1993),

with the lower values in the present study possibly attributable to the younger age of the cohort.

Hitting and shouting were common during the preschool period, becoming rarer during the school period (Table 3). Significantly more boys than girls were hit during both periods and shouted at during the preschool period. Hostility and resentment did not differ according to the gender of the study child. Domestic violence was reported for the preschool period only, with emotional domestic violence more common than being physically hurt by a partner and a partner having broken or thrown things. There were no gender differences for living in a household with domestic violence. Conflicting partnerships during

**Table 3.** Frequencies of BPD diagnosis (probable and definite) and maladaptive parenting variables shown for the total sample and by gender

Borderline diagnosis or parenting variables	Total, n (%)	Girls, n (%)	Boys, n (%)	Girls v. boys, OR (95% CI)
<b>Borderline diagnosis</b>				
Probable				
0 No	5606 (93.5)	2882 (93.5)	2724 (93.5)	
1 Yes	389 (6.5)	200 (6.5)	189 (6.5)	1.00 (0.81–1.23)
Definitely				
0 No	5995 (99.1)	3082 (99.0)	2913 (99.1)	1.13 (0.67–1.93)
1 Yes	55 (0.9)	30 (1.0)	25 (0.9)	
<b>Hitting and shouting</b>				
Preschool hitting				
0 No	3235 (56.6)	1783 (60.9)	1452 (52.1)	0.7 (0.63–0.78) <sup>a</sup>
1 Present	2479 (43.4)	1145 (39.1)	1334 (47.9)	
Preschool shouting				
0 No	3125 (54.7)	1703 (58.1)	1422 (51)	0.75 (0.68–0.83) <sup>a</sup>
1 Present	2593 (45.3)	1227 (41.9)	1366 (49)	
School hitting				
0 No	1868 (61.6)	999 (64.3)	869 (58.8)	0.79 (0.83–0.92) <sup>a</sup>
1 Present	1163 (38.4)	544 (35.7)	609 (41.2)	
School shouting				
0 No	2644 (84.7)	1349 (84.9)	1295 (84.5)	0.97 (0.8–1.81)
1 Present	477 (15.3)	240 (15.1)	237 (15.5)	
<b>Parental attitudes</b>				
Preschool hostility				
0 No	4918 (85.8)	2526 (85.9)	2392 (85.8)	0.99 (0.86– 1.15)
1 Present	812 (14.2)	416 (14.1)	396 (14.2)	
School hostility				
0 No	4595 (89.5)	2341 (89.2)	2254 (89.8)	1.06 (0.89–1.27)
1 Present	539 (10.5)	283 (10.8)	256 (10.2)	
Preschool resentment				
0 No	5011 (86.6)	2597 (87.3)	2414 (85.9)	0.89 (0.76–1.03)
1 Present	776 (13.4)	379 (12.7)	397 (14.1)	
<b>Domestic violence</b>				
Physically hurt				
0 No	5527 (93.7)	2849 (94)	2678 (93.5)	0.93 (0.75–1.14)
1 Present	369 (6.3)	183 (6.0)	186 (6.5)	
Broken or thrown				
0 No	5283 (95.7)	2724 (95.8)	2559 (95.5)	0.92 (0.71–1.2)
1 Present	240 (4.3)	119 (4.2)	121 (4.5)	
Emotional				
0 No	4828 (81.9)	2471 (81.5)	2357 (82.3)	1.05 (0.92–1.20)
1 Present	1067 (18.1)	560 (18.5)	507 (17.7)	
<b>Conflicting partnership</b>				
Preschool				
0 No	4265 (77.2)	2181 (76.7)	2084 (77.7)	1.06 (0.93–1.2)
1 Present	1262 (22.8)	663 (23.3)	599 (22.3)	
School				
0 No	3890 (80.7)	1976 (80.5)	1914 (81.0)	
1 Present	929 (19.3)	480 (19.5)	449 (19.0)	1.04 (0.90–1.2)

BPD, Borderline personality disorder; OR, odds ratio; CI, confidence interval.

<sup>a</sup> 95% CI does not include 1.



both periods did not differ according to the gender of the child.

#### ***Associations between maladaptive parenting and BPD symptoms***

Table 4 shows the associations between family adversity, maladaptive parenting and BPD probable symptoms. The Table shows the crude associations, the associations after controlling for age and gender, and the associations after controlling for age, gender, DSM-IV diagnoses and IQ.

Family adversity (one or two items; more than two items), hitting (preschool), hostility (school), partner breaking or throwing things, emotional domestic violence and conflicting partnership (preschool and school) were all significantly associated with BPD probable symptoms. After controlling for confounders, conflicting partnership (preschool and school) was no longer predictive of BPD probable symptoms. Suboptimal parenting and parent conflict led to higher odds of BPD probable symptoms after adjusting for confounders.

Table 5 shows the associations between family adversity, maladaptive parenting and BPD definite symptoms. Hitting (preschool), resentment, hostility (preschool and school), emotional domestic violence, physically hurt by partner and conflicting partnership (school) were predictive of BPD definite symptoms. After controlling for confounders, hostility (school), emotional domestic violence, physically hurt by partner and conflicting partnership (school) remained significantly predictive of BPD definite symptoms. Suboptimal parenting and parent conflict remained predictive of BPD definite symptoms after controlling for confounders.

#### ***Predictive associations between family adversity, parenting variables, potential mediators and BPD probable and definite symptoms***

The predictive associations between family adversity, parenting variables, mediators and BPD probable and definite symptoms are shown in Supplementary Table S1(A, B, C). These associations were tested according to time ordering; therefore, family adversity was considered a predictor, while Axis I DSM-IV diagnoses (DAWBA), IQ and BPD were considered outcomes of family adversity and parenting variables. Univariate analysis indicated that family adversity was predictive of suboptimal parenting, parent conflict, DSM-IV diagnosis, IQ and BPD symptoms probable and definite (Supplementary Table S1A). Suboptimal parenting and parent conflict were predictive of DSM-IV diagnoses, IQ and BPD probable and definite

symptoms (Supplementary Table S1B). DSM-IV diagnoses were predictive of BPD probable symptoms and IQ was predictive of BPD definite symptoms (Supplementary Table S1C). These findings are consistent with a pathway model in which family adversity is a precursor for suboptimal parenting and parent conflict, leading to DSM-IV diagnoses and lower IQ (child markers) culminating in BPD symptoms.

#### ***Path analysis***

The path model incorporated the family adversity, suboptimal parenting and parent conflict indices as predictors. IQ and DSM-IV diagnoses were entered as potential mediators, while gender was entered as a control. Model fit indices indicated good fit ( $\chi^2 = 11.58$ ,  $p = 0.00$ , root mean square error of approximation = 0.02, comparative fit index = 0.99). Fig. 2 shows the unstandardized and standardized (in parentheses) estimates of the direct path coefficients between the various predictor and mediating variables. Non-significant paths ( $p > 0.05$ , one-tailed) are not shown.

The direct relationships between family adversity (one or two adversities; more than two adversities), suboptimal parenting, parent conflict, DSM-IV diagnoses, IQ and BPD outcome at 11 years were significant. Direct and indirect path coefficients to the BPD outcome are shown in Table 6. The association between suboptimal parenting and BPD outcome was partially mediated by DSM-IV diagnoses and IQ at 7–8 years.

#### ***Discussion***

In line with previous research, we found that suboptimal parenting and parent conflict were more likely within families experiencing adversities, ranging from poverty and overcrowding to mental health problems (Fergusson *et al.* 2006). Family adversity was assessed in pregnancy, thereby excluding reverse-causality effects of parenting, or a challenging child, on family adversity. Family adversity had a direct impact on BPD symptoms at 11 years of age, and indirect effects via suboptimal parenting, parent conflict, poorer cognitive functioning and DSM-IV diagnoses of the child. Furthermore, there was a dose–response effect with an increase in family adversity and maladaptive parenting severity leading to increased odds of BPD symptoms. This indicates that children exposed to higher levels of family adversity and maladaptive parenting were at heightened risk of developing BPD symptoms. The direct impact of family adversity in pregnancy may be due to continued adversity throughout

**Table 4.** Associations between family adversity, maladaptive parenting and BPD probable status, showing crude associations, adjustment for age and gender, and additionally DSM-IV diagnosis and IQ

Exposure	Subgroup	BPD status, n (%)	OR (95 % CI)	OR, with controls <sup>a</sup> (95 % CI)	OR (with controls) <sup>b</sup> (95 % CI)		
Family adversity	No items	141 (5.1)					
	1 or 2 items	163 (6.7)	1.35 (1.07–1.70) <sup>c</sup>	1.35 (1.07–1.71) <sup>c</sup>	1.34 (1.01–1.77) <sup>c</sup>		
	More than 2 items	77 (11.0)	2.30 (1.72–3.08) <sup>c</sup>	2.32 (1.74–3.11) <sup>c</sup>	1.99 (1.34–2.94) <sup>c</sup>		
Hitting and shouting	Preschool	Hitting	No (3235)				
		Yes (2479)	185 (5.8)	1.31 (1.06–1.62) <sup>c</sup>	1.31 (1.06–1.62) <sup>c</sup>	1.43 (1.10–1.86) <sup>c</sup>	
	Shouting	No (3125)	185 (6.0)				
		Yes (2593)	182 (7.1)	1.20 (0.97–1.48)	1.20 (0.97–1.49)	1.22 (0.94–1.58)	
	School	Hitting	No (1868)	118 (6.4)			
			Yes (1163)	88 (7.6)	1.22 (0.91–1.62)	1.21 (0.91–1.62)	1.24 (0.88–1.74)
Shouting		No (2644)	181 (6.9)				
Yes (477)	34 (7.2)	1.04 (0.71–1.53)	1.05 (0.72–1.53)	1.11 (0.71–1.74)			
Parental attitude	Preschool	Hostility	No (4918)	298 (6.1)			
			Yes (812)	63 (7.9)	1.32 (0.98–1.75)	1.32 (0.99–1.75)	1.49 (1.07–2.08) <sup>c</sup>
		Resentment	No (5011)	314 (6.3)			
	Yes (776)	53 (6.9)	1.11(0.82–1.50)	1.11 (0.82–1.50)	1.17 (0.81–1.67)		
	School	Hostility	No (4595)	268 (5.9)			
			Yes (539)	47 (9.0)	1.58 (1.14–2.18) <sup>c</sup>	1.58 (1.14–2.19) <sup>c</sup>	1.56 (1.06–2.29) <sup>c</sup>
	Suboptimal parenting index <sup>d</sup>		1.12 (1.04–1.20) <sup>c</sup>	1.12 (1.04–1.19) <sup>c</sup>	1.13 (1.05–1.23) <sup>c</sup>		
Conflict	Domestic violence	Emotional	No (4828)	287 (6.0)			
			Yes (1067)	87 (8.3)	1.42 (1.11–1.83) <sup>c</sup>	1.43 (1.11–1.83) <sup>c</sup>	1.52 (1.12–2.06) <sup>c</sup>
	Physically hurt by partner	No (5527)	345 (6.3)				
		Yes (369)	29 (8.1)	1.31 (0.89–1.95)	1.33 (0.89–1.97)	1.58 (0.99–2.53)	
	Broken or thrown things	No (5283)	318 (6.1)				
		Yes (240)	23 (9.7)	1.66 (1.06–2.58) <sup>c</sup>	1.68 (1.08–2.63) <sup>c</sup>	1.92 (1.14–3.23) <sup>c</sup>	
	Conflicting partnership	Preschool	No (4265)	249 (5.8)			
			Yes (1262)	93 (7.4)	1.28 (1.00–1.64) <sup>c</sup>	1.28 (1.00–1.64) <sup>c</sup>	1.23 (0.90–1.67)
		School	No (3890)	253 (5.8)			
			Yes (929)	77 (8.1)	1.42 (1.09–1.86) <sup>c</sup>	1.43 (1.09–1.86) <sup>c</sup>	1.32 (0.96–1.82)
	Parent conflict index <sup>e</sup>		1.17 (1.06–1.28) <sup>c</sup>	1.17 (1.07–1.29) <sup>c</sup>	1.19 (1.06–1.34) <sup>c</sup>		

BPD, Borderline personality disorder; DSM-IV, Diagnostic and Statistical Manual of Mental Disorders, 4th edn; IQ, intelligence quotient; OR, odds ratio; CI, confidence interval.

<sup>a</sup> Controls include gender and age.

<sup>b</sup> Controls include gender, age, DSM-IV diagnosis and IQ.

<sup>c</sup> 95 % CI does not include 1.00.

<sup>d</sup> Suboptimal parenting index on a scale of 1–7.

<sup>e</sup> Parent conflict index on a scale of 1–5.

**Table 5.** Associations between maladaptive parenting and BPD definite status, showing crude associations, adjustment for age and gender, and additionally DSM-IV diagnosis and IQ

Exposure	Subgroup	BPD status, n (%)	OR (95% CI)	OR, with controls <sup>a</sup> (95% CI)	OR, with controls <sup>b</sup> (95% CI)			
Family adversity	No items							
	1 or 2 items		2.12 (1.11–4.08) <sup>c</sup>	2.13 (1.11–4.09) <sup>c</sup>	2.48 (1.01–6.08) <sup>c</sup>			
	More than 2 items		3.95 (1.88–8.32) <sup>c</sup>	3.98 (1.89–8.39) <sup>c</sup>	2.53 (0.78–8.18)			
Hitting and shouting	Preschool	Hitting	No (3235) Yes (2479)	22 (0.7) 30 (1.2)	1.79 (1.03–3.11) <sup>c</sup>	1.84 (1.06–3.21) <sup>c</sup>	1.05 (0.50–2.25)	
		Shouting	No (3125) Yes (2593)	27 (0.9) 25 (1.0)	1.12 (0.65–1.93)	1.14 (0.66–1.97)	1.17 (0.55–2.48)	
	School	Hitting	No (1868) Yes (1163)	18 (1.0) 12 (1.0)	1.07 (0.51–2.23)	1.06 (0.51–2.21)	1.26 (0.54–2.95)	
		Shouting	No (2644) Yes (477)	26 (1.0) 4 (0.8)	0.85 (0.30–2.45)	0.85 (0.30–2.45)	0.82 (0.24–2.80)	
Parental attitude	Preschool	Hostility	No (4918) Yes (812)	36 (0.7) 14 (1.7)	2.38 (1.28–4.43) <sup>c</sup>	2.38 (1.28–4.43) <sup>c</sup>	1.93 (0.81–4.64)	
		Resentment	No (5011) Yes (776)	38 (0.8) 12 (1.5)	2.06 (1.07–3.95) <sup>c</sup>	2.07 (1.08–3.98) <sup>c</sup>	2.21 (0.92–5.28)	
	School	Hostility	No (4595) Yes (539)	30 (0.7) 15 (2.8)	4.36 (2.33–8.15) <sup>c</sup>	4.34 (2.32–8.12) <sup>c</sup>	3.85 (1.69–8.78) <sup>c</sup>	
		Suboptimal parenting index <sup>d</sup>			1.29 (1.10–1.51) <sup>c</sup>	1.30 (1.11–1.52) <sup>c</sup>	1.28 (1.03–1.60) <sup>c</sup>	
	Conflict	Domestic violence	Emotional	No (4828) Yes (1067)	34 (0.7) 19 (1.8)	2.56 (1.45–4.50) <sup>c</sup>	1.96 (1.08–3.57) <sup>c</sup>	3.63 (1.71–7.73) <sup>c</sup>
			Physically hurt by partner	No (5527) Yes (369)	42 (0.8) 11 (3.0)	4.01 (2.05–7.86) <sup>c</sup>	3.02 (1.49–6.12) <sup>c</sup>	3.08 (1.17–8.10) <sup>c</sup>
Broken or thrown things		No (5283) Yes (240)	42 (0.8) 2 (0.8)	1.05 (0.25–4.36)	0.89 (0.21–3.74)	0.82 (0.11–6.21)		
		Conflicting partnership	Preschool	No (4265) Yes (1262)	38 (0.9) 6 (0.5)	0.53 (0.22–1.26)	0.43 (0.18–1.03)	0.74 (0.27–2.04)
School			No (3890) Yes (929)	26 (0.6) 16 (1.7)	2.81 (1.50–5.26) <sup>c</sup>	2.38 (1.25–4.51) <sup>c</sup>	2.89 (1.29–6.47) <sup>c</sup>	
Parent conflict composite <sup>e</sup>				1.33 (1.07–1.65) <sup>c</sup>	1.20 (0.95–1.51)	1.42 (1.06–1.91) <sup>c</sup>		

BPD, Borderline personality disorder; DSM-IV, Diagnostic and Statistical Manual of Mental Disorders, 4th edn; IQ, intelligence quotient; OR, odds ratio; CI, confidence interval.


<sup>a</sup> Controls include gender.

<sup>b</sup> Controls include gender, age, DSM-IV diagnosis and IQ.

<sup>c</sup> 95% CI does not include 1.00.

<sup>d</sup> Suboptimal parenting index on a scale of 1–7.

<sup>e</sup> Parent conflict index on a scale of 1–5.



**Fig. 2.** Final model showing unstandardized probit coefficients and standardized coefficients (in parentheses) for the direct effects of family adversity, suboptimal parenting, parent conflict, child intelligence quotient (IQ) and Diagnostic and Statistical Manual of Mental Disorders, 4th edn (DSM-IV) diagnosis ( $\chi^2 = 11.58$ ,  $p = 0.00$ , root mean square error of approximation = 0.02, comparative fit index = 0.99). Non-significant paths at the 0.05 level (one-tailed) are not shown. The Family Adversity Index (FAI) is coded into three categories: none, moderate and severe. FAI1 (one or two items) and FAI2 (more than two items) are dummy variables, with FAI (0 items) used as the reference group. -- →, Relationships (significant coefficients) for FAI2. —→, Relationships (significant coefficients) for other coefficients. 'Gender' is a nominal variable: the negative relationship indicates that male gender is a significant predictor of parenting problems. For clarity, the correlation between parenting and conflict (unstandardized coefficient 0.20, standardized coefficient 0.02,  $p = 0.00$ ) is not shown in the diagram. (⇔), Direct and indirect predictors of borderline personality disorder (BPD) are shown in Table 6.

childhood, such as social deprivation, leading to increased BPD symptoms. Alternatively, adversity in early pregnancy may lead to increased stress for the fetus, and early programming alterations of the hypothalamic–pituitary–adrenal axis (Entringer *et al.* 2009), increasing the risk of BPD symptoms.

Despite controlling for other adversities, we found that suboptimal parenting and parent conflict had significant direct associations with BPD symptoms, adding to the current research literature by providing prospective evidence for a link between maladaptive parenting and subsequent BPD symptoms in late childhood. Furthermore, there were significant indirect associations between suboptimal parenting and BPD outcome via DSM-IV diagnosis and IQ.

There is ample evidence that lower IQ is often indicative of a deleterious home environment, lacking in resources and academic encouragement (Brody & Flor, 1998; Van IJzendoorn *et al.* 2005). Therefore, maladaptive parenting is likely to contribute to poorer cognitive ability and increased BPD symptoms, as shown here. Considering the complexity of personality pathology (Tyrer *et al.* 2007), these outcomes may have various aetiological pathways. A family

environment characterized by conflict, aggression and anger directed at the child may make an impact upon the child in various ways including an alteration of internal schemata of behaviour and relationships (Westen *et al.* 2006), an exacerbation of stress responses (e.g. hypothalamic–pituitary–adrenocortical axis) (Gunnar, 1998) or an interaction with genes (Belsky & Beaver, 2011). All of these may compromise cognitive and emotional regulation (Posner *et al.* 2003). Indeed, individuals with BPD tend to display a disturbance in cognitive control processes (Posner *et al.* 2003; Rogosch & Cicchetti, 2005).

An association between IQ and increased psychotic symptoms during adolescence has been observed, curvilinear in nature, with both low and high (to a lesser extent) IQ increasing risk (Horwood *et al.* 2008). The present results suggest a more straightforward linear relationship between IQ and BPD symptoms (Supplementary Fig. S1), with high IQ possibly acting as a protective factor across the population (Batty *et al.* 2005); due to an increased ability to mobilize resources and respond appropriately in difficult situations.

Axis I (DSM-IV) diagnoses at 7–8 years were directly associated with BPD outcome at 11 years within

**Table 6.** Unstandardized probit coefficients (*B*) for the direct and indirect paths between FAI, suboptimal parenting, parental conflict, IQ and subsequent BPD outcome<sup>a</sup> at age 11 years

	Direct to BPD outcome			Indirect to BPD outcome					
				Via DSM-IV diagnosis			Via IQ		
	<i>B</i>	(s.e.)	<i>p</i> <sup>b</sup>	<i>B</i>	(s.e.)	<i>p</i>	<i>B</i>	(s.e.)	<i>p</i> <sup>b</sup>
FAI1 <sup>c</sup>	0.13 <sup>d</sup>	(0.05)	0.02	0.03	(0.002)	0.14	0.01	(0.01)	0.02
FAI2 <sup>e</sup>	0.35	(0.07)	0.000	0.01	(0.01)	0.09 <sup>b</sup>	0.03	(0.01)	0.01
Suboptimal parenting	0.053	(0.02)	0.001	0.004	(0.002)	0.08 <sup>b</sup>	0.002	(0.001)	0.07 <sup>b</sup>
Parent conflict	0.04	(0.02)	0.09 <sup>b</sup>	0.002	(0.001)	0.13	0.002	(0.001)	0.13
IQ	-0.01	(0.002)	0.01	-	-	-	-	-	-
DSM-IV	0.18	(0.10)	0.07 <sup>b</sup>						

FAI, Family Adversity Index; IQ, intelligence quotient; BPD, borderline personality disorder; DSM-IV, Diagnostic and Statistical Manual of Mental Disorders, 4th edn; s.e., standard error.

<sup>a</sup> The BPD outcome is an ordinal categorical outcome: none; probable (five or more symptoms); definite (five or more symptoms).

<sup>b</sup> The *p* value is two-tailed, therefore divided by 2 to get a significance value, as the direction of the association is clearly hypothesized.

<sup>c</sup> The FAI1 category denotes one or two items.

<sup>d</sup> A probit coefficient of 0.13 indicates that for each unit increase in FAI there is an increase of 0.13 standard deviations in the predicted Z score of the cumulative normal distribution of BPD symptoms.

<sup>e</sup> The FAI2 category denotes more than two items.

the path model. This is consistent with the 'complication model', which posits a predictive association between Axis I disorders and subsequent personality pathology (Philipsen *et al.* 2008). Our findings indicate that diagnoses of anxiety, depression, ADHD or externalizing disorders (conduct disorder, oppositional defiant disorder) were direct precursors of subsequent BPD symptoms. Both suboptimal parenting and parent conflict, to a lesser extent, were predictive of a DSM-IV Axis I diagnosis at 7–8 years. The association between suboptimal parenting and BPD via DSM diagnoses was only partial, however, and the association between parent conflict and BPD was not mediated by DSM-IV diagnoses. Thus, only a proportion of children reporting BPD at 11 years was identified by the DSM-IV diagnoses at 7–8 years.

This suggests that the DAWBA diagnoses may not capture all cases of emotional dysregulation, or the combination of internalizing and externalizing manifestations (Crawford *et al.* 2001), thought to presage the development of BPD (Crowell *et al.* 2009). While disorders such as conduct disorder and ADHD may have identified individuals more outwardly evincing the emotional dysregulation implicated in the development of BPD, other manifestations of emotional dysregulation may not have been observed by parents or teachers. Of note, male gender was significantly predictive of DSM-IV but not BPD diagnoses, suggesting that certain female typical manifestations of

emotional dysregulation (e.g. self-harm and eating disorders) may not have been identified within the DSM-IV diagnoses.

### Strengths and limitations

Study strengths include the large sample size and the assessment of family adversity before the birth of the child, precluding any reverse causality. The UK-CI-BPD was adapted from a well-validated instrument, piloted, administered by trained psychologists and showed high inter-rater reliability. The findings support the presence of a late childhood phenotype for BPD, and buttress current literature (Cohen *et al.* 2005; Chanen *et al.* 2007) by demonstrating that borderline personality symptoms, recognized in late childhood, are associated with similar risk factors to BPD diagnosed in adulthood. However, before firm conclusions can be drawn, it needs to be ascertained whether these BPD symptoms demonstrate predictive validity (Crick *et al.* 2005) and are related to BPD clinically diagnosed in adulthood.

There was substantial and selective attrition in this study. Those with more family adversity were more likely to have been lost from follow-up. Thus, the study is likely to underestimate the prevalence of BPD symptoms in late childhood (Bernstein *et al.* 1993). Despite selective drop-out, we found strong and hypothesized associations between family adversity,

suboptimal parenting and parent conflict and BPD symptoms among the remaining, less severely disadvantaged individuals. Wolke *et al.* (2009) demonstrated in simulations that even when drop-out is correlated with predictor/confounder variables, the relationships between predictors and outcome were not markedly attenuated. However, it cannot be precluded that selective drop-out had some influence on the predictive relationships reported.

Maternal hitting, shouting, hostility and resentment were based on self-report, potentially leading to an under-reporting of these factors. However, under-reporting would probably represent 'non-differential misclassification', i.e. under-reporting in both groups, therefore exerting a downward bias on our estimates of the long-term effects of suboptimal parenting, suggesting our robust estimates are conservative (underestimate effects) (Copeland *et al.* 1977). The reported rates of suboptimal parenting are still fairly high, however, suggesting a reasonable level of self-disclosure, possibly due to the anonymous, postal method of data collection. While self-report measures are regarded as less robust than observational measures, they have the benefit of capturing attitudes and behaviours across longer time spans.

Due to the very low prevalence of reported sexual abuse in this sample (0.05%), it was excluded as a predictor, potentially omitting an important experiential factor (Zanarini *et al.* 2006). Existing research, however, suggests that sexual abuse is not linked to the whole spectrum of BPD, and certain forms of BPD may be associated with maladaptive parenting other than sexual abuse (Salzman *et al.* 1993).

### *Implications and future directions*

Our results suggest that cognitive mechanisms play a direct and weak mediational role in the development of BPD symptoms. Assessing cognition via IQ supports that general cognitive ability relates to psychopathology (Batty *et al.* 2005). However, given the proposed centrality of emotional dysregulation within the BPD construct (LeGris & van Reekum, 2006), it would be prudent for future developmental studies to tap into the domain of emotional dysregulation more directly in order to clarify the pathways via which BPD symptoms develop. Though DSM-IV diagnoses at 7–8 years identified a proportion of children reporting BPD symptoms at 11 years, results suggest that there may be other precursors to BPD in mid-childhood.

In addition, the present results concur with previous studies that exposure to family adversity, suboptimal parenting and parent conflict may have numerous negative outcomes for children, including

lower cognitive ability and Axis I disorders. Further, we expand the current literature by providing prospective evidence of a link between maladaptive parenting and subsequent BPD symptoms at age 11 years, suggesting that interventions focused on improving parenting may produce wide-ranging positive effects.

We tentatively speculate that suboptimal parenting may be a marker for maternal irritable temperament (Siever & Davies, 1991), potentially exposing the child to the double jeopardy of an inherited irritable temperament (Stringaris *et al.* 2010) and suboptimal parenting, which may manifest in subsequent BPD symptoms (Crowell *et al.* 2009), including affective instability and intense inappropriate anger. Therefore, it would be desirable for future studies to ascertain whether there are prospective links between emotional/irritable temperament and later BPD symptoms. Assessing BPD symptoms in late childhood appears to be a promising avenue for understanding the development of BPD.

### **Supplementary material**

For supplementary material accompanying this paper, visit <http://dx.doi.org/10.1017/S0033291712000542>.

### **Acknowledgements**

We are extremely grateful to all the families who took part in this study, the midwives for their help in recruiting them, and the whole ALSPAC team, which includes interviewers, computer and laboratory technicians, clerical workers, research scientists, volunteers, managers, receptionists and nurses. We give special thanks to Andrea Waylen and Jeremy Horwood who helped in the conduct of the study. This article is the work of all the authors and D.W. and C.W. serve as guarantors for the content of the article. The UK Medical Research Council (grant no. 74882), the Wellcome Trust (grant no. 076467) and the University of Bristol provide core support for ALSPAC. C.W. was supported by a competitive Ph.D. scholarship funded by the University of Warwick, Department of Psychology.

### **Declaration of Interest**

None.

### **References**

- Bailey JM, Shriver A (1999). Does childhood sexual abuse cause borderline personality disorder? *Journal of Sex and Marital Therapy* 25, 45–57.



- Batty GD, Mortensen EL, Osler M (2005). Child IQ in relation to later psychiatric disorder: evidence from a Danish birth cohort study. *British Journal of Psychiatry* **187**, 180–181.
- Belsky J, Beaver KM (2011). Cumulative-genetic plasticity, parenting and adolescent self-regulation. *Journal of Child Psychology and Psychiatry* **52**, 619–626.
- Bernstein DP, Cohen P, Velez CN, Schwab-Stone M, Siever LJ, Shinsato L (1993). Prevalence and stability of the DSM-III-R personality disorders in a community-based survey of adolescents. *American Journal of Psychiatry* **150**, 1237–1243.
- Bowen E, Heron J, Waylen A, Wolke D; ALSPAC study team (2005). Domestic violence risk during and after pregnancy: findings from a British longitudinal study. *BJOG: International Journal of Obstetrics and Gynaecology* **112**, 1083–1089.
- Brody GH, Flor DL (1998). Maternal resources, parenting practices, and child competence in rural, single-parent African American families. *Child Development* **69**, 803–816.
- Chanen AM, Jovev M, Jackson HJ (2007). Adaptive functioning and psychiatric symptoms in adolescents with borderline personality disorder. *Journal of Clinical Psychiatry* **68**, 297–306.
- Chanen AM, Jovev M, McCutcheon LK, Jackson HJ, McGorry PD (2008). Borderline personality disorder in young people and the prospects for prevention and early intervention. *Current Psychiatry Reviews* **4**, 48–57.
- Cohen P, Crawford TN, Johnson JG, Kasen S (2005). The Children in the Community Study of the Developmental Course of Personality Disorder. *Journal of Personality Disorders* **19**, 466–486.
- Copeland KT, Checkoway H, McMichael AJ, Holbrook RH (1977). Bias due to misclassification in the estimation of relative risk. *American Journal of Epidemiology* **105**, 488–495.
- Crawford TN, Cohen P, Brook JS (2001). Dramatic erratic personality disorder symptoms: II. Developmental pathways from early adolescence to adulthood. *Journal of Personality Disorders* **15**, 336–350.
- Crick NR, Murray-Close D, Woods K (2005). Borderline personality features in childhood: a short-term longitudinal study. *Developmental Psychopathology* **17**, 1051–1070.
- Crowell S, Beauchaine T, Linehan M (2009). A biosocial developmental model of borderline personality: elaborating and extending Linehan's theory. *Psychological Bulletin* **135**, 495–510.
- Entringer S, Kumsta R, Hellhammer DH, Wadhwa PD, Wust S (2009). Prenatal exposure to maternal psychosocial stress and HPA axis regulation in young adults. *Hormones and Behavior* **55**, 292–298.
- Fergusson DM, Boden JM, Horwood LJ (2006). Examining the intergenerational transmission of violence in a New Zealand birth cohort. *Child Abuse and Neglect* **30**, 89–108.
- Geiger T, Crick NR (2001). A developmental psychopathology perspective on vulnerability to personality disorders. In *Vulnerability to Psychopathology: Risk Across the Lifespan* (ed. R. Ingram and J. M. Price), pp. 57–102. Guilford Press: New York.
- Golding J, Pembrey M, Jones R; ALSPAC Study Team (2001). ALSPAC: the Avon Longitudinal Study of Parents and Children, I: study methodology. *Paediatric and Perinatal Epidemiology* **15**, 74–87.
- Goodman R, Ford T, Richards H, Gatward R, Meltzer H (2000). The Development and Well-Being Assessment: description and initial validation of an integrated assessment of child and adolescent psychopathology. *Journal of Child Psychology and Psychiatry* **41**, 645–655.
- Gunnar MR (1998). Quality of early care and buffering of neuroendocrine stress reactions: potential effects on the developing human brain. *Preventive Medicine* **27**, 208–211.
- Guzder J, Paris J, Zerkowitz P, Feldman R (1999). Psychological risk factors for borderline pathology in school-age children. *Journal of the American Academy of Child and Adolescent Psychiatry* **38**, 206–212.
- Herman JL, Perry JC, van der Kolk BA (1989). Childhood trauma in borderline personality disorder. *American Journal of Psychiatry* **146**, 490–495.
- Hooley JM, Hoffman PD (1999). Expressed emotion and clinical outcome in borderline personality disorder. *American Journal of Psychiatry* **156**, 1557–1562.
- Horwood J, Salvi G, Thomas K, Hollis C, Lewis G, Menezes PA, Thompson A, Wolke D, Zammit S, Harrison G (2008). IQ and non-clinical psychotic symptoms in 12-year-olds: results from the ALSPAC birth cohort. *British Journal of Psychiatry* **193**, 185–191.
- Hyman, IA (1997). *The Case Against Spanking: How to Discipline Your Child Without Hitting*. Jossey-Bass Inc.: San Francisco.
- Johnson JG, Cohen P, Brown J, Smailes EM, Bernstein DP (1999). Childhood maltreatment increases risk for personality disorders during early adulthood. *Archives of General Psychiatry* **56**, 600–606.
- Johnson JG, Cohen P, Chen H, Kasen S, Brook JS (2006). Parenting behaviours associated with risk for offspring personality disorder during adulthood. *Archives of General Psychiatry* **63**, 579–587.
- Johnson JG, Cohen P, Smailes EM, Skodol AE, Brown J, Oldham JM (2001). Childhood verbal abuse and risk for personality disorders during adolescence and early adulthood. *Comprehensive Psychiatry* **42**, 16–23.
- Johnson JG, Smailes EM, Cohen P, Brown J, Bernstein DP (2000). Associations between four types of neglect and personality disorder symptoms during adolescence and early adulthood: findings of a community-based longitudinal study. *Journal of Personality Disorders* **14**, 171–187.
- Kitzmann GN, Holt A, Kenney E (2003). Child witness to domestic violence a meta-analytic review. *Journal of Consulting Clinical Psychology* **1**, 223–353.
- Kramer HC (2007). DSM categories and dimensions in clinical and research contexts. *International Journal of Methods in Psychiatric Research* **16** (Suppl. 1), S8–S15.
- Lee MY, Uken A, Sebold J (2007). Role of self-determined goals in predicting recidivism in domestic violence offenders. *Research on Social Work Practice* **17**, 30–41.

- LeGris J, van Reekum R** (2006). The neuropsychological correlates of borderline personality disorder and suicidal behaviour. *Canadian Journal of Psychiatry* **51**, 131–142.
- Muthén BO** (1998–2004). *Mplus Technical Appendices*. Muthén & Muthén: Los Angeles.
- Philipsen A, Limberger MF, Lieb K, Feige B, Kleindienst N, Ebner-Priemer U, Barth J, Schmahl C, Bohus M** (2008). Attention-deficit hyperactivity disorder as a potentially aggravating factor in borderline personality disorder. *British Journal of Psychiatry* **192**, 118–123.
- Posner MI, Rothbart MK, Vizueta N, Thomas KM, Levy KN, Fossella J, Silbersweig D, Stern E, Clarkin J, Kernberg O** (2003). An approach to the psychobiology of personality disorders. *Developmental Psychopathology* **15**, 1093–1106.
- Rogosch FA, Cicchetti D** (2005). Child maltreatment, attention networks, and potential precursors to borderline personality disorder. *Developmental Psychopathology* **17**, 1071–1089.
- Salzman JP, Salzman C, Wolfson AN, Albanese M, Looper J, Ostacher M, Schwartz J, Chinman G, Land W, Miyawaki E** (1993). Association between borderline personality structure and history of childhood abuse in adult volunteers. *Comprehensive Psychiatry* **34**, 254–257.
- Siever L, Davies K** (1991). A psychobiologic perspective on the personality disorders. *American Journal of Psychiatry* **148**, 1647–1658.
- Stepp SD, Pilkonis PA, Hipwell AE, Loebar R, Stouthamer-Loeber M** (2010). Stability of borderline personality disorder features in girls. *Journal of Personality Disorders* **24**, 460–472.
- Stringaris A, Maughan B, Goodman R** (2010). What's in a disruptive disorder? Temperamental antecedents of oppositional defiant disorder: findings from the Avon Longitudinal Study. *Journal of American Academy of Child and Adolescent Psychiatry* **49**, 474–483.
- Tyrer P, Coombs N, Ibrahim F, Mathilakath A, Bajaj P, Ranger M, Din R** (2007). Critical developments in the assessment of personality disorder. *British Journal of Psychiatry* **190**, 51–59.
- Van IJzendoorn MH, Juffer F, Klein Poelhuis CW** (2005). Adoption and cognitive development: a meta-analytic comparison of adopted and nonadopted children's IQ and school performance. *Psychological Bulletin* **131**, 301–316.
- Waylen A, Stallard N, Stewart-Brown S** (2008). Parenting and health in mid-childhood: a longitudinal study. *European Journal of Public Health* **18**, 300–305.
- Weaver TL, Clum GA** (1993). Early family environments and traumatic experiences associated with borderline personality disorder. *Journal of Consulting Clinical Psychology* **61**, 1068–1075.
- Wechsler D, Golombok S, Rust J** (1992). *WISC-III UK Wechsler Intelligence Scale for Children*. Psychological Corp.: Sidcup.
- Westen D, Nakash O, Thomas C, Bradley R** (2006). Clinical assessment of attachment patterns and personality disorder in adolescents and adults. *Journal of Consulting Clinical Psychology* **74**, 1065–1085.
- Wolke D, Waylen A, Samara M, Steer C, Goodman R, Ford T, Lamberts K** (2009). Selective drop-out in longitudinal studies and non-biased prediction of behaviour disorders. *British Journal of Psychiatry* **195**, 249–256.
- Zanarini MC, Frankenberg FR** (2001). Attainment and maintenance of reliability of Axis I and II disorders over the course of a longitudinal study. *Comprehensive Psychiatry* **42**, 369–374.
- Zanarini MC, Frankenburg FR, Hennen J, Reich DB, Silk KR** (2006). Prediction of the 10-year course of borderline personality disorder. *American Journal of Psychiatry* **163**, 827–832.
- Zanarini MC, Frankenburg FR, Sickel AE, Yong L** (1996). *The Diagnostic Interview for DSM-IV Personality Disorders (DIPD-IV)*. McLean Hospital: Belmont, MA.
- Zanarini MC, Horwood J, Waylen A, Wolke D** (2004). *The UK Version of the Childhood Interview for DSM-IV Borderline Personality Disorder (UK-CI-BPD)*. University of Bristol: Bristol.
- Zanarini MC, Horwood J, Wolke D, Waylen A, Fitzmaurice G, Grant BF** (2011). Prevalence of DSM-IV borderline personality disorder in two community samples: 6,330 English 11-year olds and 34,653 American adults. *Journal of Personality Disorders* **25**, 607–619.
- Zanarini MC, Skodol AE, Bender D, Dolan R, Sanislow C, Schaefer E, Morey LC, Grilo CM, Shea MT, McGlashen TH, Gunderson JG** (2000). The collaborative longitudinal personality disorders study II. Reliability of Axis I and II diagnosis. *Journal of Personality Disorders* **14**, 291–299.