

Oct4 overexpression facilitates proliferation of porcine fibroblasts and development of cloned embryos

Su Jin Kim^{2,5}, Ok Jae Koo^{3,5}, Hee Jung Park³, Joon Ho Moon², Bego Roibas da Torre², Palaksha Kanive Javaregowda², Jung Taek Kang², Sol Ji Park², Islam M. Saadeldin², Ji Yei Choi², Byeong-Chun Lee² and Goo Jang^{1,4}

Department of Theriogenology and Biotechnology, College of Veterinary Medicine and the Research Institute for Veterinary Science, Seoul National University, Seoul; Laboratory Animal Research Center, Samsung Biomedical Research Institute, Gyeonggi-do; and Emergence Center for Food-Medicine Personalized Therapy System, Advanced Institutes of Convergence Technology, Seoul National University, Gyeonggi-do, Republic of Korea

Date submitted: 16.12.2013. Date revised: 18.6.2014. Date accepted: 26.6.2014

Summary

Octamer-binding transcription factor 4 (Oct4) is a critical molecule for the self-renewal and pluripotency of embryonic stem cells. Recent reports have shown that Oct4 also controls cell-cycle progression and enhances the proliferation of various types of cells. As the high proliferation of donor fibroblasts is critical to the production of transgenic pigs, using the somatic cell nuclear transfer technique, we analysed the effect of Oct4 overexpression on the proliferation of porcine fibroblasts and embryos. Porcine endogenous Oct4 cDNA was cloned, sequenced and inserted into an expression vector. The vector was transfected into porcine fibroblasts, and a stable Oct4-overexpressed cell line was established by antibiotic selection. Oct4 expression was validated by the immunostaining of Oct4. Cell morphology was changed to sharp, and both proliferation and migration abilities were enhanced in Oct4-overexpressed cells. Real-time RT-PCR results showed that *p16*, *Bcl2* and *Myc* were upregulated in Oct4-overexpressed cells. Somatic cell nuclear transfer was performed using Oct4-overexpressed cells, and the development of Oct4 embryos was compared with that of wild-type cloned embryos. The cleavage and blastocyst formation rates were improved in the Oct4 embryos. Interestingly, blastocyst formation of the Oct4 embryos was observed as early as day 5 in culture, while blastocysts were observed from day 6 in wild-type cloned embryos. In conclusion, the overexpression of Oct4 enhanced the proliferation of both porcine fibroblasts and embryos.

Keywords: Cell proliferation, Embryos, Oct4, Porcine fibroblasts, Somatic cell nuclear transfer

Introduction

Octamer-binding transcription factor 4 (Oct4) is a transcription factor of the POU family that can be characterized as a critical molecule for modulating the self renewal and pluripotency of embryonic stem cells (Okamoto *et al.*, 1990; Mountford *et al.*, 1998; Nichols *et al.*, 1998). The precise regulation of Oct4 is very important for maintaining and differentiating embryonic stem cells; thus, both the overexpression (Hochedlinger *et al.*, 2005; Fan *et al.*, 2013) and downregulation (Pesce *et al.*, 1998; Adjaye *et al.*, 1999; Goto *et al.*, 1999; Niwa *et al.*, 2000; Fan *et al.*, 2013) of Oct4 induce a loss of pluripotency and induce differentiation. Apart from embryonic stem cells, the

¹All correspondence to: Goo Jang, Department of Theriogenology and Biotechnology, College of Veterinary Medicine, Seoul National University, 1 Gwanak-ro, Gwanak-gu, Seoul 151-742, Republic of Korea. Tel: +82 2 880 1280. Fax: +82 2 873 1269. E-mail: snujang@snu.ac.kr

²Department of Theriogenology and Biotechnology, College of Veterinary Medicine and the Research Institute for Veterinary Science, Seoul National University, Seoul 151-742, Republic of Korea.

³Laboratory Animal Research Center, Samsung Biomedical Research Institute, Gyeonggi-do 440-746, Republic of Korea.

⁴Emergence Center for Food-Medicine Personalized Therapy System, Advanced Institutes of Convergence Technology, Seoul National University, Gyeonggi-do, Republic of Korea.

⁵These authors contributed equally.

expression of Oct4 is also found in other cell types including mesenchymal stem cells (Fan *et al.*, 2013), early embryonic tissues (DeVeale *et al.*, 2013) and cancer cells (Li *et al.*, 2012), although its roles and underlying mechanisms are still unclear. Recently, several reports have shown that the expression of Oct4 controls cell-cycle progression and enhances the proliferation of the cells for both embryonic stem cells (Lee *et al.*, 2010) and other cell types (Li *et al.*, 2012; DeVeale *et al.*, 2013; Fan *et al.*, 2013).

The high proliferation of nuclear donor cells is important in transgenic animal production using the somatic cell nuclear transfer (SCNT) technique, as the transgenesis and selection procedures of the donor cells require multiple subcultures and expansion steps. Because only primary cultured cells (fibroblasts in most cases) can be used for SCNT, it is critical to maintain the highly proliferative properties of the cells that are intended for use in SCNT after gene modification. From this point of view, we wondered whether the expression of Oct4 in primary cultured porcine fibroblasts enhances proliferation, as has been suggested by previous reports. However, the role of Oct4 expression in porcine fibroblasts is still unclear.

Hence, we hypothesised that the expression of Oct4 in porcine fibroblasts might play a role in cell proliferation. In this study, we cloned porcine endogenous Oct4 cDNA and established and analysed Oct4-overexpressed porcine fibroblasts. In addition, we produced cloned embryos using the Oct4-overexpressed fibroblasts by the SCNT technique and observed the effects of Oct4 expression on the development of porcine embryos.

Materials and methods

Primary culture of porcine fibroblasts

Primary culture was performed using ear skin biopsies from neonatal piglets. The lumps of ear skin were minutely homogenised and then cultured overnight in collagenase IV (Sigma-Aldrich Corp., St. Louis, MO, USA) in a 37°C incubator. The homogenised ear tissues were washed more than three times with phosphate-buffered saline (PBS; Life Technologies, Carlsbad, CA, USA) and collected by centrifugation at 43 g, for 2 min. The collected clusters of cells were cultured in 60-mm culture dishes with Dulbecco's Modified Eagle Medium (DMEM, Life Technologies), supplemented with 10% fetal bovine serum (FBS, Life Technologies) and 1% penicillin/streptomycin (Pen/Strep; Life Technologies).

Porcine Oct4 cloning and the construction of an expression vector

For cloning the coding domain sequences (CDS) of porcine Oct4, total RNA was extracted from pig ovaries using an RNA extraction kit (Qiagen, Hilden, Germany). One microgram of total RNA was used for reverse transcription with Superscript III (Life Technologies). The CDS of the porcine Oct4 was amplified by PCR using specific primer sets (Table 1). The PCR fragments were cloned into the NT-GFP Fusion TOPO-cloning vector (Life Technologies) and sequenced for confirmation. The vectors containing porcine Oct4 CDS were digested using two restriction enzymes, *NheI* and *XhoI* (New England Bio Labs, Inc., Beverly, MA, USA), and ligated into the pIRES2 DsRed-Express2 vector (CLONTECH Laboratories Inc., Palo Alto, CA, USA). The map of the expression vector is illustrated in Figure 1A.

Transfection and establishment of a stable Oct4-overexpressed cell line

One day before transfection, fibroblasts were plated at a density of 1×10^5 cells/ml in a 35-mm culture dish and cultured overnight to achieve 50–70% confluence. Next, 1 µg of the Oct4 vector (linearised by *AflIII*; New England Bio Labs, Inc.), 3 µl of the transfection reagent (Fugene HD; Roche Inc., Switzerland) and 96 µl of DMEM were incubated for at least 20 min at 25°C in a 1.5-ml microtube and were then overlaid on a prepared porcine fibroblast. After 2 days of culture, 1 mg/ml neomycin (G418, Life Technologies) was added for another 2 weeks of culture to select the stable Oct4-expressing cells.

Immunocytochemistry validation of Oct4 expression

To determine the transfected cells that expressed the Oct4 protein, the cells were immunostained. The cells were fixed with 4% paraformaldehyde for 20 min, permeabilised with 0.1% Triton X-100 (Sigma) for 10 min and blocked with 10% normal goat serum for 1 h. The fixed cells were incubated overnight with primary antibody (Oct3/4, Santa Cruz Biotechnology, Inc., CA, USA) and then with secondary antibody (Cy3-AffiniPure Goat Anti-Mouse IgG, Jackson ImmunoResearch Laboratories, Inc., PA, USA) for 2 h. In addition, DAPI was used as a counter-stain. The stained cells were examined under ultraviolet light using a fluorescence microscope (Nikon, Japan).

Analysis of cell growth and proliferation

Cell growth was analysed using an automated cell counter (Countess[®] Cell Counter, Invitrogen). For analysis, 0.5×10^5 /ml cells were plated and cultured in 12-well plates (Falcon, NJ, USA), and

Table 1 Primer sequences used in the study

Primer name	Sequence	Reference
CDS-Oct4	F : ATGGCGGGACACCTGGCTTCCGACTT R : TCAGTTTGAATGCATGGGGGAGCCCAGA	ENSSSCT 00000001516
UTR-Oct4	F : CAAACTGAGGTGCCTGCCCTTC R : CTTAATCCCAAAGCCCTGGT	ENSSSCT 00000001516
UTR-Sox2	F : GTTCCATGGGCTCAGTGGTCAAG R : AAGCGTACCGGGTTTTTCTCCATAC	ENSSSCT 00000012883
UTR-Nanog	F : AGCCCCAGCTCCAGTTTCAGC R : AATGATCGTCACATATCTTCAGGCTGTA	ENSSSCT 00000000727
Gapdh ^a	F : ACCTGCCGTCTGGAGAAACC R : GACCATGAGGTCCACCACCCTG	ENSSSCT 00000000756
p16	F : CTGGACACTTTGGTGGTCCT R : GCGGGATCTTCTCCAGAGTT	AJ242787
p53	F : CCTCACCATCATCACTGG R : GGCTTCTTCTTTTGCCTGG	NM-213824.3
Bax	F : GCCGAAATCTTTGCTGACGG R : CGAAGGAAGTCCAGCGTCCA	AJ606301
Bcl2	F : TGGTGGTTGACTTTCTCTCC R : ATTGATGGCACTAGGGGTTT	AF216205
Cyclin D2	F : AGGAGCAGATTGAGGTCGTG R : ATATCCCGCACGTCGTAGG	NM-214088.1
Myc	F : CTGCCAAGAGGGCTAAGTTG R : AGCTTTTGCTCCTCTGCTTG	FJ882404
Gapdh ^b	F : TCTCTGCTCCCTCCCCGTTT R : TGGCAATGCACGGAACACAC	AF069649.1

^aControl for RT-PCR; ^bControl for real-time RT-PCR.

the cultured cells were collected every 24 h by 0.25% trypsin–EDTA treatment. The collected cells were resuspended in 1 ml of PBS, and 10 μ l of the sample were mixed with 10 μ l of trypan blue and then pipetted into a Countess[®] Chamber slide for analysis. Cell number results were analysed with population doubling time (PDT) calculating software (<http://www.doublingtime.com/compute.php>). To determine the cell proliferation ability, a cell migration assay was performed. Monolayers of the wild-type and Oct4-transfected cells were prepared by seeding 2×10^4 cells onto Oris[™] Cell Migration Assay kit 96-well plates (PLATYPUS Technologies, WI, USA). After 24 h, following the removal of the stopper, phase-contrast images were taken for a pre-migration reference ($t = 0$ h), and then cells were allowed to grow for 36 h to permit cell migration. After 36 h of culture, images of migrated cells were recorded. The images were analysed to establish the migration range from pre-migration ($t = 0$ h) to post-migration ($t = 36$ h), using NIH Image J freeware, and the Z factor was evaluated as previously reported (Zhang *et al.*, 1999).

Quantitative gene expression analysis

To analyse the expression of the endogenous stemness-related genes, *Oct4*, *Sox2* and *Nanog*, quantitative RT-PCR was performed as described previously (Kong *et al.*, 2009) with some modification. Briefly, total

RNA was extracted from wild-type or Oct4-transfected cells, and cDNA was synthesised and used for PCR amplification as described earlier. To distinguish the endogenous genes from those transfected with Oct4, specific primers including the untranslated region were designed and used (Table 1). For comparing the range of expression levels, Gapdh was used as a housekeeping control. For analysing gene expression related with the cell-cycle progression and apoptosis, real-time RT-PCR was used as described previously (Kinikoglu *et al.*, 2014) with some modification. For the analysis, SYBR[®] Green PCR master mix and 7300 Real-Time PCR System (Applied Biosystems, Foster City, CA, USA) were used. Details of specific primers used in the study are shown in Table 1. The expression levels of different genes were normalised to the housekeeping gene Gapdh using the $\Delta\Delta$ Ct method, as previously described (Livak and Schmittgen, 2001).

Somatic cell nuclear transfer and embryo culture

To produce Oct4-expressing porcine embryos, SCNT was performed as described in the previous report (Park *et al.*, 2014) with a slight modification. Briefly, *in vitro* matured and enucleated oocytes were placed in TCM-199, which was supplemented with 10% (v/v) FBS. Either wild-type fibroblasts or the Oct4-transfected fibroblasts were used as donor cells for SCNT. The donor cells were injected into the




Figure 1 (A) Vector construct used in the study. (B) Morphology and Oct4 expression of wild-type (Wt; Ba, Ba' and Ba'') and Oct4-overexpressed (Oct4; Bb, Bb' and Bb'') porcine fibroblasts. Cell shape was changed after transfection of Oct4 (Ba versus Bb). Wild-type and Oct4-overexpressed cells were stained with DAPI (Ba' and Bb'), and Cy3 conjugated secondary antibody was attached to Oct4 primary antibody (Oct4'; Ba'' and Bb''). Expression of Oct4 was only detected in transfected cells and localised in the nucleus. (C) The cell growth curves of wild-type and Oct4-overexpressed porcine fibroblasts. A significant difference in growth patterns was observed between two groups.

perivitelline space of the prepared oocytes and then electrically fused using an electro cell fusion generator (LF101; Nepa Gene Co., Japan). After 40 min of fusion, the artificial activation of the fused embryos was performed with a single direct current electric pulse of 1.5 kV/cm for 60 μ s using a BTX Electro Cell Manipulator 2001 (BTX Inc., San Diego, USA). The activated SCNT embryos were cultured in Porcine Zygote Medium-5 (PZM-5) (Yoshioka *et al.*, 2002) under mineral oil at 39°C in 5% CO₂, 5% O₂ and 90% N₂. Embryos were evaluated for cleavage on day 2 and for blastocyst formation on days 5 and 6.

Statistical analysis

All experiments were replicated at least three times and statistically analysed using Prism 5 software (GraphPad, La Jolla, CA, USA). Student's *t*-test was used to analyse any differences between two groups. One-way analysis of variance (ANOVA) was used to analyse any differences between three or more groups. A *P*-value less than 0.05 was considered to be statistically significant.

Results and Discussion

In this study, we cloned the CDS of porcine endogenous Oct4 and used it for overexpression in porcine fibroblasts. Several studies have used human-derived Oct4 in porcine cells to generate induced pluripotent stem cells (Kwon *et al.*, 2013; Park *et al.*, 2013; Yuan *et al.*, 2014; Zhang *et al.*, 2014). However, a few previous reports have tried to use porcine endogenous Oct4 (Liu *et al.*, 2012). The sequencing results of our cloned porcine endogenous Oct4 showed that the homologies of the amino acid sequence of porcine Oct4 and human Oct4 were 93% identical (data not shown).

After transfection and selection, the expression of Oct4 was validated with immunocytochemistry (Fig. 1B). Because Oct4 is a transcription factor, most of the cells showed Oct4-positive staining in the nucleus (Fig. 1B) as expected. Interestingly, the morphology of the fibroblasts was remarkably changed in Oct4-overexpressed cells. (Fig. 1Ba, Bb) The Oct4 cells were smaller in size with sharper and clearer margins compared to the wild-type cells. These changes in morphologies were similar to those seen in less




Figure 2 (A) Cell migrations of wild-type (Wt) and Oct4-overexpressed (Oct4) porcine fibroblasts. White circles are open areas on culture plate at 0 h (Aa and Ab) and 38 h (Aa' and Ab'). (B) Oct4-overexpressed cells showed significantly smaller open areas after cell migration, which reveals higher proliferation capacity compared with wild-type cells. The symbol (*) indicates a significant difference ($P < 0.05$).

senescent cells presented by previous reports (Bayreuther *et al.*, 1988; Cho *et al.*, 2004). The overexpression of porcine Oct4 protein also facilitated cellular doubling and proliferation. The PDT was noticeably reduced from 55.77 ± 1.32 h in the wild-type cells to 42.67 ± 0.84 h in the Oct4-overexpressed cells and, as a result, the growth curves were significantly different between the two groups (Fig. 1C). A cell migration assay was also performed to analyse the proliferation capacity of Oct4-overexpressed cells. The migrations of the cells were more active in the Oct4 cells (Fig. 2). Therefore, the migration speed of the Oct4 cells ($2.10 \times 10^6 \pm 3.67 \times 10^4 \mu\text{m}^2/\text{h}$) was significantly faster than that of the wild-type cells ($1.66 \times 10^6 \pm 2.21 \times 10^4 \mu\text{m}^2/\text{h}$). These data exhibited high reliability, evidenced not only by a CV of 2.3 and 3.0 for wild-type and Oct4-overexpressed cells, respectively, but also a Δ migration of $0.5 < Z < 1.0$.

To investigate the underlying mechanism of enhanced proliferation by Oct4 overexpression, we examined the transcription level of the cell-cycle-related genes, *p16*, *Cyclin D2*, *p53* and *Myc*, and the apoptosis-related genes, *Bax* and *Bcl2* (Fig. 3). Results showed that *p53*, *Cyclin D2* and *Bax* expression was not significantly different between the two groups. However, *p16* transcription was increased more than four-fold and *Myc* was increased about three-fold in the Oct4-overexpressed cells. Usually, in highly proliferative cells, *p16* and *p53* expression is reduced. If *p16* and *p53* expression is increased, cells are likely undergoing cell senescence and cell-cycle arrest




Figure 3 Transcriptional levels of cell cycle genes (*p16*, *p53*, *Cyclin D2* and *Myc*) and apoptosis-related genes (*Bax*, *Bcl2*) were analysed. In Oct4-overexpressed cells (Oct4), transcription levels of *p16*, *Myc* and *Bcl2* were significantly higher compared with that in wild-type cells (Wt). The symbol (*) indicates a significant difference between groups ($P < 0.05$).

(Weinberg, 1989; Raley, 1990; Hinds *et al.*, 1994; Downward, 1997; Serrano *et al.*, 1997). Thus, the upregulation of *p16* in Oct4-overexpressed cells was somewhat unexpected. However, a previous report showed that the expression of *Myc* bypassed the *p16* pathway and promoted cell proliferation even in the presence of *p16* (Alevizopoulos *et al.*, 1997). The present study also showed that the expression of *Myc* is upregulated by the expression of *Oct4*. Thus, we suggest that *Myc* is one of the key molecules

Table 2 Development of porcine cloned embryos reconstructed with wild-type (Wt) or Oct4-overexpressed (Oct4) cells

	Total	Fused (%)	Cleaved (%)	Day5		Day 6	
				BL (%)	Hatched (%)	BL (%)	Hatched (%)
Wt	249	196 (78.7)	113 (57.7) ^a	0 (0.0) ^a	–	6 (5.3)	3 (50)
Oct4	252	206 (81.7)	146 (70.9) ^b	6 (4.1) ^b	6 (100)	12 (8.2)	11 (91.7)

^{a,b}Values with different superscripts within the same column are significantly different ($P < 0.05$). BL: blastocyst.




Figure 4 Transcription level changes of endogenous *Oct4*, *Sox2* and *Nanog* in wild-type and Oct4-overexpressed porcine fibroblasts. Endogenous Oct4 expression was very weak and did not change after Oct4 vector transfection. Expression of *Sox2* and *Nanog* was reduced in Oct4-overexpressed cells compared with wild-type cells. M: 1 kb plus marker; W: wild-type cells; O: Oct4-overexpressed cells; N: blank template.

of proliferation in Oct4-overexpressed cells. On the other hand, the expression of the anti-apoptosis gene *Bcl2* was also increased about two-fold in the Oct4-overexpressed cell line. This means that the reduced apoptosis by *Bcl2* expression also contributed to faster proliferation in the Oct4-overexpressed cells.

We also analysed the expression of the endogenous stemness-related genes, *Oct4*, *Sox2* and *Nanog*, in Oct4-overexpressed cells. As shown in Fig. 4, the expression of endogenous Oct4 in porcine fibroblasts was very weak and did not change after transfection of the Oct4 vector. This means that the cellular and molecular changes found in the present study were affected by transfected Oct4 and not by the expression of endogenous *Oct4*. *Sox2* and *Nanog* are also known as molecules that modulate cell-cycle progression and promote proliferation (Zhang *et al.*, 2009; Fan *et al.*, 2013). However, the present study showed that the expression of endogenous *Sox2* and *Nanog* was downregulated (about 20 and 30%, respectively). Therefore, we conclude that enhanced proliferation in the Oct4-overexpressed cells is not related with the effect of *Sox2* and the *Nanog* pathway. Current results were similar to previous reports indicating that Oct4 overexpression induced negative feedback to *Sox2* and *Nanog* expression (Lee *et al.*, 2006b; Boer *et al.*, 2007).

The effects of Oct4 overexpression in porcine embryos were also investigated in the study. For this purpose, we produced cloned embryos reconstructed with wild-type or Oct4-overexpressed cells. Oct4 overexpression improved both the cleavage and blastocyst formation rate of cloned embryos (Table 2). As we reported earlier, the expression of *Oct4* in

cloned embryos was significantly lower compared with fertilised embryos (Lee *et al.*, 2006a). Thus, the overexpression of Oct4 in donor cells might be helpful for the proper development of cloned SCNT embryos. In addition, the higher expression of *Bcl2* in Oct4-overexpressing donor cells may affect the development of SCNT embryos. Interestingly, developmental velocity was also increased in the Oct4 group, and the formation of blastocysts could be observed as early as day 5 of culture, while blastocysts were observed from day 6 in the wild-type group. This phenomenon is similar to the result of increased PDT in Oct4-overexpressed fibroblasts in the present study. Therefore, we conclude that the overexpression of Oct4 can facilitate cell-cycle progression in both fibroblasts and cloned embryos.

In conclusion, we cloned the porcine endogenous *Oct4* gene and established a stable Oct4-overexpressed fibroblasts cell line. We found that the overexpression of Oct4 promotes the proliferation of fibroblasts. This phenomenon seems to be related to the upregulation of the *Myc* and *Bcl2* genes. The development of cloned embryos reconstructed with Oct4-overexpressed cells was improved, and blastocysts were observed as early as day 5 in culture.

Acknowledgements

This study was supported by grants from IPET (#111078–03–1–CG000), NRF (NRF-2011–0014941), Biogreen (PJ0090962012), RDA (#PJ009802), MOTIE and BK 21 plus program for veterinary science.

References

- Adjaye, J., Bolton, V. & Monk, M. (1999). Developmental expression of specific genes detected in high-quality cDNA libraries from single human preimplantation embryos. *Gene* **237**, 373–83.
- Alevizopoulos, K., Vlach, J., Hennecke, S. & Amati, B. (1997). Cyclin E and c-Myc promote cell proliferation in the presence of p16INK4a and of hypophosphorylated retinoblastoma family proteins. *EMBO J.* **16**, 5322–33.
- Bayreuther, K., Rodemann, H.P., Hommel, R., Dittmann, K., Albiez, M. & Francz, P.I. (1988). Human skin fibroblasts in vitro differentiate along a terminal cell lineage. *Proc. Natl. Acad. Sci. USA* **85**, 5112–6.
- Boer, B., Kopp, J., Mallanna, S., Desler, M., Chakravarthy, H., Wilder, P.J., Bernadt, C. & Rizzino, A. (2007). Elevating the levels of Sox2 in embryonal carcinoma cells and embryonic stem cells inhibits the expression of Sox2:Oct-3/4 target genes. *Nucleic Acids Res.* **35**, 1773–86.
- Cho, K.A., Ryu, S.J., Oh, Y.S., Park, J.H., Lee, J.W., Kim, H.P., Kim, K.T., Jang, I.S. & Park, S.C. (2004). Morphological adjustment of senescent cells by modulating caveolin-1 status. *J. Biol. Chem.* **279**, 42270–8.
- DeVeale, B., Brokhman, I., Mohseni, P., Babak, T., Yoon, C., Lin, A., Onishi, K., Tomilin, A., Pevny, L., Zandstra, P.W., Nagy, A. & van der Kooy, D. (2013). Oct4 is required ~E7.5 for proliferation in the primitive streak. *PLoS Genetics* **9**, e1003957.
- Downward, J. (1997). Cell cycle: routine role for Ras. *Curr. Biol.* **7**, R258–60.
- Fan, Y.X., Gu, C.H., Zhang, Y.L., Zhong, B.S., Wang, L.Z., Zhou, Z.R., Wang, Z.Y., Jia, R.X. & Wang, F. (2013). Oct4 and Sox2 overexpression improves the proliferation and differentiation of bone mesenchymal stem cells in Xiaomeishan porcine. *Genetics Mol. Res.* **12**, 6067–79.
- Goto, T., Adjaye, J., Rodeck, C.H. & Monk, M. (1999). Identification of genes expressed in human primordial germ cells at the time of entry of the female germ line into meiosis. *Mol. Hum. Reprod.* **5**, 851–60.
- Hinds, P.W., Dowdy, S.F., Eaton, E.N., Arnold, A. & Weinberg, R.A. (1994). Function of a human cyclin gene as an oncogene. *Proc. Natl. Acad. Sci. USA* **91**, 709–13.
- Hochedlinger, K., Yamada, Y., Beard, C. & Jaenisch, R. (2005). Ectopic expression of Oct-4 blocks progenitor-cell differentiation and causes dysplasia in epithelial tissues. *Cell* **121**, 465–77.
- Kinikoglu, B., Kong, Y.W. & Liao, E.C. (2014). Characterization of cultured multipotent zebrafish neural crest cells. *Exp. Biol. Med.* **239**, 159–68.
- Kong, Y.W., Jing, G., Yan, Z.G., Li, C. Z., Gong, N.P., Zhu, F.J., Li, D.X., Zhang, Y.R., Zheng, G.L., Wang, H.Z., Xie, L.P. & Zhang, R.Q. (2009). Cloning and characterization of Prsilkin-39, a novel matrix protein serving a dual role in the prismatic layer formation from the oyster *Pinctada fucata*. *J. Biol. Chem.* **284**, 10841–54.
- Kwon, D.J., Jeon, H., Oh, K.B., Ock, S.A., Im, G.S., Lee, S.S., Im, S.K., Lee, J.W., Oh, S.J., Park, J.K. & Hwang, S. (2013). Generation of leukemia inhibitory factor-dependent induced pluripotent stem cells from the Massachusetts General Hospital miniature pig. *BioMed Res. Int.* **2013**, 140639.
- Lee, E., Lee, S.H., Kim, S., Jeong, Y.W., Kim, J.H., Koo, O.J., Park, S.M., Hashem, M.A., Hossein, M.S., Son, H.Y., Lee, C.K., Hwang, W.S., Kang, S.K. & Lee, B.C. (2006a). Analysis of nuclear reprogramming in cloned miniature pig embryos by expression of *Oct-4* and *Oct-4* related genes. *Biochem. Biophys. Res. Commun.* **348**, 1419–28.
- Lee, J., Go, Y., Kang, I., Han, Y.M. & Kim, J. (2010). Oct-4 controls cell-cycle progression of embryonic stem cells. *Biochem. J.* **426**, 171–81.
- Lee, J., Kim, H.K., Rho, J.Y., Han, Y.M. & Kim, J. (2006b). The human OCT-4 isoforms differ in their ability to confer self-renewal. *J. Biol. Chem.* **281**, 33554–65.
- Li, C., Yan, Y., Ji, W., Bao, L., Qian, H., Chen, L., Wu, M., Chen, H., Li, Z. & Su, C. (2012). OCT4 positively regulates Survivin expression to promote cancer cell proliferation and leads to poor prognosis in esophageal squamous cell carcinoma. *PLoS One* **7**, e49693.
- Liu, K., Ji, G., Mao, J., Liu, M., Wang, L., Chen, C. & Liu, L. (2012). Generation of porcine-induced pluripotent stem cells by using OCT4 and KLF4 porcine factors. *Cell. Reprogram.* **14**, 505–13.
- Livak, K.J. & Schmittgen, T.D. (2001). Analysis of relative gene expression data using real-time quantitative PCR and the $2^{-\Delta\Delta CT}$ method. *Methods* **25**, 402–8.
- Mountford, P., Nichols, J., Zevnik, B., O'Brien, C. & Smith, A. (1998). Maintenance of pluripotential embryonic stem cells by stem cell selection. *Reprod. Fertil. Dev.* **10**, 527–33.
- Nichols, J., Zevnik, B., Anastassiadis, K., Niwa, H., Klewe-Nebenius, D., Chambers, I., Scholer, H. & Smith, A. (1998). Formation of pluripotent stem cells in the mammalian embryo depends on the POU transcription factor Oct4. *Cell* **95**, 379–91.
- Niwa, H., Miyazaki, J. & Smith, A.G. (2000). Quantitative expression of Oct-3/4 defines differentiation, dedifferentiation or self-renewal of ES cells. *Nat. Genet.* **24**, 372–6.
- Okamoto, K., Okazawa, H., Okuda, A., Sakai, M., Muramatsu, M. & Hamada, H. (1990). A novel octamer binding transcription factor is differentially expressed in mouse embryonic cells. *Cell* **60**, 461–72.
- Park, K.M., Cha, S.H., Ahn, C. & Woo, H.M. (2013). Generation of porcine induced pluripotent stem cells and evaluation of their major histocompatibility complex protein expression *in vitro*. *Vet. Res. Commun.* **37**, 293–301.
- Park, S.J., Cho, B., Koo, O.J., Kim, H., Kang, J.T., Hurh, S., Kim, S.J., Yeom, H.J., Moon, J., Lee, E.M., Choi, J.Y., Hong, J.H., Jang, G., Hwang, J.I., Yang, J., Lee, B.C. & Ahn, C. (2014). Production and characterization of soluble human TNFR1-Fc and human HO-1(HMOX1) transgenic pigs by using the F2A peptide. *Transgenic Res.* **23**, 407–19.
- Pesce, M., Gross, M.K. & Scholer, H.R. (1998). In line with our ancestors: Oct-4 and the mammalian germ. *BioEssays* **20**, 722–32.
- Ruley, H.E. (1990). Transforming collaborations between Ras and nuclear oncogenes. *Cancer Cells* **2**, 258–68.
- Serrano, M., Lin, A.W., McCurrach, M.E., Beach, D. & Lowe, S.W. (1997). Oncogenic ras provokes premature cell senescence associated with accumulation of p53 and p16INK4a. *Cell* **88**, 593–602.
- Weinberg, R.A. (1989). Oncogenes, antioncogenes, and the molecular bases of multistep carcinogenesis. *Cancer Res.* **49**, 3713–21.

- Yoshioka, K., Suzuki, C., Tanaka, A., Anas, I.M. & Iwamura, S. (2002). Birth of piglets derived from porcine zygotes cultured in a chemically defined medium. *Biol. Reprod.* **66**, 112–9.
- Yuan, Y., Lee, K., Park, K.W., Spate, L.D., Prather, R.S., Wells, K.D. & Roberts, R.M. (2014). Cell cycle synchronization of leukemia inhibitory factor (LIF)-dependent porcine-induced pluripotent stem cells and the generation of cloned embryos. *Cell Cycle* **13**, 1265–76.
- Zhang, J.H., Chung, T.D. & Oldenburg, K.R. (1999). A simple statistical parameter for use in evaluation and validation of high throughput screening assays. *J. Biomol. Screen.* **4**, 67–73.
- Zhang, X., Neganova, I., Przyborski, S., Yang, C., Cooke, M., Atkinson, S.P., Anyfantis, G., Fenyk, S., Keith, W.N., Hoare, S.F., Hughes, O., Strachan, T., Stojkovic, M., Hinds, P.W., Armstrong, L. & Lako, M. (2009). A role for NANOG in G1 to S transition in human embryonic stem cells through direct binding of CDK6 and CDC25A. *J. Cell Biol.* **184**, 67–82.
- Zhang, Y., Wei, C., Zhang, P., Li, X., Liu, T., Pu, Y., Li, Y., Cao, Z., Cao, H., Liu, Y. & Zhang, X. (2014). Efficient reprogramming of naive-like induced pluripotent stem cells from porcine adipose-derived stem cells with a feeder-independent and serum-free system. *PLoS One* **9**, e85089.