

# No-till snap bean performance and weed response following rye and vetch cover crops

Rick A. Boydston<sup>1\*</sup> and Martin M. Williams II<sup>2</sup>

<sup>1</sup>United States Department of Agriculture, Agronomist, Agricultural Research Service, 24106 N. Bunn Road, Prosser, WA 99350, USA

<sup>2</sup>United States Department of Agriculture, Ecologist, Agricultural Research Service, 1102 S. Goodwin Avenue, Urbana, IL 61801, USA

\*Corresponding author: [rick.boydston@ars.usda.gov](mailto:rick.boydston@ars.usda.gov)

Accepted 9 October 2016; First published online 5 December 2016

Research Paper

## Abstract

Fall-planted cover crops offer many benefits including weed suppressive residues in spring sown crops when controlled and left on the soil surface. However, vegetable growers have been slow to adopt direct-seeding (no-till) into cover crop residues. Field studies were conducted in 2009 and 2010 near Paterson, WA and Urbana, IL to evaluate mortality of rye and common vetch (WA) hairy vetch (IL) cover crops, weed density and biomass, and snap bean growth and yield following four cover crop control methods utilizing a roller-crimper. Rye had higher mortality than common and hairy vetch by roller-crimping, and carfentrazone applied after roller crimping only slightly increased vetch mortality. Heavy residues of rye and escaped vetch were difficult to plant into, often resulting in lower snap bean populations. Rye and hairy vetch residues suppressed final weed biomass, while common vetch reduced weed biomass 1 of 2 years. Escaped plants of both vetch species became a weed. Snap bean yields were inconsistent and often lower following cover crops compared with a fallow treatment. Being able to completely control cover crops and to plant, manage escaped weeds and mechanically harvest in the presence of heavy residues are challenges that deter vegetable growers from readily adopting these systems.

**Key words:** cover crop, cereal rye, *Secale cereal* L., common vetch, *Vicia sativa* L., cover crop termination, hairy vetch, *Vicia villosa* Roth, no-till, organic, roller-crimper, snap bean, *Phaseolus vulgaris* L., weed management, weed suppression

## Introduction

Despite the soil-building benefits of no-till, most vegetable production systems have been slower to adapt no-till systems than agronomic crops for a variety of reasons; however, production of no-till vegetable crops increased during the 1990s (Morse, 1999). Mechanical weed control (cultivation) commonly practiced in many vegetable crops is not well adapted to conservation tillage systems, where one-third or more of the soil surface remains covered with crop residues (Peigne et al., 2007). Many vegetable crops are small-seeded and difficult to plant and establish in high residue no-till fields. An alternative system, strip till, prepares a more residue-free seed bed in strips between untilled areas (Hoyt et al., 1994; Luna et al., 2012). However, such systems forfeit the weed suppressive advantages of crop residues in the seed row. Organic vegetable growers have been even slower to

develop no-till systems primarily due to limited herbicide options for terminating cover crops and managing weeds, as well as the lack of cultivation tools that operate effectively in the presence of heavy surface residues (Carr et al., 2013).

Growing cover crops that produce large quantities of residue and leaving residues on the soil surface can be an effective way to reduce soil erosion, improve water infiltration, sequester nutrients, reduce runoff of pesticides and suppress annual weeds (Morse, 1999; Rice et al., 2001; Teasdale et al., 2007; Carr et al., 2013). Mowing or using burndown herbicide are common methods of controlling cover crops for no-till planting. Mowing or flail chopping often has to be repeated and can leave uneven residues on the soil surface, including small pieces of residue that decompose quickly (Creamer and Dabney, 2002). Roller-crimpers were developed to mechanically flatten and kill cover crops, leaving a weed suppressive mulch on the soil surface (Creamer and

Dabney, 2002; Ashford and Reeves, 2003; Mischler et al., 2010a; Carr et al., 2013). Unlike mowing or flail chopping, terminating tall grass cover crops by rolling and crimping leaves residues aligned uniformly, which facilitates planting of the main crop. Cereal rye (*Secale cereal* L.) and hairy vetch (*Vicia villosa* Roth) have been successfully terminated by the roller-crimper and provide some weed suppression (Ashford and Reeves, 2003; Mirsky et al., 2009; Davis, 2010; Mischler et al. 2010a, b).

For effective annual weed suppression with cover crop residues, a dense stand with high biomass is important. However, when cover crop residues are left as surface mulch, cultivating escaped weeds can be difficult. Thus, managing weeds in a no-till system following a poor cover crop stand or with escape weeds may be more difficult than in systems without mulch.

Large-seeded vegetable crops, such as beans, cucurbits, sweet corn or transplanted vegetable crops emerge and establish through heavy plant residues better than small-seeded vegetables, making them better candidates for no-till systems (Hoyt et al., 1994; Abdul-Baki et al., 1996; Mulvaney et al., 2011; Delate et al., 2012; Kornecki et al., 2012). However, yields of transplanted tomato, zucchini and bell pepper planted into roller-crimped rye or hairy vetch were reduced 41–92% compared with a no cover crop (i.e. fallow) treatment (Leavitt et al., 2011). Poor yields were attributed in part to insufficient soil nitrogen (N) and cold soil temperatures. Tomato and eggplant transplanted into organic no-till systems resulted in crop failures (Luna et al., 2012). Whereas many vegetable crops (collards, onions and zucchini) have been transplanted into no-till systems with high residues with success (Vollmer et al., 2010; Mulvaney et al., 2011; Canali et al., 2013), obtaining good seed-to-soil contact when direct-seeding vegetables can be a challenge (Morse, 1999; Davis, 2010).

Snap bean (*Phaseolus vulgaris* L.) is an important vegetable crop grown for fresh market and processing. The majority of snap bean production in the USA is for processing (66 360 ha) (NASS, 2015). Our study areas in Illinois and Washington represent two primary production areas of processed snap beans in the Midwest and Pacific Northwest (PNW) regions. Midwestern soils typically contain higher organic matter (O.M.) and clay, whereas mineral soils low in O.M. are more frequent in the PNW vegetable-growing region. No-till planting into cover crops in the PNW region is often practiced to maintain or increase soil O.M., reduce soil erosion and improve water infiltration, whereas Midwest producers tend to be more focused on reducing pesticide runoff, reclaiming nutrients and controlling weeds.

No-till planting of snap bean into hairy vetch residues controlled by flail chopping was successful, but required a post-emergence application of sethoxydim for control of grassy weeds (Abdul-Baki and Teasdale, 1997). Yields of no-till snap bean were similar to or greater

than yields of conventionally tilled fields in studies by Skarphol et al. (1987). However, others have reported yield losses for snap bean no-till planted into cover crop residues, citing problems with inadequate crop stands, poor weed control, N immobilization or release of allelochemicals from decomposing residues (Knave and Heron, 1986; Mwaja et al., 1996; Rutledge, 1999).

The objectives of this study were to: (1) quantify the effectiveness of control methods for rye and vetch cover crops when utilizing a roller-crimper and (2) evaluate the response of snap bean and weeds to these cover crops and cover crop control methods in two primary snap bean production regions of the USA.

## Materials and methods

### Site conditions and management

Field studies were conducted in 2009 and 2010 near Paterson, WA and Urbana, IL. Cover crops were seeded with a grain drill in 18-cm rows September 15, 2008 and September 17, 2009 at Paterson and September 26, 2008 and September 28, 2009 at Urbana (Table 1). Common vetch (*Vicia sativa* L.) (Paterson) or hairy vetch (Urbana) were seeded at 76 kg ha<sup>-1</sup>, rye at 100 kg ha<sup>-1</sup> (var. 'Aroostook' at Paterson and var. 'HiRye 500' at Urbana) and a rye + vetch mixture at 56 kg ha<sup>-1</sup> of each. A fallow treatment was included as a control. Soil at the Paterson site was a Quincy sand (mixed, mesic Xeric Torripsamments) with pH 7.0 and 0.4% O.M.. Soil at Urbana was a Flanagan silt loam (Fine, smectitic, mesic Aquic Argiudolls) with pH 6.1 and 4.0% O.M. Trials were irrigated as needed with a center pivot irrigation system at Paterson and grown under natural rainfall and supplemental sprinkler irrigation at Urbana. Experiments were arranged in a split plot randomized block design with four replications. Main plots were cover crops and split plots were cover crop control methods. Main plots were 12.2 m × 12.2 m and split plots were 3 m × 12.2 m.

Four cover crop control methods were tested at each site and the timing of various field operations are listed in Table 1. To terminate cover crops, plots were roller-crimped once or twice, with the second roller crimping 5–10 days (Paterson) or 2–4 days (Urbana) after the first roller-crimping. A tractor-mounted, 3-m-wide roller-crimper, previously described by Davis (2010), was used. Briefly, the roller-crimper consisted of a 3-m-long by 0.45-m-diameter steel cylinder with 7-cm fins protruding at 90° from the cylinder in a chevron pattern; the roller weighed approximately 5000 kg when filled with water. Cover crops were roller-crimped when rye reached 95% anthesis. Rye reached this growth stage at Paterson when common vetch had fully flowered and contained small flat pods, whereas hairy vetch was <50% flowered at Urbana. Because of the difficulty planting into heavy rye residues at Urbana in 2009, glyphosate

**Table 1.** Summary of field operations at Paterson, WA and Urbana, IL for snap bean cover crop trials in 2009 and 2010.

	Paterson, WA		Urbana, IL	
	2009	2010	2009	2010
Cover crops planted	September 15, 08	September 17, 09	September 26, 08	September 28, 09
First roller-crimping	May 22	May 27	May 29	May 25
Snap bean planting	May 22	June 7 <sup>1</sup>	June 2	May 26
Burndown herbicide <sup>2</sup>	May 27	June 5	June 2	May 25
Second roller-crimping <sup>3</sup>	May 27	June 7	June 2	May 27
Snap bean harvest	July 27–31	August 23–26	July 27–31	July 19–20

<sup>1</sup> Replanted due to cool soil temperatures and poor emergence.

<sup>2</sup> Selected treatments only.

<sup>3</sup> Only one selected treatment roller-crimped twice.

was applied at 1.1 kg ae ha<sup>-1</sup> in an 18 cm band over the soon to be planted four snap bean rows April 14, 2010 at Urbana by using a bicycle sprayer with four drop nozzles spaced 76 cm apart in rye and rye + vetch plots. Rye height averaged 24 cm at the time of the glyphosate application. A third cover crop control treatment consisted of roller crimping once and broadcast application of glyphosate (rye and rye + vetch) or carfentrazone (vetch). A fourth cover crop control treatment was the same as the third treatment, but included a soil residual herbicide (s-metolachlor) tank mixed with the burndown herbicide, followed by handweeding escaped weeds. The fourth treatment served as a weed-free control. Fallow plots also were roller-crimped.

Snap bean var. 'Sahara' was planted 3–4 cm deep with a four row no-till planter in rows spaced 76 cm on May 22, 2009 and May 27, 2010 at Paterson and June 2, 2009 and May 26, 2010 at Urbana. A Kinzie no-till planter with fluted coulters, double-disc openers and rubber press wheels was used to plant snap bean at a rate of 144 300 seed ha<sup>-1</sup> at Paterson. A Monosem no-till planter with fluted coulters, double-disc openers and rubber press wheels was used at Urbana at a rate of 303 700 seed ha<sup>-1</sup>. Differences in the planting rates at the two locations were due to differences in the ability to adjust seeding density on each planter. Due to poor emergence and cool weather conditions, snap bean was replanted at Paterson on June 7, 2010.

### Data collection

Aboveground biomass of cover crops was collected May 20, 2009 and May 25, 2010 at Paterson by sampling a 1 m<sup>2</sup> quadrat in each main plot and May 20, 2009 and May 19, 2010 at Urbana by sampling a 0.25 m<sup>2</sup> quadrat in each split plot. Samples were dried at 60°C for 5 days and reweighed to determine oven-dry biomass. At approximately 2 and 4 weeks after herbicide application, cover crop mortality was visually estimated June 8 and 23, 2009 and June 23 and July 8, 2010 at Paterson on a scale of 0 = no response to 100 = all plants dead. Cover

crop mortality was visually estimated June 12 and 29, 2009 and June 9 and 28, 2010 at Urbana.

Snap bean population was determined June 8, 2009 and July 8, 2010 at Paterson and June 12, 2009 and June 4, 2010 at Urbana by counting the number of seedlings from the middle two rows of each four row plot by 6 m. Snap bean percent bloom was visually estimated July 8, 2009 and July 27, 2010 at Paterson and July 10, 2009 and June 30, 2010 at Urbana.

Weed population density was recorded June 12, 2009 and June 16, 2010 at Paterson and June 12, 2009 and June 15, 2010 at Urbana by counting all weeds by species from a 1 m<sup>2</sup> quadrat placed over the middle two rows in each split plot. Within 1 week of snap bean harvest, final biomass of weeds and escaped cover crop were determined July 24, 2009 and August 20, 2010 at Paterson and July 31, 2009 and July 15, 2010 at Urbana. Biomass of weed species and escaped cover crops was determined by clipping all escaped plants from a randomly placed 1 m<sup>2</sup> quadrat in each subplot, separated by species, dried at 65°C and weighed.

Snap bean yield was estimated July 24, 2009 and August 23, 2010 at Paterson and July 27, 2009 and July 19, 2010 at Urbana by hand harvesting all pods from 1.5 m lengths (Paterson) or 1 m lengths (Urbana) of the two middle rows of each subplot and weighing.

### Data analysis

Analysis of variance (ANOVA) was performed using PROC GLIMMIX procedure in SAS (Statistical Analysis Systems®, 2014, version 9.4, SAS Institute Inc., SAS Campus Drive, Cary, NC 27513, USA) to test for significance ( $P < 0.05$ ) of years, treatments and their interactions for the response variables recorded. Variations in cover crop species, weed species, planters, weather, soil type and irrigation led to many interactions among locations, years and other measured variables; therefore, each site was analyzed and presented separately considering blocks and years as random factors. Data were pooled across site-years when no significant year-

by-treatment interaction was detected. Mean separation was conducted using Tukey–Kramer LSMEANS ( $P = 0.05$ ). When interactions were significant, LSMEANS tests were performed separately pooling years.

## Results and discussion

### Cover crop biomass

Rye produced 8045 and 8303 kg ha<sup>-1</sup> at Paterson, WA in 2009 and 2010, respectively. Common vetch produced 20 and 56% less biomass than rye in 2009 and 2010, respectively (Table 2). Biomass of mixtures of common vetch and rye was similar to rye monoculture in both years. Rye produced 12 096 and 10 752 kg ha<sup>-1</sup> at Urbana in 2009 and 2010, respectively. Hairy vetch produced 62–64% less biomass than rye in both years (Table 2). Rye + hairy vetch produced 20% less biomass than rye monoculture in 2009, but in 2010, rye + hairy vetch and rye monoculture produced similar amount of biomass. Using a 50:50 seed mix (by weight), a barley (*Hordeum vulgare* L.)-hairy vetch mixture produced similar biomass as a barley monoculture in studies by Wayman et al. (2014) in Washington State. When testing different proportions of rye and hairy vetch (0–100%), Hayden et al. (2014) reported total shoot biomass produced in mixes was usually greater than monocultures. Sainju et al. (2005) reported greater biomass production from rye + hairy vetch mixtures when seeded at a ratio of 68:32 (by weight), than from monocultures of either species in Georgia.

### Effects of roller-crimping cover crops

Cover crop mortality was impacted by cover crop species and there was significant cover crop by year and cover crop-by-control method interactions on mortality. At both sites and years, mortality of rye was greater than common vetch or hairy vetch by roller crimping once or twice (Table 3). Leavitt et al. (2011) also reported incomplete control of hairy vetch by roller-crimping. Mischler et al. (2010b) reported hairy vetch mortality with a roller-crimper improved after early pod set. Hairy vetch was <50% flowered at Urbana when roller-crimped in these studies, which contributed to poor control. Ideally hairy vetch would have been more developed; however, the advanced growth stage of rye at Urbana complicated waiting longer to roller-crimp. Nonetheless, at the Paterson site, common vetch was in the early pod stage both years and was not controlled well by roller-crimping. Escaped vetch plants also produced seeds, which volunteered in subsequent crops. Wayman et al. (2014) reported poor control of vetch species with roller crimping in Washington.

The primary weeds at Paterson were horseweed [*Conyza canadensis* (L.) Cronq.], large crabgrass [*Digitaria sanguinalis* (L.) Scop.], hairy nightshade

**Table 2.** Dry above ground biomass of fall-planted cover crops at Paterson, WA and Urbana, IL in 2009 and 2010.

Location	Year	Common vetch	Rye	Rye + vetch
Paterson	2009	6446 b	kg ha <sup>-1</sup>	8654 a
			8045 ab	
	2010	3686 b	8303 a	8562 a
			Hairy vetch	Rye
Urbana	2009	4413 c	kg ha <sup>-1</sup>	9632 b
			12096 a	
	2010	4075 b	10799 a	12352 a

Means within a row followed by the same letter are not significantly different according to Tukey–Kramer least significant different test ( $P = 0.05$ ).

(*Solanum physalifolium* Rusby) and common lambsquarters (*Chenopodium album* L.). Early season weed density at Paterson was relatively low in all plots (<6.5 m<sup>-2</sup>) in 2009 (Table 4). Weed density was greatest in fallow plots and rye treatments (6.0–6.5 m<sup>-2</sup>), whereas plots containing common vetch averaged <1.0 m<sup>-2</sup>. However, escaped vetch became a weed and may have contributed to suppression of other weeds. Weed densities were similar among cover crop and control treatments (excluding the weed-free treatment) in 2010. Common vetch may have had less impact on weed density in 2010 due to 43% less vetch biomass produced in 2010 compared to 2009 (Table 2). Weed densities also tended to be greater in 2010 and more variable.

Final weed biomass was reduced 66–89% in snap bean following cover crops compared with the fallow treatment, which averaged 213 g m<sup>-2</sup> at Paterson in 2009 (Table 4). Escaped common vetch also was present season long and was equal to or greater than the biomass of weeds in 2009 (Table 4). In 2010, common vetch comprised a lower portion of the weed biomass, but still contributed additional competition to snap bean. Common vetch terminated by flail chopping reduced percent weed cover similar to rye and more than hairy vetch in Washington studies (Wayman et al., 2014). In 2010, only the rye treatment reduced final weed biomass by 73% compared with the fallow treatment.

At Urbana, the main weeds present were common purslane (*Portulaca oleracea* L.), ivyleaf morningglory (*Ipomoea hederacea* Jacq.), common dandelion (*Taraxacum officinale* G.H. Weber ex Wiggers) and common lambsquarters. Cover crop treatments significantly reduced early season weed density, regardless of control method (Table 5). Weed densities averaged 4.9–6.8 weeds m<sup>-2</sup>; in comparison, the fallow treatment averaged 35.8 weeds m<sup>-2</sup> (Table 5).

Final weed biomass at Urbana was reduced by more than 50% by all cover crop treatments in 2009 compared with the fallow treatment, which averaged 168 g m<sup>-2</sup>

**Table 3.** Effect of cover crop control method on cover crop mortality at Paterson, WA and Urbana, IL in 2009 and 2010.

Cover crop	Control method	Cover crop mortality			
		Paterson		Urbana	
		2009 (%)	2010	2009 (%)	2010
Vetch <sup>1</sup>	Roller-crimped 1X	51 e	40 d	39 e	25 d
	Roller-crimped 2X	60 d	54 c	49 de	34 cd
	Roll + herb <sup>2</sup>	60 d	45 cd	61 cd	53 c
	Roll + herb + HW <sup>3</sup>	58 de	70 b	66 bc	56 bc
Rye	Roller-crimped 1X	91 b	98 a	99 a	99 a
	Roller-crimped 2X	95 ab	99 a	99 a	100 a
	Roll + herb	100 a	100 a	100 a	100 a
	Roll + herb + HW	100 a	100 a	100 a	100 a
Rye + vetch	Roller-crimped 1X	76 c	75 b	36 e	41 cd
	Roller-crimped 2X	78 c	81 b	35 e	48 cd
	Roll + herb	93 ab	97 a	83 ab	79 ab
	Roll + herb + HW	94 ab	99 a	85 ab	80 ab

<sup>1</sup> Common vetch was planted at Paterson, WA and hairy vetch was planted at Urbana, IL.

<sup>2</sup> In monoculture vetch plots, the burndown herbicide was carfentrazone, whereas the herbicide used in rye or rye + vetch was glyphosate.

<sup>3</sup> HW = handweeded and also included s-metolachlor tank mixed with the burndown herbicide.

Means within a column followed by a different letter are significantly different according to Tukey–Kramer least significant different test ( $P = 0.05$ ).

(Table 5). Rye and the rye + vetch reduced weed biomass the greatest. In 2010, weed biomass was not affected by cover crop treatment, but tended to be lowest in rye or rye + vetch. Mohler and Teasdale (1993) reported greater weed emergence through hairy vetch residue compared with equal amounts of rye residue. In both years at Urbana, escaped hairy vetch biomass was greater than weed biomass and likely posed significant competition to snap bean plants.

In studies by Hayden et al. (2014), winter annual weed control in rye + hairy vetch decreased as the proportion of hairy vetch in the mix was increased. Rye reduced weed biomass in our studies in all four site-years and rye biomass was at or above the 8000 kg ha<sup>-1</sup>, reported as a threshold needed for consistent suppression of annual weeds (Mirsky et al., 2013). We suspect that part of the weed suppressiveness observed from common and hairy vetch was a result of the competition of escaped vetch with weedy species.

Snap bean population averaged only 6.4 plants m<sup>-1</sup> (84 000 plants ha<sup>-1</sup>) at Paterson both years, well below typical commercial production of 13–26 plants m<sup>-1</sup> (Table 6). The relatively low snap bean population at Paterson could be attributed, in part, to low planting rates due to planter limitations, soil type and low soil temperatures. Nevertheless, our stands were comparable to Abdul-Baki and Teasdale (1997) and Mwaja et al. (1996) who reported snap bean populations were not affected by planting into killed hairy vetch residues. In contrast, snap bean populations in our research were lowest following common vetch or rye + vetch mixtures

that were roller-crimped, but not treated with herbicides (Table 6). Regardless of treatment, common vetch reduced snap bean population because the planter had difficulty closing seed furrows in the heavy cover crop residues.

At Urbana, both year and cover crop significantly affected snap bean populations. Snap bean population averaged 17 plants m<sup>-1</sup> in 2009 and 24 plants m<sup>-1</sup> in 2010 (Table 7). Snap bean populations were lowest (9.8 plants m<sup>-1</sup>) following rye in 2009 because double-disc openers lining up on the rye row had difficulty penetrating rye roots, and when openers did penetrate, press wheels did not close the seed furrow completely in the high-residue situation. When the disc-openers were off the row, no problems with the seed furrow were observed. In 2010 at Urbana, glyphosate was applied in a narrow band over the snap bean row approximately 6 weeks prior to planting snap bean in plots containing rye and increased snap bean populations (24 plants m<sup>-1</sup> row) (Table 7). In these studies, rye was roller-crimped and snap bean planted in the same direction as rye seeding. More recently, Mirsky et al. (2013) advised against planting the main crop in the same direction as the rye seeding to avoid the problem we experienced in 2009.

In previous studies, snap bean populations direct-seeded into killed hairy vetch were 62% of those planted into conventionally tilled bare soil (Knavel and Heron, 1986). Soybean (*Glycine max*) populations were reduced by 60% of targeted plant population when planted into roller-crimped rye (Mischler et al., 2010a). Those authors reported earlier termination of rye reduced rye

**Table 4.** Early season weed density and final season weed and escaped cover crop biomass in snap bean following cover crops and control methods in 2009 and 2010 at Paterson, WA.

Cover crop	Control method	Weed density		Weed biomass		Escaped cover crop biomass	
		6/12/09	6/16/10	7/24/09	8/20/10	7/24/09	8/20/10
		(no. m <sup>-2</sup> )		(g m <sup>2</sup> )		(g m <sup>2</sup> )	
Common vetch	Roller-crimped 1X	0.6	3.0	76.7	243.2	66.1	25.1
	Roller-crimped 2X	1.1	1.3	71.2	229.7	78.5	17.5
	Roll + herb <sup>1</sup>	1.0	2.3	68.0	408.5	62.4	10.9
Rye		0.9 b			72.0 b	293.8 a	43.4 a
	Roller-crimped 1X	5.6	10.5	35.0	68.2	1.6	0.6
	Roller-crimped 2X	10.3	10.0	44.4	61.3	2.9	0.5
	Roll + herb	2.3	0.5	88.0	63.2	0	0.0
Rye + vetch		6.0 a			55.8 b	64.2 c	0.9 b
	Roller-crimped 1X	0.1	11.8	20.9	90.7	64.3	36.9
	Roller-crimped 2X	0.3	7.0	18.9	142.9	90.5	32.3
	Roll + herb	0.1	0.8	30.1	118.2	69.2	4.9
Fallow		0.2 b			23.2 b	117.3 bc	49.7 a
	Roller-crimped 1X	5.3	5.5	192.6	205.3	–	–
	Roller-crimped 2X	6.6	0.8	190.0	241.6	–	–
	Roll + herb	7.5	12.0	257.5	286.4	–	–
Main effect		6.5 a			213.3 a	244.4 ab	
	Roller-crimped 1X	–	–	87.5 b	151.8 b	–	–
	Roller-crimped 2X	–	–	81.1 b	168.9 ab	–	–
	Roll + herb	–	–	110.9 a	219.1 a	–	–
Cover crop (CC)		0.3711		0.0015		<0.0037	
Control method (CM)		0.3284		0.0027		0.2146	
Year		0.3619		0.0865		0.1106	
CC × CM		0.0116		0.2921		0.7738	
CC × year		0.7100		0.2114		0.1467	
CM × year		0.2121		0.2648		0.4085	
CC × CM × year		0.3874		0.0263		0.8259	

<sup>1</sup> In monoculture vetch plots, the burndown herbicide was carfentrazone, whereas the herbicide used in rye or rye + vetch was glyphosate.

Snap bean was planted May 22, 2009 and May 27, 2010 and replanted June 17, 2010 and harvested July 27–31, 2009 and August 23, 2010.

Means followed by a different letter within a column and within a main or simple effect are significantly different according to Tukey–Kramer least significant different test ( $P = 0.05$ ).

biomass and increased soybean plant populations relative to later termination dates. Soybean populations were reduced 30 and 17% in roller-crimped rye and hairy vetch, respectively, compared with fallow (Davis, 2010).

At Paterson, snap bean bloom in July was affected by cover crop, cover crop control method, year and all interactions between these factors. In both years, snap bean bloom was delayed most by cover crop treatments containing common vetch (Table 6). Snap bean bloom also was delayed following rye compared with fallow plots, but not to the extent as when following common vetch.

At Urbana, snap bean bloom was reduced most following cover crop treatments including rye in 2009 and was only slightly delayed when following hairy vetch (Table 7). However, in 2010, bloom was delayed most in

snap bean following hairy vetch or rye + vetch. Planting dry bean (*P. vulgaris* L.) into fall-planted rye residues delayed maturity by 2–6 days in Canadian studies (Blackshaw and Molnar, 2008) and Mwaja et al. (1996) reported delayed maturity with no-till snap bean.

At Paterson, snap bean yield was influenced by cover crop and control method; there was a significant cover crop-by-control method interaction. There were also significant year-by-control method and year-by-cover crop-by-control method interactions on snap bean yield (Table 6). Snap bean following common vetch or rye + vetch tended to yield lower even when kept weed-free compared with snap bean following rye or fallow. Snap bean following common vetch controlled by roller crimping either once or twice without a herbicide averaged only 2331–2472 kg ha<sup>-1</sup> in 2009 and 4382–5480 kg ha<sup>-1</sup> in

**Table 5.** Early-season weed density and final weed and escaped cover crop biomass in snap bean following cover crop treatments and control methods in 2009–10 at Urbana, IL.

Cover crop	Control method	Weed density		Weed biomass		Escaped cover crop biomass	
		6/12/09	6/15/10	7/31/09	7/15/10	7/24/09	7/15/10
		(no. m <sup>-2</sup> )		(g m <sup>2</sup> )		(g m <sup>2</sup> )	
Hairy vetch	Roller-crimped 1X	3.3	1.3	66.7	29.6	111	65.4
	Roller-crimped 2X	7.8	8.5	93.7	75.1	110	85.0
	Roll + herb <sup>1</sup>	3.8	5.0	86.3	13.5	152	58.1
		4.9 b		82.2 b		124 b	69.5 a
Rye	Roller-crimped 1X	1.5	12.5	29.1	24.6	0	0
	Roller-crimped 2X	0.3	15.8	0	21.2	0	0
	Roll + herb	0.8	10.3	28.5	13.2	0	0
		6.8 b		19.2 c		0 c	0 b
Rye + vetch	Roller-crimped 1X	3.3	10.8	16.5	22.2	178	35.3
	Roller-crimped 2X	3.0	10.8	29.2	20.8	159	48.8
	Roll + herb	1.8	3.8	0	8.8	189	19.8
		5.5 b		15.2 c		175 a	34.6 a
Fallow	Roller-crimped 1X	35.0	31.3	144.3	51.4	–	–
	Roller-crimped 2X	47.3	22.2	206.1	63.9	–	–
	Roll + herb	47.5	31.5	153.9	59.2	–	–
		35.8 a		168.1 a			
Cover crop (CC)		<0.0001		<0.0001		<0.0001	
Control method (CM)		0.6982		0.1051		0.8143	
Year		0.8432		0.0023		0.0610	
CC × CM		0.7280		0.3398		0.8555	
CC × year		0.0756		0.0014		<0.0001	
CM × year		0.6910		0.8247		0.0248	
CC × CM × year		0.5869		0.4847		0.3639	

<sup>1</sup> In monoculture vetch plots, the burndown herbicide was carfentrazone, whereas the herbicide used in rye or rye + vetch was glyphosate.

Snap bean was planted June 2, 2009 and May 26, 2010 and harvested July 27–31, 2009 and July 19–20, 2010.

Means followed by a different letter within a column and within a main or simple effect are significantly different according to Tukey–Kramer least significant different test ( $P = 0.05$ ).

2010 due to low plant populations and to competition from both weeds and escaped vetch.

At Urbana, both cover crop and control method influenced snap bean yield and there was a significant cover crop-by-control method interaction. Snap bean after fallow tended to yield the greatest both years at Urbana (Table 7). Snap bean yields were low (<2600 kg ha<sup>-1</sup>) following all rye containing treatments. In 2009, low snap bean yield following rye in Urbana was believed to be due in part to the low plant populations resulting from poor seed-to-soil contact from planting into heavy rye residues. However, in 2010 when rye was terminated with band-applied glyphosate prior to planting and snap bean populations were increased, snap bean yields still averaged only 2600 kg ha<sup>-2</sup> in the hand-weeded treatment. Previous studies have reported yield loss due to N immobilization from high C:N ratio of rye residues (Bottenberg et al., 1997). Rye biomass at Urbana averaged over 10 700 kg ha<sup>-1</sup> in both years. Such high levels

of rye biomass likely reduced the availability of N for snap bean in the present work.

In previous research, snap bean yielded greater in conventional tilled plots than when planted into completely controlled rye or hairy vetch (Mwaja et al., 1996). Those authors did not observe increases in disease incidence, stand reductions or problems planting no-till into glyphosate-killed cover crops, but indicated a delay in snap bean maturity following cover crops. Knavel and Heron (1986) also reported greater snap bean yield in conventional tilled plots compared to no-till planted into hairy vetch. In those studies, snap bean population was reduced due to difficulty planting into hairy vetch residues despite using a no-till planter. Recent improvements in no-till planter equipment and management of cover crop residues could improve the ability to maintain plant populations in these systems (Mirsky et al., 2013).

Snap bean was harvested by hand in the current studies, overcoming any potential machine harvest losses or

**Table 6.** Snap bean stand, percent bloom and pod yield following cover crop treatments and control methods in 2009 and 2010 at Paterson, WA.

Cover crop	Control method	Plant stand		Percent bloom		Pod yield	
		6/8/09	7/08/10	7/08/09	7/27/10	7/27/09	8/23/10
		(no. m <sup>-1</sup> row)		(%)		(kg ha <sup>-1</sup> )	
Common vetch	Roller-crimped 1X	5.5	4.0	25 de	9 d	2472 de	5480 cd
	Roller-crimped 2X	5.6	3.9	25 de	8 d	2331 e	4382 d
	Roll + herb <sup>1</sup>	5.3	5.7	26 de	8 d	2842 de	6027 cd
	Roll + herb + HW <sup>2</sup>	5.6	6.4	29 de	13 cd	5257 bcd	10660 abc
		5.3 b		–	–	–	–
Rye	Roller-crimped 1X	7.9	5.3	33 de	12 cd	3741 de	7334 bcd
	Roller-crimped 2X	8.6	6.3	49 cd	13 cd	4483 cde	7423 bcd
	Roll + herb	8.2	7.8	64 bc	19 bcd	7837 ab	10494 abc
	Roll + herb + HW	8.4	7.3	73 ab	18 bcd	8937 a	9688 abcd
		7.5 a		–	–	–	–
Rye + vetch	Roller-crimped 1X	3.6	4.5	19 e	9 d	3498 de	6511 cd
	Roller-crimped 2X	5.1	5.5	20 de	9 d	2331 de	7010 cd
	Roll + herb	6.1	5.9	30 de	12 cd	2914 de	9991 abcd
	Roll + herb + HW	6.0	5.5	34 de	18 bcd	3974 de	12982 ab
		5.3 b		–	–	–	–
Fallow	Roller-crimped 1X	7.7	8.2	100 a	24 bc	3718 de	4839 cd
	Roller-crimped 2X	7.9	7.5	100 a	27 b	3933 de	6366 cd
	Roll + herb	7.7	7.5	100 a	25 bc	4707 bcde	6745 cd
	Roll + herb + HW	8.0	7.7	100 a	36 a	7644 abc	14270 a
		7.8 a		–	–	–	–
Main effect	Roller-crimped 1X		5.8 b	–	–	–	–
	Roller-crimped 2X		6.3 ab	–	–	–	–
	Roll + herb		6.8 a	–	–	–	–
	Roll + herb + HW		6.9 a	–	–	–	–
Cover crop (CC)		0.0004		<0.0001		0.0163	
Control method (CM)		0.0032		<0.0001		<0.0001	
Year		0.1169		<0.0001		0.0888	
CC × CM		0.3251		0.0004		0.0038	
CC × year		0.2116		<0.0001		0.1487	
CM × year		0.4429		0.0465		0.0043	
CC × CM × year		0.2982		0.0017		0.0086	

<sup>1</sup> In monoculture vetch plots, the burndown herbicide was carfentrazone, whereas the herbicide used in rye or rye + vetch was glyphosate.

<sup>2</sup> HW=handweeded and also included s-metolachlor tank mixed with the burndown herbicide.

Snap bean was planted May 22, 2009 and May 27, 2010 and replanted June 17, 2010.

Means followed by a different letter within a column and within a main or simple effect are significantly different according to Tukey–Kramer least significant different test ( $P = 0.05$ ).

difficulties encountered with heavy cover crop residues or weeds. Commercially grown snap bean is harvested mechanically with large rotating reels with fingers that strip bean pods from the plants. Cover crop plant residues and escaped weeds may slow harvest and increase foreign matter in harvested product.

### Effects of addition of herbicides

Common and hairy vetch were not completely controlled by roller crimping even with application of carfentrazone (Table 3). Carfentrazone was selected to control hairy and common vetch rather than glyphosate due to reports of

incomplete control of hairy vetch with glyphosate (Shite and Worsham, 1990). For control of mixtures, glyphosate was used because carfentrazone does not control rye. After roller crimping, control of mixtures with glyphosate was greater than control of common vetch (Paterson) or hairy vetch (Urbana) monocultures with carfentrazone (Table 3). Curran et al. (2015) also reported incomplete control of hairy vetch with carfentrazone.

Weed biomass was greater when including a burndown herbicide (glyphosate or carfentrazone) with roller crimping compared with roller crimping once or twice at Paterson in 2009 and 2010 (Table 4). We speculate that including the burndown herbicide controlled more of



**Table 7.** Snap bean stand, percent bloom and pod yield following cover crop treatments and control methods in 2009 and 2010 at Urbana, IL.

Cover crop	Control method	Plant stand		Percent bloom		Pod yield	
		6/12/09	6/4/10	7/10/09	6/30/10	7/27/09	7/20/10
		(no. m <sup>-1</sup> row)		(%)		(kg ha <sup>-1</sup> )	
Hairy vetch	Roller-crimped 1X	18	21	69	15 f-h	4565	2091 ab
	Roller-crimped 2X	19	21	74	19 e-h	5581	1714 ab
	Roll + herb <sup>1</sup>	22	25	83	30 d-h	5827	2969 ab
	Roll + herb + HW <sup>2</sup>	16	23	76	32 d-h	7417	5061 a
		18.7 ab	22.3 b	75.4 a	–	5847 ab	–
Rye	Roller-crimped 1X	13	24	24	41b-f	1450	1690 ab
	Roller-crimped 2X	11	25	12	52 a-d	1123	1870 ab
	Roll + herb	5	23	26	47 a-d	1221	2215 ab
	Roll + herb + HW	10	23	26	45 a-e	1860	2600 ab
		9.8 c	24.1 ab	21.9 b	–	1414 c	–
Rye + vetch	Roller-crimped 1X	17	24	26	8 h	1221	664 b
	Roller-crimped 2X	17	23	38	12 gh	1172	812 b
	Roll + herb	16	24	50	35 c-g	2254	2526 ab
	Roll + herb + HW	16	24	34	32 d-h	2073	2477 ab
		16.6 b	23.8 ab	37.1 b	–	1680 bc	–
Fallow	Roller-crimped 1X	25	25	93	69 a	8663	4347 a
	Roller-crimped 2X	25	26	84	61 abc	7925	3994 ab
	Roll + herb	24	25	93	69 a	7286	4429 a
	Roll + herb + HW	21	26	89	67 ab	9073	4815 a
		23.7 a	25.3 a	89.4 a	–	8237 a	–
Main effect	Roller-crimped 1X	–	–	–	–	3975 b	–
	Roller-crimped 2X	–	–	–	–	3950 b	–
	Roll + herb	–	–	–	–	4147 ab	–
	Roll + herb + HW	–	–	–	–	5106 a	–
Cover crop (CC)	<0.0001		<0.0001		0.0003		
Control method (CM)	0.3747		0.0004		<0.0001		
Year	<0.0001		0.1158		0.2192		
CC × CM	0.0243		0.1217		0.0015		
CC × year	0.0004		<0.0001		0.0358		
CM × year	0.1174		0.6111		0.3050		
CC × CM × Year	0.5219		0.4103		0.7577		

<sup>1</sup> In monoculture vetch plots, the burndown herbicide was carfentrazone, whereas the herbicide used in rye or rye + vetch was glyphosate.

<sup>2</sup> HW=handweeded and also included s-metolachlor tank mixed with the burndown herbicide.

Snap bean was planted June 2, 2009 and May 26, 2010. In 2010, glyphosate was banded over rows to control rye prior to planting snap bean.

Means followed by a different letter within a column and within a main or simple effect are significantly different according to Tukey–Kramer least significant different test ( $P = 0.05$ ).

the cover crop, thereby decreasing competition of the cover crop with weeds. In previous research, both rye and hairy vetch residues left on the soil surface reduced weed density in a no-till corn system, but in the absence of a residual herbicide, weed populations increased to a severe level and residues did not always reduce final weed biomass (Teasdale et al., 1991). Weed biomass and density were reduced in soybean direct seeded into a rye cover crop terminated by roller crimping and herbicides in three of four site-years (Mischler et al., 2010a).

In 2009 at Paterson, application of glyphosate following roller crimping of rye helped prevent some of the

delay in bloom of snap bean observed when only roller crimping rye (Table 6). At Urbana, bloom of snap bean following cover crops controlled by roller crimping (once or twice) without herbicide also tended to be delayed compared to control methods including an herbicide (Table 7). Competition from escaped weeds and escaped vetch likely delayed snap bean bloom in plots where the roller-crimper was used without herbicides.

At Paterson, including herbicides to kill rye improved snap bean yield equal to snap beans following fallow, hand-weeded controls in both years (Table 6). When including an herbicide to kill hairy vetch at Urbana,

snap bean yield was comparable with snap bean after fallow (Table 7).

## Conclusions

Producing snap beans after cover crops without tillage proved challenging due to escaped vetch plants and heavy rye residue. Although a later planted crop like snap bean allows for increased production of weed-suppressive cover crop biomass in the spring, the large amount of residue complicates planting. Planters can be modified and are available to direct seed into dense cover crop residues (Mirsky et al., 2013). We used no-till planters in the present studies, but further modifications (increased weight, row cleaners, etc.) might improve the ability to obtain adequate plant populations. Short-maturity rye cultivars are available that may allow for earlier roller-crimping, although cover crop biomass and weed suppressiveness, may be sacrificed (Wells et al., 2015). Roller-crimped cover crops left on the soil surface suppress weeds, but not to the levels typically observed with residual herbicides or multiple cultivations. In addition, escaped vetch became an additional weed. Growers electing to direct-seed snap bean following these cover crops without herbicides would likely need to invest in specialized high-residue cultivation equipment for season-long control of weeds to prevent crop losses (Carr et al., 2012; Mirsky et al., 2013). Overcoming these obstacles is critical for snap bean growers to benefit from the many advantages offered by cover crops in a no-till system.

**Acknowledgements.** The authors thank AgriNorthwest for providing land and irrigation water for these studies at Paterson, WA. The authors thank Bernardo Chaves-Cordoba for statistical analysis. The authors also thank Encarnacion Rivera, Treva Anderson, Marc Seymour and Jim Moody for technical assistance.

## References

- Abdul-Baki, A.A. and Teasdale, J.R. 1997. Snap bean production in conventional tillage and in no-till hairy vetch mulch. *HortScience* 32:1191–1193.
- Abdul-Baki, A.A., Teasdale, J.R., Korcak, R.F., Chitwood, D.J., and Huettel, R.N. 1996. Fresh-market tomato production in a low-input alternative system using cover-crop mulch. *Journal of the American Society of Horticultural Science* 31:65–69.
- Ashford, D.L. and Reeves, D.W. 2003. Use of a mechanical roller-crimper as an alternative kill method for cover crops. *American Journal of Alternative Agriculture* 18:37–45.
- Blackshaw, R.E. and Molnar, L.J. 2008. Integration of conservation tillage and herbicides for sustainable dry bean production. *Weed Technology* 22:168–176.
- Bottenberg, H., Masiunas, J., Eastman, C., and Eastburn, D.M. 1997. The impact of rye cover crops on weeds, insects, and diseases in snap bean cropping systems. *Journal of Sustainable Agriculture* 9:131–155.
- Canali, S., Campanelli, G., Ciaccia, C., Leteo, F., Testani, E., and Montemurro, F. 2013. Conservation tillage strategy based on the roller crimper technology for weed control in Mediterranean vegetable organic cropping systems. *European Journal of Agronomy* 50:11–18.
- Carr, P.M., Anderson, R.L., Lawley, Y.E., Miller, P.R., and Zwinger, S.F. 2012. Organic zero-till in the northern US great plains region: opportunities and obstacles. *Renewable Agriculture and Food Systems* 27:12–20.
- Carr, P.M., Gramig, G.G., and Liebzig, M.A. 2013. Impacts of organic zero tillage systems on crops, weeds, and soil quality. *Sustainability* 5:3172–3201.
- Creamer, N.G. and Dabney, S.M. 2002. Killing cover crops mechanically: review of recent literature and assessment of new research results. *American Journal of Alternative Agriculture* 17:32–40.
- Curran, W.S., Wallace, J.M., Mirsky, S., and Crockett, B. 2015. Effectiveness of herbicides for control of hairy vetch (*Vicia villosa*) in winter wheat. *Weed Technology* 29:509–518.
- Davis, A.S. 2010. Cover-crop roller-crimper contributes to weed management in no-till soybean. *Weed Science* 58:300–309.
- Delate, L., Cwach, D., and Chase, C. 2012. Organic no-till system effects on organic soybean, corn, and tomato production and economic performance in Iowa. *Renewable Agriculture and Food Systems* 27:49–59.
- Hayden, Z.D., Ngouajio, M., and Brainard, D.C. 2014. Rye-vetch mixture proportion tradeoffs: cover crop productivity, nitrogen accumulation, and weed suppression. *Agronomy Journal* 106:904–914.
- Hoyt, G.D., Monk, D.W., and Monaco, T.J. 1994. Conservation tillage for vegetable production. *HortTechnology* 4:129–135.
- Knavel, D.E. and Heron, J.W. 1986. Response of vegetable crops to nitrogen rates in tillage systems with and without vetch and rye grass. *Journal of the American Society of Horticultural Science* 111:502–507.
- Kornecki, T.S., Arriaga, F.J., and Price, A.J. 2012. Roller type and operating speed effects on rye termination rates, soil moisture, and yield of sweet corn in a no-till system. *HortScience* 7:217–223.
- Leavitt, M.J., Sheaffer, C.C., and Wyse, D.L. 2011. Rolled winter rye and hairy vetch cover crops lower weed density but reduce vegetable yields in no-tillage organic production. *HortScience* 46:387–395.
- Luna, J.M., Mitchell, J., and Shrestha, A. 2012. Conservation tillage for organic agriculture: evolution toward a hybrid system. *Renewable Agriculture and Food Systems* 27:21–30.
- Mirsky, S., Curran, W.S., Mortensen, D.A., Ryan, M.R., and Shumway, D. 2009. Control of cereal rye with a roller/crimper as influenced by cover crop phenology. *Agronomy Journal* 101:1589–1596.
- Mirsky, S.B., Ryan, M.R., Teasdale, J.R., Curran, W.S., Reberg-Horton, C.S., Spargo, J.T., Wells, M.S., Keene, C.L., and Moyer, J.W. 2013. Overcoming weed management challenges in cover crop-based organic rotational no-till soybean production in the Eastern United States. *Weed Technology* 27:193–203.
- Mischler, R.A., Curran, W.S., Duiker, S.W., and Hyde, J. 2010a. Rolling a rye cover crop for weed suppression in no-till soybeans. *Weed Technology* 24:253–261.
- Mischler, R.A., Duiker, S.W., Curran, W.S., and Wilson, D. 2010b. Hairy vetch management for no-till organic corn production. *Agronomy Journal* 102:355–362.

- Mohler, C.L. and Teasdale, J.R.** 1993. Response of weed emergence to rate of *Vicia villosa* Roth and *Secale cereal* L. residue. *Weed Research* 33:487–499.
- Morse, R.** 1999. No-till vegetable production—Its time is now. *HortTechnology* 9:373–379.
- Mulvaney, M., Price, A., and Wood, C.W.** 2011. Cover crop residue and organic mulches provide weed control during limited-input no-till collard production. *Journal of Sustainable Agriculture* 35:312–328.
- Mwaja, V.N., Masiunas, J.B., and Eastman, C.E.** 1996. Rye (*Secale cereal* L.) and hairy vetch (*Vicia villosa* Roth) inter-crop management in fresh-market vegetables. *Journal of the American Society of Horticultural Science* 121:586–591.
- National Agricultural Statistics Service (NASS), United States Department of Agriculture** 2015. Vegetables. ISSN:1931-2857. [usda.mannlib.cornell.edu/usda/nass/vege//2010s/2015/Vege-09-04-2015.pdf](http://usda.mannlib.cornell.edu/usda/nass/vege//2010s/2015/Vege-09-04-2015.pdf) (accessed November 2, 2016).
- Peigne, J., Ball, B.C., Roger-Estrade, J., and David, C.** 2007. Is conservation tillage suitable for organic farming? A review. *Soil Use and Management* 23:129–144.
- Rice, P.J., McConnell, L.L., Heighton, L.P., Sadeghi, A.M., Isensee, A.R., Teasdale, J.R., Abdul-Baki, A.A., Harman-Fetcho, J.A., and Hapeman, C.J.** 2001. Runoff loss of pesticides and soil: a comparison between vegetative mulch and plastic mulch in vegetable production systems. *Journal of Environmental Quality* 30:1808–1821.
- Rutledge, A.D.** 1999. Experiences with conservation tillage vegetables in Tennessee. *HortTechnology* 9:366–372.
- Sainju, U.M., Whitehead, W.F., and Singh, B.P.** 2005. Biculture legume–cereal cover crops for enhanced biomass yield and carbon and nitrogen. *Agronomy Journal* 97:1403–1412.
- Shite, R.H. and Worsham, D.** 1990. Control of legume cover crops in no-till corn (*Zea mays*) and Cotton (*Gossypium hirsutum*). *Weed Technology* 4:57–62.
- Skarphol, B.J., Corey, K.A., and Meisinger, J.J.** 1987. Response of snap beans to tillage and cover crop combinations. *Journal of the American Society of Horticultural Science* 112:936–941.
- Statistical Analysis System (SAS)** 2014. Version 9.4. SAS User's Guide. Statistical Analysis Systems Institute, Cary, NC.
- Teasdale, J.R., Beste, C.E., and Potts, W.E.** 1991. Response of weeds to tillage and cover crop residue. *Weed Science* 39: 195–199.
- Teasdale, J.R., Brandsaeter, L.O., Calegari, A., and Skora Neto, F.** 2007. Cover crops and weed management. In M. K. Upadhyaya and R.E. Blackshaw (eds.). *Non-Chemical Weed Management*. CAB International, Wallingford, UK. p. 49–64.
- Vollmer, E.R., Creamer, N., Reberg-Horton, C., and Hoyt, G.** 2010. Evaluation cover crop mulches for no-till organic production of onions. *HortScience* 45:61–70.
- Wayman, S., Cogger, C., Benedict, C., Burke, I., Collins, D., and Bary, A.** 2014. The influence of cover crop variety, termination timing and termination method on mulch, weed cover and soil nitrate in reduced-tillage organic systems. *Renewable Agriculture and Food Systems*. Available on CJO2014. doi: 10.1017/S1742170514000246.
- Wells, M.S., Brinton, C.M., and Reberg-Horton, S.C.** 2015. Weed suppression and soybean yield in a no-till cover-crop mulched system as influenced by six rye cultivars. *Renewable Agriculture and Food Systems*, Available on CJO2015. doi: 10.1017/S1742170515000344.