

Vegetable Soybean Tolerance to Pyroxasulfone

Martin M. Williams II, Nicholas E. Hausman, and James L. Moody*

If registered for use on vegetable soybean, pyroxasulfone would expand the options for weed management systems in the crop. In order to determine the potential crop injury risk of pyroxasulfone on vegetable soybean, the objective of this work was to quantify vegetable soybean tolerance to pyroxasulfone applied PRE and EPOST. Twenty-one vegetable soybean and two grain-type soybean cultivars were treated with pyroxasulfone at 417 g ai ha⁻¹ (twice the recommended field use rate) PRE, EPOST, or not treated. Plant population density was unaffected by pyroxasulfone. Only low levels (<10%) of crop injury were observed within a few weeks after PRE and EPOST treatments. Soybean cultivars were not differentially affected by pyroxasulfone, as evidenced by the lack of interactions between cultivar and treatment for any crop response variable. The low amount of risk of crop injury associated with pyroxasulfone is no different for vegetable soybean cultivars grown in the US for commercial production than grain-type soybean.

Nomenclature: Bentazon; fomesafen; imazamox; linuron; pyroxasulfone; sulfentrazone; soybean, *Glycine max* (L.) Merr.

Key words: Crop injury, edamame, herbicide tolerance, minor crop.

The number of herbicides registered by the US Environmental Protection Agency for use on vegetable soybean, also known as edamame, has increased in recent years. Prior to the registration of *S*-metolachlor in 2010 (Anonymous 2010), sethoxydim appears to have been the only herbicide with a federal registration (i.e., Section 3) for use on vegetable soybean. Since then, additional herbicides have received a federal label, including bentazon (Anonymous 2015a), fomesafen (Anonymous 2014a), imazamox (Anonymous 2013c), linuron (Anonymous 2012), and trifluralin (Anonymous 2014b). Additionally, special local needs labels (i.e., Section 24c) have been issued for imazethapyr (Anonymous 2013a) and sulfentrazone (Anonymous 2013b). One consideration for herbicide registration on “minor use” crops is the need to confirm crop tolerance to the proposed herbicide. For instance, recent research showed that bentazon, fomesafen, imazamox, linuron, and sulfentrazone posed no greater risk of crop injury to vegetable soybean than they do to grain-type soybean (Williams and Nelson 2014). Such information facilitates the herbicide registration process.

Despite these advances, vegetable growers and processors need greater development of weed management systems for vegetable soybean. Recent analysis of available weed management treatments showed that a number of improvements are needed, including better crop establishment, wider use of non-chemical tactics, and registration of additional herbicides (Williams 2015a).

Pyroxasulfone is being considered for use on vegetable soybean. Pyroxasulfone is a group 15 herbicide that inhibits very long chain fatty acid (VLCFA) synthesis (Tanetani et al. 2011). The herbicide can be applied prior to crop emergence (PRE) or early postemergence (EPOST) (Anonymous 2015b). While pyroxasulfone can be applied over emerged crops, it will not control emerged weeds. Pyroxasulfone safety in grain-type soybean appears comparable to that of other VLCFA-inhibiting herbicides such as *S*-metolachlor, but pyroxasulfone is used at a lower application rate (Yamaji et al. 2014). Compared to dimethenamid and *S*-metolachlor, pyroxasulfone can provide better control of certain broadleaf weed species (Yamaji et al. 2014), including multiple herbicide-resistant waterhemp

DOI: 10.1017/wet.2017.11

* Ecologist, Agricultural Science Technician, and Biological Science Technician, United States Department of Agriculture–Agricultural Research Service, Global Change and Photosynthesis Research Unit, Urbana, IL 61801. Corresponding author’s E-mail: Martin.Williams@ars.usda.gov

(Hausman et al. 2013), a common weed in vegetable soybean production. Moreover, tank-mixing pyroxasulfone with sulfentrazone enhanced weed control compared to the herbicides applied individually (Belfry and McNaughton et al. 2015).

In order to determine if pyroxasulfone is suitable for use on vegetable soybean, crop tolerance to the herbicide needs to be confirmed. Pyroxasulfone has been registered for use on grain-type soybean for several years. This work aimed to resolve whether or not vegetable soybean tolerance to pyroxasulfone differs from that of grain-type soybean. The objective was to quantify vegetable soybean tolerance to pyroxasulfone applied PRE and EPOST.

Materials and Methods

Germplasm. Vegetable soybean cultivars were selected based on availability of seed of cultivars grown commercially in the United States. A total of 21 vegetable soybean cultivars were obtained from seven private and four public sources (Table 1). In addition, two grain-type soybean cultivars widely grown in Illinois were included as controls. The grain-type

cultivars are believed to exhibit tolerance to pyroxasulfone, given their widespread use in the state and lack of reports of injury from pyroxasulfone. Prior to planting, seed of all cultivars was characterized for 100-seed mass and germination. Germination rate was determined by incubating 100 seeds per cultivar on distilled water-moistened germination paper at 21 C and 90% relative humidity, then counting and removing seedlings daily.

Experimental Approach. Trials were conducted in a different field each year at the University of Illinois Vegetable Crop Research Farm near Urbana, Illinois. The previous year's crop was soybean. The soil was a Flanagan silt loam (fine, smectitic, mesic Aquic Argiudolls) averaging 3.1% organic matter and a pH of 6.2. Prior to planting, fields received two passes of a field cultivator. Trials were planted May 21, 2014 and May 27, 2015. Trials were kept weed free with hand-weeding as needed.

The experimental design was a split block with three replications. Main plots measured 2.5 m by 17.5 m. Subplots consisted of individual rows 2.5 m in length on 76-cm spacing. Herbicide treatments

Table 1. Source, germination, and 100-seed mass of vegetable soybean (Veg) and grain-type cultivars used in field experiments near Urbana, IL.

| Cultivar | Type | Source | Germination | 100-seed mass |
|------------------------|-------|---|-------------|---------------|
| | | | % | g |
| AGS 292 | Veg | Washington State University, Pullman, WA | 92 | 29.3 |
| BeSweet 292 | Veg | Rupp Seeds, Wauseon, OH | 90 | 28.0 |
| Butterbean | Veg | Johnny's Selected Seeds, Winslow, ME | 96 | 17.0 |
| Early Hakucho | Veg | USDA Soybean Germplasm Collection, Urbana, IL | 95 | 22.8 |
| Eda Mame Uase Chaurame | Veg | USDA Soybean Germplasm Collection, Urbana, IL | 99 | 17.9 |
| Envy | Veg | USDA Soybean Germplasm Collection, Urbana, IL | 93 | 16.0 |
| Gardensoy 24 | Veg | University of Illinois, Urbana, IL | 91 | 28.2 |
| Gardensoy 32 | Veg | University of Illinois, Urbana, IL | 98 | 26.4 |
| Gardensoy 51 | Veg | University of Illinois, Urbana, IL | 93 | 22.9 |
| IA1010 | Veg | Iowa State University, Ames, IA | 88 | 23.3 |
| JYC-2 | Veg | JYC International, Houston, TX | 83 | 34.8 |
| Misono Green | Veg | USDA Soybean Germplasm Collection, Urbana, IL | 83 | 25.8 |
| Mojo Green | Veg | Wannamaker Seeds, Saluda, NC | 86 | 23.2 |
| Okuhara Daizu | Veg | USDA Soybean Germplasm Collection, Urbana, IL | 92 | 23.0 |
| VS3 | Veg | Anonymous | 86 | 31.4 |
| VS9 | Veg | Anonymous | 94 | 25.9 |
| Sayamusume | Veg | Territorial Seed Company, Cottage Grove, OR | 88 | 40.4 |
| Sunrise | Veg | Wannamaker Seeds, Saluda, NC | 83 | 23.4 |
| Taiwame | Veg | Evergreen Seed, Anaheim, CA | 86 | 32.6 |
| Ware | Veg | USDA Soybean Germplasm Collection, Urbana, IL | 97 | 17.4 |
| WSU 729 | Veg | Washington State University, Pullman, WA | 89 | 23.8 |
| Asgrow 3253 | Grain | Asgrow, St. Louis, MO | 98 | 19.4 |
| Pioneer 93Y41 | Grain | Pioneer, Johnston, IA | 99 | 18.2 |

were assigned to main plots and soybean cultivars were assigned to subplots. Each subplot was planted with 50 seeds at a depth of 2.0 to 2.5 cm. Herbicide treatments included a nontreated control and pyroxasulfone at 417 g ha⁻¹ (2 × recommended field use rate) applied either PRE or EPOST. The PRE application was made the day of planting and the EPOST application was made when most plants had two fully emerged trifoliolate leaves (V2). All applications were made with a compressed air backpack sprayer calibrated to deliver 185 L ha⁻¹ spray volume at 275 kPa.

Data Collection. Soybean response to pyroxasulfone was quantified using five approaches. The crop seedlings were counted two weeks after planting (WAP) to determine population density. Four days later, seedlings with at least one fully emerged trifoliolate leaf, hereafter called V1 plants, were counted. The V1 plants were counted to determine if the PRE treatment delayed development of emerging seedlings. Response to pyroxasulfone was assessed visually 14 and 30 days after treatment (DAT) for the PRE application, and 7 and 14 DAT for the EPOST application. Injury relative to the nontreated control was rated on a scale from 0 (no visible symptoms) to 100 (all plants dead). Crop injury was a cumulative measure of stunting, delayed growth, and leaf damage. Five WAP, individual plant leaf area and biomass were determined. Three plants were randomly selected from each subplot and cut at the soil surface. The leaves were cut at the base of the petiole and leaf area was quantified using an area meter (LI-3100C, LI-COR, Lincoln, NE). Plant tissue (leaves and stems) was dried until constant weight to determine aboveground plant biomass.

Daily rainfall and temperature data were obtained from a weather station located within 1 km of the study sites (Illinois State Water Survey, Champaign, IL). Growing degree days (GDD) were calculated using a base temperature of 7 C.

Data Analysis. Data were found to comply with ANOVA assumptions of homogeneity of variance, based on the modified Levene's test (Neter et al. 1996). With the exception of population density and crop injury, variances were found to meet ANOVA assumptions of normality. Arcsine transformation of crop injury data improved normality. Normality of population density data could not be improved

with any transformation; therefore, non-transformed population density data were analyzed. Response variables were analyzed by ANOVA using the Proc Mixed procedure of SAS version 9.3 (SAS Institute, Cary, NC). Year and replicate nested within year were considered random effects. Herbicide treatment and cultivar were considered fixed effects. Treatment means were compared using protected, Bonferroni-corrected multiple comparisons. All effects were declared significant at $P < 0.05$.

Results and Discussion

Weather. Rainfall a few days after planting was sufficient to incorporate pyroxasulfone into the soil and facilitate seed imbibition and seedling emergence. Within two WAP, fields received 7.3 and 14.0 cm of rainfall in 2014 and 2015, respectively (Table 2). High rainfall in 2015 caused concern that pyroxasulfone applied PRE may have moved out of the seed emergence zone. Post hoc weed seedling counts were made three WAP in 2015 to determine herbicide bioavailability. Nearly weed-free conditions in the PRE treatment three WAP, compared to >60 weeds m⁻² in the nontreated control, confirmed bioavailability of pyroxasulfone.

Emergence. No interaction was observed between cultivar and treatment for population density (Table 3). Furthermore, population density was unaffected by pyroxasulfone treatment ($P = 0.278$). Across cultivars and treatments, soybean population density averaged 15.7 plants m⁻², reflecting an overall emergence rate of 58% of seed planted.

Inherent differences in emergence were observed among cultivars. For instance, population density ranged from 12.1 to 18.5 plants m⁻² among cultivars,

Table 2. Cumulative growing degree days (GDD) (base 7 C) and rainfall during field experiments conducted near Urbana, IL.

| Weeks after planting | Cumulative GDD | | Cumulative rainfall | |
|----------------------|----------------|------|---------------------|------|
| | 2014 | 2015 | 2014 | 2015 |
| | C | | cm | |
| 1 | 80 | 57 | 2.6 | 7.7 |
| 2 | 180 | 144 | 7.3 | 14.0 |
| 3 | 259 | 246 | 11.6 | 19.1 |
| 4 | 332 | 339 | 15.4 | 21.0 |
| 5 | 443 | 420 | 21.0 | 30.6 |

Table 3. Significance (P) of main effects of vegetable soybean cultivar (C), pyroxasulfone treatment (T), and their interaction (C × T) on crop response.

| Factor | Population density ^a | V1 plant no. ^b | Leaf area ^c | Plant biomass ^c |
|----------------|---------------------------------|---------------------------|------------------------|----------------------------|
| C | <0.001 | <0.001 | <0.001 | 0.011 |
| T ^d | 0.278 | <0.001 | <0.001 | <0.001 |
| C × T | 0.999 | 0.988 | 0.850 | 0.942 |

^a Determined two weeks after planting.

^b Number of plants with one or more fully expanded trifoliolate leaves, determined 18 days after planting.

^c Determined five weeks after planting.

^d Treatments at time that population density and V1 plant number were recorded were pyroxasulfone applied PRE and nontreated check; treatments at time that leaf area and plant biomass were recorded were pyroxasulfone applied PRE, pyroxasulfone applied EPOST, and nontreated check.

with grain-type cultivars having among the highest rates of emergence (data not shown). Such emergence differences would be expected given the differential germination among seed lots (Table 1). The potential for poor emergence of vegetable soybean, relative to grain-type soybean, has been reported (Williams 2015b; Zhang et al. 2013).

Crop Injury. Only low levels of crop injury were observed from pyroxasulfone at 417 g ha⁻¹ (data not shown). Mean injury 14 DAT was 10% and 8% in the PRE and EPOST treatments, respectively. Results are consistent with previous research on soybean tolerance to pyroxasulfone. Yamaji et al. (2014) reported low levels (<12%) of soybean injury from pyroxasulfone applied PRE at rates as high as 500 g ha⁻¹. In the present work, the main effect of cultivar was not significant, with P-values ranging from 0.755 to 0.996 across all evaluation periods. The few studies examining soybean cultivar sensitivity to pyroxasulfone found that up to 376 g ha⁻¹ of pyroxasulfone applied PRE resulted in <3% injury across eight grain-type cultivars commonly grown in Ontario, Canada (Belfry and Soltani et al. 2015, McNaughton et al. 2014). Visual assessment data indicate that the vegetable soybean cultivars tested were no more sensitive to pyroxasulfone applied PRE or EPOST than were the grain-type soybean cultivars.

Growth. No interactions were observed between cultivar and treatment for any of the growth response variables, including V1 plant number, leaf area, and

Table 4. Mean crop response to pyroxasulfone applied at 417 g ha⁻¹ PRE and EPOST relative to the nontreated control.

| Level | Population density ^a | V1 plant no. ^b | Leaf area ^c | Plant biomass ^c |
|------------|---------------------------------|---------------------------|-------------------------------------|----------------------------|
| | no. m ⁻² | no. m ⁻² | cm ² plant ⁻¹ | g plant ⁻¹ |
| PRE | 15.3 a | 5.2 b | 318 c | 2.39 c |
| EPOST | - ^d | - ^d | 364 b | 2.79 b |
| Nontreated | 15.9 a | 7.0 a | 424 a | 3.15 a |

^a Determined two weeks after planting.

^b Number of plants with one or more fully expanded trifoliolate leaves, determined 18 days after planting.

^c Determined five weeks after planting.

^d EPOST treatment was not applied at time of assessment.

plant biomass (Table 3). The lack of significant interactions for crop growth responses showed that soybean cultivars were not differentially affected by pyroxasulfone. Both cultivar and treatment main effects were significant (P < 0.001).

Pyroxasulfone applied PRE delayed early season growth of all cultivars. For instance, V1 plant number in the PRE treatment was delayed 26% compared to the nontreated control (Table 4). Since population density was unaffected by pyroxasulfone, reduction in V1 plant number in the PRE treatment was not due to reduced emergence, but rather was due to delayed seedling growth relative to the nontreated control. Similar reductions (approximately 25%) in plant growth due to the PRE treatment were observed for leaf area and plant biomass 5 WAP (Table 4). Pyroxasulfone applied EPOST also reduced leaf area and plant biomass 5 WAP, but to a lesser extent (approximately 12%) than it did when applied PRE.

Inherent differences in crop growth were observed among cultivars. At 5 WAP, leaf area ranged from 279 to 475 cm² per plant and plant biomass ranged from 2.25 to 3.36 g per plant (data not shown). The grain-type cultivars were among the smallest plants measured at 5 WAP. Results are consistent with those seen in previous research comparing vegetable and grain-type soybean. A comparison of 136 vegetable and 14 grain-type soybean lines showed that seedling height, leaf area, and leaf biomass at 3 WAP were larger in vegetable soybean (Williams 2015b). Differences in seedling size between soybean types was attributed to cultivar seed size. Seed of vegetable soybean cultivars used in this study were 35% heavier than were those of the grain-type cultivars, on average. Positive relationships between

soybean seed mass and various aspects of seedling growth have been reported (Burris et al. 1973; Place et al. 2011).

Implications. Pyroxasulfone does not have a negative effect on vegetable or grain-type soybean emergence or early season growth. Crop establishment was unaffected by pyroxasulfone, and visual assessment of crop injury showed minimal ($\leq 10\%$) crop response. At 5 WAP, a delay in soybean growth was seen as a result of PRE application of 417 g pyroxasulfone ha⁻¹, twice the recommended use rate for this soil type. However, vegetable soybean was no more sensitive to pyroxasulfone than was grain-type soybean. Vegetable soybean cultivars grown in the United States for commercial production have the same low risk of crop injury associated with pyroxasulfone as does grain-type soybean.

While some herbicides have become available in recent years, the total number of herbicides available for use in vegetable soybean remains limited, and weed interference continues to be a major threat to domestic production of the crop. Compared to currently registered herbicides, pyroxasulfone could provide better control over certain broadleaf species when applied alone (Hausman et al. 2013) or in combination with other herbicides (Belfry and McNaughton et al. 2015). Registration of pyroxasulfone on vegetable soybean would provide the industry with an additional tool that could be used to expand the options for weed management systems.

Acknowledgements

Mention above of any trademark, proprietary product, or vendor does not constitute a guarantee or warranty of the product by the US Department of Agriculture and does not imply its approval to the exclusion of other products or vendors that also may be suitable.

Literature Cited

- Anonymous (2010) Dual Magnum® herbicide product label 321735. Greensboro, NC: Syngenta. 53 p
- Anonymous (2012) Supplemental label, Lorox® DF herbicide. Phoenix, AZ: Tessenderlo Kerley. 2 p
- Anonymous (2013a) Special local need label, Pursuit® herbicide. Research Triangle Park, NC: BASF. 2 p
- Anonymous (2013b) Special local need label, Spartan Charge® herbicide. Philadelphia, PA: FMC. 3 p

- Anonymous (2013c) Supplemental label, Raptor® herbicide. Research Triangle Park, NC: BASF. 3 p
- Anonymous (2014a) Supplemental label, Reflex® herbicide. Greensboro, NC: Syngenta. 3 p
- Anonymous (2014b) Trefflan® HFP herbicide product label D02-047-027. Indianapolis, IN: Dow. 35 p
- Anonymous (2015a) Basagran® herbicide product label 103569-082614A. Cary, NC: Arysta. 13 p
- Anonymous (2015b) Zidua® herbicide product label NVA 2015-04-388-0072. Research Triangle Park, NC: BASF. 17 p
- Belfry KD, McNaughton KE, Sikkema PH (2015) Weed control in soybean using pyroxasulfone and sulfentrazone. *Can J Plant Sci* 95:1199–1204
- Belfry KD, Soltani N, Brown LR, Sikkema PH (2015) Tolerance of identity preserved soybean cultivars to preemergence herbicides. *Can J Plant Sci* 95:719–726
- Burris JS, Edje OT, Wahab AH (1973) Effects of seed size on seedling performance in soybeans: II. Seedling growth and photosynthesis and field performance. *Crop Sci* 13:207–210
- Hausman NE, Tranel PJ, Riechers DE, Maxwell DJ, Gonzini LC, Hager AG (2013) Responses of an HPPD inhibitor-resistant waterhemp (*Amaranthus tuberculatus*) population to soil-residual herbicides. *Weed Technol* 27:704–711
- McNaughton KE, Shropshire C, Robinson DE, Sikkema PH (2014) Soybean (*Glycine max*) tolerance to timing applications of pyroxasulfone, flumioxazin, and pyroxasulfone + flumioxazin. *Weed Technol* 28:494–500
- Neter J, Kutner MH, Nachtsheim CJ, Wasserman W (1996) *Applied Linear Statistical Models*. 4th edn. Chicago: Irwin. 1408 p
- Place GT, Reberg-Horton SC, Carter TE Jr, Smith AN (2011) Effects of soybean seed size on weed competition. *Agron J* 103:175–181
- Tanetani Y, Fujioka T, Kaku K, Shimizu T (2011) Studies on the inhibition of plant very-long-chain fatty acid elongase by a novel herbicide, pyroxasulfone. *J Pestic Sci* 36:221–228
- Williams MM 2nd (2015a) Managing weeds in commercial edamame production: current options and implications. *Weed Sci* 63:954–961
- Williams MM 2nd (2015b) Phenomorphological characterization of vegetable soybean germplasm lines for commercial production. *Crop Sci* 55:1274–1279
- Williams MM 2nd, Nelson RL (2014) Vegetable soybean tolerance to bentazon, fomesafen, imazamox, linuron, and sulfentrazone. *Weed Technol* 28:601–607
- Yamaji Y, Honda H, Kobayashi M, Hanai R, Inoue J (2014) Weed control efficacy of a novel herbicide, pyroxasulfone. *J Pestic Sci* 39:165–169
- Zhang QY, Hashemi M, Hebert SJ, Li YS (2013) Different responses of preemergence and early seedling growth to planting depth between vegetable soybean and grain soybeans. *Legume Res* 36:515–521

Received October 18, 2016, and approved February 5, 2017.

Associate Editor for this paper: Mark VanGessel, University of Delaware.