

One-year parasitological screening of stray dogs and cats in County Dublin, Ireland

A. Garcia-Campos¹, C. Power¹, J. O'Shaughnessy², C. Browne³, A. Lawlor¹,
G. McCarthy¹, E. J. O'Neill¹ and T. de Waal¹

Research Article

Cite this article: Garcia-Campos A, Power C, O'Shaughnessy J, Browne C, Lawlor A, McCarthy G, O'Neill E J, de Waal T (2019). One-year parasitological screening of stray dogs and cats in County Dublin, Ireland. *Parasitology* **146**, 746–752. <https://doi.org/10.1017/S0031182018002020>

Received: 17 September 2018
Revised: 25 October 2018
Accepted: 7 November 2018
First published online: 18 December 2018

Key words:

Ascarids; cats; *Cystoisospora*; dogs; *Giardia*; Ireland; lungworms; nematode; prevalence; protozoa

Author for correspondence:

A. Garcia-Campos,
E-mail: andres.garcia-campos@ucd.ie

¹UCD Veterinary Sciences Centre, University College Dublin, Belfield, Dublin 4, Ireland; ²Department of Agriculture, Food and the Marine, Backweston Laboratory Campus, Celbridge, Co Kildare, Ireland and ³DSPCA, Mount Venus Road, Rathfarnham, Dublin 16, Ireland

Abstract

To date, there are no recent studies identifying the prevalence of parasites of human and veterinary importance in dogs and cats in Ireland. The interaction between pets and wildlife species in the environment is an important source of parasite exposure to canids and felines, and one likely to be heightened in the stray animal population. This study aimed to establish the prevalence of endoparasites in unowned dogs and cats in County Dublin, Ireland. Feces from stray dogs ($n = 627$) and cats ($n = 289$) entering a rehoming centre were collected immediately after defecation. The main parasitic agents detected were ascarids (15.52 and 30.26%), *Cystoisospora* (3.27 and 3.69%), *Giardia* spp. (6.02 and 1.84%) and lungworms (0.64 and 2.08%), in dogs and cats respectively. Animals younger than 3 months of age were more likely to be infected with ascarids ($P < 0.001$) and *Cystoisospora* spp. ($P = 0.008$ and $P = 0.014$) than older animals. All lungworms were morphologically identified and dogs were infected with *Angiostrongylus vasorum* (0.48%) and *Crenosoma vulpis* (0.16%) whereas cats were only infected with *Aelurostrongylus abstrusus* (2.08%). This represents the first prevalence study of stray animals in Ireland. Data collected will inform the treatment and in addition, the future monitoring and control studies of parasite populations.

Introduction

In Europe, feral and semi-feral animals can play an important role in the transmission of parasites to domestic animals and humans. For example, there are concerns about the role of red foxes (*Vulpes vulpes*) in the maintenance of the life cycle of both *Echinococcus granulosus* and *E. multilocularis* in urban areas (Deplazes *et al.*, 2004; Liccioli *et al.*, 2015). In addition, their role for the transmission of these parasites to humans has been widely discussed (Hegglin and Deplazes, 2013; Hegglin *et al.*, 2015). To date, Ireland is free of both *E. granulosus* and *E. multilocularis* (Beltrán-Beck and Zancanaro, 2017) and the only species known to occur in Ireland is *E. equinus* (Hatch, 1970). However, the increase in the number of red foxes in urban and peripheric-urban areas of Co. Dublin dictates a closer interaction with humans and companion animal's habitats with the result that red foxes could play an increasing and considerable epidemiological role in other endoparasites of human and veterinary relevance. For example, a necropsy study found that 92.2 and 37.7% of red foxes harboured *Uncinaria stenocephala* and *Toxocara canis* adult worms, respectively (Wolfe *et al.*, 2001). Some years later, a survey conducted in Irish wild carnivores showed that red foxes and badgers harboured, amongst other parasitic agents, *Eucoleus aerophilus* eggs (26 and 6%, respectively) and Isospora-like oocysts (9 and 16%, respectively) (Stuart *et al.*, 2013). More recently, another study found that 39.9% of red foxes in Ireland were infected with *Angiostrongylus vasorum* and 1.3% with *Crenosoma vulpis* (McCarthy *et al.*, 2016). Notably, the percentage of foxes infected with *A. vasorum* in Co. Dublin was 66.7%.

Similar concerns have been raised as to the potential epidemiological impact foxes could pose with respect to feline parasites. One of the most important zoonotic agents that cats are responsible for is the spread of *Toxoplasma gondii* and it has been reported that one-third of the cat population in Dublin have been exposed to *T. gondii* (Juvet *et al.*, 2010). Although it is recognized that the main infective route for humans is the consumption of infected meat, infection through the ingestion of oocyst-contaminated soil and water also occurs (Elmore *et al.*, 2010). In two surveys carried out in Europe, it was estimated that more than 30% of owned cats were infected with at least one parasite species (Beugnet *et al.*, 2014; Giannelli *et al.*, 2017). In addition, *Toxocara cati*, another zoonotic agent, was identified in 19.7% of the cats tested (Beugnet *et al.*, 2014) which was very similar to the percentage of cats infected with ascarids in a follow-up study 3 years later (Giannelli *et al.*, 2017). Access to outdoors was one of the risk factors associated with parasitism in cats.

All of this information raises the question of the potential role of stray dogs and cats as potential parasite reservoirs; do they have the potential to have the same impact as wild carnivores and consequently an epidemiological impact on the transmission of endoparasites to domestic animals?

The aim of this study was to screen stray dogs and cats found roaming in County Dublin, as they presented to re-homing centres, for the presence of the most common endoparasites.

Materials and methods

Source

Fecal material (20 g approximately) was collected from unowned dogs ($n = 627$) and cats ($n = 289$) as they presented to the charity centres, Dogs Trust Ireland and The Dublin Society for Prevention of Cruelty to Animals (DSPCA) between February 2016 and June 2017. Samples were collected (by handlers) just after defecation and sent to UCD Parasitology laboratory. For each sample, the following information was recorded: the date of collection, species, age, body condition of the animal and any clinical signs and consistency of the fecal material. The consistency was categorized as formed, loose or diarrhoeic. From each sample, 0.1 g of fecal material was transferred to an eppendorf tube and stored at -20°C , the rest of the sample was subjected to routine parasitological examination as outlined below.

Fecal examination

Flotation methods

Each sample was subjected to two flotation techniques: one with saturated sugar solution and the other sulphate zinc (ZnSO_4) solution. These techniques were performed as previously described with only minor modifications (Euzéby, 1981). Briefly, 3 g of feces were suspended in 42 mL of water, filtered and 15 mL centrifuged at $1,500 \times g$ for 4 min. Pellets were then dissolved in either saturated sucrose solution ($\text{SG} = 1.3$) or ZnSO_4 solution ($\text{SG} = 1.18$), covered with cover slip and allowed to settle for 20 min. Cover slips were applied and the material observed with an Olympus BX40FA microscope (Olympus Optical Co, Ltd. Japan) using $10 \times$ and $40 \times$ Ach objectives (Olympus).

Baermann method

Detection of lungworm species was performed in all specimens using a modified Baermann technique as previously described (Euzéby, 1981). Ten grams of each fecal sample was wrapped into a piece of gauze and suspended over the mouth of a 250 mL conical beaker and immerse in lukewarm water overnight. The sediment was transferred into a watch glass and viewed using an Olympus SZ-ST stereoscope (Olympus). Any L1 larvae were then transferred to a microscope slide, stained with 10% Lugol's iodine and examined microscopically. Morphological identification of lungworm species was performed considering the following features: length, anterior and posterior extremities (Georgi and Georgi, 1991; McGarry and Morgan, 2009; Brianti *et al.*, 2014).

Modified Kinyoun staining

All samples were subjected to a Modified Kinyoun staining as described (Casemore, 1991). Briefly, a thin fecal smear was prepared onto a microscope slide, air dried and then fixed in 100% methanol and stained with Kinyoun's carbol fuchsin for 5 min. After a brief destaining step with 1% acid alcohol, samples were counterstained with 1% acid-methylene blue for 1 min and let to air dry. Samples were then examined microscopically using the $100 \times$ oil immersion objective (Olympus).

ELISA

One hundred and sixty-three samples from dogs and 85 samples from cats were randomly selected and subjected to the *Giardia* Stool Antigen Detection ELISA (IVD Research Inc., USA) following the manufacturer's instructions. Absorbance was measured at 450 nm using Ledetect 96 microplate reader (Tec GmbH, Austria). Absorbance reading of 0.08 OD and above was used as a cut-off following manufacturer's instructions.

Statistical analysis

The prevalence of each parasite was calculated as the proportion of positive animals out of the total numbers of animals examined. Animals were grouped into three age categories (younger than 3 months, between 3 months and 1 year and older than 1 year of age, respectively) determined by staff at the re-homing centre based on tooth eruption and wear (Tutt, 2006). The association between infection with a single parasite agent and animal age was analysed using Fisher's exact test (two-side, $\alpha = 0.05$) using R version 3.4.2

Results

General condition, clinical signs and stool consistency

Overall, animals were in good body condition and did not show marked signs of dehydration. Only 7 dogs had a body condition score ≤ 2 . Six dogs were coughing at the time of the clinical examination. The majority of dogs (84.04%) and cats (92.01%) had formed stools, with diarrhoea or loose feces noted in $<1\%$ of samples for each species.

Gastrointestinal nematodes

Of the 916 fecal samples, 25.8 and 36.8% were positive for the presence of endoparasites in canids and felids, respectively. The number and percentage of positive samples for each parasite are shown in Tables 1 and 2. Ascarids eggs were the most common eggs found using the saturated sugar flotation, with 15.52% of dog and 30.26% of cat samples containing eggs. Notably, 13.4% of dog feces excreted *T. canis* eggs and 2.12% *Toxascaris leonina* eggs. Only one dog was shown to be co-infected with both ascarid species. In the case of cats, only *T. cati* eggs were detected. As shown on Table 3, significantly higher ascarid prevalence was observed in dogs and cats younger than 3 months of age ($P < 0.001$ in both cases).

Uncinaria stenocephala eggs were the second most common eggs detected in 4.89% of dogs tested. Although a higher number of fecal samples (3.10%) containing *U. stenocephala* eggs was found in dogs older than 1 year of age, there was no significant association between age and carriage ($P = 0.215$). *Trichuris vulpis* eggs were identified in only 0.16% of dogs between 3 and 12 months of age. Neither *Eucoleus* spp. nor cestode eggs were identified in any of the samples examined.

Protozoan agents

Giardia spp. was the most prevalent protozoan parasite identified by ZnSO_4 flotation, particularly in dogs, with 6.02% of the samples tested excreting *Giardia* cysts. Half of these dogs which tested positive for *Giardia* spp. were over 1 year of age. However, there was no association between age and the presence of *Giardia* cysts. In the case of cats, 1.84% of the animals were positive for *Giardia* spp. by ZnSO_4 flotation. In parallel with the canine findings, more than half of the cats testing positive for *Giardia* spp. cysts were over a year old but no association

Table 1. Numbers and percentage (%) of unowned dogs positive for endoparasites at different age groups

Age group	Ascarids (n = 612)			Lungworm (n = 627)			Giardia spp.					
	<i>Toxocara canis</i>	<i>Toxascaris leonina</i>	<i>Angyostrongylus vasorum</i>	<i>Angyostrongylus vasorum</i>	<i>Crenosoma vulpis</i>	<i>Uncinaria stenocephala</i> (n = 613)	<i>Trichuris</i> spp. (n = 612)	<i>Taenia</i> spp. (n = 614)	ZnSO ₄ (n = 615)	ELISA (n = 163)	<i>Cystoisospora</i> spp. (n = 612)	<i>Cryptosporidium</i> spp. (n = 614)
<3 months	37 (6.05)	0	0	0	0	1 (0.16)	0	0	9 (1.46)	5 (3.07)	7 (1.14)	0
3–12 months	19 (3.10)	10 (1.63)	0	0	0	9 (1.47)	1 (0.16)	0	10 (1.63)	7 (4.29)	5 (0.82)	1 (0.16)
>12 months	17 (2.78)	3 (0.49)	3 (0.48)	1 (0.16)	0	19 (3.10)	0	0	18 (2.93)	15 (9.20)	4 (0.65)	0
Unknown	9 (1.47)	0	0	0	0	1 (0.16)	0	0	0	1 (0.61)	4 (0.65)	0
Total	82 (13.40)	13 (2.12)	3 (0.48)	1 (0.16)	1 (0.16)	30 (4.89)	1 (0.16)	0	37 (6.02)	28 (17.18)	20 (3.27)	1 (0.16)

Table 2. Numbers and percentage (%) of unowned cats positive for endoparasites at different age groups

Age group	Ascarids (n = 271)			Lungworm (n = 289)			Giardia spp.					
	<i>Toxocara cati</i>	<i>Toxascaris leonina</i>	<i>Aelurostrongylus abstrusus</i>	<i>Aelurostrongylus abstrusus</i>	<i>Uncinaria stenocephala</i> (n = 270)	<i>Trichuris</i> spp. (n = 271)	<i>Taenia</i> spp. (n = 271)	ZnSO ₄ (n = 271)	ELISA (n = 85)	<i>Cystoisospora</i> spp. (n = 271)	<i>Cryptosporidium</i> spp. (n = 271)	<i>Toxoplasma gondii</i> (n = 271)
<3 months	33 (12.18)	0	2 (0.69)	0	0	0	0	0	5 (5.88)	7 (2.58)	0	0
3–12 months	26 (9.59)	0	0	0	0	0	0	1 (0.37)	2 (2.35)	1 (0.37)	0	0
>12 months	18 (6.64)	0	4 (1.38)	0	0	0	0	4 (1.47)	5 (5.88)	1 (0.37)	0	2 (0.74)
Unknown	5 (1.85)	0	0	0	0	0	0	0	0	1 (0.37)	0	0
Total	82 (30.26)	0	6 (2.07)	0	0	0	0	5 (1.84)	12 (14.12)	10 (3.69)	0	2 (0.74)

Table 3. Association of animal age and presence of endoparasites

	Fisher's exact test (<i>P</i> values)	
	Dogs	Cats
Ascarids	<0.001	<0.001
Lungworms	0.378	0.493
<i>Uncinaria stenocephala</i>	0.215	NA
<i>Giardia</i> (ZnSO ₄)	0.326	0.246
<i>Giardia</i> (ELISA)	0.897	1
<i>Cystoisospora</i> spp.	0.008	0.013

NA, not applicable.

between age and cyst excretion was apparent ($P=0.246$). Interestingly, when the coproantigen ELISA was added to the screening profile for a sub-group of the samples (163 dogs and 85 cats) the percentage of animals testing positive for *Giardia* increased to 17.18% for dogs and 14.12% for cats. The number of dogs that were determined positive for *Giardia* by ELISA tended to increase with the age (3.07% for dogs younger than 3 months, 4.29% for dogs between 3 and 12 months and 9.20% for dogs older than 1 year). Cats younger than 3 months and older than 1 year showed the same prevalence (5.88% for each), but still there was no statistically significant association between *Giardia* infection and age in cats ($P=1$). Looking at the fecal consistency, only nine of the dogs and two of the cats that were positive for *Giardia* presented diarrhoea or loose feces but no significant association between *Giardia* infection and consistency of fecal material was seen in either dogs or cats ($P=0.085$ and $P=0.29$, respectively).

Cystoisospora spp. were identified in a very similar percentage of dogs and cats (3.27 and 3.69%, respectively), younger dogs and cats having the highest prevalence for both *Cystoisospora* spp. ($P=0.008$ and $P=0.013$, respectively). *Toxoplasma gondii* oocysts were only detected in two cats, *Sarcocystis* sporocysts in only two dogs and *Cryptosporidium* oocysts in only one dog.

Prevalence of lungworms

Overall four dogs (0.64%) and six cats (2.07%) were positive for lungworms using the modified Baermann technique. Three out of four L1 larvae from the infected dogs were morphologically identified as *A. vasorum*. The other larva was identified as *C. vulpis*. In cats, *A. abstrusus* was the only species of lungworm identified. All animals identified with lungworms were older than 1 year, with the exception of two kittens younger than 3 months of age. None of these animals showed clinical signs compatible with lungworm infection such as coughing or dyspnoea at the time of examination. Positive samples were detected in late summer and winter for dogs. Cats were detected to be positive for lungworms only in the months of September 2016 and May 2017.

Co-infections

Of all the animals tested, 27 dogs (4.3%) and 13 cats (4.5%) were co-infected with two parasite agents (Fig. 1). The majority of the co-infections were a combination of ascarids (22 dogs and 12 cats) with either *Cystoisospora* spp. (10 dogs and seven cats) or *Giardia* spp. (14 dogs and five cats). The dog infected with *C. vulpis* was also positive to *U. stenocephala*. Two of the cats infected with *A. abstrusus* were co-infected with either *T. cati* or *Giardia* spp.

Discussion

The aim of this study was to screen a representative number of unowned dogs and cats in order to establish the prevalence of endoparasites within this population in Co. Dublin. Over the period between February 2016 and June 2017, we were able to confirm the presence of ascarids, *U. stenocephala*, *Giardia* spp., *Cystoisospora* spp., metastrongylid parasites and occasional cases of *T. gondii*, *Sarcocystis* spp., *T. vulpis* and *Cryptosporidium* spp. In line with European reports, ascarids were the most prevalent nematodes identified in stray dogs and cats (Becker *et al.*, 2012; Waap *et al.*, 2014; Zajac *et al.*, 2017) and in owned dogs and cats with regular outdoor access (Barutzki and Schaper, 2011; Beugnet *et al.*, 2014; Knaus *et al.*, 2014; Giannelli *et al.*, 2017). In the case of *T. vulpis* the prevalence was very low compared with the 8.4 and 5% of prevalence in stray dogs found in Poland and central Italy (Paoletti *et al.*, 2015; Zajac *et al.*, 2017) but in line with another study carried out in owned dogs in Great Britain 8 years before the date of this study (Batchelor *et al.*, 2008) where only four dogs out of 4526 were positive for *T. vulpis*.

No cestodes were identified in the study. Ireland is regarded as free of *Echinococcus* spp. (Murphy *et al.*, 2012; Beltrán-Beck and Zancanaro, 2017) with the exception of *E. equinus*. Nevertheless, it should have been possible to identify other taeniid species at low prevalence rates. For example, the presence of the dog or cat flea *Ctenocephalides* spp., intermediate hosts of *Dipylidium caninum*, has been well documented in Ireland with prevalence ranging between 75 and 90% (Wall *et al.*, 1997). In addition, the prevalence of other taeniid species in Ireland has been detected at very low levels, below 0.1%, in intestinal contents of foxes (Wolfe *et al.*, 2001). This could suggest the low presence of tapeworms-infected rodents or other animal species in Co. Dublin. We believe that animals from more rural areas, where the co-habitation of dogs and livestock is closer than in Co. Dublin, would be more likely to be infected with cestodes.

The different results when cysts detection and coproantigen for *Giardia* spp. are in line with previous studies using ELISA, IFA or PCR techniques for *Giardia* diagnosis (Bouzid *et al.*, 2015). Cysts are shed intermittently, and hence repeated fecal samples are required to reduce the number of false negatives gained from infected animals. In this study, only a subset of samples were used for coproantigen thus caution should be taken for this comparison. The prevalence of *Giardia* detected by ZnSO₄ flotation dogs and cats (6.02 and 1.84%, respectively) are in line with the 5.25% of stray dogs infected in Italy (Paoletti *et al.*, 2015) and to the 2.9% of stray cats in Poland (Zajac *et al.*, 2017). When we look at the ELISA results in dogs and cats (17.18 and 14.12%, respectively) these are in line with the pool-prevalence rates calculated by a meta-analysis study worldwide (15.2% for dogs and 12% for cats) (Bouzid *et al.*, 2015). Veterinarians regard giardiasis as an important enteric pathogen in small animals as it is an agent that is not easy to cure and to eliminate for the environment (Pallant *et al.*, 2015). It is also of zoonotic importance as some assemblages of *Giardia* can infect both pets and humans. In this study, assessment of the *Giardia* assemblages was not part of the study but represents important further work in order to establish whether *Giardia* zoonotic genotypes are found in pets.

Only two cats were found to shed *T. gondii* oocysts. This is interesting as this is slightly higher than the 0.1% that was found in stray cats in Germany (Becker *et al.*, 2012) but the same percentage that was found in a 10-year study in owned cats in the same country (Raue *et al.*, 2017). Interestingly, a previous study conducted in Ireland that was looking at *T. gondii* exposure or current infection by detecting *T. gondii* IgG and IgM,




Fig. 1. Numbers of unowned dogs (A) and cats (B) co-infected with at least two endoparasites.

respectively, showed that 36.1% of stray cats were positive (Juvet *et al.*, 2010). Considering that unowned cats have more outdoor access and more chances to hunt and be in contact with intermediate hosts than indoor cats, we would have expected in the current study to identify more cats shedding *T. gondii* oocysts. It could be explained by the fact that cats usually are infected at young age and develop immunity and rarely shed oocysts again, even if re-infected.

In this study, <1% of dogs and slightly more than 2% of cats were infected with lungworms. It is possible that these values are underestimated in view of the fact that only one small volume of fecal sample was assessed. However, the prevalence of lungworm identified in our study is similar to the 1.2% of stray dogs infected in Italy (Paoletti *et al.*, 2015), the 0.97 and 0.66% of dogs positive for *A. vasorum* antigen and antibodies identified in samples in the UK (Schnyder *et al.*, 2013) and Portugal (Alho *et al.*, 2018), respectively, and for those reported in stray cats in Germany (Becker *et al.*, 2012) and to the domestic cats from Belgium (1.9%), Central-South Portugal (1.7%) and North Portugal (2.5%) (Giannelli *et al.*, 2017). It is likely that additional animals would have been identified if animals had been additionally tested using ELISA for antigen or antibody detection, capable of detecting animals in pre-patent periods (Briggs *et al.*, 2013; Schnyder *et al.*, 2015; Zottler *et al.*, 2017).

It was interesting to find that two out the eight cats infected with *A. abstrusus* were younger than 3 months of age. This finding is in line with the hypothesis of transmission of lungworms from the queen cat to suckling kittens (Brianti *et al.*, 2013; Tamponi *et al.*, 2014) as it has been identified with *Troglostrongylus brevior* and *A. abstrusus* in 8-week-old kittens. None of the animals infected showed any of the classical clinical signs attributed to lungworms. This is very important from an epidemiological point of view as stray dogs and feral cats could

become silent carriers favouring the spread of lungworms from wildlife to pets and *vice versa*.

In a survey by McCarthy *et al.* it was shown that more than 66% of red foxes in Co. Dublin were infected with *A. vasorum* in 2014 (McCarthy *et al.*, 2016). In view of our hypothesis that red foxes and unowned dogs share the same environment and potentially intermediate hosts, we would have expected to find a high prevalence of lungworm in this dog population. Although no further conclusions can be drawn, we think that one of the reasons that could explain this situation is the weather conditions. According to the records from the Irish meteorological agency Met Éireann, the period when the study was conducted was one of the driest and warmest periods for the previous 8 years (<https://www.met.ie/climate/irish-climate-monthly-summary.asp>). Although weather fluctuations have relatively minor impacts in the transmission for some parasitic agents, such as ascarids and *Trichuris* spp., it is likely to have impacted the transmission of *A. vasorum* in view of the likely impact on the important intermediate hosts, snails and slugs. This could also explain the fact that we did not see the marked seasonality of the *A. vasorum* diagnosis as described in other studies (Morgan *et al.*, 2010), although only three *A. vasorum* cases were identified overall in this study.


The significant association of young animals (puppies and kittens) to be infected with the three ascarids species found and with *Cystoisospora* spp. is in line with other studies (Barutzki and Schaper, 2013; Rauscher *et al.*, 2013). Contrary to the former parasite agents, we did not find age-specific association with the rest of the parasites targeted in this study. For *A. abstrusus*, there are some studies stating that there is an age-infection association (Traversa *et al.*, 2008; Mircean *et al.*, 2010; Capári *et al.*, 2013; Knaus *et al.*, 2014) and some others where there was not an association (Beugnet *et al.*, 2014; Giannelli *et al.*, 2017). In

this study, we consider that the number of positive samples to be too low to base a conclusion on these observations.

Considering that this study focused on unowned dogs and cats with no parasite control, a high prevalence of parasitic infection would have been expected. Nevertheless, we lack valuable information such as the time that dogs and cats have been roaming in the environment and thus exposed to parasites, their previous deworming programme and the exact localization where the animals have been retrieved makes this assessment more difficult for interpretation. Regular administration of anthelmintics significantly decreases the risk of, for example, *T. cati* infection (Beugnet *et al.*, 2014).

In conclusion, this is the first prevalence study conducted in unowned dogs and semi feral/feral cats in Ireland. The prevalence of infection with either gastrointestinal nematodes and/or protozoa agents is similar to that reported in other European countries. Considering the high prevalence of *A. vasorum* in foxes, we expected to detect a higher prevalence of lungworm infection in stray dogs. Limitations in the number of samples per animal, the laboratory techniques employed or the weather conditions could explain the results presented. The fact that these animals did not show symptoms enhances the importance of implementing monitoring programmes, control and prophylactic measures targeting this animal population in order to decrease the spillover of pathogens from silent carriers to companion animals.

Acknowledgements. The authors would like to thank all the staff of DSPCA and Dogs Trust Ireland for supplying material and the care of the animals.

Author ORCIDs.  A. Garcia-Campos 0000-0001-8887-1895.

Financial support. The study was funded, in part, by the European Scientific Counsel Companion Animal Parasites (ESCCAP) UK and Ireland.

Conflict of interest. None.

Ethical standards. Not applicable.

References

- Alho AM, Meireles J, Schnyder M, Cardoso L, Belo S, Deplazes P and de Carvalho LM (2018) *Dirofilaria immitis* and *Angiostrongylus vasorum*: the current situation of two major canine heartworms in Portugal. *Veterinary Parasitology* **252**, 120–126.
- Barutzki D and Schaper R (2011) Results of parasitological examinations of faecal samples from cats and dogs in Germany between 2003 and 2010. *Parasitology Research* **109**, 45–60.
- Barutzki D and Schaper R (2013) Age-dependant prevalence of endoparasites in young dogs and cats up to one year of age. *Parasitology Research* **112**, 119–131.
- Batchelor DJ, Tzannes S, Graham PA, Wastling JM, Pinchbeck GL and German AJ (2008) Detection of endoparasites with zoonotic potential in dogs with gastrointestinal disease in the UK. *Transboundary and Emerging Diseases* **55**, 99–104.
- Becker A-C, Rohen M, Epe C and Schnieder T (2012) Prevalence of endoparasites in stray and fostered dogs and cats in Northern Germany. *Parasitology Research* **111**, 849–857.
- Beltrán-Beck B and Zancanaro G (2017) Assessment of *Echinococcus multilocularis* surveillance reports submitted in 2017 in the context of Commission Regulation (EU) No 1152/2011. *EFSA Journal* **15**, 5051.
- Beugnet F, Bourdeau P, Chalvet-Monfray K, Cozma V, Farkas R, Guillot J, Halos L, Joachim A, Losson B, Miró G, Otranto D, Renaud M and Rinaldi L (2014) Parasites of domestic owned cats in Europe: co-infestations and risk factors. *Parasites & Vectors* **7**, 291.
- Bouziid M, Halai K, Jeffreys D and Hunter PR (2015) The prevalence of *Giardia* infection in dogs and cats, a systematic review and meta-analysis of prevalence studies from stool samples. *Veterinary Parasitology* **207**, 181–202.
- Brianti E, Gaglio G, Napoli E, Falsone L, Giannetto S, Latrofa MS, Giannelli A, Dantas-Torres F and Otranto D (2013) Evidence for direct transmission of the cat lungworm *Troglostrongylus brevior* (Strongylida: Crenosomatidae). *Parasitology* **140**, 821–824.
- Brianti E, Giannetto S, Dantas-Torres F and Otranto D (2014) Lungworms of the genus *Troglostrongylus* (strongylida: Crenosomatidae): neglected parasites for domestic cats. *Veterinary Parasitology* **202**, 104–112.
- Briggs KR, Yaros JP, Liotta JL, Lucio-Forster A, Lee AC and Bowman DD (2013) Detecting *Aelurostrongylus abstrusus*-specific IgG antibody using an immunofluorescence assay. *Journal of Feline Medicine and Surgery* **15**, 1114–1118.
- Capári B, Hamel D, Visser M, Winter R, Pfister K and Rehbein S (2013) Parasitic infections of domestic cats, *Felis catus*, in western Hungary. *Veterinary Parasitology* **192**, 33–42.
- Casemore D (1991) Laboratory methods for diagnosing cryptosporidiosis. *Journal of Clinical Pathology* **44**, 445–451.
- Deplazes P, Hegglin D, Gloor S and Romig T (2004) Wilderness in the city: the urbanization of *Echinococcus multilocularis*. *Trends in Parasitology* **20**, 77–84.
- Elmore SA, Jones JL, Conrad PA, Patton S, Lindsay DS and Dubey JP (2010) *Toxoplasma gondii*: epidemiology, feline clinical aspects, and prevention. *Trends in Parasitology* **26**, 190–196.
- Euzéby J (1981) La coprologie chez les carnivores. In *Diagnostic expérimental des helminthoses animales: travaux pratiques d'helminthologie vétérinaire, Livre 1, Informations*. Paris, France: Techniques des Services Vétérinaires, pp. 195–214.
- Georgi J and Georgi M (1991) *Canine Clinical Parasitology*. Malvern, Philadelphia: Lea & Febiger.
- Giannelli A, Capelli G, Joachim A, Hinney B, Losson B, Kirkova Z, René-Martellet M, Papadopoulos E, Farkas R, Napoli E, Brianti E, Tamponi C, Varcasia A, Margarida Alho A, Madeira de Carvalho L, Cardoso L, Maia C, Mircean V, Mihalca AD, Miró G, Schnyder M, Cantacessi C, Colella V, Cavalera MA, Latrofa MS, Annoscia G, Knaus M, Halos L, Beugnet F and Otranto D (2017) Lungworms and gastrointestinal parasites of domestic cats: a European perspective. *International Journal for Parasitology* **47**, 517–528.
- Hatch C (1970) *Echinococcus granulosus equinus* in Irish dogs. *The Veterinary Record* **86**, 632–633.
- Hegglin D and Deplazes P (2013) Control of *Echinococcus multilocularis*: strategies, feasibility and cost – benefit analyses. *International Journal for Parasitology* **43**, 327–337.
- Hegglin D, Bontadina F and Deplazes P (2015) Human-wildlife interactions and zoonotic transmission of *Echinococcus multilocularis*. *Trends in Parasitology* **31**, 167–173.
- Juvet F, Lappin MR, Brennan S and Mooney CT (2010) Prevalence of selected infectious agents in cats in Ireland. *Journal of Feline Medicine and Surgery* **12**, 476–482.
- Knaus M, Chester ST, Rosentel J, Kühnert A and Rehbein S (2014) Efficacy of a novel topical combination of fipronil, (S)-methoprene, eprinomectin and praziquantel against larval and adult stages of the cat lungworm, *Aelurostrongylus abstrusus*. *Veterinary Parasitology* **202**, 64–68.
- Liccioli S, Giroudoux P, Deplazes P and Massolo A (2015) Wilderness in the 'city' revisited: different urban shape transmission of *Echinococcus multilocularis* by altering predator and prey communities. *Trends in Parasitology* **31**, 297–305.
- McCarthy G, Ferrand M, De Waal T, Zintl A, McGrath G, Byrne W and O'Neill EJ (2016) Geographical distribution of *Angiostrongylus vasorum* in foxes (*Vulpes vulpes*) in the Republic of Ireland. *Parasitology* **143**, 588–593.
- McGarry JW and Morgan ER (2009) Identification of first-stage larvae of metastrongyles from dogs. *The Veterinary Record* **165**, 258–261.
- Mircean V, Titilincu A and Vasile C (2010) Prevalence of endoparasites in household cat (*Felis catus*) populations from Transylvania (Romania) and association with risk factors. *Veterinary Parasitology* **171**, 163–166.
- Morgan ER, Jefferies R, van Otterdijk L, McEniry RB, Allen F, Bakewell M and Shaw SE (2010) *Angiostrongylus vasorum* infection in dogs: presentation and risk factors. *Veterinary Parasitology* **173**, 255–261.
- Murphy TM, Wahlström H, Dold C, Keegan JD, McCann A, Melville J, Murphy D and McAteer W (2012) Freedom from *Echinococcus multilocularis*: an Irish perspective. *Veterinary Parasitology* **190**, 196–203.
- Pallant L, Barutzki D, Schaper R and Thompson R (2015) The epidemiology of infections with *Giardia* species and genotypes in well cared for dogs and cats in Germany. *Parasites & Vectors* **8**, 2.
- Paoletti B, Traversa D, Iorio R, De Berardinis A, Bartolini R, Salini R and Di Cesare A (2015) Zoonotic parasites in feces and fur of stray and private dogs from Italy. *Parasitology Research* **114**, 2135–2141.

- Raue K, Heuer I, Böhm C, Wolken S, Epe C and Strube C** (2017) 10-year parasitological examination results (2003 to 2012) of faecal samples from horses, ruminants, pigs, dogs, cats, rabbits and hedgehogs. *Parasitology Research* **116**, 3315–3330.
- Rauscher BA, Schäfer-Somi S, Ehling-Schulz M, Möstl K, Handl S, Hinney B, Spargser J, Schaper R and Joachim A** (2013) Control of canine endoparasites, especially *Isospora* spp., with procox in naturally infected puppies: parasitological, bacteriological and health parameters. *Open Journal of Veterinary Medicine* **3**, 121–130.
- Schnyder M, Schaper R, Bilbrough G, Morgan ER and Deplazes P** (2013) Seroepidemiological survey for canine angiostrongylosis in dogs from Germany and the UK using combined detection of *Angiostrongylus vasorum* antigen and specific antibodies. *Parasitology* **140**, 1442–1450.
- Schnyder M, Jefferies R, Schucan A, Morgan ER and Deplazes P** (2015) Comparison of coprological, immunological and molecular methods for the detection of dogs infected with *Angiostrongylus vasorum* before and after anthelmintic treatment. *Parasitology* **142**, 1270–1277.
- Stuart P, Golden O, Zintl A, De Waal T, Mulcahy G, Mccarthy E and Lawton C** (2013) A coprological survey of parasites of wild carnivores in Ireland. *Parasitology Research* **112**, 3587–3693.
- Tamponi C, Varcasia A, Brianti E, Pipia AP, Frau V, Pinna Parpaglia ML, Sanna G, Garippa G, Otranto D and Scala A** (2014) New insights on metastrongyloid lungworms infecting cats of Sardinia, Italy. *Veterinary Parasitology* **203**, 222–226.
- Traversa D, Lia RP, Iorio R, Boari A, Paradies P, Capelli G, Avolio S and Otranto D** (2008) Diagnosis and risk factors of *Aelurostrongylus abstrusus* (Nematoda, Strongylida) infection in cats from Italy. *Veterinary Parasitology* **153**, 182–186.
- Tutt C** (2006) *Small Animal Dentistry: A Manual of Techniques*, 1st Edn. Oxford, UK: Wiley-Blackwell.
- Waap H, Gomes J and Nunes T** (2014) Parasite communities in stray cat populations from Lisbon, Portugal. *Journal of Helminthology* **88**, 389–395.
- Wall R, Shaw SE and Penaliggon J** (1997) The prevalence of flea species on cats and dogs in Ireland. *Medical and Veterinary Entomology* **11**, 404–406.
- Wolfe A, Hogan S, Maguire D, Fitzpatrick C, Vaughan L, Wall D, Hayden TJ and Mulcahy G** (2001) Red foxes (*Vulpes vulpes*) in Ireland as hosts for parasites of potential zoonotic and veterinary significance. *Veterinary Record* **22**, 759–763.
- Zajac V, Wójcik-Fatla A, Sawczyn A, Cisak E, Sroka J, Kloc A, Zajac Z, Buczek A, Dutkiewicz J and Bartosik K** (2017) Stray dogs and cats as potential sources of soil contamination with zoonotic parasites. *Annals of Agricultural and Environmental Medicine* **24**, 39–43.
- Zottler E-M, Strube C and Schnyder M** (2017) Detection of specific antibodies in cats infected with the lung nematode *Aelurostrongylus abstrusus*. *Veterinary Parasitology* **235**, 75–82.