

Pushchino multibeam pulsar search — V. The bright FRB 20190203 detected at 111 MHz

S. A. Tyul'bashev,¹ V. A. Samodurov,^{2,1} A. S. Pozanenko,^{3,4,5} E. A. Brylyakova,¹ S. A. Grebenev,³ I. V. Chelovekov,³ P. Yu. Minaev,³ E. A. Isaev,⁶ and M. V. Barkov⁷

¹P.N. Lebedev Physical Institute, Russian Academy of Sciences, Astro Space Center, Pushchino Radio Astronomy Observatory, Radiotelescopnaya 1, Moscow region, Pushchino, 142290, Russia

²Graduate School of Business, HSE University

³Space Research Institute, Russian Academy of Sciences, Profsoyusnaya 84/32, Moscow, 117997, Russia

⁴National Research University Higher School of Economics, Pokrovskii bulvar 11, Moscow, 109028, Russia

⁵Institute of Physics and Technology, Institutskiy pereulok 9, Dolgoprudny, 141701, Russia

⁶Institute of Mathematical Problems of Biology, Russian Academy of Sciences, Branch of the Keldysh Institute of Applied Mathematics, Vitkevich 1a, Moscow region, Pushchino, 142290, Russia

⁷Institute of Astronomy, Russian Academy of Sciences, Pyatnitskaya 48, Moscow, 119017, Russia

Author for correspondence: S. A. Tyul'bashev, Email: serg@prao.ru.

Abstract

Since August 2014, a monitoring survey at a frequency of 111 MHz has been conducted on the Large Phased Array (LPA) radio telescope of the P.N. Lebedev Physical Institute (LPI). We report the discovery of a bright pulse having a dispersion measure (DM) equal to 134.4 ± 2 pc cm⁻³, a peak flux density (S_p) equal to 20 ± 4 Jy and a half-width (W_e) equal to 211 ± 6 ms. The excessive DM of the pulse, after taking into account the Milky Way contribution, is 114 pc cm⁻³ that indicates its extragalactic origin. Such value of DM corresponds to the luminosity distance 713 Mpc. The above parameters make the pulse to be a reliable candidate to the fast radio burst (FRB) event, and then it is the second FRB detected at such a large $\lambda \sim 2.7$ m wavelength and the first one among non-repeating FRBs. The normalized luminosity L_ν of the event, which we have designated as FRB 20190203, estimated under assumption that the whole excessive DM is determined by the intergalactic environment toward the host galaxy, is equal to $\simeq 10^{34}$ erg s⁻¹Hz⁻¹. In addition to the study of radio data we analyzed data from the quasi-simultaneous observations of the sky in the high energy (≥ 80 keV) band by the omnidirectional detector SPI/ACS aboard the INTEGRAL orbital observatory (in order to look for a possible gamma-ray counterpart of FRB 20190203). We did not detect any transient events exceeding the background at a statistically significant level. In the INTEGRAL archive, the FRB 20190203 localization region has been observed many times with with a total exposure of ~ 73.2 days. We have analyzed the data but were unable to find any reliable short gamma-ray bursts from the FRB 20190203 position. Finally we note that the observed properties of FRB 20190203 can be reproduced well in the framework of a maser synchrotron model operating in the far reverse shock (at a distance of $\sim 10^{15}$ cm) of a magnetar. However, triggering the burst requires a high conversion efficiency (at the level of 1%) of the shock wave energy into the radio emission.

Keywords: methods: data analysis; methods: observational; transients: fast radio bursts

1. Introduction

Fast radio bursts (FRBs), which are pulsed sources of intense radio emission of extragalactic origin, were discovered in 2007 in the archive data of the 64-meter telescope in Parkes (Lorimer et al. 2007). Almost all FRBs were detected at frequencies between 0.4 and 1.6 GHz, that is, in the decimeter wavelength range (see catalogs^{a,b} in Petroff et al. 2016; CHIME/FRB Collaboration et al. 2021). Externally, the bursts are similar to the pulsar pulses. They are short in duration and dispersed, that is, they come first at a high frequency and then at a low one. The median value of the observed duration of the bursts, according to the FRB catalog by Petroff et al. (2016), gets to ~ 2 ms. However, there were structures in bursts observed with a duration of less than 0.1 μ s (Majid et al. 2021).

The pulse signal, passing through the interstellar medium

in the host galaxy, the intergalactic medium, and the interstellar medium in the Milky Way, experiences scattering. Its observed duration becomes longer. The pulses with high signal-to-noise (S/N) ratios have often the characteristic FRED shape with a fast rise (stepped front) and an exponential decay.

Scattering allows us to study properties of the medium between the radiation source and us, but at the same time, scattering leads to the pulse profile smearing and decreasing in the observed S/N value of the pulse. If we are talking about observations at a certain frequency, then the further away the pulse source is, the greater the observed dispersion measure (DM) will be and the longer the delay due to scattering (τ_{sc}). If we are talking about observations at different frequencies, then the lower the frequency of observations (ν), the longer the characteristic time delay due to scattering ($\tau_{sc} \sim \nu^{-4}$) should be (Lorimer and Kramer 2004). With the difference in frequency of observations only twice, the characteristic scattering time τ_{sc} differs by more than an order of magnitude ($\sim 2^4 = 16$).

a. www.frbcat.org/

b. www.chime-frb.ca/catalog

The scattering effect cannot be compensated for and therefore leads to a deterioration in sensitivity when searching for FRBs.

In the meter wavelength range ($\nu < 300$ MHz), there were three attempts made to the moment of the full-fledged search for FRBs using new radio telescopes with wide reception bands. The Low Frequency Array (LOFAR; van Haarlem *et al.* 2013) and Murchison Widefield Array (MWA; Lonsdale *et al.* 2009) radio telescopes have been used for the search for FRBs at 45 and 182 MHz (Coenen *et al.* 2014; Karastergiou *et al.* 2015; Rowlinson *et al.* 2016). No bursts have been detected that gave upper limits of the expected number of FRBs. The problems with previous search for FRBs at 111 MHz using the Large Phased Array radio telescope of P.N. Lebedev Physical Institute (LPA LPI; Shishov *et al.* 2016) have been described by Brylyakova and Tyul'bashev (2023).

There were also a number of attempts to observe repeated FRBs, registered from different sources at high frequencies, simultaneously in the decimeter (300–800 MHz) and meter (110–180 MHz) ranges (Law *et al.* 2017; Sokolowski *et al.* 2018; Houben *et al.* 2019). The bursts were recorded in both the ranges (Chawla *et al.* 2020; Pleunis *et al.* 2021) from only one unique source FRB 20180916 (Marcote *et al.* 2020) which exhibits periodic activity. And even in this case there was no simultaneous registration for any bursts. Bursts found in decimeter range were not detected in meter range. Moreover it was noted that the burst activity was systematically delayed toward lower frequencies by about 3 days from 600 to 150 MHz.

Since the sensitivity of observations decreases due to scattering $\sim \tau_{sc}^{1/2}$, the smaller the distance to the burst source and, consequently, the smaller the scattering, the weaker bursts can be observed with all other parameters unchanged. The higher the fluctuation sensitivity of the radio telescope, which is proportional to the effective area, the more chances we have to register a burst. At the present time, the most sensitive meter-range radio telescopes in the world are LOFAR, MWA, LPA. The meter range is also claimed for Five-hundred-meter Aperture Spherical Telescope (FAST; Nan *et al.* 2011) which should have record sensitivity in both the decimeter and meter ranges. However we were unable to find any publications based on the FAST data with attempts to detect FRBs on meter waves. In this paper, we describe our search for dispersed signals in metadata obtained during technical assessment of the quality of observations carried out with the LPA LPI radio telescope.

2. LPA observations and data processing

The meridian instrument LPA LPI is a phased array consisting of 16,384 wave dipoles having one linear polarization. The input signal coming from the lines on which 128 dipoles are located goes to the amplifiers. The amplifiers have one input and four outputs. Therefore, there is a technical possibility to create four independent radio telescopes on the basis of a single antenna field (Shishov *et al.* 2016). Currently, three outputs are involved, one of which is used in the monitoring program. This radio telescope is called LPA3. It has 128 uncontrolled

(stationary) beams overlapping declinations $-9^\circ < \delta < +55^\circ$. The beams are created using the Butler matrix and provide overlap at the level of 0.405. The beam size is approximately $0.5 \times 1.0^\circ$, and its shape obeys the dependence $\sin^2(x)/x^2$. The central frequency of the observations is 110.25 MHz, the receiving frequency band is 2.5 MHz. The signal on the recorder is split into two ones and recorded onto the hard disk twice with different time–frequency resolutions. The data recorded in 6 frequency channels with a channel width of 415 kHz and a sampling time of 0.1 s are used in the “Space Weather” project (Shishov *et al.* 2016). Data recorded in 32 frequency channels with a channel width of 78 kHz and a sampling time of 12.5ms are used to search for pulsars and transients in the project Pushchino Multibeams Pulsar Search (PUMPS; Tyul'bashev *et al.* 2016; Tyul'bashev *et al.* 2018; Tyul'bashev, Kitaeva, and Tyul'basheva 2022). Data are recorded at one-hour intervals. The beginning of the recording is controlled by the atomic frequency standard. Inside the one-hour recording, the time is controlled by a quartz oscillator. Six times a day, a signal of a known temperature (calibration step) is transmitted to the input amplifiers. Using this step, the data in the frequency channels can be equalized in gain.

Quality control of observations is carried out by a special processing program. This program equalizes the gain in frequency channels, cuts all the data inside the one-hour interval into 10-s long segments (800 points in the frequency channel, $800 \times 32 = 25,600$ points in all channels) and prepares metadata which are written into files. On each segment and in each frequency channel, the baseline is subtracted and after that the parameters (metadata) are evaluated. The metadata includes the maximum and minimum values on a segment, the average and median average values on a segment, estimates of the root mean square (RMS) deviation of noise, and others. In addition, the start time of the segment and the number of the maximum point inside a segment for each frequency channel are stored (for more details, see Samodurov *et al.* 2022). If the maxima are detected inside the frequency channels at a given S/N, it is possible to estimate the time shift of these maxima between channels due to the dispersion delay and identify candidates for extraterrestrial origin. If the maxima in the frequency channels fall on one point, then there is no dispersion delay and, most likely, there is interference.

The volume of metadata recorded in the files is about 100 times smaller than the volume of input data. Further work with files is carried out using a program that, in accordance with a set of specified criteria, identifies interferences and sources of unknown nature. In total 153,729 one-hour files were processed to obtain metadata (or 17,000 square degrees every day in the interval of 3,000 days.) Three main criteria were laid down in the search program for dispersed pulses. Firstly, when passing from a high frequency to a low one, there should be a pulse shift in the frequency channels corresponding to $DM > 100$ pc cm⁻³. Secondly, in the tested segment there should be at least 4 frequency channels in which the signal has S/N > 5. For 32-channel data, the RMS deviations of noise (σ_n) in one frequency channel are approximately 1.5 Jy outside the Galactic plane in the direction to the zenith. That is, only




Figure 1. The upper panel shows the dynamic spectrum of the detected pulse. Frequency channels (in MHz) are indicated along the vertical axis. The lower panel shows the signals received for: 1) frequency channels summed up with $DM = 0 \text{ pc cm}^{-3}$; 2) channels summed up with $DM = 134.4 \text{ pc cm}^{-3}$; 3-4) channels summed up with $DM = 0 \text{ pc cm}^{-3}$ for beams above and below the detected burst; 5) base line. The lower panel along the vertical axis shows the flux density. The horizontal axis is the same for both panels and shows the time.

those sources were selected whose peak flux density (S_p), after compensation for the assumed value of DM , was higher than 7.5 Jy. Thirdly, the signal should be observed in no more than two adjacent beams. A special study of interference on LPA3 with the use of metadata showed that interference is observed in three or more beams, and most often in all beams of the radio telescope simultaneously (see Fig. 1 in Samodurov et al. 2022).

If necessary, further investigation of the sources is carried out using the original (raw) data that are stored on the server. Despite a small set of output parameters, it was based on the metadata that the first rotating radio transients (RRAT) were detected on LPA3 (Samodurov et al. 2017).

3. Discovery of FRB 20190203

Taking into account the search criteria for dispersed pulses, one pulse has been found during the metadata analysis. Figure 1 shows the dynamic spectrum and time profile of the detected pulse. The tail typical for a scattered pulse is observed in the profile. For this pulse, the maximum in frequency channels appeared first at the highest receiving frequency and consistently appeared later and later at low frequencies. The total shift of the signal peak across all frequency channels was approximately 2 s, which corresponds to $DM = 134.4 \text{ pc cm}^{-3}$. The DM value collected in the Milky Way in the direction of the




Figure 2. The same as in Fig. 1 but the red, orange and green curves shows RRAT J1005+3015 in the main beam and the beams above and below it after compensation for DM . S/N for this transient reaches 234.

pulse source is equal to 20.3 pc cm^{-3} according to the YMW16 model^c by Yao, Manchester, and Wang (2017). The DM value attributed to the Milky Way according to the NE2001 model (Cordes and Lazio 2002, 2003) differs from his value less than 2.5 pc cm^{-3} . Thus, there is an excessive $DM \approx 114 \text{ pc cm}^{-3}$ indicating the detection of the new fast radio burst (FRB).

When summing the frequency channels without compensation for DM , a source with $S/N=12$ is visible in the raw data. In the beams above and below, the source is not observed. After compensation for DM , the S/N ratio rises to 65. Such a high value of S/N excludes any possibility that the registered pulse may be a statistical fluctuation, even taking into account the long-term duration of observations and the huge number of statistical trials associated with it.

Until now, the search for dispersed pulses in the PUMPS project has been carried out up to $DM \leq 100 \text{ pc cm}^{-3}$. The estimates by Tyul'bashev, Kitaeva, and Tyul'basheva (2022) show a strong signal smearing in frequency channels due to dispersion and a catastrophic deterioration in the LPA sensitivity due to scattering when searching for periodicity in pulsars on a scale of seconds at $DM \geq 200 \text{ pc cm}^{-3}$. In Appendix 1, we demonstrate the pulses detected from three pulsars having DM significantly exceeding that of the discovered FRB. Figure 2 shows the dynamic spectrum and the time profiles of the signal in the main beam and two nearby beams for one of the similar pulses detected from the pulsar RRAT J1005+3015. These very confident detections indicate that searching for pulse signals at $DM \geq 100 \text{ pc cm}^{-3}$ is quite possible on the LPA3 radio telescope.

The fast radio burst discovered by LPA3 on February 3, 2019 (FRB 20190203) and described in detail in this paper has

c. www.atnf.csiro.au/research/pulsar/ymw16/

the following parameters:

Parameters	Values
Date	2019-02-03
Time(UT)	02:18:10.284
α_{2000}	$13^{\text{h}}40^{\text{m}}46^{\text{s}} \pm 2^{\text{m}}$
δ_{2000}	$35^{\circ}30' \pm 10'$
l	$74^{\circ}49 \pm 0^{\circ}.42$
b	$76^{\circ}56 \pm 0^{\circ}.43$
$DM, \text{pc cm}^{-3}$	134.4 ± 2.0
τ_{sc}, ms	211 ± 6
$S_{\text{peak}}, \text{Jy}$	20 ± 4
S/N	65

With the S/N ratio equal to 65, it could be expected that the FRB will be visible in one of the neighboring beams. However there is no such signal visible at the S/N level exceeding 3. It is possible to estimate the maximum shift of the FRB coordinate in declination from the coordinate of the stationary beam, based on the beam size ($0^{\circ}.5$) and shape ($\sin^2(x)/x^2$). If this FRB is located inside $\pm 5'$ of the beam coordinate, the contribution to the neighboring beams is $S/N < 3$. Thus, with a probability of about 30–35%, it is impossible to see the signal from the FRB in the neighboring beams, despite its high S/N.

The situation when a strong pulse is observed in a single beam of LPA3 is rare, but not exceptional. For example, in the paper by Tyul'bashev, Tyul'bashev, and Malofeev (2018) of the 25 discovered RRATs, four had a peak flux density exceeding 20 Jy ($S/N > 65$) and one of them, RRAT J1005+3015 (with $S_p = 28.3$ Jy), has been also observed in a single beam. We processed data for this transient obtained between August 2014 and December 2017. Its strongest pulse (see Fig. 2) had a flux density of 74 Jy and is visible in a single beam.

A two-stage calibration was carried out to determine the flux density from the found FRB. At the first step, the gain in the frequency channels was equalized using a calibration signal with a known temperature recorded in all LPA beams. In the second step, the amplitude of the found source was compared with the amplitude of the calibration source. Since the calibration source was not observed at the zenith and its coordinate did not coincide with the center of the radiation pattern, some corrections have been applied for its flux density (see some additional details in Tyul'bashev *et al.* 2019). To estimate S_p we used a calibration source 4C+34.38 (J1406+34), which has a flux density of 25 Jy. After compensation for the measured DM , we estimated the flux density of FRB 20190203 to be $S_p = 20 \pm 4$ Jy. However, the exact coordinates of the source responsible for the burst are not known. Therefore, it is impossible to correct the flux density accurately taking into account the mismatch between the centroid of the radiation pattern and coordinates of the burst. The real flux density S_p can be 1.5–2 times higher.

Also, we carried out a search for repeated bursts in the direction J1340+3530, assuming different widths for a possible

pulse (taking it equal to 1–32 points in extent, i.e. assuming that W_e varies from 12.5 to 400 ms). We used DM measured for the detected pulse and convolved the data with a model sample pulse having $\tau_{sc} = 211$ ms to obtain the maximum response (see details in the paper by Brylyakova and Tyul'bashev 2023). We also note that the scattering in the Galaxy is insignificant and is determined by the environment between the burst and the Galaxy (Kuz'min, Losovskii, and Lapaev 2007). The search was carried out in the main (central) beam, where the first pulse was detected, as well as in the beams above and below it (at the right ascension $\pm 10^{\text{m}}$ from the coordinate measured for FRB 190203). No new pulses with $S/N > 7$ were detected in the lateral beams but a pulse with $\alpha_{2000} = 13^{\text{h}}39^{\text{m}}41^{\text{s}}$ was detected in the central beam in addition to the first pulse shown in Fig. 1. This pulse, detected in the data for June 16, 2018, has the maximum $S/N = 7.2$ obtained by averaging 2 points. With further averaging, S/N begins to fall. Nevertheless, we believe that this new pulse is false (see details and discussion in Appendix 2).




Figure 3. The SPI-ACS light curve with a resolution of 50 ms in the energy range from 80 keV to 10 MeV. The vertical scale shows the number of recorded counts. The horizontal scale shows time with $T_0 = 0$ corresponding to the LPA3 registration time of FRB 20190203. The average background count rate is indicated by a black line, its deviations at the level of $\pm 1 \sigma$ are shown by red lines.

4. Search for gamma-rays with INTEGRAL

Detecting gamma-rays from any FRB remains a challenge. Several attempts have been made to detect gamma-ray bursts (GRBs) from FRB sources and there were even several reports on the possible detection of such GRBs (DeLaunay *et al.* 2016). The most reliable gamma-ray transient coinciding in both time and position with the radio burst FRB 20200428A was detected from the soft gamma repeater SGR J1935+2154 (Mereghetti *et al.* 2020) known by its exceptional X-ray and gamma-ray activity. Recently it was assumed that FRB 20190425A might be a counterpart of the second known gravitational wave event GW 190425 associated with binary neutron star merger (Moroiaru *et al.* 2023). However, this hypothesis was disfavored by further considerations (Bhardwaj *et al.* 2023; Panther *et al.* 2023; Radice *et al.* 2024; Magaña Hernandez *et al.* 2024). The hypothesis is also challenged by the detection of GRB 190425A with the SPI-ACS detector aboard INTEGRAL coinciding in time with GW 190425 and believed to be

its electromagnetic companion (Pozanenko et al. 2019). There is also some contradictory evidence of transient gamma-ray radiation from the recently discovered radio burst repeater FRB 20240114A, demonstrating extreme burst activity (Xing and Yu 2024). Note, however, that none of the five hundred FRBs detected by the Canadian Hydrogen Intensity Mapping Experiment (CHIME) were associated with any known GRBs (Curtin et al. 2023). Thus, any detection of the gamma-ray activity from observed FRBs would be useful for understanding the origin of FRBs and the mechanism of their emission.

The International Gamma-Ray Astrophysics Laboratory INTEGRAL (Winkler et al. 2003) and the Gamma-ray Burst Monitor (GBM) aboard the *Fermi* gamma-ray observatory are the most effective experiments currently in operation for studying the cosmic GRBs and other gamma-ray transients (Kuulkers et al. 2021). On INTEGRAL, GRBs are detected within the fields of view (FOV) of its main telescopes, IBIS/ISGRI ($30^\circ \times 30^\circ$ FWZR) and SPI ($25^\circ \times 25^\circ$ FWZR) (e.g., Mereghetti et al. 2003; Vianello, Götz, and Mereghetti 2009; Minaev et al. 2014; Bošnjak et al. 2014; Higgins et al. 2017; Götz et al. 2019), but also outside their FOV (so-called off-axis GRBs) — directly by the detectors of the gamma-ray telescopes (Chelovekov et al. 2019) or their massive anti-coincidence shield (first of all by SPI-ACS, e.g., Rau et al. 2005; Savchenko, Neronov, and Courvoisier 2012; Minaev et al. 2014; Biltzinger, Burgess, and Greiner 2023). An important advantage of the observatory is its ability for the persistent monitoring of the almost entire sky.

INTEGRAL allows one to search for transient gamma-ray radiation that could accompany the detected FRB, as well as for a possible gamma-ray source within the localization region of this FRB. The first task could be solved with the omnidirectional scintillation detector SPI-ACS. It registers gamma-ray photons in a single wide energy channel of 80 keV – 10 MeV with a time resolution of 50 ms. The count rate is recorded continuously throughout the spacecraft's orbit (currently 68 hours), with the exception of a few hours when it passes through the radiation belts of the Earth. At the time of the FRB registration 2019-02-03 (UT) 02:18:10.284, the observatory was at a distance of 102,200 km from the center of the Earth, in the region with a rather stable radiation background. The angle between the FRB source and the X-axis of the observatory, which coincides with the main axis of the coded-aperture telescopes, was equal to 99° . Thus, the FRB localization region was outside FOV of the telescopes. But, it was not overshadowed by the Earth disk.

4.1 Prompt gamma-ray burst

Gamma-ray emission that could accompany FRB 20190203 should be detected by SPI-ACS. However, we have not found any statistically significant excess over the background level on a time scale of 50 ms during ± 100 s from the time of the FRB registration. The 3σ upper limit for a short (< 50 ms) excess in the count rate is equal to 270 counts per bin. It corresponds, after recalculation based on the SPI-ACS calibration by Minaev and Pozanenko (2023), to the upper limit for the 10–1000 keV




Figure 4. The S/N map of the sky area in the field of view of the INTEGRAL IBIS/ISGRI telescope obtained in the 20–150 keV range on June 10, 2005 at (UT) 22:39:35.381. It contains the localization region of FRB 20190203. The exposure is 0.7 s. The lines on the map show right ascension and declination in degrees.

fluence of a short (< 50 ms) GRB equal to 1.2×10^{-7} erg cm^{-2} in the case when it has a soft energy spectrum (with spectral hardness $E_p = 20$ keV) and 4.8×10^{-8} erg cm^{-2} in the case of its hard spectrum (with $E_p = 500$ keV). The SPI-ACS light curve is shown in Fig. 3.

The source could have been registered by other telescopes aboard INTEGRAL. The most suitable for such out-of-aperture registration is the position sensitive detector ISGRI located in the focal plane of the IBIS gamma-ray telescope. ISGRI registers hard X- and gamma-rays in the 20–400 keV range. Note that many cosmic GRBs have been indeed recorded in this way (Chelovekov et al. 2019). However, our analysis of the IBIS/ISGRI data has not revealed any significant excess above the background level on the time scales of 5 ms, 10 ms or 50 ms during 100 s before the FRB 20190203 registration (GRB probably should precede the radio burst because of scattering).

4.2 Hard X-ray activity on long-term scale

During the more than 20 years of the INTEGRAL operation on orbit, the localization region for FRB 20190203 has repeatedly come into view of the observatory's main telescopes, including in the IBIS/ISGRI field of view. The total exposure for this localization region having been in the IBIS FOV was 73.244 days. A direct search for possible short GRB events within the FRB 20190203 localization region with such a long exposure is very difficult, so a special technique for this analysis was developed and used (see Appendix 3).

In total, 112 candidates to bursts exceeding the selected threshold of 10σ on a time scale of 10 ms and 15 candidates

exceeding the same threshold on a scale of 50 ms have been found in the IBIS/ISGRI detector's light curves using the mentioned technique. Unfortunately, none of these candidates has been confirmed on the related sky images, none has occurred to be a real burst. For example, Fig. 4 shows the S/N map of the sky obtained in the 20–150 keV range within the IBIS FOV for the candidate to a burst detected on the light curve on June 10, 2005 at (UT) 22:39:35.381 on a time scale of 0.01 s. The duration of the burst and the exposure used to construct the map was 0.7 s. The position of FRB 20190203 is located close to the map center and indicated by a circle. No source is found at this position at the S/N level exceeding 3, moreover there is no significant S/N excesses on the entire map. Crosses indicate the positions of the Coma cluster and several other known X-ray sources (AGNs) that happen to be present in the field. None of them appeared on the map due to the very short exposure.

The 3σ upper limit for the flux in the 20–150 keV range from a possible source at the FRB 20190203 position is equal to 360 mCrab (8×10^{-9} erg cm $^{-2}$ s $^{-1}$). This limit can be considered typical for all detected “bursts”. It is likely that all the 112+15 “bursts” found in the IBIS/ISGRI detector's light curve are connected with fluctuations or have geophysical origin.

5. Discussion

A few comments should be given regarding the discovered FRB 20190203:

1). Karastergiou et al. (2015) carried out a search for FRBs at a frequency of 145 MHz using several LOFAR stations. No FRBs have been detected and an upper estimate given for the expected number of FRBs with a flux density of $S_p > 62$ Jy for the entire sky per day was 29 events. Our observations were carried out at a frequency of 111 MHz, which is close to 145 MHz, so we can use this upper estimate to evaluate the expected number of FRBs detectable with LPA3 per day. Up to 12 FRBs per day can flare up in the 17,000–square–degree sky area covered by the daily LPA3 survey. But, taking into account the small instantaneous field-of-view of LPA3, which is only 38 square degrees, we can expect to detect less than $12 \times (38/17,000) = 0.027$ events per day or 80 FRBs for 3000 days of our observations. In fact, we detected only one event (FRB 20190203) which suggests that the detection rate is 80 times lower than expected.

Either the LOFAR upper estimate for the expected rate of FRBs was too optimistic (indirectly it is confirmed by the very small number of FRBs detected at low frequencies to date) or we need to attribute such an incredibly low LPA3 detection rate of FRBs at 111 MHz to the low sensitivity of the metadata search.

Another explanation may be related to the idea that FRBs emit in a narrow frequency range. Indeed, any upper limit of the total number of FRBs in the entire sky implies that FRB emission is generated over a wide frequency range, but if FRB intrinsic emission occurs in a narrow band (as the model by Khangulyan, Barkov, and Popov (2022) suggests), then any

estimates for a narrower range, such as that of LPA3, should be significantly smaller.

2). The trigger for signal detection in the metadata was the appearance of a peak with a flux density of > 7.5 Jy in individual frequency channels, which approximately corresponds to S/N= 5 for these channels. A trigger operating on separate channels makes it possible to isolate strong pulse interference, but is too rough to search for pulsed dispersed signals throughout the entire observation band. In a standard search for dispersed signals DM is sorted out with subsequent summation of frequency channels, which in the case of 32–channel observations on LPA3 will lead to an increase in the S/N ratio by $32^{1/2} = 5.65$ times. Furthermore, scattering widens the pulse, and additional time averaging is necessary to obtain the maximum S/N value. For the discovered FRB 20190203, averaging over 211 ms/12.5 ms= 16 points will increase the S/N ratio by a factor of 4. Thus, a standard search for pulsed dispersed signals in PUMPS data with sorting out of DM and pulse width allows bursts that are an order of magnitude weaker than those considered in this paper to be detected. We are currently developing a new FRB search program.

3). It is known that cosmic plasma scattering can differ by an order of magnitude or greater from the average value at the same DM (Bhat et al. 2004; Kuz'min, Losovskii, and Lapaev 2007). Let us estimate the expected scattering at the observed value of DM for FRB 20190203 using the formula given by Cordes and Lazio (2003):

$$\log \tau_{sc}(\mu\text{s}) = -3.59 + 0.129 \log DM + 1.02 (\log DM)^2 - 4.4 \log \nu(\text{GHz}). \quad (1)$$

Substituting the value $DM = 134.4$ pc cm $^{-3}$ and $\nu = 0.11025$ GHz, we get the expected scattering $\tau_{sc} = 330$ ms. Taking into account possible deviations of the scattering by an order of magnitude from the expected value, our estimate $\tau_{sc} = 211$ ms is reasonable close to the obtained value.

4). The distance to the burst can be determined by the observed excessive DM . We don't know the environment inside the host galaxy of the burst. If we assume that there is no contribution from the host galaxy to the observed excessive DM , then according to the YMW16 model (Yao, Manchester, and Wang 2017) the redshift z is equal to 0.17 and the luminosity distance to the burst is $d_L \simeq 700$ Mpc ($d_L \approx 10^{27.3}$ cm). This value is an upper limit for the distance to the discovered FRB. In further calculations, we will rely on this value, but we will leave the dependence of all observed and calculated values on it and introduce a normalization coefficient ($d_{L,27.3} \leq 1$). Here and further in the paper, we used the following notation in CGS units $A_a = A/10^a$.

5). At a known distance to the FRB, its luminosity can be estimated. We use the formula by Petroff, Hessels, and Lorimer (2019):

$$L = (4\pi d_L^2 S_\nu \Delta\nu)/(1+z), \quad (2)$$

where L is the isotropic equivalent of the source luminosity in the frequency band $\Delta\nu$, d_L is luminosity distance, determined taking into account the redshift z of the source. Substituting




Figure 5. The figure shows the fit of the obtained data with a logarithmic straight line (red strokes) and a logarithmic parabola (blue line). There are the observation frequency on the vertical axis and the dispersion shift relative to the infinite frequency on the horizontal axis. Black points are the dispersion shift in 32 channels obtained from the dynamic spectrum according to Fig. 1.

the measured flux $S_\nu = 20 \pm 4$ Jy, the channel width 2.5 MHz and $z \simeq 0.17$ in Eq. (2) we obtain

$$L \simeq 3 \times 10^{40} d_{L,27.3}^2 \text{ erg s}^{-1}. \quad (3)$$

According to the analysis by Cui et al. (2022) of the FRB luminosities detected in CHIME, the lower limit for the luminosity in the sample of 135 FRBs is at the level of $10^{40} - 10^{42} \text{ erg s}^{-1}$ which is consistent with our estimate for the luminosity of FRB 20190203 (Eq. 3).

At the same time, it must be remembered that the resulting estimate for the luminosity depends on the frequency band used for its evaluation. In particular, the bandwidth of CHIME is equal to 400 MHz, which is 160 times (two orders of magnitude) wider than the LPA3 reception bandwidth. Obviously, the luminosity will be higher when observing the same burst in a wider reception band. When comparing the luminosities of bursts measured by different instruments with different receiving bands and different central frequencies, it makes sense to use the luminosity normalized per unit frequency ($\Delta\nu$). In our case, we get

$$L_\nu \simeq 10^{34} d_{L,27.3}^2 \text{ erg s}^{-1} \text{ Hz}^{-1}. \quad (4)$$

Figure 16 in the paper by Petroff, Hessels, and Lorimer (2019) shows estimates of FRB normalized luminosities obtained by different telescopes except those found by CHIME. These luminosities are in the range $L_\nu \sim 10^{28} - 10^{34} \text{ erg s}^{-1} \text{ Hz}^{-1}$. The normalized luminosity of FRB 20190203 is close to the brightest bursts in the figure.

6). A short (≤ 50 ms) gamma-ray signal coinciding in time with FRB 20190203 was not detected from the FRB location at energies above 80 keV. The 3σ level for its fluence in the 10–1000 keV range was $4.8 \times 10^{-8} \text{ erg cm}^{-2}$. Assuming the signal duration to be ~ 50 ms and substituting the above distance

from the FRB source, we can estimate its maximum luminosity in gamma-rays at which it can still be detected as

$$L_\gamma \leq 6 \times 10^{49} d_{L,27.3}^2 \text{ erg s}^{-1}. \quad (5)$$

Note that such a high gamma-ray luminosity corresponds, however, to a very moderate normalized luminosity in terms of Eq. (4) $L_{\gamma,\nu} \leq 2.4 \times 10^{31} d_{L,27.3}^2 \text{ erg s}^{-1} \text{ Hz}^{-1}$.

7). Many models have been proposed to explain the origin of FRBs (Lyubarsky 2014; Metzger, Margalit, and Sironi 2019; Cordes and Chatterjee 2019; Petroff, Hessels, and Lorimer 2019; Xiao, Wang, and Dai 2021; Zhang 2022). In our opinion, the observed properties of FRB 20190203 are best explained by the model of a synchrotron maser source excited by a magnetar (Khangulyan, Barkov, and Popov 2022). For the normalized luminosity given by Eq. (4) and assuming that the radio pulse occupied 20 MHz in the frequency range, as measured in repeating bursts of FRB 20180916B by LOFAR (Pleunis et al. 2021), we have the total radio luminosity $L_r = 2 \times 10^{41} \Delta\nu_{7.3} d_{L,27.3}^2 \text{ erg s}^{-1}$, here we entered a notation of $\Delta\nu_{7.3}$ for the FRB pulse width (in 20 MHz units).

We observed a scattered burst, which became significantly wider than it was at birth. The duration of almost all FRB pulses is in the range of 1–10 ms^{d,e,f} (Petroff et al. 2016; CHIME/FRB Collaboration et al. 2021; Xu et al. 2023). If the observed pulse at birth was as narrow as the pulses observed at high frequencies, the physical power of the radio emission in vicinity of its source may be $100/dt_{-2}$ times higher, where dt_{-2} is a duration of FRB in centiseconds ($dt = dt_{-2}/100$). That is, the luminosity of radio emission from the FRB 20190203 source during this pulse could reach the value

$$L_0 \sim 2 \times 10^{43} \Delta\nu_{7.3} d_{L,27.3}^2 / dt_{-2} \text{ erg s}^{-1}, \quad (6)$$

making the discovered event one of the most powerful FRBs known. Of course, the quadratic dependence of luminosity on distance is somewhat formal, since all estimates are given for the expected value of $z = 0.17$.

Using the burst parameters, L and ν , it is possible to estimate the required power of a shock wave caused by a magnetar flare \dot{E}_β and going to infinity (see equation 19 in the paper by Khangulyan, Barkov, and Popov 2022):

$$\dot{E}_\beta \approx 6 \times 10^{44} R_{15}^{4/3} \Gamma_{\beta,3}^{4/3} \sigma_{\beta,-2}^{1/3} \text{ erg s}^{-1} \quad (7)$$

Here R_{15} is the distance from the magnetar to the front of the standing shock wave, $\sigma_{\beta,-2}$ is magnetization in the flare, and $\Gamma_{\beta,3}$ is the flare Lorentz factor. The radiation efficiency, that is, the degree of conversion of the flare energy into radiation, can be estimated by dividing the observed physical luminosity by the kinetic power of the flare ϵ_{eff} . Combining Eqs.(6) and (7) we get

$$\epsilon_{\text{eff}} = \frac{L_0}{\dot{E}_\beta} \sim 0.04 \Delta\nu_{7.3} d_{L,27.3}^2 t_{-2}^{-1} R_{15}^{-4/3} \Gamma_{\beta,3}^{-4/3} \sigma_{\beta,-2}^{-1/3} \quad (8)$$

d. www.astronomy.swin.edu.au/pulsar/frbcat/

e. www.chime-frb.ca/catalog

f. blinkverse.alkaidos.cn/

As can be seen, the degree of flare energy conversion should be high enough, at the level of a few percents, which is consistent with the parameters obtained in previous studies (Khangulyan, Barkov, and Popov 2022; Barkov and Popov 2022).

Equation (8) reproduces conversion efficiency of the kinetic energy of the shock wave to the radio pulse. The X-ray flare is probably generated within the magnetosphere of a magnetar itself, so the energy budget can be higher. Following the scenario described by Sharma, Barkov, and Lyutikov (2023, see Tabl. 1, the Large CME case), the total energy of the flare can be estimated as:

$$E_{tot,fl} \sim \sqrt{dt\dot{E}_{fl}E_{B,NS}} \sim 3 \times 10^{45} dt_{-2}^{1/2} \dot{E}_{fl,45}^{1/2} E_{B,NS,48}^{1/2} \text{ erg}, \quad (9)$$

where $E_{B,NS} \sim B_{NS}^2 R_{NS}^3 \sim 10^{48} B_{NS,15}^2 R_{NS,6}^3 \text{ erg}$ is the total energy of the magnetosphere of a magnetar. The upper limit on the power of magnetic field dissipation is

$$\dot{E}_{tot,fl} \sim \frac{E_{tot,fl}}{dt} \sim 3 \times 10^{47} dt_{-2}^{-1/2} \dot{E}_{fl,45}^{1/2} E_{B,NS,48}^{1/2} \text{ erg s}^{-1}. \quad (10)$$

The X-ray luminosity cannot exceed the total energy budget (Eq. 10), so in this scenario it should be significantly lower than the upper limit (Eq. 5) provided by X-ray observations for the maximum distance to the source ($d_{max} \simeq 700 \text{ Mpc}$). On the other hand, using the same upper limit (Eq. 5) and assuming that a fraction $\xi \leq 1$ of the total flare power was indeed emitted in hard X-rays we can estimate the minimum distance to the source $d_{min} \simeq 50\xi^{1/2} \text{ Mpc}$ explaining why such an X-ray flare has still not been detected.

8). The narrow bandwidth of LPA3 does not allow us to distinguish the linear dependence of the dispersion curve from the parabolic one. As can be seen from Fig. 5 we must have a bandwidth wider than 5 MHz in order to distinguish them.

9). Note that linear polarization of radio emission is expected in the FRB model by Khangulyan, Barkov, and Popov (2022). Unfortunately we can not measure it correctly for FRB 190203 because all wave dipoles of the LPA LPI array have single linear polarization.

10). The low localization accuracy of the found FRB 20190203 (the localization area is $60' \times 20'$ at 68% confidence) and its extreme remoteness ($d \simeq 700 \text{ Mpc}$) do not allow us to identify it with any particular host galaxy. We have found dozens of host galaxy candidates, but the number of missed galaxies (due to the known incompleteness of extragalactic catalogs) may be much higher. On the other hand, such accuracy allows us to assert that the found burst does not coincide with any previously detected FRBs^{g,h} (CHIME/FRB Collaboration *et al.* 2021; Xu *et al.* 2023).

Note finally that all our estimates in this section were made based on the assumption that the DM accumulated in the Galaxy is 20.3 pc cm^{-3} , and the DM in the host galaxy is negligible. At the same time, it is known that the DM in the Galaxy can be between 20 and 30 pc cm^{-3} (see Fig. 2 in the paper by

Cook *et al.* 2023). If we assume $DM_{host} = 100 \text{ pc cm}^{-3}$ (approximately 60% of all ATNF pulsars Manchester *et al.* 2005, in the Galaxy have $DM > 100 \text{ pc cm}^{-3}$), then $z = 0.033$, $d = 146 \text{ Mpc}$. In this case, the burst occurred close to the Galaxy, L will decrease by 25 times and the burst will become quite ordinary. We believe that only the upper estimate of the distance, which determines the maximum luminosity, is reliable. Other estimates are speculative.

6. Conclusion

We report the discovery of an intense ($S_p = 20 \pm 4 \text{ Jy}$) dispersed ($DM = 134.4 \pm 2 \text{ pc cm}^{-3}$) radio pulse at 111 MHz with the Pushchino LPA radio telescope. The pulse was detected in the direction of $\alpha = 13^{\text{h}}40^{\text{m}}46^{\text{s}} \pm 2^{\text{m}}$, $\delta = 35^{\circ}30' \pm 10'$ (epoch 2000.0) on UT 2019-02-03 02:18:10.284. It lasted $211 \pm 6 \text{ ms}$. The large measured DM indicates its extragalactic origin (it corresponds to the luminosity distance $\sim 700 \text{ Mpc}$ after subtracting the Milky Way contribution). Further analysis has shown that the pulse is likely another fast radio burst; we called it to be FRB 20190203. Repeated radio bursts from the source have not yet been detected. No also any activity has been detected in the gamma-ray range (according to INTEGRAL observations).

FRB 20190203 is the first burst of extragalactic origin discovered in the PUMPS survey. It is also the second FRB detected at such a low frequency (111 MHz), and the first one among on-off FRBs. This unique event is one of the most powerful FRBs detected so far.

Acknowledgement

SAT, EAB are grateful to the Russian Science Foundation 22-12-00236ⁱ for supporting the work in part of the FRB search with LPA. ASP, SAG, IVC, PYuM, and MVB are grateful to the Russian Science Foundation (project no. 23-12-00220^j) for partial support of the INTEGRAL observatory data reduction and analysis. VAS and ASP were supported in part through computational resources of HPC facilities at NRU HSE (project 663983).

Data Availability Statement The PUMPS survey is not finished yet. The raw data underlying this paper will be shared on reasonable request to the corresponding author. The INTEGRAL data may be obtained through the public archive of the mission^k.

References

Barkov, Maxim V., and Sergei B. Popov. 2022. Formation of periodic FRB in binary systems with eccentricity. *MNRAS* 515, no. 3 (September): 4217–4228. <https://doi.org/10.1093/mnras/stac1562>. arXiv: 2204.13489 [astro-ph.HE].

i. rscf.ru/project/22-12-00236/

j. rscf.ru/project/23-12-00220/

k. www.isdc.unige.ch

g. <https://www.chime-frb.ca/catalog/>

h. <https://blinkverse.zero2x.org/overview>

- Bhardwaj, Mohit, Antonella Palmese, Ignacio Magaña Hernandez, Virginia D'Emilio, and Soichiro Morisaki. 2023. GW190425 and FRB20190425A: Challenges for Fast Radio Bursts as Multi-Messenger Sources from Binary Neutron Star Mergers. *arXiv e-prints* (June): arXiv:2306.00948. <https://doi.org/10.48550/arXiv.2306.00948>. arXiv: 2306.00948 [astro-ph.HE].
- Bhat, N. D. Ramesh, James M. Cordes, Fernando Camilo, David J. Nice, and Duncan R. Lorimer. 2004. Multifrequency Observations of Radio Pulse Broadening and Constraints on Interstellar Electron Density Microstructure. *ApJ* 605, no. 2 (April): 759–783. <https://doi.org/10.1086/382680>. arXiv: astro-ph/0401067 [astro-ph].
- Biltzinger, Björn, J. Michael Burgess, and Jochen Greiner. 2023. Time-resolved spectral catalogue of INTEGRAL/SPI gamma-ray bursts. *A&A* 675 (July): A175. <https://doi.org/10.1051/0004-6361/202245191>.
- Bošnjak, Ž., D. Götz, L. Bouchet, S. Schanne, and B. Cordier. 2014. The spectral catalogue of INTEGRAL gamma-ray bursts. results of the joint IBIS/SPI spectral analysis. *A&A* 561 (January): A25. <https://doi.org/10.1051/0004-6361/201322256>. arXiv: 1309.3174 [astro-ph.HE].
- Brylyakova, E. A., and S. A. Tyul'bashev. 2023. On the Problems of Detecting Fast Radio Bursts with the LPA LPI. *Astronomy Reports* 67, no. 2 (February): 163–171. <https://doi.org/10.1134/S106377292302004X>. arXiv: 2305.02778 [astro-ph.HE].
- Chawla, P., B. C. Andersen, M. Bhardwaj, E. Fonseca, A. Josephy, V. M. Kaspi, D. Michilli, et al. 2020. Detection of Repeating FRB 180916.J0158+65 Down to Frequencies of 300 MHz. *ApJ* 896, no. 2 (June): L41. <https://doi.org/10.3847/2041-8213/ab96bf>. arXiv: 2004.02862 [astro-ph.HE].
- Chelovekov, I. V., S. A. Grebenev, A. S. Pozanenko, and P. Yu. Minaev. 2019. New Gamma-Ray Bursts Found in the Archival Data from the IBIS/ISGRI Telescope of the INTEGRAL Observatory. *Astronomy Letters* 45, no. 10 (December): 635–654. <https://doi.org/10.1134/S1063773719100025>. arXiv: 2005.11606 [astro-ph.HE].
- CHIME/FRB Collaboration, Mandana Amiri, Bridget C. Andersen, Kevin Bandura, Sabrina Berger, Mohit Bhardwaj, Michelle M. Boyce, et al. 2021. The First CHIME/FRB Fast Radio Burst Catalog. *ApJS* 257, no. 2 (December): 59. <https://doi.org/10.3847/1538-4365/ac33ab>. arXiv: 2106.04352 [astro-ph.HE].
- Coenen, Thijs, Joeri van Leeuwen, Jason W. T. Hessels, Ben W. Stappers, Vladislav I. Kondratiev, A. Alexov, R. P. Breton, et al. 2014. The LOFAR pilot surveys for pulsars and fast radio transients. *A&A* 570 (October): A60. <https://doi.org/10.1051/0004-6361/201424495>. arXiv: 1408.0411 [astro-ph.SR].
- Cook, Amanda M., Mohit Bhardwaj, B. M. Gaensler, Paul Scholz, Gwendolyn M. Eadie, Alex S. Hill, Victoria M. Kaspi, et al. 2023. An FRB Sent Me a DM: Constraining the Electron Column of the Milky Way Halo with Fast Radio Burst Dispersion Measures from CHIME/FRB. *ApJ* 946, no. 2 (April): 58. <https://doi.org/10.3847/1538-4357/acbbd0>. arXiv: 2301.03502 [astro-ph.GA].
- Cordes, J. M., and T. J. W. Lazio. 2002. NE2001. I. A New Model for the Galactic Distribution of Free Electrons and its Fluctuations. *arXiv e-prints* (July): astro-ph/0207156. <https://doi.org/10.48550/arXiv.astro-ph/0207156>. arXiv: astro-ph/0207156 [astro-ph.HE].
- . 2003. NE2001. II. Using Radio Propagation Data to Construct a Model for the Galactic Distribution of Free Electrons. *arXiv e-prints* (January): astro-ph/0301598. <https://doi.org/10.48550/arXiv.astro-ph/0301598>. arXiv: astro-ph/0301598 [astro-ph.HE].
- Cordes, James M., and Shami Chatterjee. 2019. Fast Radio Bursts: An Extragalactic Enigma. *ARA&A* 57 (August): 417–465. <https://doi.org/10.1146/annurev-astro-091918-104501>. arXiv: 1906.05878 [astro-ph.HE].
- Cui, Xiang-Han, Cheng-Min Zhang, Di Li, Jian-Wei Zhang, Bo Peng, Wei-Wei Zhu, Richard Strom, et al. 2022. Luminosity distribution of fast radio bursts from CHIME/FRB Catalog 1 by means of the updated Macquart relation. *Ap&SS* 367, no. 7 (July): 66. <https://doi.org/10.1007/s10509-022-04093-y>. arXiv: 2210.02290 [astro-ph.HE].
- Curtin, Alice P., Shriharsh P. Tendulkar, Alexander Josephy, Pragya Chawla, Bridget Andersen, Victoria M. Kaspi, Mohit Bhardwaj, et al. 2023. Limits on Fast Radio Burst-like Counterparts to Gamma-Ray Bursts Using CHIME/FRB. *ApJ* 954, no. 2 (September): 154. <https://doi.org/10.3847/1538-4357/ace52f>. arXiv: 2208.00803 [astro-ph.HE].
- DeLaunay, J. J., D. B. Fox, K. Murase, P. Mészáros, A. Keivani, C. Messick, M. A. Mostafá, F. Oikonomou, G. Tešić, and C. F. Turley. 2016. Discovery of a Transient Gamma-Ray Counterpart to FRB 131104. *ApJ* 832, no. 1 (November): L1. <https://doi.org/10.3847/2041-8205/832/1/L1>. arXiv: 1611.03139 [astro-ph.HE].
- Götz, Diego, Christian Gouiffès, Jérôme Rodriguez, Philippe Laurent, Elisabeth Jourdain, Jean-Pierre Roques, Sandro Mereghetti, et al. 2019. INTEGRAL results on gamma-ray bursts and polarization of hard X-ray sources. *New A Rev.* 87 (December): 101537. <https://doi.org/10.1016/j.newar.2020.101537>.
- Higgins, A. B., R. L. C. Starling, D. Götz, S. Mereghetti, K. Wiersema, T. Maccarone, J. P. Osborne, et al. 2017. Investigating the nature of the INTEGRAL gamma-ray bursts and sub-threshold triggers with Swift follow-up. *MNRAS* 470, no. 1 (September): 314–323. <https://doi.org/10.1093/mnras/stx1163>. arXiv: 1705.06744 [astro-ph.HE].
- Houben, L. J. M., L. G. Spitler, S. ter Veen, J. P. Rachen, H. Falcke, and M. Kramer. 2019. Constraints on the low frequency spectrum of FRB 121102. *A&A* 623 (March): A42. <https://doi.org/10.1051/0004-6361/201833875>. arXiv: 1902.01779 [astro-ph.HE].
- Karastergiou, A., J. Chennamangalam, W. Armour, C. Williams, B. Mort, F. Dulwich, S. Salvini, et al. 2015. Limits on fast radio bursts at 145 MHz with ARTEMIS, a real-time software backend. *MNRAS* 452, no. 2 (September): 1254–1262. <https://doi.org/10.1093/mnras/stv1306>. arXiv: 1506.03370 [astro-ph.IM].
- Khangulyan, D., Maxim V. Barkov, and S. B. Popov. 2022. Fast Radio Bursts by High-frequency Synchrotron Maser Emission Generated at the Reverse Shock of a Powerful Magnetar Flare. *ApJ* 927, no. 1 (March): 2. <https://doi.org/10.3847/1538-4357/ac4bdf>. arXiv: 2106.09858 [astro-ph.HE].
- Kuulkers, Erik, Carlo Ferrigno, Peter Kretschmar, Julia Alfonso-Garzón, Marius Baab, Angela Bazzano, Guillaume Bélanger, et al. 2021. INTEGRAL reloaded: Spacecraft, instruments and ground system. *New A Rev.* 93 (December): 101629. <https://doi.org/10.1016/j.newar.2021.101629>.
- Kuz'min, A. D., B. Ya. Losovskii, and K. A. Lapaev. 2007. Measurements of the scattering of pulsar radio emission. *Astronomy Reports* 51, no. 8 (August): 615–623. <https://doi.org/10.1134/S1063772907080021>.
- Law, C. J., M. W. Abruzzo, C. G. Bassa, G. C. Bower, S. Burke-Spolaor, B. J. Butler, T. Cantwell, et al. 2017. A Multi-telescope Campaign on FRB 121102: Implications for the FRB Population. *ApJ* 850, no. 1 (November): 76. <https://doi.org/10.3847/1538-4357/aa9700>. arXiv: 1705.07553 [astro-ph.HE].
- Lonsdale, C. J., R. J. Cappallo, M. F. Morales, F. H. Briggs, L. Benkevitch, J. D. Bowman, J. D. Bunton, et al. 2009. The Murchison Widefield Array: Design Overview. *IEEE Proceedings* 97, no. 8 (August): 1497–1506. <https://doi.org/10.1109/JPROC.2009.2017564>. arXiv: 0903.1828 [astro-ph.IM].
- Lorimer, D. R., M. Bailes, M. A. McLaughlin, D. J. Narkevic, and F. Crawford. 2007. A Bright Millisecond Radio Burst of Extragalactic Origin. *Science* 318, no. 5851 (November): 777. <https://doi.org/10.1126/science.1147532>. arXiv: 0709.4301 [astro-ph].
- Lorimer, D. R., and M. Kramer. 2004. *Handbook of Pulsar Astronomy*. Vol. 4.

- Lyubarsky, Yu. 2014. A model for fast extragalactic radio bursts. *MNRAS* 442 (July): L9–L13. <https://doi.org/10.1093/mnras/lu046>. arXiv: 1401.6674 [astro-ph.HE].
- Magaña Hernandez, Ignacio, Virginia D'Emilio, Soichiro Morisaki, Mohit Bhardwaj, and Antonella Palmese. 2024. On the association of GW190425 with its potential electromagnetic counterpart FRB 20190425A. *arXiv e-prints* (April): arXiv:2404.02496. <https://doi.org/10.48550/arXiv.2404.02496>. arXiv: 2404.02496 [astro-ph.HE].
- Majid, Walid A., Aaron B. Pearlman, Thomas A. Prince, Robert S. Wharton, Charles J. Naudet, Karishma Bansal, Liam Connor, Mohit Bhardwaj, and Shriharsh P. Tendulkar. 2021. A Bright Fast Radio Burst from FRB 20200120E with Sub-100 Nanosecond Structure. *ApJ* 919, no. 1 (September): L6. <https://doi.org/10.3847/2041-8213/ac1921>. arXiv: 2105.10987 [astro-ph.HE].
- Manchester, R. N., G. B. Hobbs, A. Teoh, and M. Hobbs. 2005. The Australia Telescope National Facility Pulsar Catalogue. *AJ* 129, no. 4 (April): 1993–2006. <https://doi.org/10.1086/428488>. arXiv: astro-ph/0412641 [astro-ph].
- Marcote, B., K. Nimmo, J. W. T. Hessels, S. P. Tendulkar, C. G. Bassa, Z. Paragi, A. Keimpema, et al. 2020. A repeating fast radio burst source localized to a nearby spiral galaxy. *Nature* 577, no. 7789 (January): 190–194. <https://doi.org/10.1038/s41586-019-1866-z>. arXiv: 2001.02222 [astro-ph.HE].
- Mereghetti, S., D. Götz, J. Borkowski, R. Walter, and H. Pedersen. 2003. The INTEGRAL Burst Alert System. *A&A* 411 (November): L291–L297. <https://doi.org/10.1051/0004-6361:20031289>. arXiv: astro-ph/0308173 [astro-ph].
- Mereghetti, S., V. Savchenko, C. Ferrigno, D. Götz, M. Rigoselli, A. Tiengo, A. Bazzano, et al. 2020. INTEGRAL Discovery of a Burst with Associated Radio Emission from the Magnetar SGR 1935+2154. *ApJ* 898, no. 2 (August): L29. <https://doi.org/10.3847/2041-8213/aba2cf>. arXiv: 2005.06335 [astro-ph.HE].
- Metzger, Brian D., Ben Margalit, and Lorenzo Sironi. 2019. Fast radio bursts as synchrotron maser emission from decelerating relativistic blast waves. *MNRAS* 485, no. 3 (May): 4091–4106. <https://doi.org/10.1093/mnras/stz700>. arXiv: 1902.01866 [astro-ph.HE].
- Minaev, P. Yu., and A. S. Pozanenko. 2023. Calibrating SPI-ACS/INTEGRAL for gamma-ray bursts and re-estimating energetics of GRB/GW 190425 in gamma-ray range. *MNRAS* 525, no. 2 (October): 2411–2421. <https://doi.org/10.1093/mnras/stad2416>. arXiv: 2308.10266 [astro-ph.HE].
- Minaev, P. Yu., A. S. Pozanenko, S. V. Molkov, and S. A. Grebenev. 2014. Catalog of short gamma-ray transients detected in the SPI/INTEGRAL experiment. *Astronomy Letters* 40, no. 5 (May): 235–267. <https://doi.org/10.1134/S106377371405003X>. arXiv: 1405.3784 [astro-ph.HE].
- Moroianu, Alexandra, Linqing Wen, Clancy W. James, Shunke Ai, Manoj Kovalam, Fiona H. Panther, and Bing Zhang. 2023. An assessment of the association between a fast radio burst and binary neutron star merger. *Nature Astronomy* 7 (May): 579–589. <https://doi.org/10.1038/s41550-023-01917-x>. arXiv: 2212.00201 [astro-ph.HE].
- Nan, Rendong, Di Li, Chengjin Jin, Qiming Wang, Lichun Zhu, Wenbai Zhu, Haiyan Zhang, Youling Yue, and Lei Qian. 2011. The Five-Hundred Aperture Spherical Radio Telescope (fast) Project. *International Journal of Modern Physics D* 20, no. 6 (January): 989–1024. <https://doi.org/10.1142/S0218271811019335>. arXiv: 1105.3794 [astro-ph.IM].
- Panther, Fiona H., Gemma E. Anderson, Shivani Bhandari, Adelle J. Goodwin, Natasha Hurley-Walker, Clancy W. James, Adela Kawka, et al. 2023. The most probable host of CHIME FRB 190425A, associated with binary neutron star merger GW190425, and a late-time transient search. *MNRAS* 519, no. 2 (February): 2235–2250. <https://doi.org/10.1093/mnras/stac3597>. arXiv: 2212.00954 [astro-ph.HE].
- Petroff, E., E. D. Barr, A. Jameson, E. F. Keane, M. Bailes, M. Kramer, V. Morello, D. Tabbara, and W. van Straten. 2016. FRBCAT: The Fast Radio Burst Catalogue. *PASA* 33 (September): e045. <https://doi.org/10.1017/pasa.2016.35>. arXiv: 1601.03547 [astro-ph.HE].
- Petroff, E., J. W. T. Hessels, and D. R. Lorimer. 2019. Fast radio bursts. *A&A Rev.* 27, no. 1 (May): 4. <https://doi.org/10.1007/s00159-019-0116-6>. arXiv: 1904.07947 [astro-ph.HE].
- Pleunis, Z., D. Michilli, C. G. Bassa, J. W. T. Hessels, A. Naidu, B. C. Andersen, P. Chawla, et al. 2021. LOFAR Detection of 110–188 MHz Emission and Frequency-dependent Activity from FRB 20180916B. *ApJ* 911, no. 1 (April): L3. <https://doi.org/10.3847/2041-8213/abec72>. arXiv: 2012.08372 [astro-ph.HE].
- Pozanenko, A. S., P. Yu. Minaev, S. A. Grebenev, and I. V. Chelovekov. 2019. Observation of the Second LIGO/Virgo Event Connected with a Binary Neutron Star Merger S190425z in the Gamma-Ray Range. *Astronomy Letters* 45, no. 11 (February): 710–727. <https://doi.org/10.1134/S1063773719110057>. arXiv: 1912.13112 [astro-ph.HE].
- Radice, David, Giacomo Ricigliano, Mukul Bhattacharya, Albino Perego, Farrukh J. Fattoyev, and Kohta Murase. 2024. What if GW190425 did not produce a black hole promptly? *MNRAS* 528, no. 4 (March): 5836–5844. <https://doi.org/10.1093/mnras/stae400>. arXiv: 2309.15195 [astro-ph.HE].
- Rau, A., A. V. Kienlin, K. Hurley, and G. G. Lichti. 2005. The 1st INTEGRAL SPI-ACS gamma-ray burst catalogue. *A&A* 438, no. 3 (August): 1175–1183. <https://doi.org/10.1051/0004-6361:20053159>. arXiv: astro-ph/0504357 [astro-ph].
- Revnivtsev, M. G., R. A. Sunyaev, D. A. Varshalovich, V. V. Zheleznyakov, A. M. Cherepashchuk, A. A. Lutovinov, E. M. Churazov, S. A. Grebenev, and M. R. Gilfanov. 2004. A Hard X-ray Survey of the Galactic-Center Region with the IBIS Telescope of the INTEGRAL Observatory: A Catalog of Sources. *Astronomy Letters* 30 (June): 382–389. <https://doi.org/10.1134/1.1764884>. arXiv: astro-ph/0402027 [astro-ph].
- Rowlinson, A., M. E. Bell, T. Murphy, C. M. Trott, N. Hurley-Walker, S. Johnston, S. J. Tingay, et al. 2016. Limits on Fast Radio Bursts and other transient sources at 182 MHz using the Murchison Widefield Array. *MNRAS* 458, no. 4 (June): 3506–3522. <https://doi.org/10.1093/mnras/stw451>. arXiv: 1602.07544 [astro-ph.HE].
- Samodurov, V. A., A. S. Pozanenko, A. E. Rodin, D. D. Churakov, D. V. Dumskij, E. A. Isaev, A. N. Kazantsev, et al. 2017. Observations of transient phenomena in bsa radio survey at 110 mhz. In *Data analytics and management in data intensive domains*, edited by Leonid Kalinichenko, Sergei O. Kuznetsov, and Yannis Manolopoulos, 130–141. Cham: Springer International Publishing.
- Samodurov, V. A., S. A. Tyul'bashev, M. O. Toropov, and S. V. Logvinenko. 2022. Detection Statistics of Pulse Signals at Declinations from +42° to +52° at the Frequency 111 MHz. *Astronomy Reports* 66, no. 4 (April): 341–347. <https://doi.org/10.1134/S1063772922040059>. arXiv: 2207.09197 [astro-ph.HE].
- Savchenko, V., A. Neronov, and T. J.-L. Courvoisier. 2012. Timing properties of gamma-ray bursts detected by SPI-ACS detector onboard INTEGRAL. *A&A* 541 (May): A122. <https://doi.org/10.1051/0004-6361/201218877>. arXiv: 1203.1344 [astro-ph.HE].
- Sharma, Praveen, Maxim V. Barkov, and Maxim Lyutikov. 2023. Relativistic coronal mass ejections from magnetars. *MNRAS* 524, no. 4 (October): 6024–6051. <https://doi.org/10.1093/mnras/stad2192>. arXiv: 2302.08848 [astro-ph.HE].
- Shishov, V. I., I. V. Chashei, V. V. Oreshko, S. V. Logvinenko, S. A. Tyul'bashev, I. A. Subaev, P. M. Svidskii, V. B. Lapshin, and R. D. Dagkesamanski. 2016. Monitoring of the turbulent solar wind with the upgraded Large Phased Array of the Lebedev Institute of Physics: First results. *Astronomy Reports* 60, no. 12 (December): 1067–1082. <https://doi.org/10.1134/S1063772916110068>.

- Sokolowski, M., N. D. R. Bhat, J. -P. Macquart, R. M. Shannon, K. W. Bannister, R. D. Ekers, D. R. Scott, et al. 2018. No Low-frequency Emission from Extremely Bright Fast Radio Bursts. *ApJ* 867, no. 1 (November): L12. <https://doi.org/10.3847/2041-8213/aae58d>. arXiv: 1810.04355 [astro-ph.HE].
- Tyul'bashev, S. A., P. Yu. Golysheva, V. S. Tyul'bashev, and I. A. Subaev. 2019. Search for Long-term Variability of Several Flat-Spectrum Sources at 111 MHz. *Astronomy Reports* 63, no. 11 (November): 920–931. <https://doi.org/10.1134/S1063772919100068>.
- Tyul'bashev, S. A., V. S. Tyul'bashev, and V. M. Malofeev. 2018. Detection of 25 new rotating radio transients at 111 MHz. *A&A* 618 (October): A70. <https://doi.org/10.1051/0004-6361/201833102>. arXiv: 1807.07565 [astro-ph.HE].
- Tyul'bashev, S. A., V. S. Tyul'bashev, V. M. Malofeev, S. V. Logvinenko, V. V. Oreshko, R. D. Dagkesamanskii, I. V. Chashei, V. I. Shishov, and N. N. Bursov. 2018. Detection of Five New RRATs at 111 MHz. *Astronomy Reports* 62, no. 1 (January): 63–71. <https://doi.org/10.1134/S1063772918010079>. arXiv: 2207.11032 [astro-ph.HE].
- Tyul'bashev, S. A., V. S. Tyul'bashev, V. V. Oreshko, and S. V. Logvinenko. 2016. Detection of new pulsars at 111 MHz. *Astronomy Reports* 60, no. 2 (February): 220–232. <https://doi.org/10.1134/S1063772916020128>. arXiv: 2208.08839 [astro-ph.HE].
- Tyul'bashev, S. A., G. E. Tyul'basheva, M. A. Kitaeva, I. L. Ovchinnikov, V. V. Oreshko, and S. V. Logvinenko. 2024. Pushchino multibeam pulsar search – IV. Detection of new pulsars at declinations $-9^\circ < \delta < +55^\circ$. *MNRAS* 528, no. 2 (February): 2220–2231. <https://doi.org/10.1093/mnras/stae070>. arXiv: 2401.10706 [astro-ph.HE].
- Tyul'bashev, Sergei A., Marina A. Kitaeva, and Gayane E. Tyul'basheva. 2022. Pushchino multibeam pulsar search – I. Targeted search of weak pulsars. *MNRAS* 517, no. 1 (November): 1112–1125. <https://doi.org/10.1093/mnras/stac2404>. arXiv: 2203.15540 [astro-ph.HE].
- van Haarlem, M. P., M. W. Wise, A. W. Gunst, G. Heald, J. P. McKean, J. W. T. Hessels, A. G. de Bruyn, et al. 2013. LOFAR: The LOw-Frequency ARray. *A&A* 556 (August): A2. <https://doi.org/10.1051/0004-6361/201220873>. arXiv: 1305.3550 [astro-ph.IM].
- Vianello, G., D. Götz, and S. Mereghetti. 2009. The updated spectral catalogue of INTEGRAL gamma-ray bursts. *A&A* 495, no. 3 (March): 1005–1032. <https://doi.org/10.1051/0004-6361:200811251>. arXiv: 0812.3349 [astro-ph].
- Winkler, C., T. J. -L. Courvoisier, G. Di Cocco, N. Gehrels, A. Giménez, S. Grebenev, W. Hermsen, et al. 2003. The INTEGRAL mission. *A&A* 411 (November): L1–L6. <https://doi.org/10.1051/0004-6361:20031288>.
- Xiao, Di, FaYin Wang, and ZiGao Dai. 2021. The physics of fast radio bursts. *Science China Physics, Mechanics, and Astronomy* 64, no. 4 (April): 249501. <https://doi.org/10.1007/s11433-020-1661-7>. arXiv: 2101.04907 [astro-ph.HE].
- Xing, Yi, and Wenfei Yu. 2024. Detection of a gamma-ray flare in the direction of FRB20240114A. *The Astronomer's Telegram*, no. 16630.
- Xu, Jiaying, Yi Feng, Di Li, Pei Wang, Yongkun Zhang, Jintao Xie, Huaxi Chen, et al. 2023. Blinkverse: A Database of Fast Radio Bursts. *Universe* 9, no. 7 (July): 330. <https://doi.org/10.3390/universe9070330>. arXiv: 2308.00336 [astro-ph.HE].
- Yao, J. M., R. N. Manchester, and N. Wang. 2017. A New Electron-density Model for Estimation of Pulsar and FRB Distances. *ApJ* 835, no. 1 (January): 29. <https://doi.org/10.3847/1538-4357/835/1/29>. arXiv: 1610.09448 [astro-ph.GA].
- Zhang, Bing. 2022. The Physics of Fast Radio Bursts. *arXiv e-prints* (December): arXiv:2212.03972. <https://doi.org/10.48550/arXiv.2212.03972>. arXiv: 2212.03972 [astro-ph.HE].

Appendix 1. Pulses detected at $DM > 130 \text{ pc/cm}^3$.

A blind search using summed power spectra showed that at $DM > 130 \text{ pc cm}^{-3}$, the LPA3 radio telescope detects 10 ordinary pulsars with periods in the range of seconds (Tyul'bashev et al. 2024). The maximum value of DM was observed for the pulsar J1950+3001 ($P = 2.7889 \text{ s}$, $DM = 249 \text{ pc cm}^{-3}$). To search for pulsed radiation from these 10 pulsars, we selected 4 of the strongest (J1921+1948, J2029+3744, J2113+4644, and J1935+1616), the S/N ratios for harmonics in the summed power spectra of which ranged from 40 to 250^1 . When searching for pulsed radiation, dispersed pulses were found in 3 of the 4 pulsars (see Fig. 6). Pulses from J2113+4644 were found in a standard search for dispersed pulses from rotating radio transients (RRAT) having a value of $100 < DM < 200 \text{ pc cm}^{-3}$.

The profiles on the bottom panel were obtained with different processing of data. For pulsar J2113+4644 (B2111+46: $P = 1.0146 \text{ s}$, $DM = 141.2 \text{ pc cm}^{-3}$, $W_e = 149 \text{ ms}$), in the lower right corner, a simple summation of channels with DM compensation is carried out. On the upper (dynamic spectrum) and lower (profiles) parts of the figure, more than half of the pulses are visible, and the scattering of strong pulses is also clearly visible. For the pulsar J1921+1948 (B1918+19: $P = 0.8210 \text{ s}$, $DM = 153.8 \text{ pc cm}^{-3}$, $W_e = 175 \text{ ms}$) averaging by 4 points was carried out, which doubled the S/N and allowed us to see the pulse profiles. Almost all of its pulses are visible for this pulsar. The pulsar J2029+3744 (B2027+37: $P = 1.2168 \text{ s}$, $DM = 190.6 \text{ pc cm}^{-3}$) is shown in the middle part of Fig. 6. With the usual averaging of points, it is poorly visible. Based on its DM value, we got a model pulse (see details in Brylyakova and Tyul'bashev 2023) and carried out convolution in each frequency channel independently. Thus, the convolution of the detected pulse with the model pulse is shown in the dynamic spectrum of the pulsar for each frequency channel. To obtain the profile, we performed the summation of channels taking into account the known DM value of the pulsar. If for the pulsars J1921+1948 and J2113+4644 pulses were observed during hundreds of days, for the pulsar J2029+3744, only one pulse with a half-width $W_e = 418 \text{ ms}$ was found during 3,000 observation sessions. For the pulsar J1935+1616 (B1933+16: $P = 0.3587 \text{ s}$, $DM = 158.5 \text{ pc cm}^{-3}$), having S/N= 95 in the first harmonic in the summed up power spectrum, pulses with $S/N > 7$ were not detected. It is possible that the time of scattering in the direction to the pulsar is higher than its period and the pulses merge.

Thus, we detect the pulses from at least 3 known pulsars. Their values of $DM > 130 \text{ pc cm}^{-3}$ and exceed DM of the found FRB 20190203, which indicates a good opportunity to search for pulses from pulsars and FRBs on LPA3 with DM up to 200 pc cm^{-3} .

Appendix 2. Checking the pulse found in the direction to J1340+3530

Dispersion smoothing at $DM = 134.4 \text{ pc cm}^{-3}$ in a frequency channel with a width of 78 kHz is equal to 65 ms or 5–6

1. <https://bsa-analytics.prao.ru/en/pulsars/>




Figure 6. The upper panel shows dynamic spectra. The vertical axis is frequencies in MHz. The lower panel shows pulse profiles. The vertical axis represents the flux density in arbitrary units (a.u.). Time is shown on the horizontal axis. The time scale is the same for the upper and lower panels. The blue color is the summation of frequency channels without DM compensation, and the red color is the summation of channels with DM compensation.

points taking into account sampling equal to 12.5 ms. That is, in our observations on LPA3, the pulse having the specified DM cannot be narrower than 5 points. Our estimate for the scattering of the discovered FRB 20190203 shows that the pulse should be wider than $211/12.5 = 16 - 17$ points. This means that if we have found a real pulse, then when averaging raw data, its S/N should grow more than 2-4 times. This is not observed in Fig. 7. The dispersion delay line is also not visible on the dynamic spectrum. At $S/N > 7$, such lines are usually visible. Therefore, we believe that the found “repeated” pulse is false. We were not able to find errors in processing or a source of interference that gives large dispersion delays. At the same time, the found pulse with such a value of DM cannot have an extraterrestrial origin.

Appendix 3. Searching for short GRBs in the IBIS/ISGRI long-term observations

To search for possible short GRBs from the FRB 20190203 localization region, during a number of episodes when it was observed by the IBIS gamma-ray telescope, we developed a special technique based on the analysis of the IBIS/ISGRI detector’s count rate history. The technique reduces the computation time required for analysis and minimizes the likelihood of a false GRB detection. The following analysis of archived data of IBIS/ISGRI has been carried out:

- At first, the standard programs from the basic package for INTEGRAL data processing (OSA version 10.2) have been applied to analyze the related IBIS/ISGRI data up to the DEAD level;
- Then we have applied the “evts_extract” program of the OSA package to receive a list of events (registered photons) adjusted for the energy and “dead” time of the detector. The list does not include events from the ISGRI “hot” pixels. The Good Time Intervals (GTI) optimal for the analysis have been selected. For each photon from this

list, there was information about the light illumination of the corresponding detector pixel (in which the photon was registered) by this source of FRB (Pixel Illumination Factor or PIF);

- Then we have constructed a light curve (count rate history) of the source in the range of 20–100 keV using events from the list with $PIF < 0.1$ to renormalize it and use as background for the light curves with $PIF > 0.2$ and $PIF > 0.4$. The light curves with time bin size of 0.01 and 0.05 s have been obtained. This procedure allowed us to clear these light curves from any flashes of geophysical origin illuminating the entire detector;
- Each of these light curves has been divided into segments containing 5,000 bins, the average value and sample variance have been calculated for each segment, and all significant (over 10 standard deviations) enhancements in the count rate were revealed and collected;
- Then the whole procedure has been repeated for the light curves divided into the new segments with an offset of 1,667 bins (to ensure that all the enhancements found near the segment edges are valid). For each of the detected enhancements (candidates to bursts), a map of the detector has been obtained, using only the photons registered during the enhancements, and compared with the map obtained during the entire observation session. If the proportion of photons from pixels containing more than 2 photons exceeded 20%, such a burst was assumed “false”, since it was more likely associated with charged particles rather than X-ray photons;
- Images of the gamma-ray sky within the IBIS field of view have been constructed for all the remaining candidates to bursts, using a coding aperture of the telescope, and then it was compared with the images obtained during the previous time intervals of the same duration in order to confirm the burst confidence and identify the burst source. An original program developed at the Space Research Institute was used to construct images at such a short time




Figure 7. The upper panel shows the dynamic spectrum of the discovered pulse. Some frequency channels in MHz are marked along the vertical axis. The next 4 panels are the pulse profiles with averaging of data on 1, 2, 4, 8 points. The maximum S/N reaches when averaging by 2 points. Along the vertical axis is the flux density in arbitrary units (a.u.). Time in seconds is shown on the horizontal axis. The time scale is given in seconds and is the same for all panels. Blue and red colors show data before and after *DM* compensation.

scale, its general principles were described by Revnivtsev et al. (2004).