

Research Article

Cite this article: Barton DP, Martelli P, Luk W, Zhu X, Shamsi S (2020). Infection of *Hexametra angusticaecoides* Chabaud & Brygoo, 1960 (Nematoda: Ascarididae) in a population of captive crested geckoes, *Correlophus ciliatus* Guichenot (Reptilia: Diplodactylidae). *Parasitology* **147**, 673–680. <https://doi.org/10.1017/S0031182020000219>

Received: 7 September 2019
Revised: 2 February 2020
Accepted: 3 February 2020
First published online: 12 February 2020


Key words:

Ascaridoidea; captive; *Hexametra*; morphology; molecular sequencing; reptile

Authors for correspondence:

Diane P. Barton, Shokoofeh Shamsi,
E-mail: dibarton@csu.edu.au
and sshamsi@csu.edu.au

Infection of *Hexametra angusticaecoides* Chabaud & Brygoo, 1960 (Nematoda: Ascarididae) in a population of captive crested geckoes, *Correlophus ciliatus* Guichenot (Reptilia: Diplodactylidae)

Diane P. Barton¹, Paolo Martelli², William Luk², Xiaocheng Zhu^{1,3} and Shokoofeh Shamsi^{1,3} 

¹School of Animal & Veterinary Sciences, Charles Sturt University, Wagga Wagga, New South Wales, Australia; ²Ocean Park Corporation, 180 Wong Chuk Hang Road, Aberdeen, Hong Kong and ³Graham Centre for Agricultural Innovation (Charles Sturt University and NSW Department of Primary Industries), Charles Sturt University, Wagga Wagga, New South Wales, Australia

Abstract

Here we report on the infection of captive crested geckos *Correlophus ciliatus* Guichenot (Reptilia: Diplodactylidae), with adults of the ascaridoid nematode, *Hexametra angusticaecoides* Chabaud & Brygoo, 1960 (Ascarididae). A population of captive crested geckoes became ill and died within a short period of time. Nematodes were recovered from the crested geckoes examined from within the coelomic cavity, penetrating various organs and migrating through subcutaneous tissues, as well as emerging through the geckos' skin. One gecko was treated with levamisole following surgical excision of nematodes from under the skin; this gecko survived. The potential source of the nematode infection in the captive geckoes is discussed. It is most likely that wild-caught Madagascan mossy geckoes, *Uroplatus sikorae* Boettger (Reptilia: Gekkonidae), introduced the infection to the colony. Molecular sequences of the nematodes are the first produced for the members of this genus. A redescription of the species and its genetic characterization based on the internal transcribed spacer sequence data is provided, suggesting some of the morphological criteria that have been used in the past to distinguish between *Hexametra* spp. may have been intraspecific morphological variations.

Introduction

Ascaridoid nematodes are well-known parasites of wild and captive reptiles. Larval stages migrate through the host tissue, potentially causing damage to organs, whereas adults in the alimentary canal often burrow into the mucosal lining (Jacobson, 2007). Mortality of reptiles due to ascaridoid infections has been reported in various captive specimens (Jacobson, 2007).

Healthy reptiles can harbour a number of parasites with no apparent ill effect; however, if placed under stress, disease can become an issue in captive reptiles (Rataj *et al.*, 2011; Reese *et al.*, 2004). A compounding factor in parasite transmission can be the mixing of host species in a tank, either consecutively or concurrently, potentially leading to a naïve host species acquiring a novel parasite that may be harmful. This is especially true if the parasite has a lack of host specificity which can also lead to the potential for transmission among species in a collection (Reese *et al.*, 2004).

Despite crested geckoes, *Correlophus ciliatus* Guichenot (Diplodactylidae), being a common pet in many parts of the world, very few parasites have been recorded from them (Brusso, 2013), and there are no records of parasites from wild crested geckoes. Crested geckoes are native to just three islands in New Caledonia (Brusso, 2013), and since their introduction to the pet trade in 1994 (Brusso, 2013), crested geckoes are now exclusively traded from captive-bred populations.

In this paper, we determine the species of nematode present in the captive crested geckoes which were ill and subsequently died, and discuss the potential sources of the infection within the gecko population. A redescription and genetic characterisation are provided for the nematodes found in the present study.

Materials and methods**History of the captive population of crested geckoes**

In 2015, 10 crested geckoes were legally sourced from Canada (from a breeder of New Caledonian species of reptiles only; $n = 4$) and Germany (from a pet shop with a variety of reptile species; $n = 6$), and subsequently kept at the Ocean Park Aquarium in Hong Kong (no permit was required). The geckoes were randomly separated into two groups of one male



Fig. 1. *Correlophus ciliatus* infected with *Hexameta angusticaecoides*. (a) Deceased gecko with adult *H. angusticaecoides* penetrating through the ventral skin surface. (b and c) Gecko with nematodes under the skin. (d) Dissected gecko with larval and immature *H. angusticaecoides* within the coelomic cavity. (e) Migrating nematode in the mouth cavity. (f) Migrating nematode in the eye socket. (g) A moribund gecko with a *H. angusticaecoides* emerging from the tympanum.

and four females each, and each group was housed in a separate terrarium. In 2016, the crested geckoes were combined into a single terrarium and a group of wild-caught Madagascan mossy geckoes, *Uroplatus sikorae* Boettger (Gekkonidae), were placed into the other. The mossy geckoes all died within a few weeks of acquisition; examination by the owner (WL) did not show any obvious signs of parasitic infection and the cause of death remains unknown. The terrarium, and its furniture, was washed with water prior to the return of the crested geckoes. The second terrarium has only ever housed crested geckoes; however, crested geckoes have been moved between the two terraria over the remaining time period for breeding purposes. All geckoes were provided with water *ad lib* and commercially available dry foods specifically formulated for geckoes (Repashy™ and Pangea™). In addition, their diet was occasionally supplemented with shop-bought crickets and roaches bred in-house.

Collection of nematodes

In the year following their return to the terrarium, the five crested geckoes contained in the terrarium that had briefly housed the Madagascan mossy geckoes became moribund and died over a short period of time. The frozen carcasses of these crested geckoes were presented to the Ocean Park Conservation Foundation Veterinary Hospital (OPCFVH), Hong Kong, in July 2018. A post-mortem examination showed that the internal organs were mildly autolysed, preventing a sound histological examination. A number of large nematodes were recovered from the coelomic cavity, under the skin and penetrating organs (liver and alimentary canal) as well as emerging from the eye socket, tympanum and skin (see Fig. 1).

One of the geckoes from the second terrarium also showed signs of infection, with two nematodes observed moving under the skin (Fig. 1b and c). The nematodes from this gecko were surgically removed and the gecko was treated with fenbendazole (25 and 50 mg/kg), then with pyrantel (10 mg/kg); fecal examination remained positive and a further two nematodes appeared under the skin were also removed surgically. The gecko was then treated with levamisole per os (10 mg/kg) as a single dose. The gecko itself was initially negatively affected by the levamisole and was inappetent and listless for 10–15 days but has subsequently recovered and appears to be in full health at the time of writing. The remaining geckoes in the second terrarium have not shown any signs of infection or illness.

Table 1. GenBank accession numbers for ITS sequences used in this study

Taxa name	GenBank accession number	Reference
<i>Ascaris lumbricoides</i>	AB571297	Arizono et al. (2010)
<i>Ascaris suum</i>	AB571302	Arizono et al. (2010)
<i>Baylisascaris devosi</i>	MH030598	Hoberg et al. (2018)
<i>Parascaris equorum</i>	MH030605	Hoberg et al. (2018)
<i>Toxascaris leonine</i>	MH030606	Hoberg et al. (2018)
<i>Seuratascaris numidica</i>	MG434690	Chen et al. (2018)
<i>Porrocaecum reticulatum</i>	MF061688	Li et al. (2018)
<i>Raphidascaris longispicula</i>	KP326545	Zhao et al. (2016)
<i>Raphidascaris lophii</i>	MH211583	Zhang et al. (2018)
<i>Acanthocheilus rotundatus</i>	MF061679	Li et al. (2018)
<i>Pseudanisakis rajae</i>	JN392470	Li et al. (2012)
<i>Toxocara canis</i>	JF837169	Chen et al. (unpublished)
<i>Toxocara cati</i>	AB571303	Arizono et al. (2010)
<i>Toxocara vitulorum</i>	KT737382	Chelladurai et al. (2015)
<i>Heterakis isolonche</i>	KM212953	Gao (unpublished)
<i>Hexameta angusticaecoides</i>	MN876031–5	Present study

Over 50 nematodes were collected from the geckoes; all of which were collected externally to the alimentary canal. However, given the state of autolysis in the geckoes at the time of post-mortem, and the positive fecal examinations, the possibility of some of the nematodes having escaped from the alimentary canal cannot be excluded. All nematodes were preserved in 90% ethanol.

Morphological identification

A small piece of the mid body of each nematode was excised for molecular study and the rest of the nematode was cleared in lactophenol for morphological examination. The anterior and




Fig. 2. Line drawing of *Hexametra angusticaecoides* found in the present study: (a and b) lateral view of the labia in a female and a male, respectively, (c) cross-section below the nerve ring in a female, (d) lateral view of the anterior end showing the nerve ring, oesophagus and short intestinal caecum, (e) lateral view of the posterior end of the male showing a pair of spicules (left spicule was broken off) and a short mucron, (f) ventral view of the male tail showing the mucron, spicules, arrangement of the postcloacal papillae and partial view of the pre-anal papillae. Note the presence of the single median pre-cloacal papilla; (g) ventral view of the posterior end of the female tail being broad, short and simple.

posterior part of each nematode was examined under a light microscope. Illustrations were made using a microscope equipped with a camera lucida. Photos were taken using an AmScope MU900 eyepiece camera. All measurements are in micrometres, unless stated otherwise. Mean measurements are given, followed by the range in parentheses. Specimens have been deposited in the Australian Helminthological Collection, South Australian Museum, Adelaide, South Australia, under accession numbers AHC48786 and AHC48787.

One male and one female were subjected to scanning electron microscopy (SEM). They were washed and dehydrated overnight in a series of graded ethanol solutions (70, 80, 90, 95% and absolute ethanol). After three additional overnight washes in absolute ethanol, the specimens were critical point dried using a tousimis Autosamdri-931 (USA). Samples were then mounted on a 12 mm carbon tab (ProSci Tech, Thuringowa, QLD, Australia) and sputter coated with gold using a K550X Sputter Coater (Quorum Technologies, UK). The specimens were examined under a JEOL (Peabody, Massachusetts, USA) NeoScope SEM with accelerating voltage set at 10 kv.

DNA extraction, PCR and sequencing

Genomic DNA was isolated from all individual nematodes using Qiagen kit and eluted into 45 μ l of water. The internal transcribed spacer (ITS) region (including ITS1, 5.8 and ITS2) was amplified using the primer sets SS1: 5'-GTTTCCGTAGGTGAACCTGCG-3' (forward) and NC2: 5'-TTAGTTTCTTTTCTCCGCT-3' (reverse), and cycling conditions in accordance with Shamsi and Suthar

(2016). An aliquot (4 μ l) of each amplicon was examined on a 1.5% w/v agarose gel, stained with GelRed™ and photographed using a gel documentation system. PCR amplicons were sent to the Australian Genome Research Facility (Queensland) for sequencing.

ITS sequences of *Hexametra angusticaecoides* Chabaud & Brygoo, 1960 (Ascarididae) were generated in the current study, while ITS sequences from additional nematode species were obtained from GenBank (Table 1). Very few molecular sequences for ascarids from reptiles are available on GenBank, thus ITS sequences were obtained for other members of the Ascarididae (Table 1). Sequences were aligned with Geneious alignment algorithm by using Geneious version 11.1.4 (Kearse *et al.*, 2012), and then were double checked with all variable sites in the original trace files for confirmation. Alignments were then truncated to 865 characters, based on the shortest sequence for ITS. *Heterakis isolonche* von Linstow, 1906 (Heterakidae), a species belonging to the superfamily Heterakoidea but also residing under the infraorder Ascaridomorpha, was used as an outgroup. Phylogenetic relationship among species was calculated by MrBayes 3.2 (Ronquist and Huelsenbeck, 2003) with the HKY + I + G model as indicated by JmodelTest 2.0 (Darriba *et al.*, 2012).

Results

Clinical findings

The geckoes presented with visible subcutaneous parasites (Fig. 1) that would occasionally emerge through the skin. The geckoes




Fig. 3. Light micrographs of *Hexametra angusticaecoides*. (a) Dorsal view of the anterior end of the larva; arrow indicates the position of the excretory pore. (b) Ventral view of the anterior extremity showing subventral lips of an adult male specimen. (c) Intestinal caecum. (d) Lateral view of the posterior end of a male (note terminal mucron). (e) Lateral view of the posterior end of a female. (f) Distal tips of alate spicules. (g and h) Dissected uteri from an immature female. Only five uterine branches were present in one specimen (image h), whereas the diagnostic feature of six uterine branches was evident in the other specimen (image g; one branch is hidden in the photo).

would sometimes lose appetite, appear uncomfortable and then die or sometimes simply be found dead with no prior evidence of distress.

Morphological findings

A total of 13 nematodes were available for examination including one larva. Descriptions of adults are based on three males and nine immature females (Figs 2–4), unless otherwise stated. None of the females contained eggs. All specimens were identified as belonging to *H. angusticaecoides* Chabaud & Brygoo, 1960.

Description: Anterior extremity with three large labia, including larger dorsal labium and two smaller subventral labia; all labia

wider than long; two papillae on dorsal labium; subventral labia with one papilla and one amphid each; labia concave on their cranial side, with minute dentigerous border, uniform in size and confined to the cranial edge of labia (Fig. 4c); inter-labia and post-labial groove absent; excretory pore almost at the same level as nerve ring, slightly posterior to it; lateral alae present between nerve ring and anterior to the oesophagus–intestinal junction; oesophagus muscular, cylindrical, slightly wider at the oesophagus–intestinal junction; short intestinal caecum present; tail short with mucron in both sexes; mucron longer in younger nematodes, becoming shorter and almost invisible in older/larger specimens, particularly in females; tail short, very broad and rounded in females, and conical, bent ventrally and papillose in




Fig. 4. Scanning electron micrographs of a male and a female *Hexametra angusticaecoides*. (a) Ventral view of the anterior part of the male parasite showing labia, excretory pore (red arrow) and an unknown feature which has not been previously described by other authors (blue arrows); (b) Magnified view of the sub-ventral labium, note the absence of inter-labia, the presence of the dentigerous ridge on the cranial edge of the labia, the presence of papilla on the same level as the labial flange, and an amphid located slightly above the flange line in the opposite direction, blue arrow corresponds with the blue arrows in image (a), red and yellow arrows pointing at the labial papilla and amphid, respectively. (c) Magnified view of the image (b), note the dentigerous ridges do not exceed the cranial edge of the labium. Also note the lobular shape of the cranial side of the labia and the presence of a pore in the centre of each lobe. (d) Dorsal view of the dorsal labium. (e) Ventral view of the posterior end of an adult male. (f) Magnified view of the precloacal papillae. (g) Close up of the median ventral precloacal papilla located superomedially to the cloaca. (h) Tip of the tail in the male specimen showing mucron and postcloacal papillae. (i) Tip of the spicule. (j) Ventral view of the female posterior extremity: Phasmid (yellow arrow), mucron (green arrow) and a minute papilla (blue arrow). (k) Magnified view of image (j). (l) High magnification view of the phasmid. (m) High magnification view of a minute papilla-like structure in the tail of a female specimen.

males; spicules almost equal and alated; six pairs of post-cloacal papillae and up to 84 pre-cloacal papillae; uterus divided into 5–6 branches (two female nematodes were dissected for the purpose of counting uterus branches, one having five uterus branches and the other six uterus branches).

Morphometrics of the specimens were as below:

Males: Total body length 46.43 mm (39.00–60.03 mm); body width 1.21 mm (1.08–1.38 mm); dorsal labium 107 (95–125) high, 168 (150–190) wide; sub-ventral labia 110 (100–130) high, 112 (100–125) wide; nerve ring 610 (480–700) from the anterior end; deirids visible in one specimen only, 700 from the anterior end; excretory pore visible in one specimen, 875 from the anterior end; oesophagus 3133 (2350–3750) long, i.e. 7% (6–8%) of the body length; intestinal caecum 650 (600–700) long; tail 207 (200–220) long (excluding mucron), i.e. 0.5% (0.4–0.5%) of the body length; tail length with mucron 247 (240–250); tail width at the level of cloaca 267 (200–300); spicules 1000 (1000–1000) and 1083 (1050–1100) long, i.e. 2% (2–3%) and 2% (2–3%), respectively, of the body length; number of pre-cloacal papillae 70 and 84 ($n = 2$).

Females: Total body length 43.40 mm (22.75–69.50 mm); body width 1.08 mm (0.65–1.70 mm); dorsal labium 107 (50–150; $n = 8$) high, 164 (100–250; $n = 8$) wide; sub-ventral labia 92 (65–150; $n = 6$) wide, 90 (50–160; $n = 6$) high; nerve ring 543 (375–800; $n = 6$) from the anterior end; excretory pore 639 (450–950; $n = 7$) from the anterior end; oesophagus 2352 (1750–3700) long, i.e. 6% (3–9%) of the body length; intestinal caecum 461 (250–650; $n = 7$) long;

tail 278 (100–700) long (excluding mucron), i.e. 1% of the body length; tail length with mucron 325 (150–700; $n = 8$); tail width at the level of cloaca 422 (220–800); vulva 16.94 mm (16.13–17.75 mm; $n = 2$) from the anterior end, i.e. 43% (34–53%; $n = 2$) of the body length.

Fourth stage larva: Total body length 13.00 mm; body width 610; nerve ring 400 from the anterior end; excretory pore 450 from the anterior end; oesophagus 1650 long, 13% of the body length; intestinal caecum 250; tail 100 long (excluding mucron), 1% of the body length; tail length with mucron 170; tail width at the level of cloaca 200.

Molecular findings

Its sequences were identical among all the males, females and larva found in the present study, confirming that only one species was involved in the infection. We found no ITS sequences for any *Hexametra* spp. in GenBank. The newly generated ITS sequences of *H. angusticaecoides* were deposited in the GenBank database under accession numbers MN876031–5. Our sequences formed a distinct group from other terrestrial ascaridoids (Fig. 5).

Discussion

This is the first report of a species of the genus *Hexametra* Travassos, 1920 from the crested gecko. Despite several reports of *Hexametra* spp. in South East Asia (Clarke *et al.*, 1970; Sprent, 1978), this is the first report of *Hexametra* from Hong




Fig. 5. Phylogenetic analysis of ITS sequences for *Hexametra angusticaecoides* found in the present study, with *Heterakis isolonche* as an outgroup. Posterior probability values >90% are indicated above the branches. Sequences obtained in this study were denoted with * symbol. MrBayes best-fit model was used for analysis as suggested by JmodelTest. Branch supports lower than 85% are not shown on the Phylogenetic tree.

Kong. Given the fact that the geckoes were captive bred and housed in a terrarium that had previously housed a different species of gecko originally from Madagascar, where *H. angusticaecoides* was originally described from, it is unlikely that the crested gecko is a natural host, hence the migration of adult nematodes and the absence of gravid females in the present study. Infertility of female nematodes may have also been due to high level of mobility observed in adult nematodes, giving them no opportunity for mating and explaining the absence of the eggs in adult females despite the presence of adult males in one of the geckoes. The high level of mobility observed among these nematodes was also an important factor in the pathogenicity and fatality caused by these parasites. Since the crested geckoes were fed on a diet specifically formulated for crested geckoes, it is unlikely that malnutrition, which is known to make hosts more vulnerable to infection with parasites and to increase the severity of the clinical signs (Urquhart *et al.*, 1996), would have played a role in the infection patterns seen in the present geckoes.

Determining the original source of the nematode infection is difficult, due primarily to the multiple sources of host animals, multiple species of hosts in the terraria and the lack of knowledge about the life cycles of most nematodes of reptilian hosts. *Hexametra angusticaecoides* is widespread in a number of reptilian species around the world (Sprent, 1978), so it is possible that the crested geckoes obtained the infection from the original captive-bred colony or pet shop from which they were acquired. However, this possibility would tend to be refuted as only geckoes within the terraria that also housed the Madagascar mossy gecko became heavily infected. This leaves two potential sources of infection: the Madagascar mossy gecko or the crickets/roaches fed to the crested geckoes. Chabaud *et al.* (1962) conducted

a series of experimental infections with eggs and larvae of *H. angusticaecoides*, using wild-caught chameleons, *Chamaeleo pardalis* (Cuvier) (Chamaeleonidae), from Madagascar with an, unfortunately, unknown infection history. Due to the successful transmission of the parasite through various insects, tadpoles, mice and chameleons, Chabaud *et al.* (1962) concluded that the 'physiological possibilities of the parasite are very broad'. They also found that in one chameleon, subcutaneous and coelomic larvae co-existed with intestinal adults. We believe that it is most likely that the Madagascar mossy gecko is the true original source of infection, as many species of *Hexametra*, including *H. angusticaecoides*, have been originally described from reptiles from Madagascar (Sprent, 1978). The fact that the Madagascar mossy geckoes that were kept in the terraria were wild caught would also support this. These geckoes died rapidly, but with no obvious signs of parasite infection. It is possible that the stress of capture and housing, in hosts that had a subclinical infection, was enough to cause the death of the hosts (Rataj *et al.*, 2011).

Host mortality due to infection with ascaridoid nematodes has previously been reported (Smith, 1999). Despite the common occurrence of *Hexametra* in lizards and snakes, there are few reports of associated mortality (Jacobson, 2007). An infection of *Hexametra boddaertii* (Baird, 1860) was found in a wild-caught snake in Argentina; the snake died after a few days despite being in apparent good health (Peichoto *et al.*, 2016). Over 100 nematodes were collected from the coelomic cavity (smaller worms) and the intestinal tract (larger worms); initial fecal examination had shown a heavy infection with ascaridoids but attempts to de-worm the snake were unsuccessful (Peichoto *et al.*, 2016). Harmful effects of ascaridoid infections are difficult to assess due to the paucity of knowledge of symptomatology in

Table 2. Measurements of male specimens of *Hexametra angusticaecoides* collected from the crested gecko, *Correlophus ciliatus*, in the present study and comparative measurements obtained from the literature

	<i>H. angusticaecoides</i>	<i>H. angusticaecoides</i>	<i>H. angusticaecoides</i>
Reference	Present study	Chabaud and Brygoo (1960)	Sprent (1978)
Host	<i>Correlophus ciliatus</i>	<i>Chamaeleo oustaleti</i>	<i>Chamaeleo</i> spp.
Body length (mm)	39–60	48	54–95
Width (maximum)	1080–1380	700	830–1500
Subventral lip length	130	190	100–140
Oesophagus length (mm)	2.35–3.75	3.2	3.5–3.9
Intestinal caecum length	600–700	400	–
Spicule length	1000–1100	1050	820–1100
Nerve ring from anterior extremity	480–700	800	490–800
Excretory pore from anterior extremity	880	920	610–950
Ejaculatory duct (mm) length	–	–	1.5–2.2
Tail length	200–220	230	210–240
Cloacal papilla pattern	70–84 pairs precloacal; single pair at level of cloaca; 2 pairs subventral and 1 or 2 pairs subdorsal postcloacal papillae; one single mid-ventral papilla located immediately anterior to cloaca	57 pairs + 1 odd precloacal; double pair at level of cloaca; 4 postcloacal	40–60 pairs precloacal; double pair at level of cloaca; 2 pairs subventral and 1 or 2 pairs subdorsal postcloacal

All measurements in micrometres unless otherwise indicated.

reptiles. Harmful effects could be due to the size of the nematode in relation to the host (competing for nutrients and causing malnutrition) and/or causing physical damage through their attachment to the mucosal lining of the stomach and/or physical damage during larval migration (Jacobson, 2007; Sprent, 1984). Clinical signs, when present, are non-specific; infected reptiles may be anorectic and may slowly lose weight (Jacobson, 2007). A captive panther chameleon, *Furcifer pardalis* (Cuvier) (Chamaeleonidae), that died suddenly without signs of illness harboured *H. angusticaecoides* in its stomach and intestine; a veiled chameleon, *Chamaeleo calyptratus* Duméril & Bibron, was euthanized after manifesting respiratory distress and had 20–30 immature *Hexametra* specimens in its coelomic cavity (Jacobson, 2007).

The fact that the terraria used in this study were not disinfected between housing of the two gecko species and the crested geckoes were returned to the terraria within a matter of days could have ensured that any nematode eggs in the terraria were able to survive and infect the new hosts. Chabaud *et al.* (1962) found that the eggs of *H. angusticaecoides* took approximately 15 days to become infective, with second stage larvae spontaneously hatching from the eggs. Additionally, the use of roaches and crickets as food items could have enabled the cycle to continue (see Chabaud *et al.*, 1962). The initial infection potentially occurred in 2016, so it would appear that the nematode is able to complete its life cycle within a captive situation, making it a potential problem in a captive colony if it became infected.

Morphological examination identified the nematodes as belonging to the genus *Hexametra* based on the keys of Hartwich (1974): female with six uterine branches, inter-labia absent and parasites of snakes and lizards. Sprent (1978) stated that an intestinal caecum can be present or absent; it was present in the current specimens. Measurements of specimens and the

presence of an intestinal caecum showed the specimens were *H. angusticaecoides*, based on the data provided previously (Chabaud and Brygoo, 1960; Sprent, 1978) (Table 2). Although most of the measurements of the specimens in the present study fall within the range of measurements provided for *H. angusticaecoides* in previous studies (Table 2), the number of precloacal papillae and the single vs a double pair of papillae at the level of the cloaca were different and might cast doubt on the specific identification of species of *Hexametra* in general. *Hexametra quadricornis* (Wedl 1862) from snakes has 50–100 precloacal papillae (Petter, 1968), but Sprent (1978) believed that this parasite was unable to develop in lizards. While in experimental infections in snakes the larvae of *H. quadricornis* did develop into adults in the body cavity, they never reached the intestine and the adult females never possessed eggs (Sprent, 1978), which is similar to the case in our study.

Additionally, during the examination of the morphology of specimens, one female was found to only possess five uterine branches, with another specimen having six (Fig. 3). In addition, despite the overall similarity in measurements, the pattern of cloacal papillae in the male was highly variable (see Table 2). This suggests that there may be a level of morphological variation in the genus that should be considered in future descriptions. This has important implications for the taxonomy of the genus and the closely related *Polydelphis* Dujardin, 1845 (Ascarididae), which are separated by the number of uterine branches (six in *Hexametra* compared to four in *Polydelphis*; Sprent, 1978). Future researchers should be aware of the potential of morphological variation and ensure genetic sequences are included to enable better systematic differentiation of this group of nematodes, especially in relation to potential issues in captive hosts.

The molecular analysis for the specimens of *H. angusticaecoides* displayed no nucleotide variability in the ITS sequences,

indicating that there was only the one species of nematode infecting the geckoes as both larval and adult stages. A comparison of ITS sequences with other species of Ascarididae from terrestrial hosts available in GenBank showed 41.2% nucleotide divergence to the closest representative, *Toxoascaris leonine* (Linstow, 1902). It must be noted, however, that there are very few full ITS sequences available for ascarid nematodes of terrestrial animals, making direct comparisons between closely related genera difficult. However, the sequences obtained here are extremely important and valuable for future research and diagnosis of infections with *H. angusticaecoides*. Specimens of *H. angusticaecoides* from the present study formed a sister group to the clade composed of other terrestrial ascaridoids, including *T. leonina*, *Baylisascaris devosi*, *Parascaris equorum*, *Ascaris lumbricoides* and *Ascaris suum* (Goeze, 1782), with the latter two taxa being considered conspecific by some authors (e.g. Leles et al., 2012; Liu et al., 2012).

In conclusion, our study reinforces the importance of application of transmission-based precautions to prevent animal and human infection with potentially deadly ascaridoid parasites. In terms of the specific identity of the nematodes, although we considered our specimens as *H. angusticaecoides*, more specimens from other members of the genus should be examined both morphologically and by molecular tools to be able to more accurately comment on the taxonomic status of these taxa.

Financial support. This study was partially funded by Charles Sturt University A516 828–66770.

Conflict of interest. None.

Ethical standards. Ethical standards were maintained during the study.

References

- Arizono N, Yoshimura Y, Tohzaka N, Yamada M, Tegoshi T, Onishi K, and Uchikawa R (2010) Ascariasis in Japan is pig-derived ascaris infecting humans? *Japanese Journal of Infectious Diseases* **63**, 447–448.
- Brusso K (2013) *Rhacodactylus ciliatus* (On-line), Animal Diversity Web. Available at https://animaldiversity.org/accounts/Rhacodactylus_ciliatus/ (Accessed 16 March 2019).
- Chabaud AG and Brygoo ER (1960) Nématodes parasites de caméléons malgaches. *Memoires de l'Institut scientifique de Madagascar. Serie A* **14**, 125–159.
- Chabaud AG, Brygoo ER and Petter AJ (1962) Parasitologie – demonstration experimentale de differents types de cycles evolutifs potentiels chez un ascaride de cameleon. *Comptes Rendus Hebdomadaires Des Seances De L Academie Des Sciences* **255**, 2320.
- Chelladurai JJ, Bader C, Snobl T, Magstadt D, Cooper V and Brewer MT (2015) *Toxocara Vitulorum* Infection in a cohort of beef calves in Iowa. *Vet Parasitol* **214**, 96–99.
- Chen H-X, Zhang K, Zhang L-P and Li L (2018) Morphological and molecular characterization of *Seuratascaris numidica* (Seurat, 1917) (Ascaridida: Ascarididae). *Acta Parasitologica* **63**, 154–159.
- Clarke M, Chang T and Cross Jr J (1970) An ascaridoid nematode from the bamboo viper, *Trimeresurus Stejnegeri* from Taiwan, Republic of China. *SEast Asian J trap Med publ Hlth* **1**, 565–567.
- Darriba D, Taboada GL, Doallo R and Posada D (2012) Jmodeltest 2: more models, new heuristics and parallel computing. *Nature Methods* **9**, 772–772.
- Hartwells G (1974) Keys to genera of the Ascaridoidea. In Anderson RC, Chabaud AG and Willmott S (eds), *CIH Keys to the Nematode Parasites of Vertebrates*, vol. 2. CAB International, UK: Commonwealth Agricultural Bureaux, pp. 1–15.
- Hoberg EP, Burek-Huntington K, Beckmen K, Camp LE and Nadler SA (2018) Transuterine infection by *Baylisascaris Transfuga*: neurological migration and fatal debilitation in sibling moose calves (*Alces alces gigas*) from Alaska. *International Journal for Parasitology-Parasites and Wildlife* **7**, 280–288.
- Jacobson E (2007) *Infectious Diseases and Pathology of Reptiles*. Boca Raton: CRC Press and Taylor & Francis Group, pp. 716.
- Kearse M, et al. (2012) Geneious basic: an integrated and extendable desktop software platform for the organization and analysis of sequence data. *Bioinformatics (Oxford, England)* **28**, 1647–1649.
- Leles D, Gardner SL, Reinhard K, Iniguez A and Araujo A (2012) Are *Ascaris lumbricoides* and *Ascaris Suum* A single species? *Parasites & Vectors* **5**. doi: 10.1186/1756-3305-5-42
- Li L, Gibson DI, Liu YY and Zhang LP (2012) Morphological and molecular study of the poorly known species *Pseudanisakis Rajae* (Yamaguti, 1941) (Nematoda: Acanthocheilidae) from elasmobranchs in the Yellow Sea and Taiwan Strait off the coast of China. *Systematic Parasitology* **81**, 115–123.
- Li L, Lu L, Nadler SA, Gibson D, Zhang L, Chen H-X, Zhao WT and Guo Y-N (2018) Molecular phylogeny and dating reveal a terrestrial origin in the early carboniferous for ascaridoid nematodes. *Systematic Biology* **67**, 888–900.
- Liu G-H, Wu C-Y, Song H-Q, Wei S-J, Xu M-J, Lin R-Q, Zhao G-H, Huang S-Y and Zhu X-Q (2012) Comparative analyses of the complete mitochondrial genomes of *Ascaris lumbricoides* and *Ascaris Suum* from humans and pigs. *Gene* **492**, 110–116.
- Peichoto ME, Sanchez MN, Lopez A, Salas M, Rivero MR, Teibler P, Toledo GdM and Tavares FL (2016) First report of parasitism by *Hexametra Boddaertii* (Nematoda: Ascaridae) in *Oxyrhopus Guibei* (Serpentes: Colubridae). *Veterinary Parasitology* **224**, 60–64.
- Petter AJ (1968) Observations on the classification and cycle of the ascarid *Hexametra quadricornis* (Wedl 1862). *Annales de Parasitologie Humaine et Comparee* **43**, 655–691.
- Rataj AV, Lindtner-Knific R, Vlahović K, Mavri U and Dovč A (2011) Parasites in pet reptiles. *Acta Veterinaria Scandinavica* **53**, 33.
- Reese DJ, Kinsella J, Zdziarski JM, Zeng Q-Y and Greiner EC (2004) Parasites in 30 captive tokay geckos, *Gekko gekko*. *Journal of Herpetological Medicine and Surgery* **14**, 21–25.
- Ronquist F and Huelsenbeck J (2003) MrBayes 3: Bayesian phylogenetic inference under mixed models. *Bioinformatics (Oxford, England)* **19**, 1572–1574.
- Shamsi S and Suthar J (2016) Occurrence of *Terranova* Larval types (Nematoda: Anisakidae) in Australian marine fish with comments on their specific identities. *PeerJ* **4**, e1722–e1722.
- Smith JW (1999) Ascaridoid nematodes and pathology of the alimentary tract and its associated organs in vertebrates, including man: a literature review. *Helminthological Abstracts* **68**, 49–96.
- Sprent JFA (1978) Ascaridoid nematodes of amphibians and reptiles – *Polydelphis*, *Travassosascaris* N.g. and *Hexametra*. *Journal of Helminthology* **52**, 355.
- Sprent J (1984) Ascaridoid nematodes. In Hoff GL, Frye FL and Jacobsen ER (eds), *Diseases of Amphibians and Reptiles*. New York: Plenum Press, pp. 219–245.
- Urquhart GM, Armour J, Duncan JL, Dunn AM and Jennings FW (1996) *Veterinary Parasitology*. Harlow, UK: Longman Group.
- Zhang K, Xu Z, Chen H-X, Guo N and Li L (2018) Anisakid and raphidascaridid nematodes (Ascaridoidea) infection in the important marine food-fish *Lophius Litulon* (Jordan) (Lophiiformes: Lophiidae). *International Journal of Food Microbiology* **284**, 105–111.
- Zhao W-T, Lu L, Chen H-X, Yang Y, Zhang L-P and Li L (2016) Ascaridoid parasites infecting in the frequently consumed marine fishes in the coastal area of China: a preliminary investigation. *Parasitology International* **65**, 87–98.