

Children's stress regulation mediates the association between prenatal maternal mood and child executive functions for boys, but not girls

REGULA NEUENSCHWANDER, KAIA HOOKENSON, URSULA BRAIN, RUTH E. GRUNAU, ANGELA M. DEVLIN, JOANNE WEINBERG, ADELE DIAMOND, AND TIM F. OBERLANDER
University of British Columbia, BC Children's Hospital Research Institute

Abstract

Prenatal exposure to maternal mood disturbances shapes children's cognitive development reflected in the critical construct of executive functions (EFs). Little is known, however, about underlying mechanisms. By examining cortisol responses in both everyday and lab challenge settings, we tested whether the child/offspring hypothalamic–pituitary–adrenal axis mediates effects of prenatal maternal mood on child EFs at age 6. In 107 Canadian children born to women with a wide range of anxious and depressive symptoms during pregnancy, we found that in boys but not girls, depressed and/or anxious prenatal maternal mood is associated with heightened diurnal cortisol levels in everyday settings, as well as heightened cortisol reactivity to a lab challenge and that this heightened reactivity was associated with poorer EFs. Among boys we also observed that cortisol reactivity but not diurnal cortisol mediated the association between depressed and/or anxious prenatal maternal mood and EFs. Depressed and/or anxious prenatal maternal mood was related to child EFs for both girls and boys. To our knowledge, this is the first study to demonstrate a mediating role for child stress regulation in the association between prenatal maternal stress-related mood disturbances and child EFs, providing evidence of a mechanism contributing to fetal programming of cognition.

Stress early in life dramatically affects the prefrontal cortex (PFC; e.g., Demir-Lira, Prado, & Booth, 2016; Noble, McCandliss, & Farah, 2007) although less is known about the ways the specific stress related to depressed or anxious prenatal maternal mood affects PFC structure and/or subsequent function during childhood (Neuenschwander & Oberlander, 2017). Given that the PFC has the highest density of glucocorticoid receptors in the primate brain and PFC glucocorticoid receptors are important in regulating PFC dopamine levels (Butts, Weinberg, Young, & Phillips, 2011; Sanchez, Young, Plotsky, & Insel, 2000), the child hypothalamic–pituitary–adrenal (HPA) axis may play a significant role in mediating effects of depressed and anxious prenatal maternal mood on child executive functions (EFs) that rely on the PFC.

In the animal literature, it is well established that the maternal and offspring HPA axis plays a crucial role in mediating prenatal stress effects on various outcomes in the offspring (Weinstock, 2008). Whereas in human studies it is

less clear if the maternal HPA axis is responsible for prenatal maternal stress effects on child outcomes (Beijers, Buitelaar, & de Weerth, 2014; Zijlmans, Riksen-Walraven, & de Weerth, 2015), there is certainly a dearth of human research examining whether altered child/offspring HPA axis activity mediates the association between prenatal maternal stress or prenatal maternal mood disturbances and child outcomes (Glover, 2015; Glover, O'Connor, & O'Donnell, 2010). The present study was undertaken in a prospective longitudinal cohort of now 6-year-olds, to examine whether child HPA axis activity under two conditions (diurnal cortisol levels across four typical days and cortisol reactivity in response to a lab challenge stress) contributes to the association between prenatal maternal stress-related mood disturbances (reflected in measures of depressed and anxious maternal mood) and child EFs (assessed with a traditional computerized task; see Figure 1).

Prenatal Maternal Mood and Fetal Programming

Pregnancy is a dramatic biological and psychological period in a woman's life that can affect health and behavior across two generations (Dunkel Schetter & Tanner, 2012; Matthews & Phillips, 2011). It is therefore not surprising that pregnancy and the postpartum period tend to heighten risk for the development or recurrence of maternal mood disorders (Leight, Fitelson, Weston, & Wisner, 2010). Prenatal maternal depres-

Support for this research was provided by a CIHR Grant MOP-89916 (to T.F.O.) and a postdoctoral fellowship from Brain Canada/NeuroDevNet (to R.N.). A. M. D. is supported by an Investigation Grant from BC Children's Hospital research Institute, University of British Columbia. We would like to thank our research assistants for data collection and all children and mothers for their participation.

Address correspondence and reprint requests to: Regula Neuenschwander, Fabrikstrasse 8 (Büro A 349), CH-3012 Bern, Switzerland; E-mail: regula.neuenschwander@psy.unibe.ch




Figure 1. Theoretical framework: Mediation model of prenatal stress exposure, hypothalamic–pituitary–adrenal (HPA) axis activity, and executive functions.

sion occurs in 10%–20% of pregnancies (Bennett, Einarson, Taddio, Koren, & Einarson, 2004; Marcus, Flynn, Blow, & Barry, 2003) and about 5% of women take a selective serotonin reuptake inhibitor (SSRI) antidepressant during pregnancy (Oberlander, Warburton, Misri, Aghajanian, & Hertzman, 2006). Furthermore, 18%–24% of women suffer from self-reported anxiety symptoms (Dennis, Falah-Hassani, & Shiri, 2017). Prenatal maternal mood disturbances have been characterized on multiple dimensions, including stress-related disorders such as anxiety and/or depression (Dunkel Schetter, 2011). For the current study, we were interested in symptom severity of such mood disturbances and not in diagnostics or the specific phenotype domain depression versus anxiety. The reason being that programming effects may not be specifically associated with a particular clinical diagnosis, but the natural range of adverse conditions or life events, which have the potential to make the mother feel generally stressed (Glover, 2011).

A well-established research literature has found that even before birth, a mother's depressed and/or anxious mood shapes her child's subsequent development of self-regulation, of which, EFs are its cognitive hallmark (e.g., Glover, 2011; Mennes, Stiers, Lagae, & van den Bergh, 2006; van den Bergh, Mulder, Mennes, & Glover, 2005). The premise that fetal experiences set pathways for health and well-being across the life span is characterized as *fetal programming* (cf. developmental origins of health and disease; Barker, 2003). This implies that fetal development is altered in a way that prepares the offspring for a particular environment it might expect to find after birth (cf. predictive adaptive response; Gluckman & Hanson, 2005). Stress regulatory systems play a pivotal role in the organism's adaptation to the demands of the external and internal environment (cf. allostasis; McEwen & Wingfield, 2003), and individual differences in

stress reactivity have been theorized to be evolutionarily selected adaptations that enable the developing organism to match its phenotype to different environmental conditions (Hostinar & Gunnar, 2013). Therefore, stress regulatory systems and the HPA axis in the offspring, in particular, may be key mechanisms underlying fetal programming and thus mediating the effects of maternal stress on child outcomes (Glover et al., 2010), although other neurocircuits, such as the dopaminergic and serotonergic systems, are likely to be involved as well (Talge, Neal, & Glover, 2007).

A series of developmental studies in animals, both rodent and nonhuman primate, established the central role of the HPA axis in mediating prenatal stress effects in both mother and offspring (for a review see Weinstock, 2008). In humans, however, empirical evidence examining *offspring* HPA activity as a mediator is still missing (for an exception see, van den Bergh, van Calster, Smits, van Huffel, & Lagae, 2008). The *maternal* HPA axis, in contrast, has been the most widely investigated biological mechanism of transmission of risk, with mixed results. A recent systematic review (Zijlmans et al., 2015), for instance, found nonsignificant associations between prenatal maternal cortisol concentrations and various child outcomes (e.g., health, cognitive, motor, behavioral, and cortisol) in as many as 76% of studies reviewed. The authors argue that apart from methodological issues (e.g., variation in statistical procedures, variation in assessment of maternal cortisol, and failure to account for confounders and moderators), additional mechanisms that may act in conjunction with, or as moderators or mediators of, the effects produced by the HPA axis should be further explored (e.g., placental functioning of the 11 β -HSD2 enzyme and altered health behaviors including eating, sleep, and exercise; Beijers et al., 2014). Nevertheless, the fact that maternal mood disturbances (anxiety, depression, and perceived stress) during

pregnancy are linked to later child outcomes, even after controlling for effects of postnatal maternal mood and other relevant prenatal and postnatal confounders, suggests that, as in animal models, some of the risk is conferred prenatally via changes in women's mood-based physiology affecting fetal neurobehavioral development.

Executive Functioning and Stress

A key assumption underlying the fetal programming hypothesis is that biological systems undergoing rapid developmental changes are especially vulnerable to organizing and disorganizing influences (Seckl & Meaney, 1993). Stress early in life, and specifically prenatal maternal stress, may have a particularly large effect on PFC structure and function because of rapid brain development during pregnancy and the high density of glucocorticoid receptors in the PFC region (Arnsten, 2009; Fuster, 2008; Sanchez et al., 2000), which are important in regulating dopamine levels (Butts et al., 2011; Sanchez et al., 2000). In particular, EFs (a set of higher order cognitive processes, such as working memory, inhibition, and cognitive flexibility, associated with PFC and integral to emerging self-regulatory behavior; Blair & Diamond, 2008; Diamond, 2013) are the first to suffer, and suffer disproportionately, under stress (Arnsten, 2009; Diamond & Ling, 2016).

In adults, acute psychosocial stress impairs EFs (e.g., Alexander, Hillier, Smith, Tivarus, & Beversdorf, 2007; Lupien, Gillin, & Hauger, 1999). Stress early in life also appears to affect PFC function (e.g., Demir-Lira et al., 2016; Noble et al., 2007). One study found that growing up under social or economic disadvantage increased young toddlers' cortisol levels, which in turn mediated the effects of poverty and parenting on EFs at the age of 3 years (Blair et al., 2011). When stressful conditions are chronic or persistent, stress response systems are under high allostatic load and adapt to the environment with over- or underactivation to an extent that impedes flexible regulation of stress physiology (McEwen & Wingfield, 2003), which underlies efficient self-regulation and optimal functioning of EFs (Ramos & Arnsten, 2007). To some extent, these effects are believed to reflect the fact that glucocorticoid levels (i.e., cortisol) modulate synaptic activity in the neural circuitry of the PFC.

The functional relationship between cortisol levels and PFC activity or EF performance has been long recognized to be curvilinear (Arnsten, 2009; de Kloet, Oitzl, & Joels, 1999; Lupien, Maheu, Tu, Fiocco, & Schramek, 2007), such that very high or very low levels of stress impair EF performance whereas moderate stress levels lead to optimal EF performance (Blair & Ursache, 2011; Cools & D'Esposito, 2011; Vijayraghavan, Wang, Birnbaum, Williams, & Arnsten, 2007). There is some evidence, however, that this may only be true for males but not females. Diamond (2011) summarized differential effects of mild stress on male versus female animals, indicating that male animals perform better on tasks dependent on the PFC when they are mildly stressed

as compared to when they are unstressed, but female animals, in contrast, perform more poorly when mildly stressed as compared to when they are unstressed.

This inverted U-shaped relationship may have important implications for beneficial effects of prenatal exposure to mild and moderate levels of maternal stress on certain child outcomes. Evidence of a nonlinear relationship between prenatal stress exposure and child outcomes is supported by DiPietro, Novak, Costigan, Atella, and Reusing (2006), who found that exposure to moderate levels of prenatal stress was associated with advanced motor and mental development in the offspring of healthy mothers with low-risk pregnancies.

Prenatal Maternal Mood Shaping Child Executive Functions

Recently, human studies have begun to examine the neurodevelopmental consequences of exposure to maternal stress during gestation. Laboratory-based measures of child neurocognitive development (behavioral measures of EFs and neurophysiological measures indexing PFC structure and activity) offer critical insights into the neural correlates (i.e., specific aspects of children's EFs including underlying structure–function relations) that may be affected by prenatal maternal stress.

Longitudinal studies indicate that prenatal stress is associated with each key component of EFs: inhibition (Buss, Davis, Hobel, & Sandman, 2011; van den Bergh, Mennes, et al., 2005; van den Bergh et al., 2006), cognitive flexibility (i.e., shifting; Mennes et al., 2006), and working memory (Buss et al., 2011; Entringer, Buss, et al., 2009; Pearson et al., 2016), as well as with reductions in PFC volume (Buss, Davis, Muftuler, Head, & Sandman, 2010) and cortical thinning in the right frontal lobes (Sandman, Buss, Head, & Davis, 2015). These findings, however, lack robustness and consistent associations between prenatal stress and EFs given that in several studies multiple tasks tapping various EF components were administered, but associations with prenatal maternal stress were only found for some of the EF tasks. For instance, some authors report impaired performance on working memory tasks (Buss et al., 2011; Pearson et al., 2016), others do not (Mennes et al., 2006; van den Bergh, Mennes, et al., 2005), and still others only find a difference between prenatally stress-exposed individuals and nonexposed individuals after hydrocortisone administration (Entringer, Buss, et al., 2009).

While it is unclear which specific aspects of children's EFs and underlying PFC structures and functions are most strongly or consistently altered following exposure to prenatal maternal stress, emerging evidence suggests that prenatal maternal stress is associated with subtle changes in EFs and the PFC in middle childhood, adolescence, and early adulthood. We build on this literature by examining whether prenatal maternal mood predicts child EFs (assessed with a task tapping all three EFs components) at 6 years. We also examined whether the functional relationship between early exposure

to stress-related mood disturbances and EF performance is curvilinear (cf. Blair & Ursache, 2011; Cools & D'Esposito, 2011; Vijayraghavan et al., 2007) such that moderate levels of stress exposure may actually enhance EFs performance.

Some of the studies reviewed above reported sex specific findings. Buss et al. (2011) found an association between maternal pregnancy-specific anxiety and inhibitory control in girls but not boys, whereas van den Bergh et al. (2006) detected impaired endogenous response inhibition in adolescent boys but not in girls. Maternal stress may thus induce sex-specific changes in the HPA axis during critical gestation periods, but to date, few human studies have addressed sex differences in child outcomes following prenatal stress exposure, and of those that have, results are mixed. Therefore, we examine whether sex moderates the effects of prenatal stress on the activity of the HPA axis in relation to EFs. Finally, indirect evidence that prenatal maternal stress affects brain development in a way that may also affect the regulation of the HPA axis in offspring (Buss et al., 2010; Entringer, Buss, et al., 2009) provides an important direction for understanding potential mechanisms.

Prenatal Maternal Mood Shaping Child Stress Regulation

Evaluating how children's stress regulation is affected by prenatal stress exposure should help elucidate our understanding of potential pathways through which the development of EFs may be affected by prenatal maternal mood disturbances. The stress system comprises two main components: the HPA axis with its end product cortisol, and the sympathetic adrenal medullary system (which is part of the autonomic nervous system) with end products epinephrine and norepinephrine. The acute secretion of glucocorticoids (corticosterone in most animals and cortisol in humans) and catecholamines (epinephrine and norepinephrine, also known as adrenaline and noradrenaline) constitute the primary agents in the chain of hormonal events triggered in response to stress. These neurochemicals act together: the catecholamines give rise to the fast "fight-or-flight response" reflected in increased heart rate and blood pressure, while the HPA axis gives rise to a slower response that mobilizes metabolic pathways; affects skeletal muscles, vascular reactivity, nervous system activity, and the immune function; and facilitates physiological and behavioral coping mechanisms. In this way, stress responses serve an adaptive survival mechanism consisting of a carefully orchestrated yet near-instantaneous sequence of hormonal changes and physiological responses enabling an individual to react quickly to threat.

However, frequent or chronic activation can result in a persisting dysregulation of the HPA axis, particularly when experienced during phases of rapid brain development such as the prenatal period, including the fetal period *ex utero* (in pre-term neonates) and during infancy (Grunau et al., 2007; Gunnar & Quevedo, 2007). Chronic stress exposure, resulting in chronic cortisol elevations, will eventually result in downregulation of

glucocorticoid receptors, and alterations in the finely tuned balance among the nervous, endocrine, and immune systems (Chrousos, 2009; Frodl & O'Keane, 2013). This can ultimately increase neuroinflammation, which itself may underlie both behavioral changes and the development of diseases or disorders later in life, including depression/anxiety, cardiovascular disease, diabetes, and autoimmune disorders (Capuron & Miller, 2011; Frodl & O'Keane, 2013; Miller & Raison, 2016). Thus, chronic stress exposure has long-term effects on physical and psychological health such as high blood pressure, increased risk of infection, arterial disease, and brain changes that may contribute to anxiety, depression, and addiction (for a general review, see Felitti & Anda, 2010; Felitti et al., 1998; McEwen, 2000).

Hyperactivation is in general suggested to be indicative of a currently stressed HPA axis (e.g., McEwen & Wingfield, 2003), whereas hypoactivation reflects reduced cortisol production, possibly due to more chronic stress that has downregulated glucocorticoid receptors and resulted in HPA dysregulation (e.g., Doom, Cicchetti, & Rogosch, 2014; Grunau et al., 2013). Moreover, exposure to chronic stress in the early years of life, when the nervous system is still developing, may result in a distinct pattern of dysregulation.

In rodents, studies have found that prenatal stress causes both an increase in basal levels and an increase in corticosterone responses to stress in the offspring, with factors such as sex, age of the offspring, nature and timing of stressor during pregnancy, and so on, playing a modulatory role (for a review, see Weinstock, 2008). In humans, there appear to be similar fetal programming effects. In Glover et al.'s (2010) review of the literature 11 studies in the last 10 years revealed an association between prenatal maternal mood or stress and some aspect of HPA axis function in the child. However, perhaps unsurprisingly, the nature of this association varied and solid replications seem to be missing. Overall, these and some additional studies indicate that prenatal stress is linked to both raised diurnal cortisol levels and increased stress responsivity in infancy (Brennan et al., 2008; Davis, Glynn, Waffarn, & Sandman, 2011; Diego et al., 2004; Grant et al., 2009; Tollenaar, Beijers, Jansen, Riksen-Walraven, & de Weerth, 2011), and early to middle childhood (Gutteling, de Weerth, & Buitelaar, 2004, 2005; O'Connor et al., 2005; Simons, Beijers, Cillessen, & de Weerth, 2015). Later in development, some studies have found either a reduced cortisol awakening response (CAR) and blunted diurnal cortisol decline in adolescents (O'Donnell et al., 2013; van den Bergh et al., 2008, however, see Vänskä et al., 2015, for an intensified CAR but nonaffected diurnal cortisol decline in 10- to 12-year-olds) or no differences in diurnal patterns but increased cortisol reactivity during the Trier Social Stress Test among young adults exposed to early stress versus an age-matched comparison group (Entringer, Kumsta, Hellhammer, Wadhwa, & Wust, 2009).

These findings provide evidence that early stress exposure is associated with elevated or hyperactivation of the HPA axis (diurnal profiles and reactivity patterns) that could, over time,

lead to adrenocortical counterregulation and hypoactivation (Miller, Chen, & Zhou, 2007). More research, however, is needed to confirm this longitudinal pattern of the HPA axis following prenatal stress exposure, where glucocorticoid receptor downregulation leads to long-term dampened stress responses.

Whether alterations in HPA axis function mediate the association between early stress exposure and altered behavioral outcomes remains a critical question (Glover et al., 2010). To the best of our knowledge, only one study (van den Bergh et al., 2008) has so far provided evidence that an altered diurnal cortisol profile in adolescent offspring associated with prenatal anxiety was underlying, thus mediating, an altered behavioral phenotype (i.e., depressive symptoms) in adolescent girls (but not in boys). Here, we investigate both diurnal cortisol levels and reactivity of the HPA axis following prenatal stress exposure, which has not been done in most previous work. Basal (diurnal) levels of HPA functioning, which follow a circadian rhythm, are important to cortisol output, which helps to maintain children's capacity to regulate their emotions and cope with stress. Reactivity of the HPA axis, in contrast, is associated with elevated cortisol levels in response to conditions that are threatening, unpredictable, and lacking in support. Thus, here, we investigate in a comprehensive way if HPA axis activity mediates the association between prenatal maternal mood and later child EFs.

Present Study

Based on prior research and theory, we hypothesize that depressed and/or anxious prenatal maternal mood is associated with poorer EFs in 6-year-olds. Further, we hypothesize that heightened HPA axis activity, reflected in two key components of the child/offspring HPA axis, namely, diurnal and reactivity cortisol levels, mediates the association of depressed and/or anxious prenatal maternal mood and children's EFs (see Figure 1). We also ask whether sex moderates the effects of prenatal maternal mood on HPA axis activity and EFs.

Method

Participants

Participants included 107 children and their mothers from southwestern Canada. A cohort of middle- to high-income pregnant women ($n = 191$) and their singleton fetuses were recruited during their second trimester (at 26 weeks gestation) from community midwifery clinics and family physician clinics in the greater Vancouver metropolitan area to examine the impact of prenatal SSRI exposure and depressed and anxious prenatal maternal mood on child development. All SSRI-treated mothers had started taking medications based on clinical need, had a diagnosis of a mood disorder, and were already taking SSRIs at the time of conception. Women in both groups (SSRI and nonexposed) had a wide range of anxious and/or depressive symptoms during pregnancy (see Table 1).

Of the original 191 mothers, 4 withdrew before the baby was born and another 4 withdrew before the end of the child's first year. At 6 years, an additional 45 children were unavailable for study (families had moved and 4 mothers had withdrawn by 3 years). At the 6-year assessment 107 children (57% girls; $M_{\text{age}} = 5;11$ years, $SD = 7$ months; $n = 44$ SSRI-exposed, $n = 63$ nonexposed) had complete EF data, complete maternal mood measures, and complete diurnal cortisol and/or complete cortisol reactivity data.

Procedure

The study was approved by the University of British Columbia Research Ethics Board and the BC Womens Hospital Research Review Committee. Informed consent was obtained from mothers at the beginning of the study. Mothers were interviewed by trained research assistants using the Hamilton Rating Scales for Depression and Anxiety (HAM-D and HAM-A; Hamilton, 1960) during the third trimester of pregnancy and again when their children were 6 years of age. At age 6 years, children came to the study center midmorning and were tested individually by trained research assistants.

Table 1. Descriptives and correlations of child and mother variables and child sex

	2	3	4	5	6	Child sex	Mean	SD	Min–Max
1. Child EFs (H&F ACC)	-.20*	-.07	-.19 [†]	.07	.03	-.09	0.68	0.23	0.05–1
2. Child cortisol reactivity (AUCg)	—	.57**	-.04	-.22*	-.09	-.01	40.52	25.90	7.48–119.93
3. Child diurnal cortisol (AUCg)		—	.09	-.20*	.03	.01	123.83	62.23	22.84–311.09
4. Prenatal maternal mood (3rd trimester)			—	.47**	.42**	.13	7.25	5.50	0–25
5. Concurrent maternal mood (6 years)				—	.33**	.12	7.73	6.16	0–27
6. Prenatal SSRI exposure					—	.19 [†]	0.41	0.49	0–1

Note: Child sex: 1 = boys, 2 = girls. Prenatal SSRI exposure: 0 = no, 1 = yes. SSRI, selective serotonin reuptake inhibitor antidepressants exposure. EFs, executive functions. H&F ACC, accuracy in Hearts & Flowers task. AUCg, area under the curve with respect to ground (raw cortisol values). [†] $p = .052$. * $p < .05$. ** $p < .001$.

The MacArthur Assessment Battery for Middle Childhood (Alkon et al., 2003) served as the lab challenge stress procedure and was administered by two research assistants blinded to exposure group status and without the parent present. This 15-min autonomic reactivity protocol for children aged 4–8 years consists of seven blocks, including the following stressors: a social task (questions about family and school); a cognitive task (number-recall task up to six digits); a physical challenge (lemon-juice taste-identification task); and an emotional challenge (two short, emotion-evoking video to elicit fear [a boy having a nightmare] and sadness [a young girl whose pet bird has died]). Age-appropriate relaxing stories were read aloud at the beginning and end of the protocol, and between challenges a quiet inactivity period separated each task. Saliva was collected before, during, and after this procedure to yield measures of children's stress reactivity cortisol. The Hearts & Flowers task (Davidson, Amso, Anderson, & Diamond, 2006) was administered to assess EFs as part of a larger battery of tasks, all of which totaled 2 hr of administration on 1 day. In the week before the lab study, the research coordinator visited children's homes and instructed mothers how to collect saliva from the children to yield measures of children's diurnal salivary cortisol.

Measures

Child EFs. Children completed the Hearts & Flowers task, a widely used computerized EFs measure that has been validated with children 4–13 years of age and with adults (previously called the Dots task; Davidson et al., 2006; Diamond, Barnett, Thomas, & Munro, 2007; Wright & Diamond, 2014; Zaitchik, Iqbal, & Carey, 2014). This task captures performance of all three components of EFs (inhibition, working memory, and cognitive flexibility [task switching]; Diamond, 2013).

In this task, the child responds to a stimulus (heart or flower) by pressing one of two locations on opposite sides of a touchscreen. There are three blocks of trials. During the first block, which consists of 12 trials, a heart appears on either the left or the right side of the screen and children are instructed to press on the same side as the heart. In the second block, which also consists of 12 trials, a flower appears on either the left or the right side of the screen and children are instructed to press on the side opposite the flower. Thus, children had to resist their prepotent response and instead press on the side opposite the stimulus. Both blocks include a short set of instructions with 4 practice trials (which if necessary can be repeated up to three times). The last block consists of 33 trials where a heart or flower appears on either side of the screen (i.e., mixed block). Children are instructed to press according to the previously learned rules. That is, they had to remember the two rules, mentally translate “same side” or “opposite side” into a left or right response on each trial, and flexibly shift between the two rules, inhibiting one to apply the other.

Stimuli were presented for 1500 ms. Responses >2000 ms were considered incorrect (inattentive) and those <250 ms,

impulsive (too fast to have been in response to the stimulus). Both responses were excluded. Outlier trials were removed by using a lower and upper threshold of 2 *SD* from the mean reaction time per trial type per block and per subject. The dependent variable used in the current analyses was the proportion of correct responses in the mixed block.

Child stress regulation. Children's HPA axis function was studied in two settings: under challenge stress (lab setting) and in an everyday (home) setting. Salivary cortisol served as a biomarker for HPA axis activity. Samples were obtained using a sterile Sorbette (Salimetrics) placed in the inside surface of the cheek for 2–3 min and stored at –20 °C until later assayed. All samples were assayed twice using a commercially available chemiluminescent technique (IBL-Hamburg) at the Technical University of Dresden (Dresden, Germany), and average values were used in analyses. The assay has a sensitivity of 0.16 ng/ml, with intra-assay and interassay coefficients of variation less than 9%.

Area under the curve with respect to ground (AUC_g; Pruessner, Kirschbaum, Meinlschmid, & Hellhammer, 2003) was calculated to assess the total hormone concentration over a specific time period (i.e., the day and challenge stress procedure). AUC_g indices were log transformed to minimize skew. Values greater than 3 *SD* above or below the mean were winsorized (2 participants or 3.4% of all values).

Diurnal cortisol. In the week before the lab challenge study, saliva was collected 3 times on 4 consecutive days to yield measures of diurnal salivary cortisol. Saliva samples were obtained upon awakening ($M_{\text{time}} = 7:34$ a.m., $SD = 0.57$ min), 20 min postawakening ($M_{\text{time}} = 7:57$ a.m., $SD = 0.57$ min), and after dinner ($M_{\text{time}} = 6:50$ p.m., $SD = 0.57$ min). For the current analyses, we calculated the mean for each time point across the 4 days. Due to missing samples, for three children the means were calculated from 3 days (one child) or 2 days (two children). The dependent variable used in the current analyses was AUC_g calculated from the three saliva samples.

Stress reactivity cortisol. To determine children's stress reactivity cortisol, five saliva samples were collected over the course of the study day when the lab challenge stress procedure was administered. The sample collected upon arrival to the lab was taken at least 15 min after the child had anything to eat or drink. Two other saliva samples were taken before the lab challenge stress procedure started (one upon completion of the Hearts & Flower task and one upon completion of the Kaufman Brief Intelligence Test). The mean of these first three saliva measures served as the baseline measure ($r_s = .54-.62$, all $p_s < .001$). The next saliva sample was collected 40 min post-stress onset and served as the reactivity measure. Given that the hypothalamic stress responses can be observed in salivary cortisol concentrations approximately 20–30 min post-stress onset and given that our stress

challenge procedure lasted 15 min, the probability that we were able to capture children's stress reactivity to the lab challenge stress procedure is relatively high. Finally, a saliva measure was collected at home in the evening (recovery; $M_{\text{time}} = 6:47$ p.m., $SD = 71$ min). For the current analyses, we replaced missing evening (recovery) values with the mean of evening cortisol values from the diurnal assessments ($n = 26$). The dependent variable used in the current analyses was AUCg calculated from the three saliva samples (baseline, reactivity, and recovery).

Depressed and/or anxious maternal mood. HAM-D and HAM-A (Hamilton, 1960) are two clinician rated scales that measure the severity of depression and anxiety symptoms in adults. The HAM-D is based on 21 items scored on a 5-point scale (9 items), a 4-point scale (11 items), or a 3-point scale (1 item) ranging from 0 = none to 4 = disabling, encompassing a range from 0 to 61. The HAM-A is based on 14 items scored on a 5-point scale (ranging from 0 = none to 4 = disabling) with a range from 0 to 56. A trained research assistant administered the scales during the third trimester of pregnancy ($M = 34.6$ weeks, $SD = 1$ week), and again at the 6-year assessment ($M_{\text{mothers' age}} = 39$ years, $SD = 5$ years). The dependent variable used in these analyses was the mean of both scales to yield a composite measure of depressed and/or anxious maternal mood (the two scales were highly correlated within each time point: $r = .89/.92$, $ps < .001$).

Missing data. Of the total 107 children, $n = 103$ had complete diurnal cortisol data (4 did not have reactivity cortisol), and $n = 104$ had complete cortisol reactivity data (3 did not have diurnal cortisol). No data were missing for EFs or maternal mood.

Statistical analyses

To test our hypotheses, multiple linear regression models were run (SPSS 20). First, to test the total effect, (i.e., our first hypothesis), we regressed child EFs on prenatal maternal mood, controlling for child age and sex, concurrent maternal mood, and prenatal SSRI antidepressant exposure (binary group status). Second, to examine whether children's stress regulation mediated the association between prenatal maternal mood and child EFs (i.e., our second hypothesis), we estimated two separate mediation models, one for each HPA activity component (i.e., stress reactivity and diurnal levels). For these analyses, we used the "indirect" macro designed for SPSS (Hayes, 2013). This is an empirical bias-corrected bootstrapping procedure, which involves obtaining 50,000 artificial samples drawn with replacement from the existing data and thereby reduces Type 1 error rates by using resampling. This approach is well suited for small sample sizes and accounts for the possibility of nonnormality and/or asymmetry for the indirect effect. In addition, contrary to the more traditional causal steps logic, which requires that both paths a

and b are statistically significant (Baron & Kenny, 1986), individual paths a and b are not required to be significant in order to determine whether M mediates the effect of X on Y (Hayes & Rockwood, in press). Parameter estimates and 95% confidence intervals (CI) for all indirect paths were derived and mediation is supported when the CIs do not contain zero.

As child sex may moderate the effects of prenatal stress on the HPA axis, we tested if our mediation models were moderated by sex (i.e., we calculated moderated mediation models where all paths are simultaneously tested for an interaction of sex with the predictor variable). All variables were mean-centered prior to creating the interaction term. Significant interactions were tested for significance in simple slopes to detect areas of significance.

In all our models we controlled for concurrent maternal mood (at 6 years of children's age) in order to isolate the effect of prenatal maternal mood on child outcomes. In addition, we controlled for prenatal SSRI exposure, and time of the day of saliva assessment where required.

Results

Preliminary analyses

Table 1 presents the descriptive data and shows zero-order correlations among the study variables as well as their correlations with sex. As expected, depressed and/or anxious prenatal maternal mood was marginally related to EFs ($r = .19$, $p = .052$); thus, the poorer the child EFs, the worse the maternal depression and/or anxiety symptoms during pregnancy. Furthermore, greater cortisol reactivity was associated with poorer EFs. Diurnal cortisol levels, however, were not significantly associated with child EFs. There were no significant associations between prenatal maternal mood and children's HPA axis activity (AUCg scores). Concurrent maternal mood, in contrast, showed small to moderate negative associations with both diurnal and reactivity cortisol levels; that is, the greater the concurrent maternal depression and/or anxiety symptoms, the lower the child cortisol levels. The two indices of HPA axis activity were highly correlated with each other. Depressed and/or anxious maternal mood showed moderate to high stability over the 6-year period and moderate correlations with prenatal SSRI antidepressants intake. Apart from these associations, prenatal SSRI antidepressants exposure was not correlated with any of the study variables, except a positive trend with child sex was found (mothers of girls were slightly more likely to be on SSRI antidepressants during pregnancy). No other significant associations with sex were observed.

Total effect of depressed and/or anxious prenatal maternal mood on child EFs

To test the first hypothesis, a multiple linear regression model was run to examine whether child EFs at 6 years was

predicted by prenatal maternal mood, while controlling for mother and child characteristics. Prenatal depressed and/or anxious maternal mood was, as hypothesized, a significant predictor of child EFs at 6 years ($\beta = -.639, p = .014$), after controlling for child age, child sex, prenatal SSRI antidepressant exposure, and concurrent maternal mood (Table 2). The quadratic term for prenatal maternal mood was also significantly related to child EFs ($\beta = .511, p = .042$), but in an unexpected way. As can be seen in Figure 2, we found a “hockey stick” relationship. At lower levels of depressed and/or anxious prenatal maternal mood, the association between child EFs and maternal mood was linear; namely, fewer correct responses on the third block of the Hearts & Flowers task was associated with increased maternal mood disturbances during pregnancy. However, when prenatal maternal mood reached a certain level of severity, this relationship plateaued and increasing mood disturbances had no further impact on EFs. The relation between prenatal maternal mood and child EFs was not moderated by sex ($\beta = -.020, p = .819$). However, in this model sex was marginally related to child EFs ($\beta = -.158, p = .059$), indicating that boys had somewhat better EFs than girls.

In sum, depressed and/or anxious prenatal maternal mood (linear effect: $\beta = -.639, p = .014$, quadratic effect: $\beta = -.511, p = .042$) predicted child EFs after controlling for child age and sex, concurrent maternal mood, and prenatal SSRI antidepressant exposure, suggesting that greater prenatal maternal depression and/or anxiety symptoms are associated with poorer child EFs.

Child HPA axis stress reactivity mediates the association of prenatal maternal mood and child EFs

To test the first mediation hypothesis, a moderated mediation model was run with cortisol stress reactivity as a mediator. The index of moderated mediation was .0053 (.0036) 95%

Table 2. Multiple linear regression predicting child EFs from prenatal maternal mood (total effect)

	<i>B</i>	β	<i>t</i>	<i>p</i>
Child age	.230	.585	7.03	<.001
Child sex	-.074	-.158	-1.91	.059
Prenatal SSRI exposure	.016	.035	0.39	.696
Prenatal maternal mood	-.027	-.639	-2.51	.014
Prenatal maternal mood ²	.001	.511	2.06	.042
Prenatal Maternal Mood \times Child Sex	-.005	-.020	-0.24	.813
Concurrent maternal mood	.000	.004	0.04	.966
<i>R</i> ²		.42		

Notes: Significant *p* values are in bold face. *B*, unstandardized beta coefficients. β , standardized beta coefficients. *R*², explained variance. Child sex: 1 = boys, 2 = girls. Prenatal SSRI exposure: 0 = no, 1 = yes. SSRI, selective serotonin reuptake inhibitor antidepressants exposure. EFs, executive functions.

CI [.000001, .0141], indicating that the mediation was moderated by sex (i.e., the CI does not contain zero). Specifically, we found a significant indirect effect of depressed and/or anxious prenatal maternal mood \rightarrow heightened cortisol reactivity \rightarrow child EFs for boys, $B = -.0053$ (.0035) 95% CI [-.0142, -.0004], but not for girls, $B = 0$ (.0011) 95% CI [-.0021, .0025]. Complete model results, including all covariates, are reported in online-only supplementary materials Table S.1. The mediation model for cortisol reactivity is depicted in Figure 3. Looking at the individual paths, path *a*₁ indicates that more depressed and/or anxious prenatal maternal mood symptoms were associated at trend level with heightened cortisol reactivity, $B = .021, p = .075$. However, path *a*₁ also indicates that there was a trend-level interaction of prenatal maternal mood with sex, $B = -.038, p = .072$.

To further explore this interaction, using *z*-standardized variables, we examined how prenatal maternal mood interacts with sex to predict cortisol reactivity to lab challenge (*z*-standardized AUCg scores, log transformed) while controlling for prenatal antidepressant exposure, concurrent maternal mood, and time of the day of saliva assessment (Figure 4). Inspection of the simple slopes revealed that for boys only cortisol reactivity depended on prenatal maternal mood (simple slope for boys: $B = .381, p < .05$): more depressed and/or anxious prenatal maternal mood symptoms were associated with increased cortisol reactivity to lab challenge. For girls, however, cortisol reactivity remained unchanged even as prenatal maternal mood symptoms worsened (simple slope for girls: $B = .022, p = .867$).

Furthermore, path *b*₁ indicates that cortisol reactivity (*z*-standardized AUCg scores, log transformed) interacted at trend level with sex to predict child EFs at 6 years, controlling for prenatal antidepressant exposure, concurrent maternal mood, and time of the day of saliva assessment, $B = .127, p = .095$ (see Figure 5). That is, for boys (simple slope: $B = -.066, p = .049$) but not for girls (simple slope: $B = .008, p = .819$), higher cortisol reactivity was associated with poorer EFs. Finally, path *c*₁ (direct effect) was significant across girls and boys ($B = -.011, p = .022$; i.e., it was not moderated by sex).

In sum, these findings indicate that depressed and/or anxious prenatal maternal mood was negatively associated with child EFs via heightened cortisol reactivity, but only for boys. Prenatal maternal mood, however, had a significant direct effect on child EFs for both girls and boys.

Child diurnal HPA axis activity mediates the association of prenatal maternal mood and child EFs

To test the second mediation hypothesis, we again ran a moderated mediation model with diurnal cortisol levels as a mediator. The index of moderated mediation was $-.0023$ (.0035) 95% CI [-.0048, .0087], indicating that this mediation was not moderated by sex (i.e., the CI contains zero). There was neither an indirect effect for boys, $B = -.0022$ (.0034) 95% CI [-.0086, .0047], nor for girls, $B = 0$




Figure 2. Prenatal maternal mood predicts child executive functions (EFs; accuracy in Hearts & Flowers task) at 6 years while controlling for child sex, child age, prenatal antidepressant exposure, and concurrent maternal mood (linear effect: $\beta = -.639, p = .014$, quadratic effect: $\beta = .511, p = .042$). At lower levels of depressed and/or anxious prenatal maternal mood, the association between child EFs and maternal mood was linear, namely, fewer correct responses on the third block of the Hearts & Flowers task was associated with increased mood disturbances. However, when prenatal maternal mood reached a certain level of severity, this relation disappeared, and increasing mood disturbances had no ongoing impact on EFs. This may have implications for interventions. Namely, among highly depressed/anxious women, it may be crucial to know that improvements in maternal mood alone may not directly translate into better child outcomes.

(.0008) 95% CI [-.0014, .0020]. Complete model results, including all covariates, are reported in online-only Supplementary Table S.2. The mediation model for diurnal cortisol is depicted in Figure 6. Looking at the individual paths, path a_2 indicates that more depressed and/or anxious prenatal maternal mood symptoms were associated with heightened diurnal cortisol levels, $B = .023, p = .029$. However, path a_2 also shows that there was a significant interaction of prenatal maternal mood and sex, $B = -.52, p = .008$.

To further understand this interaction, using z -standardized variables, we examined how prenatal maternal mood interacts with sex to predict diurnal cortisol levels (z -standardized AUCg scores, log transformed) while controlling for prenatal antidepressant exposure, concurrent maternal mood, and time of the day of saliva assessment (Figure 7).

For boys (simple slope: $B = .492, p = .002$), but not for girls (simple slope: $B = -.027, p = .833$), diurnal cortisol levels depended on prenatal maternal mood. Namely, more depressed and/or anxious prenatal maternal mood symptoms were associated with higher diurnal cortisol levels.

Path b_2 was not significant, $B = -.008, p = .870$ (also no significant interaction with sex), indicating that children’s diurnal cortisol levels were not associated with EFs performance. Finally, path c_2 (direct effect) was significant, $B = -.014, p = .008$, across girls and boys.

Taken together, among boys we observed that heightened cortisol reactivity but not diurnal cortisol levels mediated the association between depressed and/or anxious prenatal maternal mood and EFs. However, it is important to note that prenatal maternal mood was also associated with EFs in girls as




Figure 3. Bootstrapping results testing the mediation model. Unstandardized estimates. Dashed lines indicate marginally significant paths ($p < .10$). Significant indirect effects for boys, $B = -.0053$ (.0035) 95% CI [-.0142, -.0004], but not for girls, $B = 0$ (.0011) 95% CI [-.0021, .0025]. Controlling for time of the day of saliva assessment, prenatal antidepressant exposure, and concurrent maternal mood.




Figure 4. Prenatal maternal mood interacts with sex to predict cortisol reactivity to lab challenge (z -standardized AUCg scores, log transformed) while controlling for prenatal antidepressant exposure, concurrent maternal mood, and time of the day of saliva assessment. For boys only, cortisol reactivity depended on prenatal maternal mood (simple slope for boys: $B = .381$, $p < .05$): more depressed/ anxious prenatal maternal mood was associated with increased cortisol reactivity to lab challenge. For girls, however, cortisol reactivity was not associated with exposure to prenatal maternal mood (simple slope for girls: $B = .022$, $p = .867$).

indicated by the significant total effect and both significant direct effects in the diurnal cortisol model and the cortisol reactivity model.

Discussion

Prenatal exposure to maternal mood disturbances has been reported to shape a child's subsequent development of EFs. However, specific mechanisms that link the prenatal environment with postnatal outcomes require further investigation.




Figure 5. Cortisol reactivity (z -standardized AUCg scores, log transformed) interacts with sex to predict child executive functions (EFs; accuracy in Hearts & Flowers task) at 6 years, while controlling for prenatal antidepressant exposure, concurrent maternal mood, and time of the day of saliva assessment. For boys only, EFs depended on cortisol reactivity: higher cortisol reactivity was associated with poorer EFs (simple slope for boys: $B = -.066$, $p = .049$). For girls, in contrast, cortisol reactivity was not associated with EFs (simple slope for girls: $B = .008$, $p = .819$).

By examining children's cortisol responses under both unstressed conditions and following stress, this study sought to determine the role of the child/offspring HPA axis in mediating effects of prenatal maternal mood on child EFs at age 6 years. We found that in boys but not girls, depressed and/or anxious prenatal maternal mood is associated with both heightened diurnal cortisol levels and heightened cortisol reactivity to a lab challenge, and that heightened reactivity is associated with poorer EFs. Further, among boys, cortisol reactivity but not diurnal cortisol levels mediated the association between depressed and/or anxious prenatal maternal mood and EFs. The total effect of depressed and/or anxious prenatal maternal mood on child EFs, however, was significant for both girls and boys. Specifically, depressed and/or anxious prenatal maternal mood predicted child EFs up to a point and then asymptoted so that even worse prenatal maternal mood symptoms had no further impact on child EFs. To the best of our knowledge, these are the first findings that demonstrate a role for child stress regulation in mediating the relationship between prenatal maternal stress-related mood disturbances and child EFs, thereby providing insight into potential pathways through which EF development (at least in boys) may be affected by depressed and/or anxious prenatal maternal mood.

Total effect on executive functions

Given the central role of EFs for child development and health, these findings highlight the importance of incorporating the prenatal period into our models of EF development. In line with the notion of equifinality (Cicchetti & Rogosch, 1996), a central tenet of developmental psychopathology denoting that there are many developmental pathways that may lead to the same outcome, our findings expand on the child development literature showing that EFs are malleable, and context-specific experiences matter both at home and at school (for reviews, see Diamond & Lee, 2011; Hughes, 2011; Ling, Kelly, & Diamond, 2016). Specifically, our findings add to the emerging literature within the field of prenatal maternal stress showing that depressed and/or anxious prenatal maternal mood is associated with subtle changes in EFs and frontal lobe structures in middle childhood, adolescence, and early adulthood (e.g., Buss et al., 2011; van den Bergh, Mennes, et al., 2005).

Contrary to our hypothesis, we were not able to detect an inverted U-shaped relation between prenatal stress and EFs (Arnsten, 2009). In our sample, the data were best represented by a regression line that flattened when depressed and/or anxious prenatal maternal mood reached a certain level of severity. This may have implications for interventions. Namely, among highly depressed and/or anxious women, it may be crucial to know that improvements in maternal mood alone may not directly translate into better child outcomes. Possibly, beyond mood itself, related but less prominent maternal characteristics of mood disorders such as maternal cognitive disturbances may also shape child outcomes (Snyder, 2013).




Figure 6. Bootstrapping results testing the mediation model. Unstandardized estimates. Dotted lines indicate nonignificant paths. No indirect effect. Controlling for time of the day of saliva assessment, prenatal antidepressant exposure, and concurrent maternal mood.

Clearly, replication of our findings with another cohort and/or other EF tasks is important. Future research is also needed to address the question when and how moderate stress/cortisol levels experienced during the prenatal period may lead to optimal EF performance.

Mediation through child HPA axis stress reactivity

Our findings may also reflect multifinality, the other central tenet of psychopathology, which refers to the notion that any one process may function differently across systems or within individuals (Cicchetti & Rogosch, 1996). Thus, not all children are affected equally by prenatal maternal stress or mood disturbances. Our finding indicating that cortisol

reactivity but not diurnal cortisol levels mediated the effect of prenatal maternal mood on EFs for boys only suggests that there probably are different mechanisms underlying the effects of exposure to depressed/ anxious prenatal maternal mood symptoms on EFs for girls (e.g., immune system and inflammation, such as the pro-inflammatory cytokines, Coussons-Read, Okun, & Nettles, 2007; epigenetics, Oberlander et al., 2008; autonomic nervous system responses, Surland et al., 2017; dopaminergic systems, Zhang, Chretien, Meaney, & Gratton, 2005).

Within the literature of developmental effects of prenatal exposure to substance abuse (e.g., nicotine), animal studies have demonstrated sex differences, with males generally showing higher vulnerability than females (e.g., Shacka, Fennell, & Robinson, 1997). Similarly, among humans, substance-exposed boys showed greater cognitive deficits than exposed girls (Moe & Slining, 2001). In contrast, a study showing that altered HPA activity played a mechanistic role in fetal programming of child behavioral outcomes found evidence for mediated effects of prenatal anxiety on depressive symptoms for 14- to 15-year-old girls but not boys (van den Bergh et al., 2008). The authors argue that maternal anxiety may induce sex-specific changes in the HPA axis during critical gestation periods, which only become apparent post-puberty, when the HPA axis has reached its full maturation. Alternatively, they argue, it may also be plausible that only during puberty do females become more vulnerable to depressive symptoms because of crucial sex differences in gonadal hormones. It will be interesting to see what we find when the children in our study reach puberty. As well, future research needs to explore further the possibility that the timing of the exposure to maternal depression and anxiety as well as the amount of sex hormones in the developing fetus may be relevant for sex differences in the outcome (de Bruijn, van Bakel, & van Baar, 2009; Glover & Hill, 2012).

Prenatal stress has been associated with HPA axis dysregulation in infants, adolescents, and adults (reviewed in Glover et al., 2010). Our findings add to and expand on this literature as our study is one of the very few showing that prenatal




Figure 7. Prenatal maternal mood interacts with sex to predict diurnal cortisol levels (z-standardized AUCg scores, log transformed) while controlling for prenatal antidepressant exposure, concurrent maternal mood, and time of the day of saliva assessment. For boys only, diurnal cortisol levels depended on prenatal maternal mood: more depressed and/or anxious prenatal maternal mood was associated with higher diurnal cortisol levels (simple slope for boys: $B = .492, p = .002$). For girls, however, diurnal cortisol levels were not associated with exposure to prenatal maternal mood (simple slope for girls: $B = -.027, p = .833$).

maternal mood is associated with both aspects of children's HPA axis activity (heightened diurnal cortisol and heightened stress reactivity levels). Whether and how sex moderates the effects of prenatal stress or maternal mood on the development of the HPA axis remains a critical question. Contrary to our results, prior research has found that both prenatal maternal anxiety and prenatal exposure to synthetic glucocorticoids increased stress reactivity in girls only (Alexander et al., 2012; de Bruijn et al., 2009). Whether HPA axis dysregulation appears to be sex specific during specific developmental periods (i.e., if it appears early on and again later in girls or whether it continues to be true only for boys in our sample) remains unknown. Future studies analyzing how sex moderates the relationship between prenatal maternal depression and/or anxiety symptoms and development of the HPA axis are necessary to reconcile inconsistent findings in the literature.

Apart from sex, genetics have been shown to be powerful moderators of prenatal maternal exposures. A recent study (Buchmann et al., 2014), for example, shows that in 19-year-old adolescents exposed to prenatal maternal stress, only carriers of the dopamine D4 receptor gene (*DRD4*) seven-repeat allele were found to have altered (i.e., attenuated) cortisol secretion during the Trier Social Stress Test. These results suggest that prenatal maternal stress may only affect the HPA axis of carriers of certain "risk alleles" (the *DRD4* 7r allele has been shown to be a "risk allele" for externalizing problems, particularly in the presence of environmental adversity; Bakermans-Kranenburg & van IJzendoorn, 2006). In our own cohort, we found that among children whose mothers exhibited some degree of depressive-anxious affect during pregnancy, a child's EFs at age 6 varied depending on the child's serotonin transporter (*SLC6A4*) genotype and his/her mother's mood (Weikum et al., 2013). The EFs of children with at least one short allele of the gene were well preserved even if their mothers reported many depressive symptoms (i.e., they showed resilience), whereas the EFs of children with two long forms of the *SLC6A4* gene appeared to be very sensitive to their mothers' mood. If their mothers were more symptomatic, these children showed worse EFs than those in any other group; but if their mothers were happier, these children's EFs were better than those of any other group. Thus, context counts whereby children of more depressed mothers with one or two short alleles of the *SLC6A4* gene showed the better EFs, but given a mother who was not symptomatic, children with two long alleles showed the best EFs. Another intriguing example of multifinality or "biological sensitivity to context"/ "differential susceptibility" (Ellis, Boyce, Belsky, Bakermans-Kranenburg, & van IJzendoorn, 2011) was provided by Pluess et al. (2011), who showed that the association between maternal anxiety during pregnancy and negative emotionality in early infancy was only significant in infants carrying one or more copies of the serotonin transporter linked polymorphic repeat (*5-HTTLPR*) short allele but not in those homozygous for the long allele. In this way, the *5-HTTLPR* allelic variations

might increase vulnerability to adverse environmental influences as early as the fetal period for some infants. Furthermore, in a study of infants born very preterm and exposed to the stress of multiple procedures during fetal life ex-utero, Chau et al. (2014) showed that variation in the catechol-O-methyltransferase (*COMT*) genotype moderated the relationship between early stress and internalizing behaviors at 18 months corrected age. The idea of biological sensitivity to context/differential susceptibility offers a critical perspective that allows us to move our thinking about fetal programming from "invariant" developmental outcomes associated with early adverse exposure to a perspective that outcomes in childhood represent interactions between biological (a child's genotype) and contextual (maternal mood) variables enabling both positive and negative outcomes.

Executive functions and cortisol responses

Physiological arousal or children's cortisol responses and EF skills are not linked in a fixed linear pattern, but are more accurately represented in a complex dynamic interplay. In our sample, a heightened HPA axis reactivity but not a heightened diurnal HPA axis profile was associated with poorer EFs. These findings are mostly in line with a small number of studies with children suggesting that heightened cortisol reactivity and heightened baseline cortisol levels are negatively associated with effortful control and EF performance (Blair et al., 2011; Donzella, Gunnar, Krueger, & Alwin, 2000; Wagner et al., 2015). In contrast, other studies have found that cortisol reactivity was associated positively with preschoolers' effortful control and EFs (Blair, Granger, & Peters Razza, 2005; Spinrad et al., 2009). More nuanced analyses suggest, however, that the interplay between physiological arousal/cortisol responses and EF skills is even more complex. Blair and Berry (2017), for instance, recently reported that young children with relatively low cortisol levels, showing moderate fluctuations in their cortisol levels over the first 4 years of their lives, tended to show better EF performance at 5 years of age than did children with either highly stable or highly variable temporal profiles. Recent research also suggests that heightened cortisol reactivity may promote EF skills in nurturing contexts, but undermine them in less resourced or less supportive contexts (probably because the children are too stressed). In a study of kindergarteners, cortisol responses interacted with family income to predict EFs (Obradović, Portilla, & Ballard, 2016): heightened cortisol levels were associated with better EFs in children from higher income families and with worse EFs in children from lower income families. It is an open question if children in our sample experienced a less resourced or less supportive context and therefore their heightened HPA axis reactivity was associated with worse EFs. Clearly context matters, and research is needed to clarify under which conditions heightened cortisol reactivity and/or heightened baseline cortisol levels are adaptive versus maladaptive and may therefore represent a plasticity versus risk factor.

It is somewhat surprising that we only found a relationship between heightened cortisol reactivity and lower EFs for boys and not girls. Based partly on the animal literature, researchers have emphasized that it is important to examine potential differential sex effects when investigating relationships between stress reactivity and child outcomes (Spinrad et al., 2009). In line with our findings, Donzella et al. (2000) reported that all but one of the children who showed an increase in cortisol in response to a competition were male. These boys were described by teachers as more surgent and lower in effortful control. One possible mechanism for these sex differences may be that boys and girls may respond to stress differently (Stroud, Salovey, & Epel, 2002). Specifically, females are more likely to respond to challenge by focusing more on managing interpersonal relationships (building alliances and calming others) whereas males, when faced with stress, tend to engage more in risky behaviors (Booth, Granger, & Shirtcliff, 2008). As a case in point, Shirtcliff, Granger, Booth, and Johnson (2005) found that lower basal cortisol levels predicted higher externalizing problems for boys, but not for girls. In addition to these behavioral mechanisms, we may not have been able to detect a relationship between cortisol reactivity and EFs for girls because females are more likely to show increases in autonomic responses to stress as opposed to males who are more likely to show increased cortisol to stress (Kudielka & Kirschbaum, 2005). Furthermore, bodies of females evidence stress in different ways than do the bodies of males (even though the levels of stress might be comparable; Kudielka & Kirschbaum, 2005). Thus, boys' and girls' stress responses may have different implications for their social functioning and possibly also for their cognitive functioning. Future research may explore implications of these sex-specific associations in the context of prenatal stress exposure.

Limitations and future directions

It is important to note several limitations of the present study. First, as in all human studies investigating prenatal stress or mood disturbances, where randomizing exposure is not possible, we cannot speak to the causal effects of prenatal maternal mood on child behavior. However, as we took care to control statistically for postnatal maternal mood (as commonly done in this field), our findings support the concept of fetal programming, namely, that some of the risk is conferred prenatally from mother to child. Nevertheless, potential confounding by genetic influences cannot be ruled out completely. Rice et al. (2010), for instance, leveraged *in vitro* fertilization to tease apart genetic and putative prenatal stress effects on offspring development and reported results suggesting offspring attention-deficit/hyperactivity disorder (ADHD) may be due to shared inheritance of characteristics, rather than prenatal stress effects of the *in utero* environment. It remains questionable, however, to what extent these findings translate to EFs deficits following prenatal stress exposure. Cognitive deficits associated with ADHD are usually

thought of in terms of poor EFs (Johnson, 2015), but as other crucial neuropsychological factors also contribute to the complex symptoms of this developmental disorder (Pauli-Pott & Becker, 2011), caution is warranted when translating ADHD findings to EFs outcomes.

Second, and related to the issue of causality, we recognize that by using mediational models, the mediator is presumed to cause the outcome and not vice versa (cf. reverse causation; Cole & Maxwell, 2003). However, in our case, the timing of the measurement of the mediator (HPA axis activity) and outcome (EF performance) was not optimal; that is, the two variables were measured at the same time point (at 6 years). However, given our strongly theoretically based hypotheses, reverse causal effects appear unlikely. Nevertheless, future work with our cohort where we will be able to measure the outcome *after* the mediator will improve the strength of our conclusions.

Third and finally, it is important to bear in mind that no measurement of EFs is perfect. Likewise, cortisol is an imperfect measure of level of stress, especially in females (Kudielka & Kirschbaum, 2005). Furthermore, the same experience is differentially stressful to different persons; and the cognitive testing (EFs tests and Kaufman Brief Intelligence Test) may have been stressful for some children but not others, hence providing an imperfect baseline. The fact, however, that we assessed baseline cortisol levels with three saliva measures (one upon arrival to the lab) strengthens our conclusions. As to the recovery saliva measure taken at home after the lab visit, we are not aware of any other study applying such a measure in combination with lab-based stress reactivity. Unfortunately, no perceived stress state of child concurrent with time of home recovery sample was assessed, so we cannot determine the emotional/arousal state the child was in when this last saliva sample was taken. Based on previous research showing lower stress reactivity evoked in home-based studies compared to childcare- and/or lab-based studies (Alkon et al., 2003), we are though inclined to assume that our recovery saliva measure is not only a reliable indicator of stress recovery but also may actually maximize ecological validity of this study.

Despite these limitations, this study makes innovative contributions to the field of the developmental origins of psychopathology. It is a first step in the attempt not only to better understand how prenatal maternal mood shapes children's EF development but also to illuminate child/offspring HPA axis reactivity as a prominent mechanism in fetal programming. Future research may identify other mechanisms (e.g., autonomic nervous system responses, epigenetics, dopaminergic systems, or immune system and inflammation, such as the pro-inflammatory cytokines) through which children's, and maybe in particular girls', EFs are affected by prenatal maternal mood.

Clearly, more work needs to be done to better understand how sex moderates the effects of prenatal stress on the development of stress responsivity and regulation. Future work may also gain critical insight by investigating whether altered EFs act as a mediator between early stress exposure and

subsequent child outcomes. A recent study (Pearson et al., 2016), reported that prenatal anxiety was associated with impaired working memory in the offspring at age 8, which then mediated the effect of prenatal anxiety on math grades at age 16. Such findings may allow us to understand how exposure to prenatal stress affects children's functioning across different developmental domains.

Finally, research is needed to examine interactions of prenatal stress with postnatal environment and the child's genetic profile. Despite the popularity of the fetal programming model, it is also important to acknowledge that such effects are not necessarily permanent as animal research has shown that young animals show remarkable neuronal resilience if the stress is discontinued (cf. McEwen & Morrison, 2013). Thus, while in these analyses we focused on prenatal maternal mood exposure and its associations with child stress regulation and EFs (i.e., while controlling for postnatal maternal mood), it is important to bear in mind that postnatal factors (e.g., postnatal maternal mood, parenting, secure attachment, and socioeconomic status) are important as well and may reverse the effects of prenatal stress exposure (or exacerbate them). The congruence between the demands of prenatal and postnatal environments may be a crucial factor. For instance, Sandman, Davis, and Glynn (2012) reported accelerated motor and mental development during the first year of life among infants whose mothers experienced congruent

levels of depressive symptoms during and after pregnancy, even when the levels of symptoms were relatively high and the prenatal and postnatal environments were unfavorable. In this sense, prenatal environments prepare the fetus for postnatal life and confer an adaptive advantage for critical survival functions during early development. Furthermore, maternal prenatal and postnatal mental health problems may be differentially associated with later outcomes in the offspring. For instance, Vänskä et al. (2015) showed that both maternal prenatal and postnatal mental health problems predicted children's later stress regulation, but in unique ways (maternal prenatal mental health problems predicted an intensified CAR, whereas mothers' early postpartum mental health problems predicted a reduced CAR). Future research will hopefully explore in more detail the combination of early life stress, genetics, and ongoing stress that may ultimately determine an individual's responsiveness to stress and the vulnerability or resilience in developmental outcomes, such as EFs and psychiatric disorders (e.g., depression; Charney & Manji, 2004).

Supplementary Material

To view the supplementary material for this article, please visit <https://doi.org/10.1017/S095457941800041X>.

References

- Alexander, J. K., Hillier, A., Smith, R. M., Tivarus, M. E., & Beversdorf, D. Q. (2007). Beta-adrenergic modulation of cognitive flexibility during stress. *Journal of Cognitive Neuroscience*, *19*, 468–478. doi:10.1162/jocn.2007.19.3.468
- Alexander, N., Rosenlocher, F., Stalder, T., Linke, J., Distler, W., Morgner, J., & Kirschbaum, C. (2012). Impact of antenatal synthetic glucocorticoid exposure on endocrine stress reactivity in term-born children. *Journal of Clinical Endocrinology and Metabolism*, *97*, 3538–3544. doi:10.1210/jc.2012-1970
- Alkon, A., Goldstein, L. H., Smider, N., Essex, M. J., Kupfer, D. J., & Boyce, W. T. (2003). Developmental and contextual influences on autonomic reactivity in young children. *Developmental Psychobiology*, *42*, 64–78. doi:10.1002/dev.10082
- Arnsten, A. F. T. (2009). Stress signalling pathways that impair prefrontal cortex structure and function. *Nature Review Neuroscience*, *10*, 410–422. doi:10.1038/nrn2648
- Bakermans-Kranenburg, M. J., & van IJzendoorn, M. H. (2006). Gene-environment interaction of the dopamine D4 receptor (DRD4) and observed maternal insensitivity predicting externalizing behavior in preschoolers. *Developmental Psychobiology*, *48*, 406–409. doi:10.1002/dev.20152
- Barker, D. J. (2003). The developmental origins of adult disease. *European Journal of Epidemiology*, *18*, 733–736. doi:10.1023/A:1025388901248
- Baron, R. M., & Kenny, D. A. (1986). The moderator–mediator variable distinction in social psychological research: Conceptual, strategic, and statistical considerations. *Journal of Personality and Social Psychology*, *51*, 1173–1182. doi:10.1037/0022-3514.51.6.1173
- Beijers, R., Buitelaar, J. K., & de Weerth, C. (2014). Mechanisms underlying the effects of prenatal psychosocial stress on child outcomes: Beyond the HPA axis. *European Child and Adolescent Psychiatry*, *23*, 943–956. doi:10.1007/s00787-014-0566-3
- Bennett, H. A., Einarson, A., Taddio, A., Koren, G., & Einarson, T. R. (2004). Prevalence of depression during pregnancy: Systematic review. *Obstetrics and Gynecology*, *103*, 698–709. doi:10.1097/01.AOG.0000116689.75396.5f
- Blair, C., & Berry, D. J. (2017). Moderate within-person variability in cortisol is related to executive function in early childhood. *Psychoneuroendocrinology*, *81*, 88–95. doi:10.1016/j.psyneuen.2017.03.026
- Blair, C., & Diamond, A. (2008). Biological processes in prevention and intervention: The promotion of self-regulation as a means of preventing school failure. *Development and Psychopathology*, *20*, 899–911. doi:10.1017/S0954579408000436
- Blair, C., Granger, D., & Peters Razza, R. (2005). Cortisol reactivity is positively related to executive function in preschool children attending Head Start. *Child Development*, *76*, 554–567. doi:10.1111/j.1467-8624.2005.00863.x
- Blair, C., Granger, D. A., Willoughby, M., Mills-Koonce, R., Cox, M., Greenberg, M. T., . . . Family Life Project Investigators. (2011). Salivary cortisol mediates effects of poverty and parenting on executive functions in early childhood. *Child Development*, *82*, 1970–1984. doi:10.1111/j.1467-8624.2011.01643.x
- Blair, C., & Ursache, A. (2011). A bidirectional model of executive functions and self-regulation. In K. D. Vohs & R. F. Baumeister (Eds.), *Handbook of self-regulation: Research, theory, and applications* (2nd ed., pp. 300–320). New York: Guilford Press.
- Booth, A., Granger, D. A., & Shirtcliff, E. A. (2008). Gender- and age-related differences in the association between social relationship quality and trait levels of salivary cortisol. *Journal of Research on Adolescence*, *18*, 239–260. doi:10.1111/j.1532-7795.2008.00559.x
- Brennan, P. A., Pargas, R., Walker, E. F., Green, P., Newport, D. J., & Stowe, Z. (2008). Maternal depression and infant cortisol: Influences of timing, comorbidity and treatment. *Journal of Child Psychology and Psychiatry*, *49*, 1099–1107. doi:10.1111/j.1469-7610.2008.01914.x
- Buchmann, A. F., Zohsel, K., Blomeyer, D., Hohm, E., Hohmann, S., Jenßen-Steinmetz, C., . . . Laucht, M. (2014). Interaction between prenatal stress and dopamine D4 receptor genotype in predicting aggression and cortisol levels in young adults. *Psychopharmacology*, *231*, 3089–3097. doi:10.1007/s00213-014-3484-7
- Buss, C., Davis, E. P., Hobel, C. J., & Sandman, C. A. (2011). Maternal pregnancy-specific anxiety is associated with child executive function

- at 6-9 years age. *Stress*, 14, 665-676. doi:10.3109/10253890.2011.623250
- Buss, C., Davis, E. P., Muftuler, L. T., Head, K., & Sandman, C. A. (2010). High pregnancy anxiety during mid-gestation is associated with decreased gray matter density in 6-9-year-old children. *Psychoneuroendocrinology*, 35, 141-153. doi:10.1016/j.psyneuen.2009.07.010
- Butts, K. A., Weinberg, J., Young, A. H., & Phillips, A. G. (2011). Glucocorticoid receptors in the prefrontal cortex regulate stress-evoked dopamine efflux and aspects of executive function. *Proceedings of the National Academy of Sciences of the United States of America*, 108, 18459-18464. doi:10.1073/pnas.1111746108
- Capuron, L., & Miller, A. H. (2011). Immune system to brain signaling: Neuropsychopharmacological implications. *Pharmacology and Therapeutics*, 130, 226-238. doi:10.1016/j.pharmthera.2011.01.014
- Charney, D. S., & Manji, H. K. (2004). Life stress, genes, and depression: Multiple pathways lead to increased risk and new opportunities for intervention. *Science Signaling*, 2004, re5. doi:10.1126/stke.2252004re5
- Chau, C. M. Y., Ranger, M., Sulistyoningrum, D., Devlin, A. M., Oberlander, T. F., & Grunau, R. E. (2014). Neonatal pain and COMTVal158Met genotype in relation to serotonin transporter (SLC6A4) promoter methylation in very preterm children at school age. *Frontiers in Behavioral Neuroscience*, 8, 409. doi:10.3389/fnbeh.2014.00409
- Chrousos, G. P. (2009). Stress and disorders of the stress system. *Nature Reviews Endocrinology*, 5, 374-381. doi:10.1038/nrendo.2009.106
- Cicchetti, D., & Rogosch, F. A. (1996). Equifinality and multifinality in developmental psychopathology. *Development and Psychopathology*, 8, 597-600. doi:10.1017/S0954579400007318
- Cole, D. A., & Maxwell, S. E. (2003). Testing mediational models with longitudinal data: Questions and tips in the use of structural equation modeling. *Journal of Abnormal Psychology*, 112, 558-577. doi:10.1037/0021-843X.112.4.558
- Cools, R., & D'Esposito, M. (2011). Inverted-U shaped dopamine actions on human working memory and cognitive control. *Biological Psychiatry*, 69, e113-e125. doi:10.1016/j.biopsych.2011.03.028
- Coussons-Read, M. E., Okun, M. L., & Nettles, C. D. (2007). Psychosocial stress increases inflammatory markers and alters cytokine production across pregnancy. *Brain, Behavior, and Immunity*, 21, 343-350. doi:10.1016/j.bbi.2006.08.006
- Davidson, M. C., Amso, D., Anderson, L. C., & Diamond, A. (2006). Development of cognitive control and executive functions from 4 to 13 years: Evidence from manipulations of memory, inhibition, and task switching. *Neuropsychologia*, 44, 2037-2078. doi:10.1016/j.neuropsychologia.2006.02.006
- Davis, E. P., Glynn, L. M., Waffarn, F., & Sandman, C. A. (2011). Prenatal maternal stress programs infant stress regulation. *Journal of Child Psychology and Psychiatry*, 52, 119-129. doi:10.1111/j.1469-7610.2010.02314.x
- de Bruijn, A. T., van Bakel, H. J., & van Baar, A. L. (2009). Sex differences in the relation between prenatal maternal emotional complaints and child outcome. *Early Human Development*, 85, 319-324. doi:10.1016/j.earlhumdev.2008.12.009
- de Kloet, E. R., Oitzl, M. S., & Joëls, M. (1999). Stress and cognition: Are corticosteroids good or bad guys? *Trends in Neurosciences*, 22, 422-426.
- Demir-Lira, Ö. E., Prado, J., & Booth, J. R. (2016). Neural correlates of math gains vary depending on parental socioeconomic status (SES). *Frontiers in Psychology*, 7, 1-12. doi:10.3389/fpsyg.2016.00892
- Dennis, C.-L., Falah-Hassani, K., & Shiri, R. (2017). Prevalence of antenatal and postnatal anxiety: Systematic review and meta-analysis. *British Journal of Psychiatry*, 210, 315-323. doi:10.1192/bjp.bp.116.187179
- Diamond, A. (2011). Biological and social influences on cognitive control processes dependent on prefrontal cortex. *Progress in Brain Research*, 189, 319-339. doi:10.1016/B978-0-444-53884-0.00032-4
- Diamond, A. (2013). Executive functions. *Annual Review of Psychology*, 64, 135-168. doi:10.1146/annurev-psych-113011-143750
- Diamond, A., Barnett, W. S., Thomas, J., & Munro, S. (2007). Preschool program improves cognitive control. *Science*, 318, 1387-1388. doi:10.1126/science.1151148
- Diamond, A., & Lee, K. (2011). Interventions shown to aid executive function development in children 4 to 12 years old. *Science*, 333, 959-964. doi:10.1126/science.1204529
- Diamond, A., & Ling, D. S. (2016). Conclusions about interventions, programs, and approaches for improving executive functions that appear justified and those that, despite much hype, do not. *Developmental Cognitive Neuroscience*, 18, 34-48. doi:10.1016/j.dcn.2015.11.005
- Diego, M. A., Field, T., Hernandez-Reif, M., Cullen, C., Schanberg, S., & Kuhn, C. (2004). Prepartum, postpartum, and chronic depression effects on newborns. *Psychiatry*, 67, 63-80. doi:10.1521/psyc.67.1.63.31251
- DiPietro, J. A., Novak, M. F., Costigan, K. A., Atella, L. D., & Reusing, S. P. (2006). Maternal psychological distress during pregnancy in relation to child development at age two. *Child Development*, 77, 573-587. doi:10.1111/j.1467-8624.2006.00891.x
- Donzella, B., Gunnar, M. R., Krueger, W. K., & Alwin, J. (2000). Cortisol and vagal tone responses to competitive challenge in preschoolers: Associations with temperament. *Developmental Psychobiology*, 37, 209-220. doi:10.1002/1098-2302(2000)37:4<209::AID-DEV1>3.0.CO;2-S
- Doom, J. R., Cicchetti, D., & Rogosch, F. A. (2014). Longitudinal patterns of cortisol regulation differ in maltreated and nonmaltreated children. *Journal of the American Academy of Child & Adolescent Psychiatry*, 53, 1206-1215. doi:10.1016/j.jaac.2014.08.006
- Dunkel Schetter, C. (2011). Psychological science on pregnancy: Stress processes, biopsychosocial models, and emerging research issues. *Annual Review of Psychology*, 62, 531-558. doi:10.1146/annurev.psych.031809.130727
- Dunkel Schetter, C., & Tanner, L. (2012). Anxiety, depression and stress in pregnancy: Implications for mothers, children, research, and practice. *Current Opinion in Psychiatry*, 25, 141-148. doi:10.1097/YCO.0b013e3283503680
- Ellis, B. J., Boyce, W. T., Belsky, J., Bakermans-Kranenburg, M. J., & van IJzendoorn, M. H. (2011). Differential susceptibility to the environment: An evolutionary-neurodevelopmental theory. *Development and Psychopathology*, 23, 7-28. doi:10.1017/S0954579410000611
- Entringer, S., Buss, C., Kumsta, R., Hellhammer, D. H., Wadhwa, P. D., & Wuest, S. (2009). Prenatal psychosocial stress exposure is associated with subsequent working memory performance in young women. *Behavioral Neuroscience*, 123, 886-893. doi:10.1037/a0016265
- Entringer, S., Kumsta, R., Hellhammer, D. H., Wadhwa, P. D., & Wust, S. (2009). Prenatal exposure to maternal psychosocial stress and HPA axis regulation in young adults. *Hormones and Behavior*, 55, 292-298. doi:10.1016/j.yhbeh.2008.11.006
- Felitti, V. J., & Anda, R. F. (2010). The relationship of adverse childhood experiences to adult medical disease, psychiatric disorders and sexual behavior: Implications for healthcare. In R. A. Lanius, E. Vermetten, & C. Pain (Eds.), *The impact of early life trauma on health and disease* (pp. 77-87). Cambridge: Cambridge University Press.
- Felitti, V. J., Anda, R. F., Nordenberg, D., Williamson, D. F., Spitz, A. M., Edwards, V., . . . Marks, J. S. (1998). Relationship of childhood abuse and household dysfunction to many of the leading causes of death in adults: The adverse childhood experiences (ACE) study. *American Journal of Preventive Medicine*, 14, 245-258. doi:10.1016/S0749-3797(98)00017-8
- Frodl, T., & O'Keane, V. (2013). How does the brain deal with cumulative stress? A review with focus on developmental stress, HPA axis function and hippocampal structure in humans. *Neurobiology of Disease*, 52, 24-37. doi:10.1016/j.nbd.2012.03.012
- Fuster, J. M. (2008). *The prefrontal cortex* (4th ed.). London: Academic Press.
- Glover, V. (2011). Annual Research Review: Prenatal stress and the origins of psychopathology: An evolutionary perspective. *Journal of Child Psychology and Psychiatry*, 52, 356-367. doi:10.1111/j.1469-7610.2011.02371.x
- Glover, V. (2015). Prenatal stress and its effects on the fetus and the child: Possible underlying biological mechanisms. In M. C. Antonelli (Ed.), *Perinatal programming of neurodevelopment* (pp. 269-283). New York: Springer.
- Glover, V., & Hill, J. (2012). Sex differences in the programming effects of prenatal stress on psychopathology and stress responses: An evolutionary perspective. *Physiology and Behavior*, 106, 736-740. doi:10.1016/j.physbeh.2012.02.011
- Glover, V., O'Connor, T. G., & O'Donnell, K. (2010). Prenatal stress and the programming of the HPA axis. *Neuroscience and Biobehavioral Reviews*, 35, 17-22. doi:10.1016/j.neubiorev.2009.11.008
- Gluckman, P., & Hanson, M. (2005). *The fetal matrix: Evolution, development and disease*. New York: Cambridge University Press.
- Grant, K. A., McMahon, C., Austin, M. P., Reilly, N., Leader, L., & Ali, S. (2009). Maternal prenatal anxiety, postnatal caregiving and infants' cortisol responses to the still-face procedure. *Developmental Psychobiology*, 51, 625-637. doi:10.1002/dev.20397

- Grunau, R. E., Cepeda, I. L., Chau, C. M. Y., Brummelte, S., Weinberg, J., Lavoie, P., . . . Turvey, S. E. (2013). Neonatal pain-related stress and NFKBIA genotype are associated with altered cortisol levels in preterm boys at school age. *PLOS ONE*, *8*, e73926. doi:10.1371/journal.pone.0073926
- Grunau, R. E., Haley, D. W., Whitfield, M. F., Weinberg, J., Yu, W., & Thiesen, P. (2007). Altered basal cortisol levels at 3, 6, 8 and 18 months in infants born extremely low gestational age. *Journal of Pediatrics*, *150*, 151–156. doi:10.1016/j.jpeds.2006.10.053
- Gunnar, M. R., & Quevedo, L. (2007). The neurobiology of stress and development. *Annual Review of Psychology*, *58*, 145–173. doi:10.1146/annurev.psych.58.110405.085605
- Gutting, B. M., de Weerth, C., & Buitelaar, J. K. (2004). Maternal prenatal stress and 4–6 year old children's salivary cortisol concentrations pre- and post-vaccination. *Stress*, *7*, 257–260. doi:10.1080/10253890500044521
- Gutting, B. M., de Weerth, C., & Buitelaar, J. K. (2005). Prenatal stress and children's cortisol reaction to the first day of school. *Psychoneuroendocrinology*, *30*, 541–549. doi:10.1016/j.psneuen.2005.01.002
- Hamilton, M. (1960). A rating scale for depression. *Journal of Neurology, Neurosurgery & Psychiatry*, *23*, 56–62. doi:10.1136/jnnp.23.1.56
- Hayes, A. F. (2013). *Introduction to mediation, moderation, and conditional process analysis: A regression-based approach*. New York: Guilford Press.
- Hayes, A. F., & Rockwood, N. J. (in press). Regression-based statistical mediation and moderation analysis in clinical research: Observations, recommendations, and implementation. *Behaviour Research and Therapy*. doi:10.1016/j.brat.2016.11.00
- Hostinar, C. E., & Gunnar, M. (2013). The developmental effects of early life stress: An overview of current theoretical frameworks. *Current Directions in Psychological Science*, *22*, 400–406. doi:10.1177/0963721413488889
- Hughes, C. (2011). Changes and challenges in 20 years of research into the development of executive functions. *Infant and Child Development*, *20*, 251–271. doi:10.1002/icd.736
- Johnson, A. C. (2015). Developmental pathways to attention-deficit/hyperactivity disorder and disruptive behavior disorders: Investigating the impact of the stress response on executive functioning. *Clinical Psychology Review*, *36*, 1–12. doi:10.1016/j.cpr.2014.12.001
- Kudielka, B. M., & Kirschbaum, C. (2005). Sex differences in HPA axis responses to stress: A review. *Biological Psychology*, *69*, 113–132. doi:10.1016/j.biopsycho.2004.11.009
- Leight, K. L., Fitelson, E. M., Weston, C. A., & Wisner, K. L. (2010). Childbirth and mental disorders. *International Review of Psychiatry*, *22*, 453–471. doi:10.3109/09540261.2010.514600
- Ling, D. S., Kelly, M. K., & Diamond, A. (2016). Human-animal interaction and the development of cognitive control (executive functions). In L. S. Freund, S. McCune, L. Esposito, N. R. Gee, & P. McCardle (Eds.), *Social neuroscience of human-animal interaction*. New York: American Psychological Association Press.
- Lupien, S. J., Gillin, C. J., & Hauger, R. L. (1999). Working memory is more sensitive than declarative memory to the acute effects of corticosteroids: A dose-response study in humans. *Behavioral Neuroscience*, *113*, 420–430. doi:10.1037/0735-7044.113.3.420
- Lupien, S. J., Maheu, F., Tu, M., Fiocco, A., & Schramek, T. E. (2007). The effects of stress and stress hormones on human cognition: Implications for the field of brain and cognition. *Brain and Cognition*, *65*, 209–237. doi:10.1016/j.bandc.2007.02.007
- Marcus, S. M., Flynn, H. A., Blow, F. C., & Barry, K. L. (2003). Depressive symptoms among pregnant women screened in obstetrics settings. *Journal of Women's Health*, *12*, 373–380. doi:10.1089/154099903765448880
- Mathews, S. G., & Phillips, D. I. (2011). Minireview: Transgenerational inheritance of the stress response: A new frontier in stress research. *Endocrinology*, *151*, 7–13. doi:10.1210/en.2009-0916
- McEwen, B. S. (2000). Allostasis and allostatic load: Implications for neuropsychopharmacology. *Neuropsychopharmacology*, *22*, 108–124. doi:10.1016/S0893-133X(99)00129-3
- McEwen, B. S., & Morrison, J. H. (2013). The brain on stress: Vulnerability and plasticity of the prefrontal cortex over the life course. *Neuron*, *79*, 16–29. doi:10.1016/j.neuron.2013.06.028
- McEwen, B. S., & Wingfield, J. C. (2003). The concept of allostasis in biology and biomedicine. *Hormones & Behavior*, *43*, 2–16. doi:10.1016/S0018-506X(02)00024-7
- Mennes, M., Stiers, P., Lagae, L., & van den Bergh, B. R. H. (2006). Long-term cognitive sequelae of antenatal maternal anxiety: Involvement of the orbitofrontal cortex. *Neuroscience and Biobehavioral Reviews*, *30*, 1078–1086. doi:10.1016/j.neubiorev.2006.04.003
- Miller, A. H., & Raison, C. L. (2016). The role of inflammation in depression: From evolutionary imperative to modern treatment target. *Nature Reviews Immunology*, *16*, 22–34. doi:10.1038/nri.2015.5
- Miller, G. E., Chen, E., & Zhou, E. S. (2007). If it goes up, must it come down? Chronic stress and the hypothalamic-pituitary-adrenocortical axis in humans. *Psychological Bulletin*, *133*, 25–45. doi:10.1037/0033-2909.133.1.25
- Moe, V., & Slining, K. (2001). Children prenatally exposed to substances: Gender-related differences in outcome from infancy to 3 years of age. *Infant Mental Health Journal*, *22*, 334–350. doi.org/10.1002/imhj.1005
- Neuenschwander, R., & Oberlander, T. F. (2017). Developmental origins of self-regulation: Prenatal maternal stress and psychobiological development during childhood. In K. Deater-Deckard & R. K. Panneton (Eds.), *Parental stress and early child development: Adaptive and maladaptive outcomes* (pp. 127–156). New York: Springer.
- Noble, K. G., McCandliss, B. D., & Farah, M. J. (2007). Socioeconomic gradients predict individual differences in neurocognitive abilities. *Developmental Science*, *10*, 464–480. doi:10.1111/j.1467-7687.2007.00600.x
- Oberlander, T. F., Warburton, W., Misri, S., Aghajanian, J., & Hertzman, C. (2006). Neonatal outcomes after prenatal exposure to selective serotonin reuptake inhibitor antidepressants and maternal depression using population-based linked health data. *Archives of General Psychiatry*, *63*, 898–906. doi:10.1001/archpsyc.63.8.898
- Oberlander, T. F., Weinberg, J., Papsdorf, M., Grunau, R., Misri, S., & Devlin, A. M. (2008). Prenatal exposure to maternal depression, neonatal methylation of human glucocorticoid receptor gene (NR3C1) and infant cortisol stress responses. *Epigenetics*, *3*, 97–106. doi:10.4161/epi.3.2.6034
- Obradović, J., Portilla, X. A., & Ballard, P. J. (2016). Biological sensitivity to family income: Differential effects on early executive functioning. *Child Development*, *87*, 374–384. doi:10.1111/cdev.12475
- O'Connor, T. G., Ben-Shlomo, Y., Heron, J., Golding, J., Adams, D., & Glover, V. (2005). Prenatal anxiety predicts individual differences in cortisol in pre-adolescent children. *Biological Psychiatry*, *58*, 211–217. doi:10.1016/j.biopsycho.2005.03.032
- O'Donnell, K. J., Glover, V., Jenkins, J., Browne, D., Ben-Shlomo, Y., Golding, J., & O'Connor, T. G. (2013). Prenatal maternal mood is associated with altered diurnal cortisol in adolescence. *Psychoneuroendocrinology*, *38*, 1630–1638. doi:10.1016/j.psneuen.2013.01.008
- Pauli-Pott, U., & Becker, K. (2011). Neuropsychological basic deficits in pre-schoolers at risk for ADHD: A meta-analysis. *Clinical Psychology Review*, *31*, 626–637. doi:10.1016/j.cpr.2011.02.005
- Pearson, R. M., Bornstein, M. H., Cordero, M., Scerif, G., Mahedy, L., Evans, J., . . . Stein, A. (2016). Maternal perinatal mental health and offspring academic achievement at age 16: The mediating role of childhood executive function. *Journal of Child Psychology and Psychiatry*, *57*, 491–501. doi:10.1111/jcpp.12483
- Pluess, M., Velders, F. P., Belsky, J., van IJzendoorn, M. H., Bakermans-Kranenburg, M. J., Jaddoe, V. W. V., . . . Tiemeier, H. (2011). Serotonin transporter polymorphism moderates effects of prenatal maternal anxiety on infant negative emotionality. *Biological Psychiatry*, *69*, 520–525. doi:10.1016/j.biopsycho.2010.10.006
- Pruessner, J. C., Kirschbaum, C., Meinlschmid, G., & Hellhammer, D. H. (2003). Two formulas for computation of the area under the curve represent measures of total hormone concentration versus time-dependent change. *Psychoneuroendocrinology*, *28*, 916–931. doi:10.1016/S0306-4530(02)00108-7
- Ramos, B. P., & Arnsten, A. F. (2007). Adrenergic pharmacology and cognition: Focus on the prefrontal cortex. *Pharmacology and Therapeutics*, *113*, 523–536. doi:10.1016/j.pharmthera.2006.11.006
- Rice, F., Harold, G., Boivin, J., van den Bree, M., Hay, D. F., & Thapar, A. (2010). The links between prenatal stress and offspring development and psychopathology: Disentangling environmental and inherited influences. *Psychological Medicine*, *40*, 335–345. doi:10.1017/S0033291709005911
- Sanchez, M. M., Young, L. J., Plotsky, P. M., & Insel, T. R. (2000). Distribution of corticosteroid receptors in the rhesus brain. *Journal of Neuroscience*, *20*, 4657–4658.
- Sandman, C. A., Buss, C., Head, K., & Davis, E. P. (2015). Fetal exposure to maternal depressive symptoms is associated with cortical thickness in late childhood. *Biological Psychiatry*, *77*, 324–334. doi:10.1016/j.biopsych.2014.06.025

- Sandman, C. A., Davis, E., & Glynn, L. (2012). Prescient human fetuses thrive. *Psychological Science*, *23*, 93–100. doi:10.1177/0956797611422073
- Seckl, J. R., & Meaney, M. J. (1993). Early life events and later development of ischaemic heart disease. *Lancet*, *342*, 1236. doi:10.1016/0140-6736(93)92215-F
- Shacka, J. J., Fennell, O. B., & Robinson, S. E. (1997). Prenatal nicotine sex-dependently alters agonist-induced locomotion and stereotypy. *Neurotoxicology and Teratology*, *19*, 467–476. doi:10.1016/S0892-0362(97)00063-9
- Shirtcliff, E. A., Granger, D. A., Booth, A., & Johnson, D. (2005). Low salivary cortisol levels and externalizing behavior problems in youth. *Development and Psychopathology*, *17*, 167–184. doi:10.1017/S09545794050050091
- Simons, S. S. H., Beijers, R., Cillessen, A. H. N., & de Weerth, C. (2015). Development of the cortisol circadian rhythm in the light of stress early in life. *Psychoneuroendocrinology*, *62*, 292–300. doi:10.1016/j.psyneuen.2015.08.024
- Snyder, H. R. (2013). Major depressive disorder is associated with broad impairments on neuropsychological measures of executive function: A meta-analysis and review. *Psychological Bulletin*, *139*, 81–132. doi:10.1037/a0028727
- Spinrad, T. L., Eisenberg, N., Granger, D. A., Eggum, N. D., Sallquist, J., Haugen, R., . . . Hofer, C. (2009). Individual differences in preschoolers' salivary cortisol and alpha-amylase reactivity: Relations to temperament and maladjustment. *Hormones and Behavior*, *56*, 133–139. doi:10.1016/j.yhbeh.2009.03.020
- Stroud, L. R., Salovey, P., & Epel, E. S. (2002). Sex differences in stress responses: Social rejection versus achievement stress. *Biological Psychiatry*, *52*, 318–327. doi:10.1016/S0006-3223(02)01333-1
- Suurland, J., van der Heijden, K. B., Smaling, H. J. A., Huijbregts, S. C. J., van Goozen, S. H. M., & Swaab, H. (2017). Infant autonomic nervous system response and recovery: Associations with maternal risk status and infant emotion regulation. *Development and Psychopathology*, *29*, 759–773. doi:10.1017/S0954579416000456
- Talge, N. M., Neal, C., & Glover, V. (2007). Antenatal maternal stress and long-term effects on child neurodevelopment: How and why? *Journal of Child Psychology and Psychiatry and Allied Disciplines*, *48*, 245–261. doi:10.1111/j.1469-7610.2006.01714.x
- Tollenaar, M. S., Beijers, R., Jansen, J., Riksen-Walraven, J. M. A., & de Weerth, C. (2011). Maternal prenatal stress and cortisol reactivity to stressors in human infants. *Stress*, *14*, 53–65. doi:10.3109/10253890.2010.499485
- van den Bergh, B. R. H., Mennes, M., Oosterlaan, J., Stevens, V., Stiers, P., Marcoen, A., & Lagae, L. (2005). High antenatal maternal anxiety is related to impulsivity during performance on cognitive tasks in 14- and 15-year-olds. *Neuroscience and Biobehavioral Reviews*, *29*, 259–269. doi:10.1016/j.neubiorev.2004.10.010
- van den Bergh, B. R. H., Mennes, M., Stevens, V., van der Meere, J., Bögner, N., Stiers, P., . . . Lagae, L. (2006). ADHD deficit as measured in adolescent boys with a continuous performance task is related to antenatal maternal anxiety. *Pediatric Research*, *59*, 78–82. doi:10.1203/01.pdr.0000191143.75673.52
- van den Bergh, B. R. H., Mulder, E. J., Mennes, M., & Glover, V. (2005). Antenatal maternal anxiety and stress and the neurobehavioural development of the fetus and child: Links and possible mechanisms: A review. *Neuroscience and Biobehavioral Reviews*, *29*, 237–258. doi:10.1016/j.neubiorev.2004.10.007
- van den Bergh, B. R. H., van Calster, B., Smits, T., van Huffel, S., & Lagae, L. (2008). Antenatal maternal anxiety is related to HPA-axis dysregulation and self-reported depressive symptoms in adolescence: A prospective study on the fetal origins of depressed mood. *Neuropsychopharmacology*, *33*, 536–545. doi:10.1038/sj.npp.1301450
- Vänskä, M., Punamäki, R.-L., Lindblom, J. K., Tolvanen, A., Flykt, M., Unkila-Kallio, L., . . . Tiitinen, A. (2015). Timing of early maternal mental health and child cortisol regulation. *Infant and Child Development*, *25*, 461–483. doi:10.1002/icd.1948
- Vijayraghavan, S., Wang, M., Birnbaum, S. G., Williams, G. V., & Arnsten, A. F. T. (2007). Inverted-U dopamine D1 receptor actions on prefrontal neurons engaged in working memory. *Nature Neuroscience*, *10*, 376–384. doi:10.1038/nn1846
- Wagner, S., Cepeda, I. L., Krieger, D., Maggi, S., D'Angiulli, A., Weinberg, J., & Grunau, R. E. (2015). Higher cortisol is associated with poorer executive functioning in preschool children: The role of parenting stress, parent coping and quality of daycare. *Child Neuropsychology*, *3*, 1–17.
- Weikum, W. M., Brain, U., Chau, C. M. Y., Grunau, R. E., Boyce, T., Diamond, A., & Oberlander, T. F. (2013). Prenatal serotonin reuptake inhibitor (SRI) antidepressant exposure and serotonin transporter promoter genotype (SLC6A4) influence executive functions at 6 years of age. *Frontiers in Cellular Neuroscience*, *7*, 1–12. doi:10.3389/fncel.2013.00180
- Weinstock, M. (2008). The long-term behavioural consequences of prenatal stress. *Neuroscience and Biobehavioral Reviews*, *32*, 1073–1086. doi:10.1016/j.neubiorev.2008.03.002
- Wright, A., & Diamond, A. (2014). An effect of inhibitory load in children while keeping working memory load constant. *Frontiers in Psychology*, *5*, 213. doi:10.3389/fpsyg.2014.00213
- Zaitchik, D., Iqbal, Y., & Carey, S. (2014). The effect of executive function on biological reasoning in young children: An individual differences study. *Child Development*, *85*, 160–175. doi:10.1111/cdev.12145
- Zhang, T. Y., Chretien, P., Meaney, M. J., & Gratton, A. (2005). Influence of naturally occurring variations in maternal care on prepulse inhibition of acoustic startle and the medial prefrontal cortical dopamine response to stress in adult rats. *Journal of Neuroscience*, *25*, 1493–1502. doi:10.1523/JNEUROSCI.3293-04.2005
- Zijlmans, M. A., Riksen-Walraven, J. M., & de Weerth, C. (2015). Associations between maternal prenatal cortisol concentrations and child outcomes: A systematic review. *Neuroscience and Biobehavioral Reviews*, *53*, 1–24. doi:10.1016/j.neubiorev.2015.02.015