



Preston J. Boyer¹ , Joshua A. Yell², Jennifer G. Andrews³ and Michael D. Seckeler³ 

Original Article

Cite this article: Boyer PJ, Yell JA, Andrews JG, and Seckeler MD (2020) Anxiety reduction after pre-procedure meetings in patients with CHD. *Cardiology in the Young* 30: 991–994. doi: [10.1017/S1047951120001407](https://doi.org/10.1017/S1047951120001407)

Received: 11 March 2020
Accepted: 10 May 2020
First published online: 5 June 2020

Keywords:

CHD; cardiac catheterisation; anxiety; three-dimensional printing; education of patients

Author for correspondence:

Michael D. Seckeler, MD, MSc, Department of Pediatrics (Cardiology), The University of Arizona, 1501 N. Campbell Ave, PO Box 245073, Tucson, AZ 85724, USA. Tel: +1 520 626 6508; Fax: +1 520 626 6571. E-mail: mseckeler@peds.arizona.edu

¹University of Arizona, Department of Pediatrics, Tucson, AZ, USA; ²University of Arizona, College of Medicine, Tucson, AZ, USA and ³University of Arizona, Department of Pediatrics, Division of Cardiology, Tucson, AZ, USA

Abstract

Background: Cardiac catheterisations for CHD produce anxiety for patients and families. Current strategies to mitigate anxiety and explain complex anatomy include pre-procedure meetings and educational tools (cardiac diagrams, echocardiograms, imaging, and angiography). More recently, three-dimensionally printed patient-specific models can be added to the armamentarium. The purpose of this study was to evaluate the efficacy of pre-procedure meetings and of different educational tools to reduce patient and parent anxiety before a catheterisation. **Methods:** Prospective study of patients ≥ 18 and parents of patients < 18 scheduled for clinically indicated catheterisations. Patients completed online surveys before and after meeting with the interventional cardiologist, who was blinded to study participation. Both the pre- and post-meeting surveys measured anxiety using the State-Trait Anxiety Inventory. In addition, the post-meeting survey evaluated the subjective value (from 1 to 4) of individual educational tools: physician discussion, cardiac diagrams, echocardiograms, prior imaging, angiograms and three-dimensionally printed cardiac models. Data were compared using paired t-tests. **Results:** Twenty-three patients consented to participate, 16 had complete data for evaluation. Mean State-Trait Anxiety Inventory scores were abnormally elevated at baseline and decreased into the normal range after the pre-procedure meeting (39.8 versus 31, $p = 0.008$). Physician discussion, angiograms, and three-dimensional models were reported to be most effective at increasing understanding and reducing anxiety. **Conclusion:** In this pilot study, we have found that pre-catheterisation meetings produce a measurable decrease in patient and family anxiety before a procedure. Discussions of the procedure, angiograms, and three-dimensionally printed cardiac models were the most effective educational tools.

Cardiac catheterisations have long been recognised as a significant source of anxiety for patients and families. While the procedures are less invasive than surgery, cardiac catheterisations still incite a great deal of fear. Uncertainty of the outcome and fear of unfavourable findings, pain, death, and the need for further procedures all contribute to increased patient anxiety prior to a catheterisation procedure.^{1–5} This is especially true for children and adults with CHD.^{5–7} The burden of CHD often includes multiple high-risk procedures, frequent and extended hospitalisations, social isolation, and economic instability. Due to the complexity, chronicity, and uncertainty that accompanies living with CHD, many of these patients and their families experience elevated levels of anxiety at baseline.^{8–10} These baseline symptoms of anxiety and depression can be exacerbated by an upcoming cardiac catheterisation, which has been shown to increase morbidity and mortality associated with procedural interventions.¹¹ Previous studies in the general adult population have examined multiple interventions targeted at lowering the patient and family's anxiety prior to a cardiac catheterisation, ranging from educational approaches to sensory and therapy-based approaches.^{12–14} These interventions often take place during a pre-procedure meeting to obtain informed consent, which alone has been shown to improve levels of anxiety.¹⁵ There is still a paucity of literature regarding the effectiveness of these interventions on children and adults with CHD undergoing cardiac catheterisations. Additionally, while there are a variety of educational tools that have been proposed and studied in isolation in the adult population, there are limited data regarding the comparative effectiveness of different educational tools and methods. In recent years, as part of an increase in personalised medicine, patient-specific three-dimensional models have increasingly been utilised across many medical specialties. A unique advantage of these models is the ability to generate a physical representation of complex anatomy which can highlight the anatomic region of interest for the healthcare provider and the patient. The benefits of these models have been evidenced in pre-procedure planning, resident and medical education, as well as patient education.^{16–19} The purpose of this study was to evaluate patient and family anxiety before and after a pre-procedure meeting and qualitatively assesses the educational methods used in discussing complex anatomy and procedures.

Material and methods

Ethical approval was obtained through the University of Arizona Institutional Review Board. Parents of children with CHD and adults with CHD who were scheduled to undergo clinically indicated cardiac catheterisations from October 2017 through March 2019 were invited to participate in the study. Standard practice at our institution for congenital cardiac catheterisations includes an in-person pre-procedural meeting with the attending congenital interventional cardiologist. During these meetings, the procedure is discussed in detail, including the patient's individual anatomy, the reason for the procedure, and any anticipated interventions, and informed consent is obtained. During each meeting, the interventionalist uses cardiac diagrams as well as any available additional imaging, including echocardiograms, MRI or CT images, angiograms (from the patient's prior cardiac catheterisations or those of similar anatomy), and a three-dimensionally printed model of the patient's heart defect. Parents of patients <18 and patients ≥18 who provided informed consent to participate completed pre- and post-meeting online surveys to evaluate their state of anxiety before and after the meeting, and to examine their opinion about the educational methods used. Data collected included demographics, severity of CHD (simple, moderate, severe, or complex single ventricle), and whether the planned procedure was diagnostic or interventional. The attending interventionalist who conducted the meetings was blinded to study participation.

Both the pre- and post-meeting surveys measured anxiety using the six-item Short Form State-Trait Anxiety Inventory.^{20,21} The State-Trait Anxiety Inventory scores range from 20 to 80, with higher scores indicating higher levels of anxiety. A normal score is defined as 34–36 for non-psychiatric patients, with scores above 38 suggesting significantly elevated anxiety.^{22,23} In addition, the post-meeting survey evaluated the subjective value (from 1 to 4, least to most) of each educational tool for increasing understanding and reducing anxiety. Exclusion criteria included: emergent cardiac catheterisations, patient refusal of a pre-catheterisation meeting, no appropriate three-dimensionally printed model, inability to complete the English language survey, incomplete survey data, and refusal to participate in the study. Data were compared using paired t-tests.

Results

During the study period, there were 135 congenital cardiac catheterisations performed at our institution. Of these, 23 met inclusion criteria and consented to participate in the study and 16 completed both the pre- and post-meeting surveys (Fig 1). Demographics are summarised in Table 1. The mean State-Trait Anxiety Inventory score was markedly elevated before the meeting and returned to the normal range after the meeting (39.8 versus 31.0, $p = 0.008$) (Fig 2). There were no significant differences in State-Trait Anxiety Inventory scores or a score reduction between different CHD complexities. There were no differences in State-Trait Anxiety Inventory score based on age, race, or the primary respondent (patient versus parent).

Subjective assessment of the educational tools for increasing understanding and reducing anxiety showed that physician description, angiograms, and three-dimensionally printed models were the most well-received (3.87 ± 0.34 , 3.75 ± 0.44 , and 3.71 ± 0.77 , respectively), while cardiac diagrams, echocardiograms, and CT/MRI scans were rated lower (3.59 ± 0.63 , 3.43 ± 0.94 , and 3.20 ± 1.10 , respectively) (Fig 3).

Table 1. Demographics of survey patients

Characteristic	
Age, mean \pm sd, (y)	36.8 \pm 12.3
Female, n (%)	7 (43.8)
CHD severity, n (%):	
Simple	5 (31.3)
Moderate	2 (12.5)
Complex	3 (18.8)
Complex single ventricle	5 (31.3)
Procedure type, n (%):	
Diagnostic	1 (6.3)
Interventional	13 (81.3)
Unknown	3 (12.5)
Race/Ethnicity, n (%):	
Hispanic	7 (43.8)
Non-Hispanic White	6 (37.5)
Other	3 (18.8)

sd=standard deviation; y=years.




Figure 1. Flow diagram for study enrollment.




Figure 2. Change in anxiety from pre- to post-meeting. State-Trait Anxiety Inventory scores for anxiety levels from pre- and post-meeting surveys for the entire study group. Individual responses are shown in dashed grey, and the overall mean is shown in bold black. The normal range of anxiety levels in a non-psychiatric population is shown between the dotted lines.

Discussion

In this pilot study of children and adults with CHD undergoing cardiac catheterisation, we found that patients and their families experience a high level of anxiety prior to a cardiac catheterisation and that interventions including a pre-procedure meeting with the interventional cardiologist and three-dimensionally printed models help to reduce their anxiety to normal levels. Recognising the




Figure 3. Average subjective rating of each teaching method. The y-axis shows the average patient rating for each of the six teaching methods evaluated.

degree of anxiety surrounding, these procedures as well as the utility of three-dimensionally printed models to explain complex CHD anatomy will be important as patients with CHD undergo higher numbers of increasingly complex interventions.

The notion of informed consent gives interventional cardiologists opportunities not only to ensure patient's agreement with planned interventions but also to educate and prepare patients for their upcoming procedure. This has been implemented in a variety of ways in the fields of adult and paediatric cardiology, from paper handouts to multimedia presentations.^{7,12,24,25} In our experience, it is common for paediatric cardiologists to offer patients and families some type of in-person engagement prior to the procedure. It has previously been demonstrated that patients experience notably high levels of anxiety prior to meeting with a member of the paediatric cardiology team, levels even above that which would be expected for the anticipation of an upcoming procedure.⁵ It is important to know if the pre-procedural face-to-face informational meetings are effective at reducing the elevated anxiety experienced by patients and families. In addition, there are a number of educational tools that can be used to help explain complex anatomy and pathophysiology, including diagrams, echocardiograms, CT/MR imaging, and angiography. More recently, three-dimensionally printed, patient-specific models can be added to the armamentarium, which have been subjectively well-received by patients and physicians as educational tools that can aid comprehension.^{26–28} Better understanding of the effect of pre-procedural meetings on patient/parent anxiety, as well as increased awareness of the effectiveness of different teaching methods can help interventional cardiologists ensure patient preparedness for procedures and promote improved outcomes.

Anxiety experienced by patients and parents in this population is important to consider as part of their overall health and quality of life. Patients with CHD and their families already experience higher levels of stress, anxiety, and depression than the general population.^{10,29} As our data confirm, baseline anxiety is likely to be further exacerbated in anticipation of a cardiac catheterisation. Higher levels of anxiety have been associated with increased morbidity and mortality for patients undergoing cardiac procedures and have a significant negative impact on the quality of life for

these patients and their families.^{6,11,30} As patients with CHD are able to live longer and healthier lives, and as advances in the field of interventional cardiology continue, these patients are expected to undergo an increasing number of cardiac catheterisations throughout their lifetime. It is important to know what tools are effective in reducing the anxiety associated with these procedures.

In addition to answering questions and obtaining informed consent, an important aspect of a pre-catheterisation meeting is the education for patients and families regarding not only the intended procedure but also the patient's anatomy and physiology. Patients with CHD often have complex anatomy which can be difficult to describe with traditional two-dimensional representations. However, better understanding of their heart disease, especially the anatomy, increases the overall well-being of patients with CHD.³¹ The increasing incorporation of three-dimensionally printed models into medicine is a perfect medium to educate patients with CHD and their families. Our study found that families prefer verbal and tactile educational tools, including three-dimensionally printed models, over standard cardiac imaging. While echocardiography and advanced cardiac imaging are familiar to and useful for cardiologists, the current study suggests that their utility as educational tools for patients is limited.

This study has several limitations. The sample size is small, which may limit the generalisability of the results. However, pre-procedural meetings for cardiac catheterisations are already common among congenital cardiologists, and our study adds important objective data to support the ongoing practice of such meetings. The data were obtained through survey responses, which could be subject to response bias, recall bias, and selection bias; however, the State-Trait Anxiety Inventory assessment is a well-validated tool that is commonly used to evaluate patient anxiety and is designed to minimise bias from self-reported data.

In conclusion, patients with CHD and their families experience abnormally high levels of anxiety prior to cardiac catheterisations. Pre-procedure meetings utilising a variety of educational tools, including three-dimensionally printed models, can measurably reduce this anxiety. Expanding this practice among congenital cardiologists, an important step towards personalised medicine could lead to reduced peri-procedural anxiety and improved outcomes for this complex patient population.

Acknowledgements. None.

Financial support. Dr Seckeler received funding from the Sarver Heart Center Congenital Heart Disease Education Grant and the University of Arizona College of Medicine Vernon & Virginia Furrow Award for Medical Education Research. All other authors report no sources of funding from public, commercial, or not-for-profit sectors.

Conflicts of interest. None.

Ethical standards. The authors assert that all procedures contributing to this work comply with the ethical standards of the relevant national guidelines on human experimentation (United States Title 45 Code of Federal Regulations, Part 46 [45 CFR 46]) and with the Helsinki Declaration of 1975, as revised in 2008, and has been approved by the institutional committees (University of Arizona Institutional Review Board).

References

- Gallagher R, Trotter R, Donoghue J. Preprocedural concerns and anxiety assessment in patients undergoing coronary angiography and percutaneous coronary interventions. *Eur J Cardiovasc Nurs* 2010; 9: 38–44.
- Trotter R, Gallagher R, Donoghue J. Anxiety in patients undergoing percutaneous coronary interventions. *Hear Lung J Acute Crit Care* 2011; 40: 185–192.
- Delewi R, Vlastra W, Rohling WJ, et al. Anxiety levels of patients undergoing coronary procedures in the catheterization laboratory. *Int J Cardiol* 2017; 228: 926–930.
- Pederson C. Effect of imagery on children's pain and anxiety during cardiac catheterization. *J Pediatr Nurs* 1995; 10: 365–374.
- Kobayashi D, Turner DR, Forbes TJ, Aggarwal S. Parental anxiety among children undergoing cardiac catheterisation. *Cardiol Young* 2018; 28: 315–321.
- Üzger A, Başpınar O, Bülbül F, Yavuz S, Kiliç M. Evaluation of depression and anxiety in parents of children undergoing cardiac catheterization. *Türk Kardiyol Dern Ars* 2015; 43: 536–541.
- Rigatelli G, Magro B, Ferro S, et al. Education, and obtaining of informed consent, using multimedia before adults with congenitally malformed hearts are submitted to transcatheter interventions. *Cardiol Young* 2009; 19: 60–63.
- Gupta S, Giuffre RM, Crawford S, Waters J. Covert fears, anxiety and depression in congenital heart disease. *Cardiol Young* 1998; 8: 491–499.
- Gupta S, Mitchell I, Michael Giuffre R, Crawford S. Covert fears and anxiety in asthma and congenital heart disease. *Child Care Health Dev* 2001; 27: 335–348.
- Kovacs AH, Saidi AS, Kuhl EA, et al. Depression and anxiety in adult congenital heart disease: predictors and prevalence. *Int J Cardiol* 2009; 137: 158–164.
- Celano CM, Millstein RA, Bedoya CA, Healy BC, Roest AM, Huffman JC. Association between anxiety and mortality in patients with coronary artery disease: a meta-analysis. *Am Heart J* 2015; 170: 1105–1115.
- Flory J, Emanuel E. Interventions to improve research in informed consent for research. *J Am Med Assoc* 2004; 292: 1593–1601.
- Carroll DL, Malecki-Ketchell A, Astin F. Non-pharmacological interventions to reduce psychological distress in patients undergoing diagnostic cardiac catheterization: a rapid review. *Eur J Cardiovasc Nurs* 2017; 16: 92–103.
- Kendall PC, Williams L, Pechacek TF, Graham LE, Shisslak C, Herzoff N. Cognitive-behavioral and patient education interventions in cardiac catheterization procedures: the Palo Alto medical psychology project. *J Consult Clin Psychol* 1979; 47: 49–58.
- Freeman WR, Pichard AD, Smith H. Effect of informed consent and educational background on patient knowledge, anxiety, and subjective responses to cardiac catheterization. *Cathet Cardiovasc Diagn* 1981; 7: 119–134.
- Teishima J, Takayama Y, Iwaguro S, et al. Usefulness of personalized three-dimensional printed model on the satisfaction of preoperative education for patients undergoing robot-assisted partial nephrectomy and their families. *Int Urol Nephrol* 2018; 50: 1061–1066.
- Bernhard JC, Isotani S, Matsugasumi T, et al. Personalized 3D printed model of kidney and tumor anatomy: a useful tool for patient education. *World J Urol* 2016; 34: 337–345.
- Loke YH, Harahsheh AS, Krieger A, Olivieri LJ. Usage of 3D models of tetralogy of Fallot for medical education: impact on learning congenital heart disease. *BMC Med Educ* 2017; 17: 1–8.
- Jones TW, Seckeler MD. Use of 3D models of vascular rings and slings to improve resident education. *Congenit Heart Dis* 2017; 12: 578–582.
- Marteau TM, Bekker H. The development of a six-item short-form of the state scale of the Spielberger State-Trait Anxiety Inventory (STAI). *Br J Clin Psychol* 1992; 31: 301–306.
- Spielberger CD. *The State-Trait Anxiety Inventory [Test Manual]*. Consulting Psychologists Press, Palo Alto, California, 1970.
- Julian LJ. Measures of anxiety. *Arthritis Care Res* 2011; 63: 1–11.
- Knight RG, Waal Manning HJ, Spears GF. Some norms and reliability data for the State-Trait Anxiety Inventory and the Zung Self-Rating depression scale. *Br J Clin Psychol* 1983; 22: 245–249.
- Wu KL, Chen SR, Ko WC, et al. The effectiveness of an accessibility-enhanced multimedia informational educational programme in reducing anxiety and increasing satisfaction of patients undergoing cardiac catheterisation. *J Clin Nurs* 2014; 23: 2063–2073.
- Harkness K, Morrow L, Smith K, Kiczula M, Arthur HM. The effect of early education on patient anxiety while waiting for elective cardiac catheterization. *Eur J Cardiovasc Nurs* 2003; 2: 113–121.
- Biglino G, Capelli C, Wray J, et al. 3D-manufactured patient-specific models of congenital heart defects for communication in clinical practice: feasibility and acceptability. *BMJ Open* 2015; 5: e007165.
- Biglino G, Koniordou D, Gasparini M, et al. Piloting the use of patient-specific cardiac models as a novel tool to facilitate communication during clinical consultations. *Pediatr Cardiol* 2017; 38: 813–818.
- Lau I, Sun Z. Three-dimensional printing in congenital heart disease: a systematic review. *J Med Radiat Sci* 2018; 65: 226–236.
- Uzark K, Jones K. Parenting stress and children with heart disease. *J Pediatr Heal Care* 2003; 17: 163–168.
- Lawoko S, Soares JFF. Quality of life among parents of children with congenital heart disease, parents of children with other diseases and parents of healthy children. *Qual Life Res* 2003; 12: 655–666.
- Wang Q, Hay M, Clarke D, Menahem S. Associations between knowledge of disease, depression and anxiety, social support, sense of coherence and optimism with health-related quality of life in an ambulatory sample of adolescents with heart disease. *Cardiol Young* 2014; 24: 126–133.