



# Linear independence of series related to the Thue–Morse sequence along powers

Michael Coons and Yohei Tachiya

*Abstract.* The Thue–Morse sequence  $\{t(n)\}_{n \geq 0}$  is the indicator function of the parity of the number of ones in the binary expansion of nonnegative integers  $n$ , where  $t(n) = 1$  (resp.  $= 0$ ) if the binary expansion of  $n$  has an odd (resp. even) number of ones. In this paper, we generalize a recent result of E. Miyanohara by showing that, for a fixed Pisot or Salem number  $\beta > \sqrt{\varphi} = 1.272019\dots$ , the set of the numbers

$$1, \sum_{n \geq 1} \frac{t(n)}{\beta^n}, \sum_{n \geq 1} \frac{t(n^2)}{\beta^n}, \dots, \sum_{n \geq 1} \frac{t(n^k)}{\beta^n}, \dots$$

is linearly independent over the field  $\mathbb{Q}(\beta)$ , where  $\varphi := (1 + \sqrt{5})/2$  is the golden ratio. Our result yields that for any integer  $k \geq 1$  and for any  $a_1, a_2, \dots, a_k \in \mathbb{Q}(\beta)$ , not all zero, the sequence  $\{a_1 t(n) + a_2 t(n^2) + \dots + a_k t(n^k)\}_{n \geq 1}$  cannot be eventually periodic.

## 1 Introduction

Let  $s_2(n)$  denote the number of ones in the binary expansion of  $n$ . The *Thue–Morse sequence*  $\mathbf{t} = \{t(n)\}_{n \geq 0}$  is defined, for  $n \geq 0$ , by  $t(n) = 1$  if  $s_2(n)$  is odd, and  $t(n) = 0$  if  $s_2(n)$  is even. The Thue–Morse sequence is paradigmatic in the areas of complexity and symbolic dynamics, and as such is an object of current interest in a variety of areas. While the sequence goes back at least to the 1851 paper of Prouhet [6], its interest in the context of complexity is usually attributed to Thue [8, 9] who showed that  $\mathbf{t}$  is overlap-free; that is, viewing  $\mathbf{t}$  as a one-sided infinite word  $t(0)t(1)t(2)\dots$ , it contains no subwords of the form  $awawa$ , where  $a \in \{0, 1\}$  and  $w$  is a finite binary word, possibly empty (cf. [1, p. 15]). This shows that  $\mathbf{t}$  contains no three consecutive identical subwords (namely,  $\mathbf{t}$  is cube-free) and consequently  $\mathbf{t}$  is nonperiodic. Thus, we find that the number  $\sum_{n \geq 1} t(n)b^{-n}$  is irrational for any integer  $b \geq 2$ , but more strongly, a result of Mahler [3] provides the transcendence of  $\sum_{n \geq 1} t(n)\alpha^{-n}$  for any algebraic number  $\alpha$  with  $|\alpha| > 1$ .

On the other hand, the Thue–Morse sequence along powers has also been studied by several authors. In 2007, Moshe [5] investigated the subword complexity of the sequence along squares  $\{t(n^2)\}_{n \geq 1}$  (see Section 4 for details) and solved

---

Received by the editors December 20, 2023; revised February 27, 2024; accepted February 29, 2024.

Published online on Cambridge Core March 6, 2024.

This work was supported by the Research Institute for Mathematical Sciences, an International Joint Usage/Research Center located in Kyoto University. This research was partly supported by JSPS KAKENHI Grant Number JP22K03263.

AMS subject classification: 11J72, 11A63, 11B85.

Keywords: Thue–Morse sequence, linear independence, Pisot number, Salem number.



a problem of Allouche and Shallit [1, p. 350] answering that every finite word  $a_0 a_1 \cdots a_{m-1}$  ( $a_i \in \{0, 1\}$ ) of length  $m$  appears in the infinite word  $t(0)t(1)t(4)\cdots$ . Moreover, this result was generalized by Drmota, Mauduit, and Rivat [2] who proved that the sequence  $\{t(n^2)\}_{n \geq 1}$  is normal (to the base 2); that is, for any  $m \geq 1$  and any  $a_0 a_1 \cdots a_{m-1}$  ( $a_i \in \{0, 1\}$ ), we have

$$\lim_{N \rightarrow \infty} \frac{\#\{n < N \mid t(n^2) = a_0, t((n+1)^2) = a_1, \dots, t((n+m-1)^2) = a_{m-1}\}}{N} = \frac{1}{2^m}.$$

As a consequence, the number  $\sum_{n \geq 1} t(n^2)2^{-n}$  is normal in base 2. Recently, Spiegelhofer [7] proved that the sequence along cubes  $\{t(n^3)\}_{n \geq 1}$  is simply normal;

$$\lim_{N \rightarrow \infty} \frac{\#\{n < N \mid t(n^3) = 0\}}{N} = \frac{1}{2}.$$

In this direction, it is expected that the sequence  $\{t(P(n))\}_{n \geq 0}$  is normal for any nonnegative integer-valued polynomial  $P$  of degree at least 3; however, it is unsolved (cf. [2, Conjecture 1]).

A Pisot (resp. Salem) number is an algebraic integer  $\beta > 1$  whose Galois conjugates other than  $\beta$  have moduli less than 1 (resp. less than or equal to 1 and at least one conjugate lies on the unit circle). Recently, Miyanochara [4] showed that if  $\beta$  is a Pisot or Salem number with  $\beta > 2$ , then the number  $\sum_{n \geq 1} t(n^2)\beta^{-n}$  does not belong to the field  $\mathbb{Q}(\beta)$ . Note that his method depends on elementary arguments without the use of the normality of  $\{t(n^2)\}_{n \geq 0}$ . In this paper, we generalize Miyanochara’s results by proving the following theorem. Throughout the paper,  $\varphi := (1 + \sqrt{5})/2$  denotes the golden ratio.

**Theorem 1.1** *Let  $\beta$  be a Pisot or Salem number with  $\beta > \sqrt{\varphi} = 1.272019 \dots$ . Then, for any integer  $k \geq 1$ , the  $k + 1$  numbers*

$$1, \sum_{n \geq 1} \frac{t(n)}{\beta^n}, \sum_{n \geq 1} \frac{t(n^2)}{\beta^n}, \dots, \sum_{n \geq 1} \frac{t(n^k)}{\beta^n}$$

*are linearly independent over the field  $\mathbb{Q}(\beta)$ . In particular, for any integer  $k \geq 1$ , the number  $\sum_{n \geq 1} t(n^k)\beta^{-n}$  does not belong to  $\mathbb{Q}(\beta)$ .*

It should be noted that all Pisot numbers are covered in Theorem 1.1, since the smallest Pisot number is the plastic ratio  $\rho = 1.324717 \dots$ . On the other hand, all Salem numbers are not; for example, the smallest known Salem number is  $\lambda = 1.176280 \dots$  (see Sect. 4 for details). Let  $\beta$  be as in Theorem 1.1. Then, as an immediate corollary of Theorem 1.1, for any nontrivial  $\mathbb{Q}(\beta)$ -linear combination of the Thue–Morse sequence along powers

$$(1.1) \quad s(n) := a_1 t(n) + a_2 t(n^2) + \dots + a_k t(n^k)$$

the number  $\sum_{n \geq 1} s(n)\beta^{-n}$  does not belong to  $\mathbb{Q}(\beta)$ , and hence the sequence  $\{s(n)\}_{n \geq 0}$  cannot be eventually periodic.

The present paper is organized as follows. In Section 2, for each integer  $r = 1, 2, \dots, k$ , we will investigate the appearance of zeros in the sequences defined

by the difference of  $t(n^r)$  and a certain shift  $t((n + n_0)^r)$ . This observation makes it possible to find a good rational approximation to

$$\xi := \sum_{n \geq 1} s(n)\beta^{-n},$$

where the sequence  $\{s(n)\}_{n \geq 1}$  is defined in (1.1). Section 3 is devoted to the proof of Theorem 1.1. In the last Section 4, we will give remarks and further questions related to our results.

## 2 Some useful equalities and inequalities

The first lemma allows us to optimize our choice of rational approximations.

**Lemma 2.1** *For any integer  $k \geq 2$ , there exist positive integers  $m$  and  $n$  such that the system of simultaneous congruences*

$$(2.1) \quad \begin{aligned} X &\equiv 2^{2m-1} - 1 \pmod{2^{2m}}, \\ 3^{k-1}X &\equiv 2^{2n} - 1 \pmod{2^{2n+1}} \end{aligned}$$

has an integer solution.

**Proof** For an even integer  $k \geq 2$ , clearly  $x = 1$  is a solution of (2.1) with  $m = n = 1$ . Thus, let  $k = 2\ell + 1 \geq 3$  be odd. Then there exist an integer  $s \geq 3$  and  $a \in \{0, 1\}$  satisfying

$$3^{k-1} = 9^\ell \equiv 1 + 2^s + a2^{s+1} \pmod{2^{s+2}}.$$

If  $s = 2u + 1 \geq 3$  is odd, we set  $x := 2^{2u+1} - 1 + (1 - a)2^{2u+2}$ , so that

$$\begin{aligned} 3^{k-1}x &\equiv (1 + 2^{2u+1} + a2^{2u+2})(2^{2u+1} - 1 + (1 - a)2^{2u+2}) \pmod{2^{2u+3}} \\ &\equiv 2^{2u+2} - 1 \pmod{2^{2u+3}}, \end{aligned}$$

and hence the integer  $x$  is a solution of (2.1) with  $m = n = u + 1$ . If  $s = 2u \geq 4$  is even, then setting  $x := 2^{2u+1} - 1 + 2^{2u+2}$ , we obtain

$$3^{k-1}x \equiv (1 + 2^{2u})(-1) \equiv 2^{2u} - 1 \pmod{2^{2u+1}},$$

and the integer  $x$  is a solution of (2.1) with  $m = u + 1$  and  $n = u$ . Lemma 2.1 is proved. ■

Let  $x \geq 1$  be an integer solution of the system of congruences (2.1). Then the binary expansions of the integers  $x$  and  $x3^{k-1}$  have the forms

$$(x)_2 = w_1 0 \overbrace{11 \cdots 1}^{2m-1} \quad \text{and} \quad (x3^{k-1})_2 = w_2 0 \overbrace{11 \cdots 1}^{2n},$$

respectively, for some binary words  $w_1$  and  $w_2$ , and hence we obtain the equalities

$$(2.2) \quad t(1 + x) = t(x) \quad \text{and} \quad t(1 + x3^{k-1}) = 1 - t(x3^{k-1}).$$

Let  $k \geq 2$  be an integer, and let  $v(k) := v_2(k)$  be the 2-adic valuation of  $k$ . Fix positive integers  $m, n, x$  (depending only on  $k$ ) as in Lemma 2.1. Since  $k2^{-v(k)}$  is an odd integer, the congruence

$$(2.3) \quad k2^{-v(k)} Y \equiv x \pmod{2^{2m+2n+1}}$$

has an integer solution. Let  $y \leq 2^{2m+2n+1}$  be the least positive integer solution of (2.3).

The remaining lemmas give us quantitative information about our approximations, as well as allowing us to optimize the range of  $\beta$  in Theorem 1.1. Throughout the paper, let  $N$  be a sufficiently large integer. In the following Lemmas 2.2 and 2.3, we investigate the Thue–Morse values of the integers

$$(2.4) \quad (y2^{kN-v(k)+\delta} + j)^k = \sum_{\ell=0}^k \binom{k}{\ell} y^\ell j^{k-\ell} \cdot 2^{(kN-v(k)+\delta)\ell}, \quad \delta \in \{0, 1\}$$

for  $j = 0, 1, \dots, 2^N + 4$  and  $j = 2^N + 2h$  ( $h = 3, 4, \dots, 2^{N-3}$ ). Define

$$A_{\ell,j} := \binom{k}{\ell} y^\ell j^{k-\ell}, \quad \ell = 0, 1, \dots, k, \quad j = 1, 2, \dots, 2^N + 2^{N-2}.$$

When  $1 \leq \ell \leq k$ , we have

$$(2.5) \quad A_{\ell,j} \leq 2^k \cdot (2^{2m+2n+1})^k \cdot (5 \cdot 2^{N-2})^{k-1} < 2^{kN-v(k)+\delta}$$

since  $N$  is sufficiently large. Hence, using (2.4) and (2.5), we obtain

$$(2.6) \quad \begin{aligned} t((y2^{kN-v(k)+\delta} + j)^k) &\equiv t(A_{0,j} + A_{1,j}2^{kN-v(k)+\delta}) + \sum_{\ell=2}^k t(A_{\ell,j}2^{(kN-v(k)+\delta)\ell}) \\ &\equiv t(j^k + zj^{k-1}2^{kN+\delta}) + \sum_{\ell=2}^k t(A_{\ell,j}) \pmod{2}, \end{aligned}$$

where  $z := k2^{-v(k)}y$ . Note that the integers  $A_{\ell,j}$  are independent of  $\delta$ .

**Lemma 2.2** For every integer  $j = 0, 1, \dots, 2^N - 1$  and  $j = 2^N + 2h$  ( $h = 0, 1, \dots, 2^{N-3}$ ), we have

$$(2.7) \quad t((y2^{kN-v(k)} + j)^k) = t((y2^{kN-v(k)+1} + j)^k).$$

**Proof** Let  $\delta \in \{0, 1\}$ . For  $j = 0, 1, \dots, 2^N - 1$ , we have

$$(2.8) \quad j^k < 2^{kN} \quad \text{and} \quad 2^{kN} \mid zj^{k-1}2^{kN+\delta},$$

and moreover, for  $j = 2^N + 2h$  ( $h = 0, 1, \dots, 2^{N-3}$ ),

$$(2.9) \quad j^k \leq (5 \cdot 2^{N-2})^k < 2^{kN+k-1} \quad \text{and} \quad 2^{kN+k-1} \mid z(2^{N-1} + h)^{k-1}2^{kN+k-1+\delta},$$

since  $2^{kN+k-1} \mid zj^{k-1}2^{kN+\delta}$ . Hence, by (2.8) and (2.9),

$$(2.10) \quad t(j^k + zj^{k-1}2^{kN+\delta}) \equiv t(j^k) + t(zj^{k-1}) \pmod{2},$$

so that (2.6) and (2.10) yield

$$(2.11) \quad t\left((y2^{kN-v(k)+\delta} + j)^k\right) \equiv t(j^k) + t(zj^{k-1}) + \sum_{\ell=2}^k t(A_{\ell,j}) \pmod{2}$$

for  $j = 0, 1, \dots, 2^N - 1$  and  $j = 2^N + 2h$  ( $h = 0, 1, \dots, 2^{N-3}$ ). Thus, Lemma 2.2 is proved since the right-hand side in (2.11) is independent of  $\delta$ . ■

Next, we show that the equality (2.7) also holds for  $j = 2^N + 3$ , but not for  $j = 2^N + 1$ .

**Lemma 2.3** *We have*

$$t\left((y2^{kN-v(k)} + 2^N + 1)^k\right) \neq t\left((y2^{kN-v(k)+1} + 2^N + 1)^k\right)$$

and

$$t\left((y2^{kN-v(k)} + 2^N + 3)^k\right) = t\left((y2^{kN-v(k)+1} + 2^N + 3)^k\right).$$

**Proof** Let  $\delta \in \{0, 1\}$  and  $j := 2^N + i$  ( $i = 1, 3$ ). Then we have

$$\begin{aligned} j^k + zj^{k-1}2^{kN+\delta} &= \sum_{\ell=0}^k \binom{k}{\ell} i^{k-\ell} \cdot 2^{N\ell} + \sum_{\ell=0}^{k-1} \binom{k-1}{\ell} zi^{k-1-\ell} \cdot 2^{N(\ell+k)+\delta} \\ &= \sum_{\ell=0}^{k-1} \binom{k}{\ell} i^{k-\ell} \cdot 2^{N\ell} + (1 + zi^{k-1}2^\delta) \cdot 2^{kN} \\ (2.12) \quad &+ \sum_{\ell=k+1}^{2k-1} \binom{k-1}{\ell-k} zi^{2k-\ell-1} \cdot 2^{N\ell+\delta}. \end{aligned}$$

Since the integers

$$B_{\ell,i} := \binom{k}{\ell} i^{k-\ell}, \quad 1 + zi^{k-1}2^\delta, \quad C_{\ell,i} := \binom{k-1}{\ell-k} zi^{2k-\ell-1}$$

are independent of  $N$ , it follows from (2.12) that

$$(2.13) \quad t\left(j^k + zj^{k-1}2^{kN+\delta}\right) \equiv \sum_{\ell=0}^{k-1} t(B_{\ell,i}) + t(1 + zi^{k-1}2^\delta) + \sum_{\ell=k+1}^{2k-1} t(C_{\ell,i}) \pmod{2}.$$

Moreover, combining Lemma 2.1 with (2.3), we obtain

$$\begin{aligned} zi^{k-1} &= k2^{-v(k)} yi^{k-1} \equiv xi^{k-1} \pmod{2^{2m+2n+1}} \\ &\equiv \begin{cases} 2^{2m-1} - 1 \pmod{2^{2m}}, & \text{if } i = 1, \\ 2^{2n} - 1 \pmod{2^{2n+1}}, & \text{if } i = 3, \end{cases} \end{aligned}$$

so that by (2.2)

$$(2.14) \quad t(1 + zi^{k-1}) = \begin{cases} t(zi^{k-1}), & \text{if } i = 1, \\ 1 - t(zi^{k-1}), & \text{if } i = 3. \end{cases}$$

Thus, by (2.6), (2.13), and (2.14), we obtain

$$\begin{aligned} t\left((y2^{kN-v(k)+1} + j)^k\right) - t\left((y2^{kN-v(k)} + j)^k\right) & \\ \equiv t\left(j^k + zj^{k-1}2^{kN+1}\right) - t\left(j^k + zj^{k-1}2^{kN}\right) & \\ \equiv t(1 + 2zi^{k-1}) - t(1 + zi^{k-1}) & \\ \equiv 1 + t(zi^{k-1}) - \begin{cases} t(zi^{k-1}), & \text{if } i = 1, \\ 1 - t(zi^{k-1}), & \text{if } i = 3, \end{cases} & \\ \equiv \begin{cases} 1, & \text{if } j = 2^N + 1, \\ 0, & \text{if } j = 2^N + 3, \end{cases} \pmod{2}, & \end{aligned}$$

which finishes the proof of the Lemma 2.3. ■

Define

$$\lambda := 1 + \frac{1}{2(k-1)} > 1,$$

and let  $[\alpha]$  denote the integral part of the real number  $\alpha$ .

**Lemma 2.4** For every integer  $r = 1, \dots, k - 1$  and  $j = 0, 1, \dots, 2^{\lfloor \lambda N \rfloor}$ , we have

$$(2.15) \quad t\left((y2^{kN-v(k)} + j)^r\right) = t\left((y2^{kN-v(k)+1} + j)^r\right).$$

**Proof** Let  $\delta \in \{0, 1\}$  and  $r, j$  be fixed integers as in the lemma. Then we have

$$(y2^{kN-v(k)+\delta} + j)^r = \sum_{\ell=0}^r \binom{r}{\ell} y^\ell j^{r-\ell} \cdot 2^{(kN-v(k)+\delta)\ell}.$$

Since

$$D_{\ell,j} := \binom{r}{\ell} y^\ell j^{r-\ell} \leq 2^{k-1} \cdot (2^{2m+2n+1})^{k-1} \cdot (2^{\lambda N})^{k-1} < 2^{kN-v(k)+\delta},$$

we obtain

$$t\left((y2^{kN-v(k)+\delta} + j)^r\right) \equiv \sum_{\ell=0}^r t(D_{\ell,j}) \pmod{2},$$

which is independent of  $\delta$ . Lemma 2.4 is proved. ■

### 3 Proof of Theorem 1.1

Let  $\beta$  be a Pisot or Salem number with  $\beta > \sqrt{\varphi} = 1.272019 \dots$ . Suppose to the contrary that there exist an integer  $k \geq 1$  and algebraic numbers  $a_0, a_1, \dots, a_k \in \mathbb{Q}(\beta)$ , not all zero, such that

$$(3.1) \quad a_0 + a_1 \sum_{n \geq 1} \frac{t(n)}{\beta^n} + a_2 \sum_{n \geq 1} \frac{t(n^2)}{\beta^n} + \dots + a_k \sum_{n \geq 1} \frac{t(n^k)}{\beta^n} = 0.$$

We may assume that  $a_0, a_1, \dots, a_k \in \mathbb{Z}[\beta]$  and  $a_k \neq 0$ . As mentioned in Section 1, the number  $\sum_{n \geq 1} t(n)\alpha^{-n}$  is transcendental for any algebraic number  $\alpha$  with  $|\alpha| > 1$ , and thus we have  $k \geq 2$ . Define the sequence  $\{s(n)\}_{n \geq 1}$  by

$$s(n) := a_1 t(n) + a_2 t(n^2) + \dots + a_k t(n^k), \quad n \geq 1,$$

and

$$\xi := \sum_{n \geq 1} \frac{s(n)}{\beta^n}.$$

Note that  $\{s(n)\}_{n \geq 1}$  is bounded and the number  $\xi = -a_0 \in \mathbb{Z}[\beta]$  by (3.1). Let  $v(k), y$  be as in Section 2, and let  $N$  be a sufficiently large integer such that Lemmas 2.2–2.4 all hold. For convenience, let

$$\kappa(N) := kN - v(k).$$

Define the algebraic integers  $p_N, q_N \in \mathbb{Z}[\beta]$  by

$$p_N := (\beta^{y2^{\kappa(N)}} - 1) \sum_{n=1}^{y2^{\kappa(N)}-1} s(n)\beta^{y2^{\kappa(N)}-n} + \sum_{n=y2^{\kappa(N)}}^{y2^{\kappa(N)+1}-1} s(n)\beta^{y2^{\kappa(N)+1}-n}$$

and  $q_N := (\beta^{y2^{\kappa(N)}} - 1)\beta^{y2^{\kappa(N)}}$ , respectively. Then, we obtain

$$\begin{aligned} \frac{p_N}{q_N} &= \sum_{n=1}^{y2^{\kappa(N)}-1} \frac{s(n)}{\beta^n} + \frac{\beta^{y2^{\kappa(N)}}}{\beta^{y2^{\kappa(N)}} - 1} \cdot \sum_{n=y2^{\kappa(N)}}^{y2^{\kappa(N)+1}-1} \frac{s(n)}{\beta^n} \\ &= \sum_{n=1}^{y2^{\kappa(N)}-1} \frac{s(n)}{\beta^n} + \left( 1 + \frac{1}{\beta^{y2^{\kappa(N)}}} + \left( \frac{1}{\beta^{y2^{\kappa(N)}}} \right)^2 + \dots \right) \sum_{n=y2^{\kappa(N)}}^{y2^{\kappa(N)+1}-1} \frac{s(n)}{\beta^n} \\ &= \sum_{n=1}^{y2^{\kappa(N)+1}-1} \frac{s(n)}{\beta^n} + \frac{1}{\beta^{y2^{\kappa(N)}}} \sum_{n=y2^{\kappa(N)}}^{y2^{\kappa(N)+1}-1} \frac{s(n)}{\beta^n} + O\left(\left(\frac{1}{\beta^{y2^{\kappa(N)}}}\right)^2\right) \cdot O\left(\frac{1}{\beta^{y2^{\kappa(N)}}}\right) \\ (3.2) \quad &= \sum_{n=1}^{y2^{\kappa(N)+1}-1} \frac{s(n)}{\beta^n} + \sum_{j=0}^{y2^{\kappa(N)}-1} \frac{s(y2^{\kappa(N)} + j)}{\beta^{y2^{\kappa(N)+1+j}}} + O\left(\frac{1}{\beta^{3y2^{\kappa(N)}}}\right). \end{aligned}$$

By the equalities (2.7) and (2.15) with  $j = 0, 1, \dots, 2^N$ , we have

$$(3.3) \quad s(y2^{\kappa(N)} + j) = s(y2^{\kappa(N)+1} + j), \quad j = 0, 1, \dots, 2^N.$$

Hence, by (3.2) and (3.3),

$$(3.4) \quad \frac{p_N}{q_N} = \sum_{n=1}^{y2^{\kappa(N)+1}+2^N} \frac{s(n)}{\beta^n} + \sum_{j=2^N+1}^{2^{\lfloor \lambda N \rfloor}} \frac{s(y2^{\kappa(N)} + j)}{\beta^{y2^{\kappa(N)+1+j}}} + O\left(\frac{1}{\beta^{y2^{\kappa(N)+1+2^{\lfloor \lambda N \rfloor}}}}\right),$$

where we used  $1 < \lambda < 2 \leq k$  in the big  $O$  notation. Therefore, by using (3.4) and the equalities (2.15) with  $j = 2^N + 1, \dots, 2^{\lfloor \lambda N \rfloor}$ , we obtain

$$(3.5) \quad \xi - \frac{p_N}{q_N} = a_k \sum_{j=2^N+1}^{2^{\lfloor \lambda N \rfloor}} \frac{u(j)}{\beta^{y2^{\kappa(N)+1}+j}} + O\left(\frac{1}{\beta^{y2^{\kappa(N)+1}+2^{\lfloor \lambda N \rfloor}}}\right),$$

where

$$u(j) := t\left((y2^{\kappa(N)+1} + j)^k\right) - t\left((y2^{\kappa(N)} + j)^k\right), \quad j \geq 2^N + 1.$$

Note that  $|u(j)| \leq 1$  since  $u(j) \in \{-1, 0, 1\}$  for every integer  $j$ . Moreover, by the definition of  $q_N$ , we have  $q_N \leq \beta^{y2^{\kappa(N)+1}}$ , and so by (3.5),

$$(3.6) \quad q_N \xi - p_N = O\left(\frac{1}{\beta^{2^N}}\right).$$

On the other hand, applying Lemmas 2.2 and 2.3, we obtain

$$(3.7) \quad \begin{aligned} \left| \sum_{j=2^N+1}^{2^{\lfloor \lambda N \rfloor}} \frac{u(j)}{\beta^j} \right| &\geq \frac{1}{\beta^{2^N+1}} - \sum_{j=2^N+5}^{2^{\lfloor \lambda N \rfloor}} \frac{|u(j)|}{\beta^j} \\ &= \frac{1}{\beta^{2^N+1}} - \sum_{\substack{j \geq 2^N+5 \\ j:\text{odd}}} \frac{1}{\beta^j} - \sum_{\substack{j \geq 2^N+2^{N-2}+1 \\ j:\text{even}}} \frac{1}{\beta^j} \\ &\geq \frac{1}{\beta^{2^N+1}} - \frac{\beta^2}{\beta^2 - 1} \left( \frac{1}{\beta^{2^N+5}} + \frac{1}{\beta^{5 \cdot 2^{N-2}+2}} \right) \\ &= \frac{\beta^4 - \beta^2 - 1}{\beta^3(\beta^2 - 1)} \cdot \frac{1}{\beta^{2^N}} + O\left(\frac{1}{\beta^{5 \cdot 2^{N-2}}}\right), \end{aligned}$$

and hence, by (3.5) and (3.7),

$$\begin{aligned} \beta^{y2^{\kappa(N)+1}} \left| \xi - \frac{p_N}{q_N} \right| &\geq |a_k| \cdot \left| \sum_{j=2^N+1}^{2^{\lfloor \lambda N \rfloor}} \frac{u(j)}{\beta^j} \right| + O\left(\frac{1}{\beta^{2^{\lfloor \lambda N \rfloor}}}\right) \\ &= |a_k| \cdot \frac{\beta^4 - \beta^2 - 1}{\beta^3(\beta^2 - 1)} \cdot \frac{1}{\beta^{2^N}} + O\left(\frac{1}{\beta^{5 \cdot 2^{N-2}}}\right) + O\left(\frac{1}{\beta^{2^{\lfloor \lambda N \rfloor}}}\right). \end{aligned}$$

Thus, noting that  $\beta > \sqrt{\varphi}$ ,  $a_k \neq 0$ , and that both  $5 \cdot 2^{N-2} - 2^N$  and  $2^{\lfloor \lambda N \rfloor} - 2^N$  tend to infinity as  $n \rightarrow \infty$ , we obtain

$$(3.8) \quad \left| \xi - \frac{p_N}{q_N} \right| > 0.$$

Combining (3.6) and (3.8), there is a positive constant  $c_1$  such that

$$(3.9) \quad 0 < |q_N \xi - p_N| < \frac{c_1}{\beta^{2^N}}$$

for every sufficiently large  $N$ .



Now, we complete the proof. When  $\beta$  is a rational integer, (3.9) is clearly impossible for large  $N$ , since  $q_N \xi - p_N$  is a nonzero rational integer. So, suppose not, and let  $\beta =: \beta_1, \beta_2, \dots, \beta_d$  ( $d \geq 2$ ) be the Galois conjugates over  $\mathbb{Q}$  of  $\beta$ . Since  $\xi = -a_0 \in \mathbb{Z}[\beta]$ , there exist rational integers  $A_0, A_1, \dots, A_{d-1}$  such that  $\xi = \sum_{i=0}^{d-1} A_i \beta^i$ . Define the polynomial over  $\mathbb{Z}[\beta]$

$$F_N(X) := q_N(X)\xi(X) - p_N(X),$$

where

$$p_N(X) := (X^{y^{2^{\kappa(N)}}} - 1) \sum_{n=1}^{y^{2^{\kappa(N)}}-1} s(n)X^{y^{2^{\kappa(N)}}-n} + \sum_{n=y^{2^{\kappa(N)}}}^{y^{2^{\kappa(N)}}+1-1} s(n)X^{y^{2^{\kappa(N)}}+1-n},$$

$$q_N(X) := (X^{y^{2^{\kappa(N)}}} - 1)X^{y^{2^{\kappa(N)}}},$$

$$\xi(X) := \sum_{i=0}^{d-1} A_i X^i.$$

Note that  $p_N(\beta) = p_N$ ,  $q_N(\beta) = q_N$ , and  $\xi(\beta) = \xi$ . By (3.9),

$$(3.10) \quad 0 < |F_N(\beta)| = |q_N \xi - p_N| < \frac{c_1}{\beta^{2^N}}.$$

Moreover, since  $\beta$  is a Pisot or Salem number, we have  $|\beta_i| \leq 1$  ( $i = 2, 3, \dots, d$ ), and so by the definitions of  $p_N(X), q_N(X), \xi(X)$ , there exists a positive constant  $c_2$  independent of  $N$  such that

$$(3.11) \quad 0 < |F_N(\beta_i)| \leq c_2 \cdot y^{2^{\kappa(N)}}, \quad i = 2, 3, \dots, d,$$

where the first inequality follows since  $F_N(\beta_i)$  are the Galois conjugates of  $F_N(\beta) \neq 0$ . Therefore, considering the norm over  $\mathbb{Q}(\beta)/\mathbb{Q}$  of the algebraic integer  $F_N(\beta)$ , we obtain, by (3.10) and (3.11),

$$1 \leq |N_{\mathbb{Q}(\beta)/\mathbb{Q}} F_N(\beta)| = \prod_{i=1}^{d-1} |F_N(\beta_i)| < \frac{c_1 c_2^{d-1} \cdot (y^{2^{\kappa(N)}})^d}{\beta^{2^N}},$$

which is impossible for sufficiently large  $N$ , since  $\kappa(N) = O(N) = o(2^N)$ . The proof of Theorem 1.1 is now complete.

### 4 Concluding remarks and further questions

As there is no known nontrivial lower bound on Salem numbers, for our Theorem 1.1 to apply to all Salem numbers, we would need our result to be valid for  $\beta > 1$ . It seems very unlikely that the type of optimization that we have done here could be carried out to reach that range. A similar approach could increase the range a bit, but a new idea is probably necessary to get the full range of possible  $\beta$ . Here, our proof works for Pisot and Salem numbers, but it seems reasonable to conjecture that Theorem 1.1 holds for any algebraic number  $\beta$  with  $|\beta| > 1$ , though without more information on the structure of the sequences  $\{t(n^k)\}_{n \geq 0}$  this seems out of reach at the moment.

When we first started our investigation, we wanted to show that the three numbers

$$1, \sum_{n \geq 1} \frac{t(n)}{b^n}, \sum_{n \geq 1} \frac{t(n^2)}{b^n}$$

are linearly independent over  $\mathbb{Q}$  for any positive integer  $b \geq 2$ . Considering this question, two properties of the Thue–Morse sequence stood out to us. First, the sequence  $\{t(n)\}_{n \geq 0}$  is produced by a finite automaton (see [1, Section 5.1]), so it is not a very complicated sequence. Second, the sequence  $\{t(n^2)\}_{n \geq 0}$  is extremely complicated – as we mentioned in Section 1, the sequence  $\{t(n^2)\}_{n \geq 0}$  is normal; that is, all  $2^m$  patterns of finite subwords of length  $m$  occur with frequency  $2^{-m}$ . A result of Wall [10, Corollary 1, p. 15] states that if  $\xi$  is normal and  $q_1 \neq 0$  and  $q_2$  are rational numbers, then  $q_1 \xi + q_2$  is also normal, which implies the  $\mathbb{Q}$ -linear independence of the above three numbers when  $b = 2$ . It seems reasonable to think that for any rational numbers  $q_1 \neq 0$  and  $q_2$ , the number

$$q_2 + q_1 \sum_{n \geq 1} \frac{t(n)}{b^n}$$

must have a “not very complicated” base- $b$  expansion. In fact, this is the case since  $\{t(n)\}_{n \geq 0}$  is produced by a finite automaton (see [1, Section 13.1]). There is a gap in the literature regarding sequences and numbers that fall in between automatic and normal. We make explicit a question that would be a first step in this direction.

Recall, for any sequence  $f$  taking values in  $\{0, 1\}$ , we let  $p_f(m)$  denote the (subword) complexity of  $f$  as a one-sided infinite word. In particular,  $p_f(m)$  counts the number of distinct blocks of length  $m$  in  $f$ . So, for example,  $f$  is eventually periodic if and only if  $p_f(m)$  is uniformly bounded, and if  $f$  is 2-normal, then  $p_f(m) = 2^m$  for all  $m$ , since every binary word of any length appears in a 2-normal binary word. The entropy of  $f$  is the limit,  $h(f) := \lim_{m \rightarrow \infty} (\log p_f(m))/m \in [0, \log 2]$ . Considering the Thue–Morse sequence (or word)  $\mathbf{t}$ , since  $\mathbf{t}$  is generated by a finite automaton, we have that  $p_{\mathbf{t}}(m) = O(m)$ , so  $h(\mathbf{t}) = 0$ . In the present paper, we considered the sequences  $\mathbf{t}^k := \{t(n^k)\}_{n \geq 0}$ . Moshe [5] established that  $p_{\mathbf{t}^k}(m) \geq 2^{m/2^{k-2}}$  for any  $k \geq 2$ ; that is, that  $h(\mathbf{t}^k) \geq (\log 2)/2^{k-2} > 0$ . The question about numbers with lower complexity seems immediate.

**Question 4.1** For a real number  $\xi$ , let  $p_\xi(m, b)$  be the number of  $b$ -ary words of length  $m$  appearing in the base- $b$  expansion of  $\xi$ . Is it true that for any two rational numbers  $q_1 \neq 0$  and  $q_2$ , we have  $p_{q_1 \xi + q_2}(m, b) = O(p_\xi(m, b))$ ?

**Acknowledgements** The authors would like to thank Eiji Miyanohara for his helpful comments on an earlier version of this paper, and in particular, for pointing us toward the results of [1, Section 13.1]. This work began during the first author’s visit to Hirosaki University. Michael Coons thanks Hirosaki University for their support, and their staff and students for their outstanding hospitality.

## References

- [1] J.-P. Allouche and J. Shallit, *Automatic sequences: theory, applications, generalizations*, Cambridge University Press, Cambridge, 2003.
- [2] M. Drmota, C. Mauduit, and J. Rivat, *Normality along squares*. J. Eur. Math. Soc. 21(2019), no. 2, 507–548.
- [3] K. Mahler, *Arithmetische Eigenschaften der Lösungen einer Klasse von Funktionalgleichungen*. Math. Ann. 101(1929), 342–366. Corrigendum 103(1930), 532.
- [4] E. Miyahara, *An arithmetical property of the real numbers generated by Thue–Morse sequence along squares*. Acta Math. Hungar. 170(2023), 430–435.
- [5] Y. Moshe, *On the subword complexity of Thue–Morse polynomial extractions*. Theor. Comput. Sci. 389(2007), 318–329.
- [6] E. Prouhet, *Mémoire Sur quelques relations entre les puissances des nombres*. C. R. Acad. Sci. Paris Sér. I 33(1851), 225.
- [7] L. Spiegelhofer, *Thue–Morse along the sequence of cubes*. Preprint, 2023, [arXiv:2308.09498](https://arxiv.org/abs/2308.09498) [math.NT].
- [8] A. Thue, *Über unendliche Zeichenreihen*. Norske vid. Selsk. Skr. Mat. Nat. Kl. 7(1906), 1–22. Reprinted in: T. Nagell (Ed.), Selected mathematical papers of Axel Thue, Universitetsforlaget, Oslo, 1977, pp. 139–158.
- [9] A. Thue, *Über die gegenseitige Lage gleicher Teile gewisser Zeichenreihen*. Norske vid. Selsk. Skr. Mat. Nat. Kl. 1(1912), 1–67. Reprinted in: T. Nagell (Ed.), Selected mathematical papers of Axel Thue, Universitetsforlaget, Oslo, 1977, pp. 413–478.
- [10] D. D. Wall, *Normal numbers*. Ph.D. Thesis, University of California, Berkeley, 1950.

*Department of Mathematics and Statistics, California State University, Chico, CA 95929, United States*  
*e-mail:* [mjcoons@csuchico.edu](mailto:mjcoons@csuchico.edu)

*Graduate School of Science and Technology, Hirosaki University, Hirosaki 036-8561, Japan*  
*e-mail:* [tachiya@hirosaki-u.ac.jp](mailto:tachiya@hirosaki-u.ac.jp)