

Longitudinal Study of Sustained Attention in Outpatients with Bipolar Disorder

Alexandrea L. Harmell,^{1,2} Brent T. Mausbach,¹ Raeanne C. Moore,^{1,3} Colin A. Depp,^{1,3} Dilip V. Jeste,^{1,3}
AND Barton W. Palmer^{1,3,4}

¹University of California, San Diego, Department of Psychiatry, La Jolla, California

²San Diego State University, Department of Psychology, San Diego, California

³The Sam and Rose Stein Institute for Research on Aging, University of California San Diego, La Jolla, California

⁴Veterans Medical Research Foundation, VASDHS, San Diego, California

(RECEIVED June 18, 2013; FINAL REVISION December 6, 2013; ACCEPTED December 9, 2013)

Abstract

Individuals with bipolar disorder (BD) may exhibit attentional deficits, however, the extent of impairment and long-term fluctuations in performance in attention are relatively unknown. We investigated the relationship between sustained attention and affective symptoms over time among BD patients. We also examined whether global differences in attentional capacity differed among BD *versus* normal comparison (NC) subjects. Participants included 106 outpatients with BD and 66 NC subjects who were administered symptom rating scales and a measure of sustained attention (Continuous Performance Test- Identical Pairs). Measures were repeated 6, 12, and 26 weeks post-baseline. Compared to NC subjects, participants with BD showed impairment in sustained attention across time. Within patient increases in manic symptoms were associated with increased false alarms; both manic and depressive symptoms were associated with worse discrimination. Neither manic nor depressive symptoms were related to hit rates. Our results indicate that the ability to inhibit a response to near miss stimuli (i.e., those that are close to but not identical to the target) is globally impaired among BD patients relative to NC subjects, as well as state-dependent, covarying with affective symptoms. Psychosocial interventions requiring high levels of attentional capacity may need to be adapted according to patients' current symptomatology. (*JINS*, 2014, 20, 230–237)

Keywords: Neuropsychology, Intra-individual variability, Cognition, Continuous performance test, Mood disorder, Bipolar disorder

INTRODUCTION

Although bipolar disorder (BD) is associated with deficits in several cognitive domains, there is some evidence suggesting that impairment in sustained attention may be particularly characteristic of the disorder. Several cross sectional studies have demonstrated that individuals with BD exhibit deficits in sustained attention during both manic (Clark, Iversen, & Goodwin, 2001; Najt et al., 2005) and depressive episodes (Borkowska & Rybakowski, 2001). More recently, however, it is becoming increasingly recognized that neurocognitive deficits in sustained attention may partially persist during periods of euthymia (Latalova, Prasko, Diveky, & Velartova, 2011; Pattanayak, Sagar, & Mehta, 2012), and may even be present among unaffected family members of patients with

BD (Sepede et al., 2012). Such findings raise the question of whether deficits in sustained attention may be a trait-like feature of BD, rather than a secondary effect of the psychopathologic symptoms on test performance.

Studies using Continuous Performance Tests (CPT) that reveal deficits in attention during euthymic phases of the disorder offer preliminary evidence that sustained attention/vigilance may be a stable trait of the disorder (Ancín et al., 2010; Clark, Kempton, Scarnà, Grasby, & Goodwin, 2005; Walshe et al., 2012). Some researchers have suggested that sustained attention may be one of the *most* sensitive markers of cognitive impairment in BD as it is unrelated to medication status and even partially present in patients with a good functional recovery (Clark & Goodwin, 2004). Further investigation of sustained attention over time may aid in better understanding the degree to which sustained attention is a stable trait-like deficit in BD.

Despite a growing literature on sustained attention in BD, there remains a paucity of literature specifically examining

Correspondence and reprint requests to: Barton W. Palmer, Division of Geriatric Psychiatry 116A-1, University of California, San Diego, 8950 Villa La Jolla Drive Suite B101, La Jolla, CA 92037. E-mail: bpalmer@ucsd.edu

the stability of sustained attention within individual patients. Exploring within-person stability of sustained attention among people with BD may provide further information to researchers and clinicians about potential barriers and/or the most effective timing for psychotherapeutic interventions. Additionally, knowing whether individual fluctuations in attention are associated with symptom exacerbation may be helpful in explaining functional outcomes in the disorder as impaired sustained attention likely impacts social activities as well as school and work performance (Goodwin & Jamison, 1990).

The aim of the present report was to examine level and stability of sustained attention among outpatients with BD, and the degree to which this stability is affected by concurrent severity or fluctuations in psychopathologic symptoms. Subjects were repeatedly assessed with a measure of sustained attention at baseline, and 6-, 12-, and 26-week follow-up evaluations, regardless of current symptomatology. We first compared level of sustained attention among patients relative to normal comparison (NC) subjects. We hypothesized that the BD group would have worse sustained attention performance relative to the NC group. [We also ran a sub-analysis at baseline to compare sustained attention among euthymic patients and NC subjects.] In regard to stability over time, we hypothesized that, within patients, having more severe manic or depressive symptoms would result in significantly worse sustained attention than when less symptomatic.

METHODS

Participants

Subjects included 106 adult outpatients with bipolar disorder (BD) (84 Type I and 22 Type II) and 66 psychiatrically healthy normal comparison (NC) subjects. Data in the present report were collected as part of subjects' participation in an ongoing study of capacity for informed consent among people with BD. Some ($n = 91$) of the subjects provided data for a prior report in which we examined the stability of overall neurocognitive functioning (Depp, Savla, Vergel de Dios, Mausbach, & Palmer, 2012), but the present analyses represent our first examination of the sustained attention data.

Patient participants were recruited through a variety of sources including the University of California, San Diego (UCSD) and Veterans Affairs San Diego Healthcare System outpatient psychiatry services, the San Diego County Adult and Older Adult Mental Health Services, as well as through San Diego area Board and Care facilities. The NC subjects were recruited through flyers posted in newspapers, online, or in the community.

Inclusion criteria included: (a) DSM-IV diagnosis of BD or, if NC, absence of a major neuropsychiatric disorder, and (b) English fluency. Participants were excluded from the study if they: (a) met diagnostic criteria for substance dependence (per DSM-IV criteria) within the 2 months before enrollment, (b) had a known diagnosis of dementia or

other neurological or medical conditions known to affect cognitive functioning, or (c) had medical problems that would interfere with the participant's ability to complete the study assessments. Diagnostic status of patients and NC subjects was established with the Structured Clinical Interview for the DSM-IV-TR Axis I Disorders (SCID-I; First, Spitzer, Gibbon, & Williams, 2002) administered by a trained research assistant and subsequently reviewed by a board certified psychiatrist. Approval for this study was provided by the UCSD Human Research Protections Program and all participants provided written informed consent before enrollment.

Measures

Sociodemographic information

Information about each subject's age, education, gender, and ethnicity was collected *via* interview and/or review of available records. Premorbid verbal cognitive functioning was estimated with the Word Reading subtest from the Wide Range Achievement Test – 4th edition (WRAT-4) (Wilkinson & Robertson, 2006).

Psychiatric background

Clinical history and status information was determined as part of the SCID-I administration including age of onset and current affective episode type (if any).

Severity of psychopathology

Severity of manic symptoms was evaluated at each study visit (baseline, 6, 12, and 26 weeks) with the Young Mania Rating Scale (YMRS; Young, Biggs, Ziegler, & Meyer, 1978). The presence and severity of depressive symptoms was evaluated at each study visit using the Montgomery-Asberg Depression Rating Scale (MADRS; Montgomery & Asberg, 1979).

Sustained Attention

Sustained attention was assessed at each visit in both NC and BD participants using the Continuous Performance Test-Identical Pairs (CPT-IP) from the MATRICS Consensus Cognitive Battery (Nuechterlein & Green, 2006). This task measures attention with increased memory load using computer-generated two, three, and four-digit stimuli. The software for this test provides separate results for the two, three, and four-digit stimuli, but does not provide an index of overall performance. Rather than analyze all three conditions in parallel (and thus risk inflated Type I error due to multiple comparisons), we focused our analyses on the results for the four-digit trials, as this block of trials requires the highest cognitive demand in sustained attention. However, to determine whether task complexity affected the pattern of results, we also conducted secondary analyses in which we examined the pattern of results for our key comparisons when substituting the simplest (two-digit) condition scores.

In the four-digit condition, participants are shown a rapid sequence of four-digit numbers (150 trials) presented on a

computer screen, and are instructed to press a mouse key any time two numbers in a row are identical. Each of the 150 trials consists of presenting a four-digit number. Responses to trials were scored as either correct detections/hits, false alarms (stimulus presented was very similar but not identical to that of the preceding trial), or random errors (responses to randomly distributed fillers). D -prime (d'), a statistic applied in signal detection theory, was used to measure each individuals' ability to discriminate signal from background noise (this was calculated as the difference between z -transformations of hits minus false alarms). As prior studies have indicated that both hits and false alarms may be sensitive to fluctuations in depression and mania (Snyder, Nussbaum, & Robins, 2006), it appeared prudent to include both in the present analyses, along with d' .

Data Analysis

All statistical analyses were performed using SPSS software version 20. Independent t -tests were performed to compare BD and NC groups on quantitative demographic variables. Pearson chi-square tests were used for categorical variables such as gender and race.

Mixed model analyses were performed to compare mean sustained attention scores of the BD and NC groups across time. In these analyses, random intercepts were modeled and diagnostic group (BD or NC) were entered as fixed effects. Hits, false alarms, and d' were used separately as the dependent variable.

In addition, a sub-analysis was conducted by using independent t -tests to compare CPT-IP performances of euthymic patients (per SCID-I) at baseline and NC subjects.

Next, we restricted the analysis to only the BD group. First, within and between person variables were created for each mood measure (YMRS and MDRS). For the within-person scores, each participant's YMRS or MDRS score at a given study visit was expressed in terms of deviation from the mean of his or her YMRS or MDRS scores across all study visits. Development of this variable isolates within-person change in symptoms so that we can correlate change in symptoms to change in attention. For the between-person scores, we first calculated each participant's mean YMRS or MDRS score across all study visits, and then expressed that score in terms of deviation from the mean of all patients. This variable allows us to examine average (long-term) symptom levels and their relation to the attention intercepts across time. These within and between variables were entered into a linear mixed model analysis with random intercepts and age, education, and either YMRS within and between or MADRS within and between total scores entered as the fixed effects. Separate analyses were conducted for each of the CPT-IP outcome variables: hits, false alarms, and d' to determine whether within the BD group, age, education, and symptoms explained within or between differences in CPT-IP performance. Number of random errors was also explored, however due to the limited variability between patients (across all visits, 90.6% of patients had less than 4 random errors), it was not a primary focus of this study. Significance was

defined as $p < .05$, two tailed, for all analyses. In an effort to estimate the effect sizes for all significant statistical analyses, we computed pseudo R -square statistics consistent with the approach outlined by Singer and Willet (2003). Specifically, this statistic was calculated by squaring the correlation between the actual CPT-IP scores and the model predicted CPT-IP scores. Cohen's d effect sizes were calculated for independent t -tests.

RESULTS

Demographic and Clinical Characteristics:

As shown in Table 1, relative to the NC group, the BD group was older (45.4 years vs. 39.6) had slightly fewer years of education (14.6 years vs. 15.6), and had differences in ethnic composition ($\chi^2(4) = 16.38; p = .003$). However, there were no significant differences between the two groups in gender (40.7% vs. 40.5%) or estimated premorbid verbal ability (WRAT-4 Reading; scaled score of 96.4 vs. 98.4).

As expected, the BD group endorsed significantly more severe manic and depressive symptoms relative to the NC group (although even in the BD group the baseline scores were, on average, in the mild range of severity). Also, per the SCID-I, within the month preceding the baseline evaluation, 62% of patients were euthymic, 25% experienced a major depressive episode, 8% a hypomanic episode, and 5% had met criteria for a mixed or a manic episode. In terms of current medication status, 71.7% of patients were taking a mood stabilizer, 37.7% an anti-depressant, and 42.7% were taking an anti-psychotic. Some participants were taking a combination of more than one of these drug classes including 2.8% of patients on both antipsychotics and anti-depressants, 14.2% on both anti-depressants and mood stabilizers, 16.4% on both mood stabilizers and anti-psychotics, and 16.0% on all three.

All participants with at least one baseline visit were included in the analyses. Within the patient group, 87% ($n = 92$) of patients had at least 2 CPT-IP scores from two different time-points [67% ($n = 71$) had at least three scores and 36% ($n = 38$) of patients had CPT-IP scores from all four visits]. In the NC group, 82% ($n = 54$) had at least 2 CPT-IP scores from two different time-points [76% ($n = 50$) had at least three scores and 46% ($n = 30$) of NC subjects had CPT-IP scores from all four visits].

Between Group Differences in Sustained Attention

Within the entire sample, compared to NC subjects, participants with BD performed significantly worse across all time points on d' [estimate = -0.504 ; $SE = 0.133$; $t = -3.795$; $p < .001$, 95% confidence interval (CI) ($-.766, -.241$)], had significantly lower hit rates [estimate = -2.178 ; $SE = 0.905$; $t = -2.407$; $p = .017$, 95% CI ($-3.964, .392$)], and exhibited a significantly higher rate of false alarms on the CPT-IP four-digit task [estimate = 1.735 ; $SE = 0.478$; $t = 3.630$; $p < .001$, 95% CI ($.791, 2.678$)]. These predictors accounted

Table 1. Demographic characteristics, severity of symptoms, and overall cognitive functioning by diagnosis

	NC group (<i>n</i> = 66)	Bipolar disorder (<i>n</i> = 106)	<i>t</i> or χ^2	<i>df</i> or <i>N</i>	<i>p</i>
Age (years)	39.6 (14.5)	45.4 (11.7)	2.76	170	.006
Education (years)	15.6 (2.1)	14.6 (2.3)	-2.86	170	.005
Gender (% women)	40.5%	40.7%	3.20	1	.074
Ethnicity (%)			16.38	4	.003
Caucasian	59.1%	66.0%			
African American	6.1%	7.5%			
Hispanic/Latino	7.6%	14.2%			
Asian	16.7%	0.9%			
Other	10.5%	11.4%			
WRAT scaled score (at baseline)	98.4 (11.8)	96.4 (15.4)	-0.83	137	.393
YMRS (at baseline)	1.4 (1.8)	8.0 (7.7)	6.75	168	< .001
MADRS (at baseline)	1.2 (2.1)	14.0 (9.9)	10.25	169	< .001
Age of onset (years)	NA	20.0 (9.1)		<i>N</i> = 84	
% Bipolar I	NA	79.2%			
D-prime rate (at baseline)	1.9 (0.9)	1.5 (0.9)	3.25	156	.001
Hit rate (at baseline)	22.1 (6.1)	19.6 (6.5)	2.39	156	.018
False alarm rate (at baseline)	4.3 (3.2)	6.0 (4.3)	-2.55	156	.012

Note. Values represent means (and *SDs*) or percentages (where indicated).

YMRS = Young Mania Rating Scale; MADRS = Montgomery-Asberg Depression Rating Scale; WRAT = Wide Range Achievement Test – 4th edition.

for 6.9%, 3.0%, and 5.2% of the variance in *d'*, hit rate, and false alarm scores respectively.

Due to the fact that the BD and NC groups differed in terms of mean age and education, we performed additional statistical analyses after removing participants whose age or education values were outside of the range of either group (age ≥ 23 and ≤ 72 ; years of education ≥ 9 and ≤ 17). This restriction led to a subsample of 39 NC subjects and 91 patients with BD (mean age 44.4 vs. 45.1 years and mean years of education 14.7 vs. 14.2 years, respectively). The pattern of differences between NC subjects and BD in this subsample (see Table 2) was consistent with our original analyses. Specifically, relative to NC subjects, the BD group performed significantly worse on *d'* [estimate = -0.522; *SE* = 0.160; *t* = -3.261; *p* = .001; 95% CI (-.840, -.205)], had lower hit rates [estimate = -2.251;

SE = 1.130; *t* = -1.991; *p* = .049; 95% CI (-4.489, -.014)], and had significantly higher false alarm rates [estimate = 1.740; *SE* = 0.592; *t* = 2.938; *p* = .004; 95% CI (.568, 2.912)].

After substituting the two-digit task scores, and using the demographically matched subsample, the BD group showed significant impairment across all measures of sustained attention relative to NC subjects including *d'* [estimate = -0.446; *SE* = 0.122; *t* = -3.667; *p* < .001; 95% CI (-.687, -.205)], hits [estimate = -1.656; *SE* = 0.619; *t* = -2.678; *p* = .008; 95% CI (-2.880, -.432)], and false alarms [estimate = .660; *SE* = 0.233; *t* = 2.837; *p* = .005; 95% CI (.199, 1.122)].

In our other sub-analysis just looking at baseline visit, euthymic patients had significantly lower *d'* [*t*(117) = 3.556; *p* = .001] and hit rates [*t*(117) = 2.647; *p* = .009] on the CPT-IP when compared to NC subjects. Moreover, euthymic patients showed higher rates of false alarms relative to NC subjects [*t*(117) = -2.507; *p* = .014]. Cohen's *d* effect sizes were .65, .49, and .46, respectively, suggesting effects in the medium range. These results were generally consistent using the age and education matched sample, however false alarms did not quite reach statistical significance [*d'* *t*(82) = 3.105; *p* = .003; hits *t*(82) = 2.470; *p* = .016; false alarms *t*(82) = -1.943; *p* = .055]. Effect sizes in this latter analysis were comparable with those using the full euthymic sample (Cohen's *d* = .67, .54, and .44, respectively).

Table 2. Group differences in sustained attention between age and education matched patients with bipolar disorder (*n* = 91) and normal comparison subjects (*n* = 39) (parameter estimates, standard errors, *t*-values, and *p*-values)

	Estimate	<i>SE</i>	<i>t</i> -Value	<i>p</i> -Value
D-prime				
Intercept	2.050	.133	15.354	.000
Bipolar group	-.522	.160	-3.261	.001
Hits				
Intercept	22.639	.942	24.029	.000
Bipolar group	-2.251	1.130	-1.991	.049
False alarms				
Intercept	4.199	.493	8.515	.000
Bipolar group	1.740	.592	2.938	.004

Within-Patient Impact of Affective Symptoms on Sustained Attention over Time

Adjusting for age and education, we examined the impact of change in affective symptoms on sustained attention within the BD group. As shown in Table 3, a negative relationship was found between severity of manic symptoms (YMRS) and

Table 3. Associations of sustained attention and mania within and between bipolar disorder subjects (parameter estimates, standard errors, *t*-values, and *p*-values)

Mania symptom model	Estimate	SE	<i>t</i> -Value	<i>p</i> -Value
D-prime				
Intercept	.601	.539	1.114	.268
Age	-.021	.006	-3.493	.001
Education (in years)	.133	.032	4.130	.000
YMRS total_within	-.016	.008	-2.092	.038
YMRS total_between	.012	.012	.997	.321
Hits				
Intercept	14.257	4.184	3.408	.001
Age	-0.101	.047	-2.147	.034
Education (in years)	0.757	.250	3.024	.003
YMRS total_within	.051	.056	.904	.367
YMRS total_between	.137	.091	1.496	.137
False alarms				
Intercept	6.414	2.111	3.038	.003
Age	.067	.024	2.814	.006
Education (in years)	-.256	.126	-2.027	.045
YMRS total_within	.100	.044	2.304	.022
YMRS total_between	.047	.047	1.002	.319

YMRS = Young Mania Rating Scale.

the ability to accurately discriminate between target and non-target stimuli (*d'*) in that increases in manic symptoms were related to decreases in *d'* [estimate = -.016; *SE* = 0.008; *t* = -2.092; *p* = .038; 95% CI (-.031, -.000)]. Based on the pseudo R-squared statistic, results of this model indicate that approximately 20.0% of the variance in *d'* scores was accounted for by the predictors in our model. In addition, a relationship also emerged between manic symptoms and false alarms showing that as manic symptoms increased, a patient's rate of false alarms also increased [estimate = .100; *SE* = 0.044; *t* = 2.304; *p* = .022; 95% CI = (.014, .186)] (in respect to pseudo R-squared, 8.8% of the variance was explained). However, no significant association was found between manic symptoms and hit rate [estimate = .051; *SE* = 0.056; *t* = 0.904; *p* = .367; 95% CI = (-.060, .162)].

As shown in Table 4, changes in depressive symptoms over time were not significantly associated with change in hits or false alarms. However, increases in depressive symptoms appeared to be associated with decreases in discriminability [*d'* estimate = -0.011; *SE* = 0.004; *t* = -2.394; *p* = .018; 95% CI = (-.020, -.002)]. Specifically, pseudo R-square statistic indicated that 19.6% of the variance in *d'* scores was accounted for by the predictors used in our model.

When the within BD analyses were performed with the scores from the two-digit CPT-IP task, neither manic [*d'* *t*(208.907) = -1.878; *p* = .062; hits *t*(-1.780) = 209.444; *p* = .077; false alarms *t*(219.684) = .233; *p* = .816] nor depressive symptoms [*d'* *t*(211.084) = -1.011; *p* = .313; hits *t*(211.593) = -.772; *p* = .441; false alarms *t*(222.602) = -.755; *p* = .451] were associated with significant changes on any of the sustained attention outcome variables.

Table 4. Associations of sustained attention and depression within and between bipolar disorder subjects (parameter estimates, standard errors, *t*-values, and *p*-values)

Depression symptom model	Estimate	SE	<i>t</i> -Value	<i>p</i> -Value
D-Prime				
Intercept	0.755	0.522	1.446	.151
Age	-0.021	0.006	-3.467	.001
Education (in years)	0.123	0.031	3.975	.000
MADRS total_within	-.011	.004	-2.394	.018
MADRS total_between	.000	.008	.010	.992
Hits				
Intercept	15.994	4.079	3.921	.000
Age	-0.101	.047	-2.117	.037
Education (in years)	0.636	.241	2.645	.010
MADRS total_within	-.061	.034	-1.791	.075
MADRS total_between	.008	.064	.134	.894
False alarms				
Intercept	6.806	2.047	3.325	.001
Age	.068	.024	2.834	.006
Education (in years)	-.285	.121	-2.362	.020
MADRS total_within	.015	.027	.576	.565
MADRS total_between	.029	.033	.879	.381

MADRS = Montgomery-Asberg Depression Rating Scale.

Between-Patient Impact of Affective Symptoms on Sustained Attention over Time

Between-patient mania and depression measures, which represent the individual's overall mean across the total number of assessments completed, were not significant predictors of sustained attention measures in any analyses (see Tables 3 and 4).

DISCUSSION

This study investigated sustained attention, assessed at multiple time points using the Continuous Performance Test-Identical Pairs (CPT-IP), in a sample of 106 patients with BD and 66 NC subjects. Results of this study suggest that compared to NC subjects, participants with BD show sustained attention impairment across time. These results remained, even after performing additional analyses restricting the two groups to comparable age and education ranges. Of note, patients with BD also performed worse across all measures of sustained attention on the two-digit CPT-IP task relative to NC subjects. However, patient performance did not appear to be affected by exacerbation in mood symptoms on the two-digit task. Use of four- rather than two-digit stimuli may increase cognitive load, thereby taxing a greater proportion of limited working memory resources (Bucks & Seljos, 1994; van Merriënboer & Sweller, 2005). In our other sub-analysis, it was also found that euthymic patients performed significantly worse on all three CPT-IP outcomes relative to NC subjects, signifying that attentional deficits may represent a partially trait-like marker of dysfunction in patients with BD (Clark & Goodwin, 2004).

Our results also indicate that certain aspects of sustained attention, namely discrimination and false alarms, are potentially sensitive to within-person mood variability. At the group level (comparing between subjects), severity of manic or depressive symptoms did not impact performance on attention. However, within patients, those who had increases in either manic or depressive symptoms over time demonstrated decreases in the ability to discriminate between target and non-target stimuli. Together these findings provide support to the theory that sustained attention may be both trait and state-like in regard to mood symptoms.

Attentional impairments are among the most common behavioral manifestations of brain dysfunction, and underlie many higher order cognitive deficits (Glisky, 2007; Sarter, Givens, & Bruno, 2001; Shi et al., 2012). Sustained attention, specifically, requires the selection of salient sensory information for future processing, and must be maintained over time. Waxing and waning attention is a common cognitive symptom of major affective disorders, such as BD or Major Depressive Disorder. Previous research has indicated that the quality of sustained attentional impairment varies as a function of affective state, in that more false alarms are observed in patients who are in a manic state and fewer hits are observed in patients with depressive symptomatology (Snyder et al., 2006). Our results are consistent with these findings in the patients exhibiting more severe mania, but not in those exhibiting depressive symptoms. The reason for this discrepancy in findings is unclear; however one possible explanation for this difference is that our sample, as a whole, endorsed only mild levels of depressive symptoms. A study that looked cross-sectionally within more mildly depressed participants also did not find significant associations between hit rates and depression, but did find significant associations between mania and false alarms (Swann, Anderson, Dougherty, & Moeller, 2001). Therefore, it is possible that subsyndromal depression does not impact sustained attention in patients with BD as much as subsyndromal mania.

The finding that increases in manic symptoms were associated with higher false alarms rates may be attributed to impulsivity (Riccio, Reynolds, Lowe, & Moore, 2002). Specifically, as patients become more impulsive they may have a lower threshold for responding to near miss stimuli (i.e., those that are close to but not identical to the target). That possibility noted, however, it is unclear why such a lowering of response threshold did not lead to a parallel increase in hit rates.

Increases in false alarm rates when manic are of particular interest because they may represent an inability for manic patients to discriminate between subtle differences in visual information. Alternatively, the tendency for increased false alarms may represent difficulty inhibiting a response when a near target stimulus is presented. Deficits in response inhibition have frequently been noted in BD on the CPT test (Quraishi & Frangou, 2002), as well as performance on the Stroop Color Word Interference Task (Pompei et al., 2011; Stefanopoulou et al., 2009). Furthermore, Glosser, Deutsch, Cole, Corwin and Saykin (1998) and Corwin, Peselow,

Feenan, Rotrosen, and Fieve (1990) have previously reported on the impact that elevations in mood can have on producing more liberal response bias. This liberal response bias may be a result of distractibility, impulsive behavior, increased levels of arousal, or a combination of all of these symptoms that are typically characteristic of mania.

The ability to allocate attentional resources to relevant sensory information while simultaneously ignoring non-relevant information can have a significant impact on everyday functioning and quality of life. For example, in a sample of chronically hospitalized patients with schizophrenia, Silverstein, Schenkel, Valone, and Nuernberger (1998) demonstrated that 50% of change from a conversation skills training program could be predicted by measures of sustained attention and verbal learning. Another study found that improvement on a measure of social problem solving skills could be linked back to sustained attention and cognitive flexibility in outpatients with schizophrenia (Ucok et al., 2006).

One limitation of the present study is that sustained attention findings are based solely on performance on the CPT-IP. Whether the same pattern of results would generalize to other measures of sustained attention is unknown.

A potential confound in comparing the NC and BD groups is that only the latter were receiving psychotropic medications. Even the comparisons within the BD group may be complicated by the wide range of pharmacological treatments that patients with BD typically receive. There is some evidence that stimulant medications may enhance performance on the CPT task (Tucha et al., 2006), but only four participants in our study reported taking stimulants at the time of testing. Many of our participants were taking mood stabilizers, but these do not appear to have substantial effects on attention (Quraishi & Frangou, 2002; Wingen, Kuypers, van de Ven, Formisano, & Ramaekers, 2008). While benzodiazepines have been known to impair processing speed, previous research reported no significant correlation between performance on attentional tasks and benzodiazepine use (Clark et al., 2001). Combined, these studies seem to provide some assurance that the deficits in sustained attention in the BD group are not primarily attributable to medication effects (Clark & Goodwin, 2004; Goswami et al., 2009; Phillips, Travis, Fagiolini, & Kupfer, 2008).

The preceding conclusion does not necessarily suggest that within patient changes in medications would have no effect on sustained attention, particularly if such a change resulted in exacerbation or improvement in symptoms; it only suggests that there is not a strong relationship between type or specific dose of medications on sustained attention when comparing between patients. (In fact, in the present study, the association between symptom severity and sustained attention was only present when examined within subjects, not when comparing between patients.)

In addition to the aforementioned limitations, there are some methodological issues that deserve mention. The study did not include a measure of effort level, so it is conceivable that BD related performance deficits partially reflect suboptimal motivation rather than attention skills, *per se*. We also did not screen participants for ADHD which may be

co-morbid with BD. [The actual co-morbidity of ADHD and BD continues to remain controversial as some studies suggest that the two disorders share overlap in diagnostic symptoms and are sometimes difficult to distinguish from one another (Youngstrom, Arnold, & Frazier, 2010).] Furthermore, although those with recent (within 2 months) substance dependence were screened out, we did not exclude people with alcohol or substance abuse. Questions about abuse were included as part of the SCID-I interviews; only five patients met criteria for current alcohol abuse, one of whom also met criteria for substance abuse. But these values are based largely on self-report so may underestimate the true levels. Therefore, it is possible that some of the attentional effects attributed to BD in our results may further reflect effects of undetected co-morbid disorders.

It is also possible that the generalizability of our results may be restricted as the BD participants in this study, as a whole, endorsed relatively mild levels of manic and depressive symptoms and reported above average education levels. Finally, the proportions of specific ethnic minority participants (particularly Latino and Asian American) differed between the NC and BD groups. The potential effects of ethnicity/culture on CPT performance are currently unknown. Future research is needed to explore group differences in CPT performance in various minority populations.

A major strength of this study is that it included measures of sustained attention and symptomatology over time rather than the use of only cross-sectional or static indices of attentional capacity. Additionally, we evaluated both within- and between-patient performance, which allowed us to examine intra-individual variability as well as group differences. In a previous longitudinal study of overall cognitive functioning using a subset of participants from this sample, it was concluded that manic and depressive symptoms had little effect on overall level of cognitive impairment (Depp, Savla, et al., 2012). Therefore, it may be the case that there is something unique about sustained attention in isolation that is more susceptible to symptom fluctuations than is overall cognitive functioning. In general, patients with bipolar disorder have exhibited more intra-individual variability in neurocognitive functioning, as a whole, relative to patients with schizophrenia and NC subjects (Depp et al., 2008); sustained attention appears to be one of the domains consistent with this variable pattern.

One practical implication of these findings is that psychosocial interventions for BD, particularly those requiring high levels of sustained attention, may need to be adjusted according to a patient's current symptom severity. Our findings suggest that when patients with BD are symptomatic, their ability to pay attention may be compromised. Therefore, it may be helpful for clinicians providing psychotherapy or rehabilitation services to take this into consideration and to evaluate whether patients are attending to and understand material presented. Cognitive supports (e.g., repetition of information, multi-modal presentation) may be useful to aid in the successful translation of instructions (e.g., medication dosing) to action. Moreover, there is some evidence that

attention/vigilance deficits in BD may adversely impact everyday functioning (reviewed in Depp, Mausbach, et al., 2012); investigation of the benefits of cognitive training interventions specifically targeting sustained attention would be a welcome addition to the field.

ACKNOWLEDGMENTS

This work was supported, in part, by National Institutes of Health (grants MH64722, T32MH019934, 5P30MH066248) and by the Department of Veterans Affairs. The authors have no conflicts of interest to report.

REFERENCES

- Ancín, I., Santos, J.L., Teijeira, C., Sánchez-Morla, E.M., Bescós, M.J., Argudo, I., & Cabranes-Díaz, J.A. (2010). Sustained attention as a potential endophenotype for bipolar disorder. *Acta Psychiatrica Scandinavica*, *122*(3), 235–245.
- Backs, R.W., & Seljos, K.A. (1994). Metabolic and cardiorespiratory measures of mental effort: The effects of level of difficulty in a working memory task. *International Journal of Psychophysiology*, *16*(1), 57–68.
- Borkowska, A., & Rybakowski, J.K. (2001). Neuropsychological frontal lobe tests indicate that bipolar depressed patients are more impaired than unipolar. *Bipolar Disorders*, *3*, 88–94.
- Clark, L., & Goodwin, G.M. (2004). State- and trait-related deficits in sustained attention in bipolar disorder. *European Archives of Psychiatry and Clinical Neuroscience*, *254*(2), 61–68.
- Clark, L., Kempton, M.J., Scarnà, A., Grasby, P.M., & Goodwin, G.M. (2005). Sustained attention-deficit confirmed in euthymic bipolar disorder but not in first-degree relatives of bipolar patients or euthymic unipolar depression. *Biological Psychiatry*, *15*, 57(2), 183–187.
- Clark, L., Iversen, S.D., & Goodwin, G.M. (2001). A neuropsychological investigation of prefrontal cortex involvement in acute mania. *American Journal of Psychiatry*, *158*, 1605–1611.
- Corwin, J., Peselow, E., Feenan, K., Rotrosen, J., & Fieve, R. (1990). Disorders of decision in affective disease: An effect of b-Adrenergic Dysfunction? *Biological Psychiatry*, *27*, 813–833.
- Depp, C.A., Mausbach, B.T., Harmell, A.L., Savla, G.N., Bowie, C.R., Harvey, P.D., & Patterson, T.L. (2012). Meta-analysis of the association between cognitive abilities and everyday functioning in bipolar disorder. *Bipolar Disorders*, *14*(3), 217–226.
- Depp, C.A., Savla, G.N., Moore, D.J., Palmer, B.W., Stricker, J.L., Lebowitz, B.D., & Jeste, D.V. (2008). Short-term course of neuropsychological abilities in middle-aged and older adults with bipolar disorder. *Bipolar Disorders*, *10*(6), 684–690.
- Depp, C.A., Savla, G.N., Vergel de Dios, L.A., Mausbach, B.T., & Palmer, B.W. (2012). Affective symptoms and intra-individual variability in the short-term course of cognitive functioning in bipolar disorder. *Psychological Medicine*, *42*, 1409–1416.
- First, M.B., Spitzer, R.L., Gibbon, M., & Williams, J.B.W. (2002). *Structured clinical interview for DSM-IV-TR Axis I disorders, research version, patient edition*. (SCID-I/P) New York: Biometrics Research, New York State Psychiatric Institute.
- Glisky, E.L. (2007). Changes in cognitive function in human aging. In: D. R. Riddle, (Ed.), *Brain aging: Models, methods, and*

- mechanisms*. Boca Raton, FL: CRC Press. Chapter 1. Retrieved from: <http://www.ncbi.nlm.nih.gov/books/NBK3885/>
- Glosser, G., Deutsch, G.K., Cole, L.C., Corwin, J., & Saykin, A.J. (1998). Differential lateralization of memory discrimination and response bias in temporal lobe epilepsy patients. *Journal International Neuropsychology Society*, 4(5), 502–511.
- Goodwin, F.K., & Jamison, K.R. (1990). *Manic-depressive illness*. New York: Oxford University Press.
- Goswami, U., Sharma, A., Varma, A., Gulrajani, C., Ferrier, I.N., Young, A.H., ... Moore, P.B. (2009). The neurocognitive performance of drug-free and medicated euthymic bipolar patients do not differ. *Acta Psychiatrica Scandinavica*, 120, 456–463.
- Latalova, K., Prasko, J., Diveky, T., & Velartova, H. (2011). Cognitive impairment in bipolar disorder. *Biomedical papers of the medical faculty at the University Palacky Olomouc, Czech Republic*, 155(1), 19–26.
- Montgomery, S.A., & Asberg, M. (1979). A new depression scale designed to be sensitive to change. *British Journal of Psychiatry*, 134, 382–389.
- Najt, P., Glahn, D., Bearden, C.E., Hatch, J.P., Monkul, E.S., Kaur, S., ... Soares, J.C. (2005). Attention deficits in bipolar disorder: A comparison based on the Continuous Performance Test. *Neuroscience Letters*, 379, 122–126.
- Nuechterlein, K.H., & Green, M.F. (2006). *MCCB: MATRICS consensus cognitive battery manual*. Los Angeles, CA: MATRICS Assessment, Inc.
- Pattanayak, R.D., Sagar, R., & Mehta, M. (2012). Neuropsychological performance in euthymic Indian patients with bipolar disorder type I: Correlation between quality of life and global functioning. *Psychiatry Clinical Neuroscience*, 66(7), 553–563.
- Phillips, M.L., Travis, M.J., Fagiolini, A., & Kupfer, D.J. (2008). Medication effects in neuroimaging studies of bipolar disorder. *The American Journal of Psychiatry*, 165(3), 313.
- Pompei, F., Jogia, J., Tatarelli, R., Girardi, P., Rubia, K., Kumari, V., & Frangou, S. (2011). Familial and disease specific abnormalities in the neural correlates of the Stroop Task in Bipolar Disorder. *Neuroimage*, 56(3), 1677–1684.
- Quraishi, S., & Frangou, S. (2002). *Neuropsychology of bipolar disorder: A review Journal of Affective Disorders*, 72(3), 209, 226.
- Riccio, C.A., Reynolds, C.R., Lowe, P., & Moore, J.J. (2002). The Continuous Performance Test: A window on the neural substrates for attention? *Archives of Clinical Neuropsychology*, 17, 235–272.
- Sarter, M., Givens, B., & Bruno, J.P. (2001). The cognitive neuroscience of sustained attention: Where top-down meets bottom-up. *Brain Research Reviews*, 35, 146–160.
- Sepede, G., De Berardis, D., Campanella, D., Perrucci, M.G., Ferretti, A., Serroni, N., & Gambi, F. (2012). Impaired sustained attention in euthymic bipolar disorder patients and non-affected relatives: An fMRI study. *Bipolar Disorders*, 4(7), 764–779.
- Shi, T., Li, X., Song, J., Zhao, N., Sun, C., Xia, W., ... Tomoda, A. (2012). EEG characteristics and visual cognitive function of children with attention deficit hyperactivity disorder (ADHD). *Brain & Development*, 34(10), 806–811.
- Silverstein, S.M., Schenkel, L.S., Valone, C., & Nuernberger, S.W. (1998). Cognitive deficits and psychiatric rehabilitation outcomes in schizophrenia. *Psychiatric Quarterly*, 69(3), 169–191.
- Singer, J.D., & Willet, J.B. (2003). *Applied longitudinal data analysis modeling change and event occurrence*. New York: Oxford University Press.
- Snyder, P.J., Nussbaum, P.D., & Robins, D.L. (2006). *Clinical neuropsychology: A pocket handbook for assessment* (2nd ed., pp. 572–606). Washington, DC: American Psychological Association.
- Stefanopoulou, E., Manoharan, A., Landau, S., Geddes, J.R., Goodwin, G., & Frangou, S. (2009). Cognitive functioning in patients with affective disorders and schizophrenia: A meta-analysis. *International Review of Psychiatry*, 21, 336–356.
- Swann, A.C., Anderson, J.C., Dougherty, D.M., & Moeller, F.G. (2001). Measurement of inter-episode impulsivity in bipolar disorder. *Psychiatry Research*, 101(2), 195–197.
- Tucha, O., Mecklinger, L., Laufkötter, R., Klein, H.E., Walitza, S., & Lange, K.W. (2006). Methylphenidate-induced improvements of various measures of attention in adults with attention-deficit hyperactivity disorder. *Journal Neural Transmission*, 113, 1575–1592.
- Ucok, A., Cakir, S., Duman, Z.C., Dişçigil, A., Kandemir, P., & Atli, H. (2006). Cognitive predictors of skill acquisition on social problem solving in patients with schizophrenia. *European Archives of Psychiatry and Clinical Neuroscience*, 256(6), 388–394.
- Van Merriënboer, J.J., & Sweller, J. (2005). Cognitive load theory and complex learning: Recent developments and future directions. *Educational Psychology Review*, 17(2), 147–177.
- Walshe, M., Schulze, K.K., Stahl, D., Hall, M.H., Chaddock, C., Morris, R., ... Kravariti, E. (2012). Sustained attention in bipolar I disorder patients with familial psychosis and their first degree relatives. *Psychiatry Research*, 199(1), 70–73.
- Wilkinson, G.S., & Robertson, G.J. (2006). *Wide Range Achievement Test 4 (WRAT4)*. Odessa, FL: Psychological Assessment Resources, Inc.
- Wingen, M., Kuypers, K.P., van de Ven, V., Formisano, E., & Ramaekers, J.G. (2008). Sustained attention and serotonin: A pharmaco-fMRI study. *Human Psychopharmacology*, 23(3), 221–230.
- Young, R.C., Biggs, J.T., Ziegler, V.E., & Meyer, D.A. (1978). A rating scale for mania: Reliability, validity and sensitivity. *British Journal Psychiatry*, 133, 429–435.
- Youngstrom, E.A., Arnold, L.E., & Frazier, T.W. (2010). Bipolar and ADHD comorbidity: Both artifact and outgrowth of shared mechanisms. *Clinical Psychology (New York)*, 17(4), 350–359.