

Effects of morphodynamics and across-shore physical gradients on benthic macroinfauna on two sandy beaches in south-eastern Brazil

JOAQUIM N.S. SANTOS¹, RAFAELA DE S. GOMES², RUAN MANAGNA VASCONCELLOS³,
DÉBORA DE SOUZA SILVA¹ AND FRANCISCO GERSON ARAÚJO²

¹Instituto Federal do Norte de Minas Gerais—IFNMG, Fazenda Meio Pé da Serra, s/n°. BR 367, km 278, Caixa Postal 11, CEP 39600-000, Araçuaí, MG, Brasil, ²Laboratório de Ecologia de Peixes, Universidade Federal Rural do Rio de Janeiro, BR 465, km 7, CEP 23890-000 Seropédica, RJ, Brasil, ³Instituto Federal de Educação, Ciência e Tecnologia do Espírito Santo—IFES, Campus Colatina. Avenida Arino Gomes Leal, 1700, Bairro Santa Margarida, CEP 29.700-660, Colatina, ES, Brasil

We examined the benthic macroinfauna at three strata during two seasons (winter and summer) on two sandy beaches (dissipative and reflective) in south-eastern Brazil. The hypothesis raised is that effects of morphodynamics and zonation determine the structure of macroinfauna, with dissipative beaches having comparatively higher species richness and density than the reflective beaches. Flamengo beach (dissipative) had higher species richness but lower density compared to Grumari beach (reflective). A high dissimilarity in assemblage structure (91.75%) was detected between the two beaches. Zonation in the occurrence of macroinfauna was detected for the two beaches in the two examined seasons. At Grumari beach, Emerita brasiliensis occurred mainly in stratum 1 (intertidal swept zone) while Saccocirus sp. occurred in stratum 2 (infralittoral at 0.5 m depth), whereas at Flamengo beach E. brasiliensis and Enoploides sp. had the highest density in stratum 1 (intertidal swept zone) whereas Scolelepis goodbodyi and Donax uncinata dominated in stratum 3 (infralittoral at 1.0 m depth). Scolelepis goodbodyi, Dispio uncinata, Enoploides sp., Nematoda and Trileptium sp. were associated with higher a Dean parameter and content of organic matter at Flamengo beach. In contrast, Hastula sp., Donax sp., Pisionidens indica, Hemipodus californiensis, Saccocirus sp. and Phyllocididae were associated with the higher wave period and grain size of Grumari beach. The hypothesis that macroinfauna structure differs between the beaches and strata was confirmed, with the dissipative beaches having comparatively higher richness but lower density than the reflective beaches.

Keywords: benthic invertebrates, surf zones, wave exposure, benthic community, beaches

Submitted 15 August 2013; accepted 31 December 2013; first published online 31 January 2014

INTRODUCTION

Sandy beaches are dynamic systems where the continued dynamism of environmental variables determined by winds and the tidal movements on the sandy substrate influence the spatial and temporal distribution of benthic organisms (Brown & McLachlan, 1990). Short & Wright (1983) suggested that the morphodynamic state is a useful concept to classify sandy beaches in an objective way, because it refers to the depositional forms of sandy beaches and their hydrodynamic processes, namely the interactions between wave height, sediment granulometry and tide ranges. Moreover, the geological setting can influence the morphological development and beach states (Jackson *et al.*, 2005). According to the intensity of these factors, sandy beaches can be classified into two extreme states: dissipative and reflective (Defeo & McLachlan, 2005). The dissipative state is characterized by low beach slope, smooth profile, sediment with high content of organic matter and great water retention, with waves breaking and dissipating energy within an extensive area. On the

other hand, the reflective state is characterized by high beach slope, coarse sediment, low content of organic matter and strong wave exposure.

The benthic macroinfauna of sandy beaches has low richness and diversity and high dominance of a few species. Such characteristics are closely related to the morphodynamic state of the beach, with richness and diversity increasing from the reflective to the dissipative state (Brown & McLachlan, 1990). Several macroinfaunal species are tolerant to a wide spectrum of morphodynamic conditions, keeping stable population in both energy states (Velooso *et al.*, 2006). Moreover, the vertical distribution of the organisms (zonation) is a sequential species replacement, in which all or some of the species have restricted ranges of distribution (McLachlan & Jaramillo, 1995; Brazeiro & Defeo, 1996; Neves *et al.*, 2007). Four biological zones have been considered on sandy beaches according to the hydrodynamic state and interstitial water contents (Salvat, 1964): (1) dry sandy zone, at the top of the beach; (2) retention zone, slightly wet zone with water retention at ebb tide; (3) resurge zone, where water flows in and out of the sediment; and (4) saturation zone, a permanently water-saturated zone.

Seasonal changes in the density of macroinfaunal populations are common and closely associated with the

Corresponding author:
F.G. Araújo
Email: gerson@ufrjr.br

reproductive dynamic of species and with fluctuation in environmental variables, resulting in different macrofauna structures between winter and summer (Souza & Gianuca, 1995). Moreover, sandy beach urbanization is an increasing phenomenon that contributes to decreasing composition and density of benthic organisms by affecting sediment characteristics and consequently the macrofaunal community (Velooso *et al.*, 2006; Defeo *et al.*, 2009). Main threats are linked with the social and economic use of beaches such as, sand nourishment to counter erosion, the mechanical cleaning of beaches and disturbance by tourist pressure. The enrichment of coastal waters with nutrients has very likely exerted a significant influence on the macrofauna in recent years.

This study aimed to assess the effect of morphodynamics and zonation on the macrofauna assemblages during two seasons (winter 2005 and summer 2006), considering the permanently subaquatic infralittoral zone and the intertidal swept zone on two sandy beaches of south-eastern Brazil. The hypothesis raised is that morphodynamics determine the composition and relative abundance of macrofauna, with the dissipative beaches having comparatively higher richness and density than the reflective beaches, and that the structure assemblage changes according to morphodynamics and zonation.

MATERIALS AND METHODS

Study area

Flamengo beach ($22^{\circ}55'S$ $43^{\circ}10'W$) is a dissipative beach in the outer zone of Guanabara Bay (Vasconcellos *et al.*, 2010),

while Grumari beach ($23^{\circ}03'S$ $43^{\circ}32'W$) is a reflective oceanic beach (Velooso *et al.*, 2003) (Figure 1).

Flamengo beach is located near to the sea connection in the south-western part of Guanabara Bay, which is surrounded by one of the largest metropolitan areas in Brazil. The beach is approximately 1800 m long and has low amplitude semi-diurnal tides with 0.7 m of mean range. It is characterized by median to fine sand grains size with mean diameter of 0.20 mm and gentle slope profile (Vasconcellos *et al.*, 2011).

Grumari beach is located in an unprotected coastal area where the action of waves is most striking. The beach is approximately 2500 m long. It is characterized by medium to coarse sand, with mean grain size of 0.43 mm, tide range of approximately 1 m, and intertidal slope varying from 1/5.29 to 1/17.82 (Petraco *et al.*, 2003; Velooso *et al.*, 2006).

Field sampling and laboratory procedures

Sampling was conducted according to a systematic design in which three transects (fixed 30 m apart) during two seasons (winter 2005 and summer 2006) were performed. In each transect, three sampling points were defined (strata 1, 2 and 3), distributed at the spread washing zone (stratum 1), at 0.5 m depth (stratum 2) and at 1 m depth (stratum 3) (Figure 1). Samples were collected at each point for sediment analysis during ebb tide, with tide height ranging between 0.0 and 1.3 m.

Sediment samples were taken with a PVC 'corer' (50 cm long, 10 cm diameter) with a collecting area of 0.00785 m² at a depth of 15 cm. For biological samples (macrofauna), two samples and two replicates were collected at each stratum, totalling 36 samples (3 transects \times 3 strata \times 2




Fig. 1. Study area with indications of the two beaches on the coastline of Rio de Janeiro State (1, Flamengo beach; 2, Grumari beach) and the design of quantitative samplings of the sediment.

samples \times 2 replicates). Sediment samples for particle-size and organic matter (g.dm^{-3}) were determined in 18 samples per season on each beach (3 transects \times 3 strata \times 2 replicates).

The collected sediment was placed in plastic bags, then weighed (precision of 0.01 g) and dried at 80°C in a stove. A sub-sample of 300 g was collected for the particle-size analysis in sieves of different mesh size over 15 min., according to Suguio (1973). Six sieves of respectively 2 mm, 1 mm, 0.5 mm, 0.25 mm, 0.09 mm and 0.06 mm mesh size were used, which corresponded to granule fractions of very coarse, coarse, medium, fine, very fine and finest. Organic matter content (g.dm^{-3}) was obtained by the volumetric method of potassium bichromate (Walkey & Black, 1943) modified by Frattinni & Kalckmam (1967).

Biological samples were initially screened in plastic trays (80 cm \times 40 cm \times 7 cm) using tap water for removal of the largest specimens, than sieved through a 0.5 mm mesh and examined under light stereo microscope for identification of the smallest specimens. All identified specimens were preserved in 70% ethanol solution. Vouchers specimens were deposited in the macroinfauna collection of the Laboratory of Fish Ecology, Universidade Federal Rural do Rio de Janeiro.

Morphodynamics measurements

On each sampling occasion, the wave height and period were estimated. Wave height was estimated by measuring the height of the waves with graduated poles against the horizon. The wave period (T) was estimated with a chronometer and corresponded to 1/10 of the real time between eleven wave consecutive crests from a fixed point in the surf zone. Three replicates were performed to obtain an average measurement.

Classification of the morphodynamic state of each beach was assessed through the adimensional Dean parameter, Ω (Short & Wright, 1983). The Dean parameter was determined according to the following equation:

$$\Omega = H_b / (W_s \times T),$$

where H_b is the mean wave height (cm); W_s is the velocity of settling sediment (cm.s^{-1}), obtained through the mean sediment diameter, considering a density of 2.65 and a temperature of 22°C (Gibbs *et al.*, 1971) and T is the time interval (period, in seconds) between two consecutive waves. Thus, reflective beaches have $\Omega < 2$, intermediary beaches have $2 < \Omega < 5$, and dissipative beaches have $\Omega > 5$.

Data processing

The mean granules size was determined from each granulometric fraction weight retained in each sieve, using the software Sysgran 3.0 (Camargo, 2006). A multifactorial design (nested ANOVA) was used to compare differences in sediment granules and Dean parameter between seasons (fixed factor), beaches (fixed factor) and strata (nested in beaches). Wave height and period were compared between beaches and seasons, using factorial ANOVA. These analyses were performed using the software Statistica, v.8.0 (StatSoft, 2007).

Macroinfauna assemblage structure was compared between the two beaches and among the three strata using analysis of similarity (ANOSIM). A similarity matrix for these biological samples was obtained using the Bray–Curtis index with the numerical data undergoing four-root transformation to balance the rare and abundant species (Clarke & Warwick, 2001). A cluster analysis on the similarity matrix using the group average method was performed to assess eventual spatial (strata and beaches) and seasonal patterns. The species that most contributed to similarity with the groups were determined by similarity percentage—SIMPER. These analyses were performed using the software PRIMER, v.6.0 (Clarke & Warwick, 2001). Canonical correspondence analysis (CCA) was performed to assess environmental influences on the macroinfauna assemblages by using CANOCO v.4.5 (ter Braak & Šmilauer, 2002) on fourth-root transformed data.

RESULTS

Physical characteristics

SEDIMENT GRANULES

At Flamengo beach, the granule diameter ranged from 0.11 to 0.33 mm in winter, and from 0.12 to 0.23 mm in summer, with no significant difference between seasons ($F = 0.386$; $P = 0.54$) (Table 1). On the other hand, Grumari beach had coarser sediment with granule diameter ranging from 0.29 to 0.66 mm in winter and from 0.35 to 0.54 mm in summer, with no significant differences between seasons ($F = 0.377$; $P = 0.54$) (Table 1). However, a clear difference in the mean granule diameter between the two beaches (Flamengo = 0.17 ± 0.05 mm; Grumari = 0.46 ± 0.08 mm; mean \pm standard deviation) was detected ($F = 277.06$; $P = 0.0001$).

WAVE HEIGHT

At Flamengo beach, wave height ranged from 20 to 80 cm in winter and from 10 to 70 cm in summer, with no significant

Table 1. Means (\pm standard error) for environmental variables in Flamengo and Grumari beaches by seasons.

Environmental variables	Flamengo		Grumari	
	Winter	Summer	Winter	Summer
Wave height (cm)	60.83 \pm 3.55	61.33 \pm 6.79	97.78 \pm 4.39	80.56 \pm 3.52
Wave period (s)	6.83 \pm 0.46	6.83 \pm 0.33	12.67 \pm 0.27	13.33 \pm 0.69
Dean parameter (Ω)	5.07 \pm 0.01	5.01 \pm 0.13	0.97 \pm 0.09	0.80 \pm 0.04
Grain size (mm)	0.16 \pm 0.01	0.17 \pm 0.01	0.47 \pm 0.03	0.45 \pm 0.02
Organic matter (g.dm^{-3})	0.19 \pm 0.01	0.21 \pm 0.01	0.12 \pm 0.01	0.13 \pm 0.01
Morphodynamic state	Dissipative		Reflective	

difference between seasons ($F = 0.116$; $P = 0.736$) (Table 1). Grumari beach had comparatively higher wave height, ranging from 70 to 130 cm in winter, and from 40 to 90 cm in summer, with significant between-seasons differences ($F = 22.52$; $P = 0.001$) (Table 1). Significant differences ($F = 21.57$; $P = 0.0001$) were also detected between the two beaches (Flamengo = 61.6 ± 21.9 cm; Grumari = 83.7 ± 21.3 cm).

WAVE PERIOD

No significant difference was found for the wave period on Flamengo beach between seasons ($F = 0.00$; $P = 1.0$), with values ranging from 4 to 10 s in winter, and from 5 to 9 s in summer (Table 1). On Grumari beach, the wave period ranged from 11 to 14 s in winter and from 9 to 19 s in summer, with no significant difference between seasons ($F = 1.42$; $P = 0.243$) (Table 1). Significant differences in the wave mean period were found between the two beaches ($F = 175.64$; $P = 0.0001$) with comparatively higher values for Grumari beach (13.0 ± 2.1 s) compared with Flamengo beach (6.8 ± 1.6 s).

DEAN PARAMETER (Ω)

At Flamengo beach, the Dean parameter was >5 , therefore, this beach was classified as dissipative. On the other hand, Grumari beach had a Dean parameter <1 and was classified as a reflective beach. At both beaches, no significant differences for Dean parameter were detected between seasons ($F = 4.0$; $P = 0.054$) and strata ($F = 2.0$; $P = 0.083$) (Table 1), although a significant difference was found between the two beaches ($F = 2253.0$; $P = 0.0001$).

ORGANIC MATTER (g.dm^{-3})

The content of organic matter in sediment differed significantly between seasons and beaches (Table 1). Significantly higher values were recorded in summer compared with the winter at both Flamengo (t -test = -5.75 ; $P = 0.0001$) and Grumari (t -test = -3.46 ; $P = 0.003$) beaches. There was a clear difference between the beaches, with significant higher

values for Flamengo beach ($0.201 \pm 0.005 \text{ g.dm}^{-3}$) compared with Grumari beach ($0.125 \pm 0.003 \text{ g.dm}^{-3}$) (t -test = 20.64 ; $P < 0.0001$).

Beach comparisons

A total of 12 taxa was recorded at the two examined beaches, five of them at species level, five at genus level, one at family level and one at phylum level. Polychaeta and Nematoda were the numerically dominant groups (Table 2).

At Flamengo beach, eight taxa were recorded: (1) Polychaeta: *Dispio uncinata* Hartman, 1951; *Scolecopsis goodbodyi* Jones, 1962; and *Pisionidens indica* (Aiyar & Alikuhni, 1940); (2) Crustacea Decapoda: *Emerita brasiliensis* Schmitt, 1935; (3) two Nematoda genus: *Trileptium* sp. and *Enoploides* sp., (4) specimens of Nemertea; and (5) Mollusca (*Donax* sp.). *Enoploides* sp. and *E. brasiliensis* were the numerically dominant taxa in winter (43.4% and 21.1% of the total number of individuals, respectively) and in the summer (57.3% and 19.3%, respectively) (Table 2). *Emerita brasiliensis* made the highest contribution to biomass, with 93.2% of the total weight in the winter and 95.6% in the summer (Table 2). The highest density (individuals.m⁻²) was recorded in summer, except for *Trileptium* sp. and Nemertea, with the highest contributions from *Enoploides* sp. (3386.4 individuals.m⁻²) and *E. brasiliensis* (1139.4 individuals.m⁻²) (Table 3). The highest Margalef richness was recorded in winter (0.879) (Table 3).

At Grumari beach, a total of seven taxa was recorded: (1) Polychaeta: *Saccocirus* sp.; specimens of the *Phyllodocidae* family; *P. indica*; and *Hemipodus californiensis* (Hartman, 1938); (2) *Emerita brasiliensis*; and (3) Mollusca: *Donax* sp. and *Hastula* sp. In winter, the numerically dominant groups were *Saccocirus* sp. (84.9% of the total number of individuals) and *E. brasiliensis* (8.67%), and a similar pattern was also recorded in summer, when *Saccocirus* sp. reached 98.5% of total number of individuals (Table 2). *Emerita brasiliensis* made the greatest contribution to biomass, with 94.5% of the total weight in winter, and 85.58% in summer (Table 2).

Table 2. Percentage and absolute (in parentheses) values for number and biomass (weight) of macroinfauna species in Flamengo and Grumari beaches by seasons. Seasonal comparisons for absolute values (P -values) according to Student's t -test also indicated. N, number of individuals.

Flamengo	Number		P -values	Weight (%)		P -values
	Winter	Summer		Winter	Summer	
<i>Enoploides</i> sp.	43.42 (N = 399)	57.34 (N = 957)	0.001	<0.001 (g)	<0.001 (g)	–
<i>E. brasiliensis</i>	21.11 (194)	19.29 (322)	0.02	93.23 (44.62)	95.60 (74.10)	0.002
<i>Trileptium</i> sp.	9.58 (88)	3.83 (64)	0.05	<0.001	<0.001	–
<i>D. uncinata</i>	9.38 (86)	6.35 (106)	0.004	4.13 (2.98)	1.86 (2.44)	0.09
<i>S. goodbodyi</i>	8.27 (76)	9.47 (158)	0.002	2.06 (1.99)	2.42 (2.87)	0.06
Nemertea	8.05 (74)	2.99 (51)	0.05	<0.001	<0.001	–
<i>P. indica</i>	–	0.72 (12)	–	–	0.13	–
<i>Donax</i> sp.	0.22 (2)	–	–	0.58	–	–
Grumari						
<i>Saccocirus</i> sp.	84.99 (3.743)	98.46 (52.254)	0.0001	0.83 (0.67)	5.52 (6.90)	0.0001
Phyllodocidae	5.54 (244)	0.36 (194)	0.05	0.04 (0.62)	0.60 (0.43)	0.08
<i>E. brasiliensis</i>	8.67 (382)	0.98 (522)	0.04	94.54 (52.72)	85.58 (93.96)	0.0001
<i>P. indica</i>	0.45 (20)	0.09 (49)	0.05	0.36 (0.21)	0.46 (0.52)	0.04
<i>H. californiensis</i>	–	0.02 (13)	–	–	0.93	–
<i>Donax</i> sp.	0.34 (15)	0.07 (37)	0.003	4.03	6.25	0.02
<i>Hastula</i> sp.	–	0.01 (4)	–	–	0.66	–

Table 3. Densities (individuals.m⁻²), richness and total number of individuals in Flamengo and Grumari beaches by seasons. Seasonal comparisons for species densities (*P*-values) according to Student's *t*-test also indicated.

Taxa	Flamengo			Grumari		
	Winter	Summer	<i>P</i> -values	Winter	Summer	<i>P</i> -values
<i>Enoploides</i> sp.	1415.4	3386.4	0.001	–	–	–
<i>E. brasiliensis</i>	686.5	1139.4	0.01	1351.73	1847.13	0.02
<i>Trileptium</i> sp.	311.4	226.5	0.04	–	–	–
<i>D. uncinata</i>	304.3	375.0	0.05	–	–	–
<i>S. goodbodyi</i>	268.9	559.1	0.03	–	–	–
Nemertea	261.9	176.9	0.06	–	–	–
<i>P. indica</i>	–	42.5	–	70.77	176.93	0.05
<i>Saccocirus</i> sp.	–	–	–	13244.87	184904.0	0.0001
Phyllodocidae	–	–	–	863.41	686.48	0.04
<i>H. californiensis</i>	–	–	–	–	46.00	–
<i>Donax</i> sp.	7.1	–	–	53.08	130.93	0.04
<i>Hastula</i> sp.	–	–	–	–	14.15	–
Total density (ind.m ²)	3255.5	5904.7		15583.9	187806.1	
Total number of individuals	920	1669		24374	53084	
Numbers of taxa	7	7		5	7	
Margalef richness (seasons)	0.879	0.809		0.477	0.552	
Margalef richness (beaches)	0.8907			0.5475		

The highest densities were recorded in summer, except for Phyllodocidae, with *Saccocirus* sp. having 184,904 individuals.m⁻² and *E. brasiliensis* 1847.1 individuals.m⁻² (Table 3). The highest Margalef richness (0.552) was recorded in summer.

The highest densities were recorded at Grumari beach (187,806.1 individuals.m⁻²), while the highest Margalef richness was at Flamengo beach (0.891) (Table 3). Three taxa were common at the two beaches: *P. indica*, *E. brasiliensis* and *Donax* sp. *Scolecopsis goodbodyi*, *D. uncinata*, *Trileptium* sp. and *Enoploides* sp. were recorded exclusively on Flamengo beach, while *Saccocirus* sp., *H. californiensis*, *Hastula* sp. and Phyllodocidae occurred on Grumari beach only.

Cluster analysis on macroinfauna density showed a clear separation between the two beaches, which were clustered in different groups (Figure 2). ANOSIM used to compare assemblage structure indicated the macroinfauna differed between the two beaches (*R* global = 0.835, *P* < 0.001). The SIMPER analysis showed a high dissimilarity between the two

beaches (91.75%), with *Saccocirus* sp., *Enoploides* sp., *D. uncinata*, *S. goodbodyi* and *E. brasiliensis* explaining 63.70% of the difference between the two beaches (Table 4). Species characteristics for Flamengo beach had 55.54% of average similarity, with the greatest contribution from *D. uncinata*, *S. goodbodyi* and *Enoploides* sp. At Grumari beach, the characteristic species that explained most of the 37.90% average similarity were *Saccocirus* sp., Phyllodocidae and *E. brasiliensis* (Table 4).

Environmental variables and macroinfauna assemblages

The first two axes of CCA explained 92.8% of the total variance of the species–environment correlation. The first axis revealed a well-defined spatial pattern and separated the two beaches (Figure 3), whereas the second axis showed a slight separation along the strata for each beach. According to the


**Fig. 2.** Dendrogram from cluster analysis on the density of the macroinfauna of Flamengo (FLA) and Grumari (GRU) beaches.

Table 4. Species contribution for the average within similarity in Flamengo and Grumari beaches according to SIMPER analysis. Dissimilarity also indicated.

Similarity	Flamengo	Grumari	Flamengo × Grumari	
	55.44	37.90	Dissimilarity	91.75
<i>D. uncinata</i>	25.05	–	<i>Saccocirus</i> sp.	17.95
<i>S. goodbodyi</i>	23.77	–	<i>Enoploides</i> sp.	12.47
<i>Enoploides</i> sp.	19.72	–	<i>D. uncinata</i>	11.23
Nemertea	11.79	–	<i>S. goodbodyi</i>	11.22
<i>E. brasiliensis</i>	9.94	23.37	<i>E. brasiliensis</i>	10.82
<i>Saccocirus</i> sp.	–	32.44	Phyllodocidae	9.10
Phyllodocidae	–	23.78	Nemertea	7.52
<i>P. indica</i>	–	10.31	<i>Trileptium</i> sp.	6.89
<i>Donax</i> sp.	–	8.86	<i>P. indica</i>	5.56

CCA, *S. goodbodyi*, *D. uncinata*, *Enoploides* sp., Nematoda and *Trileptium* sp. were closely associated with higher values of organic matter and the Dean parameter of Flamengo beach. In contrast, *Hastula* sp., *Donax* sp., *P. indica*, *H. californiensis*, *Saccocirus* sp. and Phyllodocidae were associated with the higher values of wave period and grain size of Grumari beach. *Emerita brasiliensis* appeared near the centre of the diagram, suggesting a wide spatial distribution across the two beaches (Figure 3).

Strata comparisons

FLAMENGO BEACH

According to ANOSIM, different structures of the macroinfauna were found among the three examined strata ($R_{\text{global}} = 0.424$, $P = 0.03$). A spatially differentiated pattern suggesting zonation of the macroinfaunal community was found by cluster analysis on the density data (Figure 4A).




Fig. 3. Ordination diagram (triplet) from canonical correspondence analysis of macroinfauna species abundance, environmental variables and seasons. Species code: Sgo, *Scolecis goodbodyi*; Dun, *Dispio uncinata*; Eno, *Enoploides* sp.; Nem, Nematoda; Tri, *Trileptium* sp.; Has, *Hastula* sp.; Don, *Donax* sp.; Pin, *Pisionidens indica*; Hca, *Hemipodus californiensis*; Sac, *Saccocirus* sp.; Eme, *Emerita brasiliensis*; Phy, Phyllodocidae. Environmental variables code: OrgMat, organic matter; Dean, Dean parameter; Wp, wave period; Grain, grain size; Wh, wave height. Samples code (beach, season and stratum), e.g. FS1, Flamengo beach, summer, stratum 1; GW3, Grumari beach, winter, stratum 3.

According to ANOSIM, strata 1 and 2 did not differ significantly ($R = 0.168$, $P = 0.02$), with SIMPER analysis indicating as characteristic species from stratum 1 the taxa *Enoploides* sp. and *E. brasiliensis*, which contributed 40.57% and 30.02% to the average similarity, respectively (Table 5), whereas *Enoploides* sp., *D. uncinata* and *S. goodbodyi* were the species with greatest contribution to the within similarity in stratum 2 (Table 5). Dissimilarity between these two strata was only 44.27%, with the greatest contributions from *Enoploides* sp. (34.22%) and *E. brasiliensis* (23.46%) (Table 5).

ANOSIM indicated significant difference in structure assemblages between strata 1 and 3 ($R = 0.704$, $P = 0.001$), with SIMPER analysis determining that the Polychaeta *S. goodbodyi* and *D. uncinata* had the greatest contribution to within similarity in stratum 3. Average dissimilarity between these two strata was 70.45%, with *Enoploides* sp. (31.83%) and *E. brasiliensis* (27.12%) having the greatest contribution (Table 5).

Strata 2 and 3 differed significantly in structure assemblage, although with some indication of species overlapping because of the intermediary R -value (ANOSIM, $R_{\text{global}} = 0.418$, $P = 0.01$). Average dissimilarity between these two strata was 54.69%, with greatest contribution from *Enoploides* sp. (30.61%) and *E. brasiliensis* (18%) (Table 5).

Seasonally, the macroinfauna structure differed slightly between summer and winter for the pooled strata ($R_{\text{global}} = 0.361$, $P = 0.01$). Only stratum 1 had no significant difference in assemblage structure ($R = 0.232$, $P = 0.02$) between seasons, with low percentage of dissimilarity (31.53%) and the greatest contribution from *Enoploides* sp. (33.46%) (Table 6).

GRUMARI BEACH

The assemblage structure had significant differences between the three examined strata according to ANOSIM ($R_{\text{global}} = 0.41$, $P = 0.01$) and cluster analysis (Figure 4B), indicating differences in assemblage macroinfauna structure between the three strata.

Strata 1 and 2 did not differ in assemblage structure ($R_{\text{global}} = 0.128$, $P = 0.03$) according to ANOSIM (Table 5). In stratum 1, *E. brasiliensis* and *Saccocirus* sp. were the species to contribute most to within similarity, with 39.22% and 37.65% of the average similarity, respectively (Table 6). In stratum 2, *Saccocirus* sp. (47.42%) had the greatest contribution to within similarity. Average dissimilarity between these two strata was only 50.86%, with the greatest contribution from *Saccocirus* sp. (41.55%) (Table 6).

A significant difference in assemblage structure was found between strata 1 and 3 ($R = 0.689$, $P = 0.01$), with SIMPER analysis indicating *Saccocirus* sp. and *E. brasiliensis* as having the greatest contribution to average similarity in stratum 3 (Table 6). Average dissimilarity between these two strata (81.70%) was mainly due to the contribution from *Saccocirus* sp. (38.52%) and *E. brasiliensis* (26.13%) (Table 6).

Strata 2 and 3 differed significantly in assemblage structure according to ANOSIM ($R_{\text{global}} = 0.445$, $P = 0.01$). Average dissimilarity between strata 2 and 3 was 71.97%, with the greatest contribution from *Saccocirus* sp. (37.28%) (Table 6).

Seasonally, a slight difference was found for assemblage structure between the summer and winter for the pooled strata ($R_{\text{global}} = 0.463$, $P = 0.01$). However, stratum 3 did not differ between seasons ($R_{\text{global}} = 0.201$, $P = 0.42$) (Table 6). In strata 1 and 2, *Saccocirus* sp. made the greatest




Fig. 4. Dendrogram from cluster analysis of the density of the macrofauna in Flamengo (A) and Grumari (B) beaches, with samples coded by strata. E1, intertidal swept zone; E2, infralittoral at 0.5 m depth; E3, infralittoral at 1.0 m depth.

Table 5. Species contribution for the average within similarity by strata in Flamengo and Grumari beaches according to SIMPER analysis. Dissimilarity also indicated. Strata: 1, spread washing zone; 2, infralittoral at 0.5 m depth; 3, infralittoral at 1.0 depth.

	Stratum			Dissimilarity		
	1	2	3	(1 × 2)	(1 × 3)	(2 × 3)
Flamengo						
Taxa / similarity	62.60	59.31	51.40	44.27	70.45	54.69
<i>Enoploides</i> sp.	40.57	20.56	–	34.22	31.83	30.61
<i>E. brasiliensis</i>	30.02	13.88	–	23.46	27.12	18.00
<i>D. uncinata</i>	9.71	20.77	36.23	–	5.76	10.13
<i>Trileptium</i> sp.	8.54	12.84	–	11.33	10.14	13.22
<i>S. goodbodyi</i>	–	20.43	49.89	12.27	12.64	11.27
Nemertea	8.41	11.52	5.63	8.90	7.83	12.74
Grumari						
Taxa / similarity	56.23	50.00	35.90	50.86	81.70	71.97
<i>E. brasiliensis</i>	39.22	23.27	–	18.03	26.13	19.00
<i>Saccocirus</i> sp.	37.65	47.42	–	41.55	38.52	37.28
Phyllodocidae	21.35	13.86	22.91	14.99	13.28	13.67
<i>P. indica</i>	–	11.70	37.94	10.51	8.53	12.30
<i>Donax</i> sp.	–	–	37.75	7.59	7.85	11.37

contributions (60.99% and 37.57%, respectively) for average dissimilarity between seasons (Table 6).

DISCUSSION

The macrofauna assemblages structure in the two studied sandy beaches were closely associated to the morphodynamic

Table 6. *R* and *P*-values for seasonal (winter × summer) comparisons of macrofauna assemblage structure in each stratum at Flamengo and Grumari beaches according to ANOSIM. Average dissimilarity for significant differences also indicated.

Stratum (winter × summer)	<i>R</i>	<i>P</i> -values	Average dissimilarity
Flamengo			
1	0.232	0.02	31.53
2	0.109	0.18	–
3	–0.069	0.71	–
Grumari			
1	0.635	0.004	52.96
2	0.304	0.013	53.13
3	–0.02	0.42	–

states. High grain size and wave periods at Grumari beach determined the predominance of the *Saccocirrus* sp., whereas high organic matter at Flamengo beach favoured high abundances of *Enoploides* sp. Grain size often is coarser at reflective rather than at dissipative beaches as a result of intense wave action at the former (McLachlan, 1990; Defeo *et al.*, 1992; McArdle & McLachlan, 1992; Borzone *et al.*, 1996). Close association between high abundances of *Saccocirrus* with coarse sands and high wave period was also reported by Di Domenico *et al.* (2009) for reflective beaches in southern Brazil, suggesting that exposed beaches provide an ideal environment for the occurrence of *Saccocirrus* species. Conversely, the predominance of fine sediment particles at Flamengo beach favoured high abundances of *Enoploides* sp. Maria *et al.* (2013) found that the sediment characteristics influence the nematode community distribution, among them *Enoploides* sp., and these findings coincide with the results of the present study.

Emerita brasiliensis was abundant in both beaches irrespective of the morphodynamic state. Veloso *et al.* (2003) also did not detect significant differences in densities of *E. brasiliensis* on 15 exposed beaches, both dissipative and reflective, along the coast of Rio de Janeiro. Therefore, it is reasonable to suppose that *E. brasiliensis* has a high ability to occur in different environmental conditions, and that the morphodynamic state seems to play a secondary role in determining its distribution on sandy beaches.

Our findings pointed to a higher richness at the dissipative Flamengo beach, which is in accordance with the expectation that the morphodynamic state is the main factor determining macroinfaunal richness on sandy beaches (McLachlan, 1990; Defeo *et al.*, 1992; Borzone *et al.*, 1996), with an increasing trend from the reflective to the dissipative state. However, lower densities at Flamengo beach could be associated with high anthropogenic influences on this beach, and the allochthonous material that arrives into the sea may affect the macroinfauna community. The higher density of *Saccocirrus* sp. at Grumari beach is not in accordance with Dexter (1992), who reported Polychaeta preferring sheltered beaches. On the other hand, a higher diversity of these animals in areas with poorly-selected sediments with biogenic contributions (shell fragments) rather than in muddy or sandy-muddy sediments was found by Villora-Moreno (1997). *Saccocirrus* sp. was not recorded in previous studies at Grumari beach (Veloso *et al.*, 2003, 2006). Differences between the findings of Veloso *et al.* (2003, 2006) and this study could be due to the larger sampling size in the present study, which also incorporates a permanently-submersed zone (strata 2 and 3), while those studies covered only the spread washing zone.

A clear zonation was found for the macroinfauna assemblage along the three strata on the two beaches, with more evident differences at Grumari beach. The most abundant taxon in stratum 1 (spread washing zone) was *E. brasiliensis* in both seasons, with comparatively higher values in the summer. According to Defeo & Cardoso (2002) this suspensivorous-filtering crustacean is found in the tidal zones of reflective and dissipative beaches, with comparatively higher values on the reflective beaches. In the most studied Rio de Janeiro sandy beaches, *E. brasiliensis* occurs in high densities, mainly in those with a reflective state (Veloso *et al.*, 1997, 2006; Cardoso *et al.*, 2003), and our results corroborate those findings, since the highest density of *E. brasiliensis* was

recorded for Grumari beach. Fernandes & Soares-Gomes (2006) found that *E. brasiliensis* persists on beaches with higher sediment thrust caused by water pressure, which favours its suspensivorous-filtering habit. Veloso & Cardoso (1999), Gianuca, (1985) and Petracco *et al.* (2003) found higher densities of *E. brasiliensis* during spring and summer, which partially corroborates our findings of highest density during the summer.

At Grumari beach, the predominance of *Saccocirrus* sp. in stratum 2, and *P. indica* and *Donax* sp. in stratum 3 suggests a clear pattern of vertical zonation for this reflective beach. According to several studies (e.g. Hill & Hunter, 1976; Leber, 1982; Knott *et al.*, 1983) the species composition between the tidal and infralittoral zones differs because of the micro-habitat selection and hydrodynamic state that determines the stability of the communities. Jaramillo (1987) and McLachlan *et al.* (1993) found that macroinfaunal diversity and density increase with depth, which is in accordance with the findings of the present study.

In Flamengo beach, a high similarity between strata 1 and 2 was most influenced by the common occurrence of *Enoploides* sp. (Nematoda) and *E. brasiliensis* taxa, which had the highest density in summer. Giere (1993) observed that the tidal zone offers favourable conditions for Nematoda occurrence, and Rodriguez *et al.* (2001) found that this group of animals has the greatest density in dissipative sandy beaches. A similar pattern was observed in the present study, with the great density of *Enoploides* sp. at Flamengo beach, which coincided with the greatest content of organic matter. According to Fleeger & Decho (1987), the density of Nematoda is related to the quantity of organic matter in the sediment. In stratum 3, *S. goodbodyi* and *D. uncinata* were the most representative taxa. *Scolecopsis goodbodyi* is recognized as a dominant species in beaches of intermediary to dissipative state (Souza & Gianuca, 1995; Borzone *et al.*, 1996, 2003; Barros *et al.*, 2001). Souza & Gianuca (1995) associated the highest density of this species to the recruitment season in the summer.

In this study, we found that the dissipative beach had a different macroinfauna composition and higher richness compared to the reflective beach, and that this latter beach had a comparatively higher density. Furthermore, the macroinfauna also differed between strata, mainly between the spread washing zone (stratum 1) and the deeper infralittoral zone (stratum 3). Therefore, the hypothesis of different macroinfauna composition between the beaches and strata was confirmed, with the dissipative beaches having comparatively higher richness but lower density than the reflective beaches.

ACKNOWLEDGEMENTS

We thank Dr Paulo Cesar de Paiva (UFRJ) for helping in Polychaeta identification. We also thank Ana Paula Penha Guedes, Antonio Gomes da Cruz-Filho and Benjamin Carvalho T. Pinto for helping in fieldwork.

FINANCIAL SUPPORT

The authors are grateful to the FAPERJ—Fundação Carlos Chagas de Amparo à Pesquisa do Estado do Rio de Janeiro

and CNPq—Conselho Nacional de Desenvolvimento Científico Tecnológico, for financial support.

REFERENCES

- Barros F., Borzone C.A. and Rosso S.** (2001) Macrofauna of six beaches near Guaratiba bay, Southern Brazil. *Brazilian Archives of Biology and Technology* 44, 351–364.
- Borzone C.A., Melo S.G., Rezende K.V., Vale R. and Krul R.** (2003) Macrobenthic intertidal communities from wave to tide dominated beach environments. A case study in two Brazilian beaches. *Journal of Coastal Research* 35, 472–480.
- Borzone C.A., Souza J.R.B. and Soares A.G.** (1996) Morphodynamic influence on the structure of inter and subtidal macrofaunal communities of subtropical sandy beaches. *Revista Chilena de História Natural* 69, 565–577.
- Brazeiro A. and Defeo O.** (1996) Macrofauna zonation in macrotidal sandy beaches: is it possible to identify patterns in such variable environments? *Estuarine, Coastal and Shelf Science* 42, 523–536.
- Brown A.C. and McLachlan A.** (1990) *Ecology of sandy shores*. Amsterdam: Elsevier.
- Camargo M.G.** (2006) SYSGRAN: Um sistema de código aberto para análises granulométricas do sedimento. *Revista Brasileira de Geociências* 36, 371–378.
- Cardoso R.S., Veloso V.G. and Caetano C.H.S.** (2003) Life history of *Emerita brasiliensis* (Decapoda: Hippidae) on two beaches with different morphodynamic characteristics. *Journal of Coastal Research* 35, 392–401.
- Clarke K.R. and Warwick R.M.** (2001) *Changes in marine communities: an approach to statistical analysis and interpretation*. 2nd edition Plymouth: Primer E.
- Defeo O. and Cardoso R.S.** (2002) Macroecology of population dynamics and life history traits of the mole crab *Emerita brasiliensis* in Atlantic sandy beaches of South America. *Marine Ecology Progress Series* 239, 169–179.
- Defeo O. and McLachlan A.** (2005) Patterns, processes and regulatory mechanisms in sandy beach macrofauna: a multi scale analysis. *Marine Ecology Progress Series* 295, 1–20.
- Defeo O., McLachlan A., Shoeman D.S., Schilancher A., Dugan J., Jones A., Lastra M. and Scapini F.** (2009) Threats to sandy beach ecosystems: a review. *Estuarine, Coastal and Shelf Science* 81, 1–12.
- Defeo O., Ortiz E. and Castilla J.C.** (1992) Growth mortality and recruitment of the yellow clam *Mesodesma mactroides* on Uruguayan beaches. *Marine Biology* 114, 429–437.
- Dexter D.M.** (1992) Sandy beach community structure: the role of exposure and latitude. *Journal of Biogeography* 19, 59–66.
- Di Domenico M., Lana P.C. and Garraffoni A.R.S.** (2009) Distribution patterns of interstitial polychaetes in sandy beaches of southern Brazil. *Marine Ecology* 30, 47–62.
- Fernandes R.S.R. and Soares-Gomes A.** (2006) Community structure of macrobenthos in two tropical sandy beaches with different morphodynamic features, Rio de Janeiro, Brazil. *Marine Ecology* 27, 160–169.
- Fleeger J.W. and Decho A.W.** (1987) Spatial variability of interstitial meiofauna: a review. *Stygologia* 3, 35–48.
- Frattini C.T.A. and Kalkman R.E.** (1967) Correlação entre alguns métodos de determinação do carbono. *Pesquisa Agropecuária Brasileira* 2, 259–261.
- Gianuca N.M.** (1985) *The ecology of a sandy beach in Southern Brazil*. PhD thesis. University of Southampton, UK.
- Gibbs R.J., Matthews M.D. and Link D.A.** (1971) The relationship between sphere size and settling velocity. *Journal of Sedimentary Petrology* 41, 7–18.
- Giere O.** (1993) *Meiobenthology, the microscopic fauna in aquatic sediments*. Berlin: Springer.
- Hill G.M. and Hunter R.E.** (1976) Interaction of biological and geological process in the beach and nearshore, northern Padre Island, Texas. In Davis Jr R.A. and Ethington R.L. (eds) *Beach and nearshores sedimentation*. Tulsa, OK: Society of Economic Paleontologists and Mineralogists, pp. 169–187.
- Jackson D.W.T., Cooper J.A.G. and Rio L.D.** (2005) Geological control of beach morphodynamic state. *Marine Geology* 216, 297–314.
- Jaramillo E.** (1987) Sandy beach macrofauna from the Chilean coast: zonation patterns and zoogeography. *Vie et Milieu* 37, 165–174.
- Knott D.M., Calder D.R. and VanDolah R.F.** (1983) Macrobenthos of sandy beach and nearshore environments at Murrells Inlet, South Carolina, U.S.A. *Estuarine, Coastal and Shelf Science* 16, 573–590.
- Leber K.M.** (1982) Seasonality of macroinvertebrates on a temperate, high wave energy sandy beach. *Bulletin of Marine Science* 32, 86–98.
- Maria T.F., Paiva P., Vanreusel A. and Esteves A.M.** (2013) The relationship between sandy beach nematodes and environmental characteristics in two Brazilian sandy beaches (Guanabara Bay, Rio de Janeiro). *Anais da Academia Brasileira de Ciências* 85, 25–270.
- McArdle S. and McLachlan A.** (1992) Sand beach ecology: swash factors relevant to the macrofauna. *Journal of Coastal Research* 8, 398–407.
- McLachlan A.** (1990) Dissipative beaches and macrofauna communities on exposed intertidal sands. *Journal of Coastal Research* 6, 57–71.
- McLachlan A. and Jaramillo E.** (1995) Zonation on sandy beaches. *Oceanography and Marine Biology: an Annual Review* 33, 305–335.
- McLachlan A., Jaramillo E., Donn T.E. and Wessels F.** (1993) Sandy beach macrofauna communities and their control by the physical environment: a geographical comparison. *Journal of Coastal Research* 15, 27–38.
- Neves L.P., Da Silva P.R. and Bemvenuti C.E.** (2007) Zonation of benthic macrofauna on Cassino Beach, southern most Brazil. *Brazilian Journal of Oceanography* 55, 293–307.
- Petracco M., Veloso V.G. and Cardoso R.S.** (2003) Population dynamics and secondary production of *Emerita brasiliensis* (Crustacea: Hippidae) at Prainha beach, Brazil. *Marine Ecology* 24, 231–245.
- Rodriguez J.G., Lopez J. and Jaramillo E.** (2001) Estructura comunitaria de la meiofauna intermareal en un gradiente de tipos morfodinámicos de playas arenosas en el sur de Chile. *Revista Chilena de Historia Natural* 74, 885–897.
- Salvat B.** (1964) Les conditions hydrodynamiques interstitielles des sédiments meubles intertidaux et la repartition verticale de la faune endogée. *Cahiers de Recherche de l'Académie de Science de Paris* 259, 43–56.
- Short A.D. and Wright L.D.** (1983) Physical variability of sandy beaches. In McLachlan A. and Erasmus T. (eds) *Sandy beaches as ecosystems. 1st International Symposium on Sandy Beaches, Port Elizabeth, South Africa, 17–21 January 1983*. The Hague: W. Junk, pp. 133–144.
- Souza J.R.B. and Gianuca N.M.** (1995) Zonation and seasonal variation of the intertidal macrofauna on a sandy beach of Paraná State, Brazil. *Scientia Marina* 59, 103–111.
- StatSoft Inc.** (2007) *STATISTICA (data analysis software system)*, version 8.0. Available at: www.statsoft.com (accessed 9 January 2014).
- Suguio K.** (1973) *Introdução a sedimentologia*. São Paulo: EDUSP.

- Ter Braak C.F.J. and Šmilauer P.** (2002) *CANOCO reference manual and CanoDraw for Windows user's guide: software for canonical community ordination* (version 4.5). Ithaca, NY: Microcomputer Power.
- Vasconcellos R.M., Araújo F.G., Santos J.N.S. and Silva M.A.** (2011) Diel seasonality in fish biodiversity in a sandy beach in south-eastern Brazil. *Journal of the Marine Biological Association of the United Kingdom* 91, 1337–1344.
- Veloso V.G., Silva E.S., Caetano C.H.S. and Cardoso R.S.** (2006) Comparisons between the macroinfauna of urbanized and protected beaches in Rio de Janeiro State, Brazil. *Biological Conservation* 27, 510–515.
- Veloso V.G., Caetano C.H.S. and Cardoso R.S.** (2003) Composition, structure and zonation of intertidal macroinfauna in relation to physical factors in microtidal sandy beaches in Rio de Janeiro state, Brazil. *Scientia Marina* 67, 393–402.
- Veloso V.G. and Cardoso R.S.** (1999) Population biology of the mole crab *Emerita brasiliensis* (Decapoda: Hippidae) at Fora beach, Brazil. *Journal of Crustacean Biology* 19, 147–153.
- Veloso V.G., Cardoso R.S. and Fonseca D.B.** (1997) Spatio-temporal characterization of intertidal macrofauna community at Prainha Beach, State of Rio de Janeiro. *Oecologia Brasiliensis* 3, 213–225.
- Villora-Moreno S.** (1997) Environmental heterogeneity and the biodiversity of interstitial polychaeta. *Bulletin of Marine Science* 60, 494–501.
- and
- Walkey A. and Black J.A.** (1943) An examination of the Degtjareff method for determining soil organic matter, and proposed modification of the chronic acid titration method. *Soil Science* 37, 29–38.

Correspondence should be addressed to:

F.G. Araújo
Laboratório de Ecologia de Peixes, Universidade Federal Rural do Rio de Janeiro, BR 465, km 7, CEP 23890-000 Seropédica RJ, Brasil
email: gereson@ufrj.br