

Vaccination of eels (*Anguilla japonica* and *Anguilla anguilla*) against *Anguillicola crassus* with irradiated L₃

K. KNOPF^{1*} and R. LUCIUS²

¹ Leibniz-Institute of Freshwater Ecology and Inland Fisheries, Müggelseedamm 310, 12587 Berlin, Germany

² Department of Molecular Parasitology, Humboldt-University Berlin, Philippstraße 13, Haus 14, 10115 Berlin, Germany

(Received 22 November 2007; revised 27 December 2007; accepted 3 January 2008; first published online 27 February 2008)

SUMMARY

The original host of the swimbladder nematode *Anguillicola crassus*, the Japanese eel (*Anguilla japonica*) and the recently colonized European eel (*Anguilla anguilla*) were immunized with 40 irradiated (500 Gy) 3rd-stage larvae (L₃) of this parasite and challenged with an infection of 40 normal L₃. The immunization induced a significant reduction of the number of adult worms developing from the challenge infection in *A. japonica*, but not in *A. anguilla*. The induced resistance (calculated using the relation of the number of adult worms in immunized eels and in non-immunized control eels) in *A. japonica* was 87.3% ± 30.4%. Following a single infection, the percentage of adult worms found in *A. japonica* was lower as compared to *A. anguilla*, and the few adult worms were much smaller, revealing a lower susceptibility of *A. japonica* to *A. crassus* in comparison to *A. anguilla*. Both eel species developed an antibody response against *A. crassus*, but the level of antibody responses was not positively correlated with the protection against infection, suggesting that the antibody response is not a key element in resistance of eels against *A. crassus*. This study suggests that the original host of *A. crassus* is able to mount efficient protective immune responses against its parasite, whereas the newly acquired host seems to lack this ability.

Key words: *Anguillicola crassus*, *Anguilla japonica*, *Anguilla anguilla*, irradiated L₃, vaccination, resistance, antibody response.

INTRODUCTION

Anguillicola crassus is a nematode that develops in the swimbladder of eels. This parasite originates from East Asia where it is a parasite of the Japanese eel (*Anguilla japonica*). Introduced to Europe about 25 years ago and a few years later to Northern America, *A. crassus* spread within stocks of the endemic European eel (*Anguilla anguilla*) and American eel (*Anguilla rostrata*), respectively. Whereas little is known about the epidemiology of *A. crassus* in Northern America, a number of studies documented the successful and fast spread of *A. crassus* over almost all of Europe, where it became one of the most prevalent parasites of *A. anguilla* (Sures *et al.* 1999; Sures and Streit, 2001; Kirk, 2003).

So far, neither field studies nor experiments revealed evidence for a protective immunity against *A. crassus* in *A. anguilla* (Knopf, 2006). However, a recent experimental study on the infectivity of *A. crassus* in *A. japonica* and *A. anguilla* showed that the original host *A. japonica* is less susceptible and obviously possesses more effective defence

mechanisms against this parasite compared to the newly acquired host *A. anguilla* (Knopf and Mahnke, 2004). A single infection with thirty 3rd-stage larvae (L₃) of *A. crassus* resulted in an approximately 3 times higher recovery rate in *A. anguilla* compared to *A. japonica*, with a 10 times higher wet weight of parasites in *A. anguilla*. Only 27% of the recovered worms became adult in *A. japonica*, but 94% of the worms reached maturity in *A. anguilla* during a 98-day experiment. The fact that dead, encapsulated and necrotic larvae (almost 60% of the number of recovered parasites) were only found in *A. japonica* suggested the presence of protective immune effector mechanisms in the original host. However, such differences in susceptibility could theoretically also be related to factors other than immune responses, such as physiological differences or lack of certain stimuli that trigger the development of the parasites.

In an attempt to establish and compare protective immunity in both eel species, we used an approach taken earlier in animal models of filariasis. In these nematode infections a long-lasting and nearly complete immunity can be induced by vaccination with irradiation attenuated L₃, and detailed protocols have been worked out for the rodent filaria *Acanthocheilonema viteae* (see Schrempf-Eppstein *et al.* 1997 for references). These reports stimulated us to compare the effect of an irradiated vaccine in *A. japonica*

* Corresponding author: Leibniz-Institute of Freshwater Ecology and Inland Fisheries, Müggelseedamm 310, 12587 Berlin, Germany. Tel: +49 30 64181637. Fax: +49 30 64181663. E-mail: klaus.knopf@igb-berlin.de

and *A. anguilla*, with the goal to find out whether possible differences in immune response exist that could play a role in the different susceptibility towards *A. crassus* of the two eel species.

MATERIALS AND METHODS

Assessment of an optimal irradiation dose

Attenuation of infective larvae (L₃) by irradiation has been shown in a variety of filariae parasitizing in mammals (Lucius *et al.* 1991) but, to our knowledge, has not been reported for parasitic nematodes of fish. Therefore, an orienting experiment was performed to assess the effect of ¹³⁵Cs irradiation on L₃ of *A. crassus*. It was expected that a certain dose of irradiation would stop the development of the parasite without killing it.

L₃ isolated from their intermediate host were irradiated by exposing to a ¹³⁵Cs radiation source. Doses of 25 L₃, irradiated with 0, 175, 350 and 525 Gy were applied to groups of 6 *A. anguilla* by oral administration, using a 1-ml syringe fitted with a 1.5 mm diameter plastic tubing.

The eels were maintained at a water temperature of 23 °C, killed after 70 days and the swimbladder was examined for larvae and adults of *A. crassus*. Because L₃ and 4th-stage larvae (L₄) cannot be distinguished from each other perfectly by means of light microscopy (Blanc *et al.* 1992), larvae with a body length exceeding 1.5 mm were counted as L₄, according to Knopf *et al.* (1998).

Experimental design

Groups each of 16 *A. japonica* and *A. anguilla* were vaccinated with 1 dose of 40 irradiation (¹³⁵Cs, 500 Gy) attenuated L₃ of *A. crassus*, and after 5 weeks the eels were challenged with 40 L₃. The L₃ were counted in a round-bottomed 98-well plate and suspended in approximately 100 µl of RPMI-1640 medium, Hepes modification (Sigma-Aldrich, Taufkirchen, Germany). This suspension was introduced into the stomach of each eel as described above.

Challenge control groups each of 16 eels of both species were sham treated by peroral administration of medium and challenged with 40 normal L₃. Medium control groups each of 16 eels of both species were sham infected twice with medium to monitor potential changes of the antibody response due to factors not related to the *A. crassus* infection. The irradiation control groups, also consisting of 16 eels of each species, were treated with 40 irradiated L₃ without subsequent infection to test for a possible development of the L₃ after the irradiation.

At dissection (12 weeks p.i.) living and dead/encapsulated larvae in the swimbladder wall and adults of *A. crassus* in the swimbladder lumen were counted.

Male and female adult worms were individually weighed. Nematodes showing no reaction to mechanical stimulation were considered dead. The presence of *A. crassus* eggs/2nd-stage larvae (L₂) in the swimbladder lumen was considered as evidence of reproduction of the nematodes.

The experiment was split into 2 consecutive parts (A and B), each performed with 8 eels per treatment group. Eels treated in part B of the experiment were bled by caudal vein puncture at -5, 0, 4, 8 and 12 weeks post-infection (p.i.).

Source and maintenance of eels

Anguilla anguilla were obtained from a commercial eel farm known to be free of *A. crassus*. The absence of *A. crassus* was confirmed by necroscopy of 15 eels. *Anguilla japonica* were imported as glass-eels from Japan and raised in a recirculation system free of *A. crassus*. For the experiment eels were kept individually in aerated 40-l compartments of 200 l tanks, equipped with a polypropylene tube serving as a hiding-place. Water temperature was maintained at 23 °C. The eels were allowed to feed *ad libitum* on pellet food. Prior to the experiment fish were allowed to acclimatize for 2 weeks.

Parasites

L₃ of *A. crassus* were obtained according to the method described by Knopf *et al.* (1998). Briefly, 2nd-stage larvae (L₂) collected from the swimbladder lumen of naturally infected eels were fed to planktonic copepods serving as intermediate hosts. After 14 days at 20 °C, L₃ were isolated from the intermediate hosts by the tissue potter method described by Haenen *et al.* (1994) and stored in RPMI-1640 medium containing 100 U ml⁻¹ penicillin and 100 µg ml⁻¹ streptomycin at 4 °C until use.

Enzyme-linked immunosorbent assay (ELISA)

Preliminary tests with monoclonal antibodies (mAbs) raised against *A. anguilla* immunoglobulin (Ig) heavy and light chain (WEI 1 and WEI2, van der Heijden *et al.* 1995) revealed that *A. japonica* Ig are not recognized by WEI1, and WEI2 showed only a very weak reaction with *A. japonica* Ig. In contrast, polyclonal antibodies against *A. anguilla* Ig (Buchmann *et al.* 1992) also showed an appreciable reaction to *A. japonica* Ig. We used the most sensitive detection system for each species, namely the polyclonal antibodies to detect *A. japonica* Ig, and WEI1 for *A. anguilla* Ig. To allow a limited comparability of the results, the intensity of the antibody responses was expressed relative to the antibody content at the beginning of the experiment.

Crude antigen extracts from complete L₃ and from the body wall of adult *A. crassus* were prepared by sonication on ice in a 10-fold amount of sarcosyl-TE-buffer (10 mM Tris, 1 mM EDTA, 2% N-lauroyl-sarcosine-sodium salt, pH 8.0) and centrifuged for 20 min at 16 000 g. The supernatant was stored at -70 °C until use.

Polystyrene microtitre plates (Nunc, Kamstrup, Denmark) were coated with the crude antigen extracts in a concentration of 1.5 µg · ml⁻¹ in carbonate buffer (10 mM Na₂CO₃, 35 mM NaHCO₃, pH 9.6) overnight at 4 °C. Wells were washed 3 times with PBS containing 0.05% (v/v) Tween 20 (PBS-T), blocked with 1% (w/v) non-fat dry milk (Bio-Rad Laboratories, USA) in PBS for 3 h at 20 °C and washed 4 times with deionised water. After drying at 37 °C the plates were sealed with plastic tape and stored at -70 °C until use.

Eel sera were tested in triplicate at a dilution of 1:100 in PBS + 1% dry milk and incubated for 1 h at 37 °C. Antibodies of *A. japonica* were detected with polyclonal rabbit anti-eel Ig (Buchmann *et al.* 1992) in a concentration of 1:1000 in PBS + 1% dry milk followed by incubation with horseradish peroxidase conjugated sheep anti-rabbit IgG (AP311, The Binding Site, England) in a concentration of 1:2000 in PBS + 1% dry milk. Antibodies of *A. anguilla* were detected with a monoclonal mouse anti-eel Ig (WEI 1, van der Heijden *et al.* 1995) diluted 1:500 in PBS + 1% dry milk followed by incubation with sheep anti-mouse IgG conjugated with horseradish peroxidase (AP271, The Binding Site, England) diluted 1:1000 in PBS + 1% dry milk.

Incubation with the secondary and tertiary antibodies was for 45 min at 37 °C, and subsequently the wells were washed 3 times with PBS-T. The substrate reaction with TMB (3,3',5,5'-Tetra-Methyl-Benzidine, Sigma) was stopped after 15 min with 2N H₂SO₄. The absorbance was read at 492 nm with a plate reader (Genios, Tecan, Männedorf, Switzerland).

Statistical evaluation

Differences between groups were evaluated with the Mann-Whitney-U-Test. Statistical analysis of sequentially measured values within a group were analysed with the Friedman-Test and the Wilcoxon-Test. Fisher's exact test was used to check if the number of eels with eggs/L₂ of *A. crassus* in the swimbladder lumen differed significantly between immunized eels with a challenge infection and the challenge control group and between the host species. Spearman's rank correlation coefficient was used to detect a link between the number of retrieved adult worms and the intensity of the antibody response. Significance was accepted when $P < 0.05$. Statistical analyses were performed with SPSS 9.0 (SPSS Inc., Chicago, Illinois).




Fig. 1. Effect of different doses of radiation (¹³⁵Cs) on the larval development of *Anguillicola crassus* in *Anguilla anguilla* at a water temperature of 23 °C. Shown are the percentage of L₃, L₄, and adult worms retrieved at 70 days p.i. ($n = 6$).

The resistance of immunized animals was calculated as follows, with the number of adult worms:

$$\text{resistance (\%)} = \left(1 - \frac{\text{worm no. of immunized eels}}{\text{mean worm no. of challenge control group}} \right) \times 100$$

RESULTS

Assessment of optimal irradiation dose

Irradiation of L₃ of *A. crassus* resulted in a dose-dependent inhibition of the larval development (Fig. 1). Whilst 78.0% ± 20.7% of the worms retrieved at day 70 p.i. became adult in the non-irradiated control group, irradiation with 175 Gy and 350 Gy reduced the percentage of adults to 42.5% ± 27.1% and 2.8% ± 6.8%, respectively. Irradiation with 525 Gy still allowed the development from L₃ to L₄, but development to the adult stage was completely stopped.

The recovery rates (including L₃, L₄ and adults) were 23.3% ± 21.2%, 19.5% ± 13.8%, and 18.0% ± 10.4% for the worms irradiated with 175 Gy, 350 Gy, and 525 Gy, respectively, being not significantly different from the non-irradiated control with a recovery rate of 15.6% ± 16.8%. These data show that the irradiation attenuated the development without immediately killing the L₃. Based on these results an irradiation dose of 500 Gy was chosen for further experiments.

Worm recovery in immunized eels and control eels

In part A ($n = 8$) of the experiment, immunization of *A. japonica* with irradiated L₃ induced 96.8% ± 9.2% protection, based on the number of adult worms developing from the challenge infection ($P < 0.05$). In part B ($n = 8$) of the experiment a similar trend, but no statistically significant difference, was

Table 1. Effect of immunization of *Anguilla anguilla* and *Anguilla japonica* with 40 irradiated L₃ of *Anguillicola crassus*(The experiment was split into 2 independent parts A and B. Data presented are mean values \pm s.d.)

Trial A

	<i>Anguilla japonica</i>			<i>Anguilla anguilla</i>		
	Imm. & inf.	Imm.	Inf.	Imm. & inf.	Imm.	Inf.
Immunization	40 L3 irr.	40 L3 irr.		40 L3 irr.	40 L3 irr.	
Challenge infection	40 L3		40 L3	40 L3		40 L3
Number of adult worms (mean \pm s.d.)	0.1 \pm 0.4	—	3.9 \pm 4.4*	6.6 \pm 5.7	0.9 \pm 1.0	5.6 \pm 4.5
Ratio male: female worms	1: 0.0		1: 0.8	1: 0.8		1: 0.5
Recovery rate (% retrieved alive)	5.1 \pm 4.7	7.5 \pm 5.0	17.6 \pm 13.3*	20.9 \pm 14.5	6.9 \pm 5.5	18.1 \pm 13.7
Adult worms (% of retrieved worms)	0.4 \pm 1.3	—	27.2 \pm 27.2*	29.7 \pm 24.2	13.7 \pm 10.4	76.5 \pm 20.8*
Dead larvae (% of retrieved worms)	83.6 \pm 14.0	71.3 \pm 17.5	49.2 \pm 33.2	28.9 \pm 35.5	55.4 \pm 17.5	1.0 \pm 2.5*
Reproduction of <i>A. crassus</i> (% eels with worm eggs)	—	—	42.9	75.0	—	62.5
Protection (mean \pm s.d.)	96.8 \pm 9.2					
Surviving eels	8/8	8/8	7/8	8/8	8/8	8/8

Trial B

	<i>Anguilla japonica</i>			<i>Anguilla anguilla</i>		
	Imm. & inf.	Imm.	Inf.	Imm. & inf.	Imm.	Inf.
Immunization	40 L3 irr.	40 L3 irr.		40 L3 irr.	40 L3 irr.	
Challenge infection	40 L3		40 L3	40 L3		40 L3
Number of adult worms (mean \pm s.d.)	0.3 \pm 0.5	—	1.1 \pm 2.1	13.1 \pm 11.9	0.4 \pm 0.7	18.3 \pm 6.9
Ratio male: female worms	1: 1.0		1: 0.8	1: 0.8		1: 0.8
Recovery rate (% retrieved alive)	10.2 \pm 7.3	4.1 \pm 5.0	13.4 \pm 7.3	30.0 \pm 20.8	13.4 \pm 9.5	66.3 \pm 16.6*
Adult worms (% of retrieved worms)	3.5 \pm 6.6	—	12.7 \pm 25.9*	51.5 \pm 28.8	9.4 \pm 18.6	67.2 \pm 18.0
Dead larvae (% of retrieved worms)	52.7 \pm 30.9	67.0 \pm 29.6	48.4 \pm 29.1*	9.6 \pm 12.6	16.4 \pm 24.5	0.7 \pm 2.0*
Reproduction of <i>A. crassus</i> (% eels with worm eggs)	—	—	—	75.0	—	100.0
Protection (mean \pm s.d.)	77.8 \pm 41.1					
Surviving eels	8/8	8/8	8/8	8/8	8/8	8/8

* Significant difference (U-test, $P < 0.05$) between Imm. & inf. and Inf.

observed (Table 1). Combining the results from both parts of the experiment ($n = 16$) revealed a significant reduction of the number of adult worms in immunized *A. japonica* ($P < 0.05$), implying a resistance of $87.3\% \pm 30.4\%$. In contrast, immunization of *A. anguilla* with irradiated L₃ had no effect on the number of adult worms developing from the challenge infection (Table 1).

In *A. japonica* only immunized with irradiated L₃ no adult worms were recovered, whereas in *A. anguilla* 1.7% of the irradiated L₃ had developed to adult worms. These worms were very small (females weighing 3.9 ± 4.1 mg, 1 male weighing 0.2 mg) compared to adult worms which had developed in *A. anguilla* within the same time from normal L₃ (Fig. 2). The total burden of living worms (L₃, L₄, adults) in the immunization control group in *A. anguilla* was about one third to one fourth compared to the challenge control group (Table 1).

The wet weight of adult worms from immunized and challenge infected *A. anguilla* did not

significantly differ from the wet weight of challenge control worms (Fig. 2). In contrast, the few adult *A. crassus* found in the immunized and challenge infected *A. japonica* tended to be smaller than the challenge control worms, but due to the low number of adult worms found in the immunized *A. japonica* this difference could not be shown to be statistically significant (Fig. 2).

The sex ratio of the adult worms was similar in both host species and in immunized versus non-immunized eels (Table 1). The number of eels with eggs/L₂ of *A. crassus* in their swimbladder lumen was similar in *A. anguilla* immunized with a challenge infection and the challenge control group. In *A. japonica* 3 of 15 specimens of the challenge control group harboured eggs/L₂ of *A. crassus*, while none of the immunized eels contained eggs/L₂.

In the challenge control groups the percentage of adult *A. crassus* recovered was significantly lower in *A. japonica* compared to *A. anguilla* ($19.5\% \pm 26.6\%$ versus $71.5\% \pm 19.3\%$, respectively), and the




Fig. 2. Mean wet weight of (A) female and (B) male *Anguillicola crassus* from experimentally infected *Anguilla japonica* and *Anguilla anguilla* kept at a water temperature of 23 °C and examined 12 weeks p.i. Error bars indicate s.d.; numbers within or on the columns indicate sample sizes. Columns with superscripts in common are not different from one another ($P > 0.05$). Grey columns, normally infected eels (challenge control group); white columns, immunized and challenge-infected eels.

wet weight of adult worms was significantly lower in *A. japonica* than in *A. anguilla* (Fig. 2). The proportion of dead/encapsulated larvae from all worms retrieved was significantly higher in *A. japonica* ($48.8\% \pm 29.9\%$) than in *A. anguilla* ($0.8\% \pm 2.2\%$), and the percentage of eels containing eggs/ L_2 of *A. crassus* in the swimbladder lumen was significantly lower in *A. japonica* compared to *A. anguilla*.

Antibody response of immunized eels and control eels

To study the course of the antibody response, sera obtained from eels in part B of the experiment were tested by ELISA with total soluble antigens of L_3 and body wall soluble antigens of adult worms. The qualitative course of the antibody responses detected in both eel species was similar for both crude antigens. However, there were significant differences between host species (Fig. 3).

In immunized and challenge-infected *A. japonica* the first antibody responses were detected 1 month after the challenge infection, i.e. 2 months after the first antigen contact. The antibody response rose slightly until the end of the experiment. *A. japonica* of the challenge control group also reacted 2 months after the first antigen contact, i.e. 2 months after the challenge infection and had a slightly rising antibody response. In immunized *A. japonica* without challenge infection, antibody responses were also detectable 2 months after the immunization. Antibody responses against body wall antigens remained low, while antibodies against L_3 antigens rose slightly. Sera from the *A. japonica* control group that was neither immunized nor challenge infected showed no reaction with the *Anguillicola* antigen preparations.

In immunized and challenge-infected *A. anguilla* first antibody responses (2 of 8 eels) were also detected 2 months after the first antigen contact (i.e. 1 month after the challenge infection). Much in contrast to *A. japonica*, the antibody response then increased drastically and reached a high level at the end of the experiment. *A. anguilla* of the challenge control group showed a relatively weak, but significant reaction 2 months after the challenge infection, but the antibody levels did not rise. No antibody response was detected in immunized *A. anguilla* without challenge infection, and in the control group that was neither immunized nor challenge infected.

The level of antibody responses in immunized *A. japonica* was not correlated with protection against challenge infection. However, in immunized *A. anguilla* a positive correlation was found between the level of antibody responses against the larval antigen preparation and the number of adult worms (week 8: $r = 0.802$, $P = 0.017$; week 12: $r = 0.786$, $P = 0.021$).

DISCUSSION

Our study shows that eels can be successfully vaccinated against *A. crassus* by application of irradiated L_3 , provided that the animals belong to a host species that is able to mount a protective immune response. Whereas *A. japonica* could be protected by vaccination with attenuated L_3 , the newly acquired host species *A. anguilla* could not restrict the worm burden deriving from challenge infection. These data suggest that the original host can restrict the burden of its parasite by immune responses, whereas *A. anguilla* cannot. Therefore, it is likely that the recent spread of *A. crassus* in Europe was facilitated by an immunologically determined susceptibility of *A. anguilla*.

Our data show that gamma irradiation (^{135}Cs) is a useful method to obtain attenuated L_3 of *A. crassus*. Compared with L_3 of the filarial nematode *Acanthocheilonema viteae*, which is almost completely




Fig. 3. ELISA study of antibody responses of immunized and non-immunized *Anguilla japonica* (A, B) and *Anguilla anguilla* (C, D) against antigens of the adult worm body wall (A, C) and somatic L₃ antigens (B, D) of *Anguillicola crassus* during the course of a challenge infection. Shown is the serum antibody content relative to the start of the experiment. Arrowheads indicate the date of immunization (imm.) and infection (inf.), error bars indicate s.d., and asterisks significantly increased antibody contents. ■, immunized; ●, infected; ▲, immunized and infected; ▼, not immunized, not infected.

attenuated with a dose of 350 Gy (Lucius *et al.* 1991; Schrepf-Eppstein *et al.* 1997), attenuation of L₃ of *A. crassus* requires a higher dose of radiation. Irradiation with 525 Gy still allowed the larval development from L₃ to L₄, but interfered with the further development to adults. This prompted us to reduce the level of irradiation to 500 Gy in the vaccination experiment, as we anticipated that a slightly better larval growth could induce a better immunity against the challenge infection. However, the vaccination experiment showed that a few 500 Gy-irradiated L₃ of the irradiation control group reached maturity in the European eel, such that the original slightly higher irradiation dose of 525 Gy can be considered optimal for attenuation of *A. crassus*.

In *A. japonica*, the original host of *A. crassus*, immunization with irradiated L₃ of *A. crassus* resulted in a significantly reduced number of adult worms developing from a subsequent challenge infection compared to the challenge control group only infected with normal L₃, indicating that immunization

induced partial resistance. In the new host *A. anguilla*, immunization with irradiated L₃ had obviously no effect on the number of adult worms developing from the challenge infection, providing no evidence for an induced resistance. Although the basis of the immunity induced by irradiated nematode L₃ has not been completely elucidated, work in rodent models suggests that adaptive immunity induced by irradiated filarial L₃ requires IgE and eosinophils, and furthermore depends on activation of Toll-like receptor 4 (TLR4) (Abraham *et al.* 2004; Kerepesi *et al.* 2005). Therefore, the differences in reactivity between *A. japonica* and *A. anguilla* could be caused by a multitude of factors involved in such adaptive immune responses. Among others, variation in MHC genes or differences in cytokine regulation might determine the host qualities of eel species.

The lower susceptibility of *A. japonica* to *A. crassus* in comparison to *A. anguilla*, which has previously been demonstrated by experimental infection (Knopf and Mahnke, 2004), could be confirmed with the

present experiment. Following a single infection, the percentage of adult worms found in *A. japonica* was lower as compared to *A. anguilla*, and the few adult worms were much smaller in *A. japonica* compared to *A. anguilla*. As published data prove that *A. crassus* has the potential to reach a similar size in both host species (Kuwahara *et al.* 1974; Moravec and Taraschewski, 1988) our data indicate that the worms' development is retarded in the original host, *A. japonica*, as compared to *A. anguilla*. This might indicate poorer living conditions for *A. crassus* in *A. japonica* as compared to *A. anguilla*. Such differences in growth conditions for *A. crassus* could be due to adaptive immune responses developing during the infection, but also to stronger innate immune responses of *A. japonica* as, for example, attacking neutrophils.

Comparison of the present experiment with a similar study on infection of various rodent species with the filarial nematode *A. viteae* reveals, as an interesting parallel, that in both cases the highest degree of protection was observed in the original host (Schrempf-Eppstein *et al.* 1997). A second interesting parallel pertains to the role of antibody responses in protection. It has been assumed that L₃ of *A. crassus* can be killed by antibody-mediated mechanisms (Nielsen *et al.* 1999; Knopf *et al.* 2000), but hitherto there is no proof for a role of antibody responses in immune protection neither in *A. japonica* nor in *A. anguilla*. In the present study, the level of antibody responses against the larval antigen preparation in vaccinated *A. anguilla* was positively correlated with the number of adult worms developing from the challenge infection, suggesting that the antibody response measured is more a marker for susceptibility than for resistance. This is intriguing, as antibody-mediated cellular cytotoxicity is also regarded as an important mechanism of protection in rodent infections with filarial nematodes (Abraham *et al.* 2004). However, a recent vaccination study with recombinant tropomyosin of *A. viteae* revealed also an inverse correlation between protection and antibody responses and suggested T cell-mediated immune effector mechanisms (Hartmann *et al.* 2006). The same might hold true for the infection of *A. japonica* with *A. crassus*. It is, however, possible that the use of other antigen preparations, e.g. from L₄, or other experimental conditions would reveal protective antibody-mediated immune mechanisms.

It is suggestive that comparative experiments with *A. japonica* and *A. anguilla*, that differ significantly in their susceptibility to *A. crassus*, might be a key for further insights into immune effector mechanisms of fish against a nematode parasite. Moreover, comparison between immune mechanisms of hosts as different as mammals and fish might help to determine common denominators of protection against nematode parasites.

We are most grateful to Professor Kazuo Ogawa, University of Tokyo, for organizing the shipment of the Japanese glass eels. We thank Dr Kurt Buchmann, University of Copenhagen, and Dr Jan Rombout, Wageningen University, for providing the antibodies against eel immunoglobulin. We also thank Dr Heidi Hecker-Kia, Deutsches Rheuma-Forschungszentrum Berlin, who enabled us to use the radiation source, and Eric Eckmann for technical support.

REFERENCES

- Abraham, D., Leon, O., Schnyder-Candrian, S., Wang, C. C., Galioto, A. M., Kerepesi, L. A., Lee, J. J. and Lustigman, S.** (2004). Immunoglobulin E and eosinophil-dependent protective immunity to larval *Onchocerca volvulus* in mice immunized with irradiated larvae. *Infection and Immunity* **72**, 810–817.
- Blanc, G., Bonneau, S., Biagiatti, S. and Petter, A. J.** (1992). Description of the larval stages of *Anguillicola crassus* (Nematoda, Dracunculoidea) using light and scanning electron microscopy. *Aquatic Living Resources* **5**, 307–318.
- Buchmann, K., Østergaard, L. and Glamann, J.** (1992). Affinity purification of antigen-specific serum immunoglobulin from the European eel (*Anguilla anguilla*). *Scandinavian Journal of Immunology* **36**, 89–97.
- Haenen, O. L. M., van Wijngaarden, T. A. M. and Borgsteede, F. H. M.** (1994). An improved method for the production of infective third-stage juveniles of *Anguillicola crassus*. *Aquaculture* **123**, 163–165.
- Hartmann, S., Sereda, M. J., Sollwedel, A., Kalinna, B. and Lucius, R.** (2006). A nematode allergen elicits protection under specific conditions. *Vaccine* **24**, 3581–3590.
- Kerepesi, L. A., Leon, O., Lustigman, S. and Abraham, D.** (2005). Protective immunity to the larval stages of *Onchocerca volvulus* is dependent on Toll-like receptor 4. *Infection and Immunity* **73**, 8291–8297.
- Kirk, R. S.** (2003). The impact of *Anguillicola crassus* on European eels. *Fisheries Management and Ecology* **10**, 385–394.
- Knopf, K.** (2006). The swimbladder nematode *Anguillicola crassus* in the European eel *Anguilla anguilla* and the Japanese eel *Anguilla japonica*: differences in susceptibility and immunity between a recently colonized host and the original host. *Journal of Helminthology* **80**, 129–136.
- Knopf, K. and Mahnke, M.** (2004). Differences in susceptibility of the European eel (*Anguilla anguilla*) and the Japanese eel (*Anguilla japonica*) to the swimbladder nematode *Anguillicola crassus*. *Parasitology* **129**, 491–496.
- Knopf, K., Naser, K., van der Heijden, M. H. T. and Taraschewski, H.** (2000). Humoral immune response of European eel *Anguilla anguilla* experimentally infected with *Anguillicola crassus*. *Diseases of Aquatic Organisms* **42**, 61–69.
- Knopf, K., Würtz, J., Sures, B. and Taraschewski, H.** (1998). Impact of low water temperature on the development of *Anguillicola crassus* in the final host

- Anguilla anguilla*. *Diseases of Aquatic Organisms* **33**, 143–149.
- Kuwahara, A., Niimi, A. and Itagaki, H.** (1974). Studies on a nematode parasitic in the air bladder of the eel. I. Description of *Anguillicola crassa* n. sp. (Philometridae, Anguillicolidae). *Japanese Journal of Parasitology* **23**, 275–279.
- Lucius, R., Textor, G., Kern, A. and Kirsten, C.** (1991). *Acanthocheilonema viteae*: vaccination of jirds with irradiation-attenuated stage-3 larvae and with exported larval antigens. *Experimental Parasitology* **73**, 184–196.
- Moravec, F. and Taraschewski, H.** (1988). Revision of the genus *Anguillicola* Yamaguti, 1935 (Nematoda: Anguillicolidae) of the swimbladder of eels, including descriptions of two new species, *A. novaezelandiae* sp. n. and *A. papernai* sp. n. *Folia Parasitologica* **35**, 125–146.
- Nielsen, M. E.** (1999). An enhanced humoral immune response against the swimbladder nematode, *Anguillicola crassus*, in the Japanese eel, *Anguilla japonica*, compared with the European eel, *A. anguilla*. *Journal of Helminthology* **73**, 227–232.
- Schrempf-Eppstein, B., Kern, A., Textor, G. and Lucius, R.** (1997). *Acanthocheilonema viteae*: vaccination with irradiated L3 induces resistance in three species of rodents (*Meriones unguiculatus*, *Mastomys coucha*, *Mesocricetus auratus*). *Tropical Medicine and International Health* **2**, 104–110.
- Sures, B., Knopf, K., Würtz, J. and Hirt, J.** (1999). Richness and diversity of parasite communities in European eels *Anguilla anguilla* of the River Rhine, Germany, with special reference to helminth parasites. *Parasitology* **119**, 323–330.
- Sures, B. and Streit, B.** (2001). Eel parasite diversity and intermediate host abundance in the River Rhine, Germany. *Parasitology* **123**, 185–191.
- van der Heijden, M. H. T., Rooijackers, J. B. M. A., Booms, G. H. R., Rombout, J. H. W. M. and Boon, J. H.** (1995). Production, characterization and applicability of monoclonal antibodies to European eel (*Anguilla anguilla* L., 1758) immunoglobulin. *Veterinary Immunology and Immunopathology* **45**, 151–164.