

Malaria pigment accelerates MTT – formazan exocytosis in human endothelial cells

Sarah D'Alessandro¹, Yolanda Corbett², Silvia Parapini¹, Federica Perego³, Loredana Cavicchini¹, Lucia Signorini¹, Serena Delbue¹, Carla Perego³, Pasquale Ferrante¹, Donatella Taramelli³ and Nicoletta Basilico¹

Research Article

Cite this article: D'Alessandro S *et al.* (2019). Malaria pigment accelerates MTT – formazan exocytosis in human endothelial cells. *Parasitology* **146**, 399–406. <https://doi.org/10.1017/S0031182018001579>

Received: 18 June 2018
Revised: 2 August 2018
Accepted: 22 August 2018
First published online: 1 October 2018

Key words:

Endothelial cells; haemozoin; malaria; MTT reduction

Author for correspondence:

Nicoletta Basilico, E-mail: nicoletta.basilico@unimi.it

¹Dipartimento di Scienze Biomediche, Chirurgiche e Odontoiatriche, Università degli Studi di Milano, Milano, Italy; ²Dipartimento di Bioscienze, Università degli Studi di Milano, Milano, Italy and ³Dipartimento di Scienze Farmacologiche e Biomolecolari, Università degli Studi di Milano, Milano, Italy

Abstract

Haemozoin is a by-product of haemoglobin digestion by intraerythrocytic malaria parasites, which induces immunologic responses on different tissues, including endothelial cells. In the present paper, the incubation of human microvascular endothelial cells with haemozoin significantly inhibited MTT reduction, a measure of cytotoxicity, without increasing the release of cytoplasmic lactate dehydrogenase. Moreover, haemozoin did not induce apoptosis or cell cycle arrest nor decreased the number of live cells, suggesting that cells viability itself was not affected and that the inhibition of MTT reduction was only apparent and probably due to accelerated MTT-formazan exocytosis. After 30 min of MTT addition, a significant increase in the % of cells exocytosing MTT formazan crystals was observed in haemozoin-treated cells compared with control cells. Such an effect was partially reversed by the addition of genistein, an inhibitor of MTT-formazan exocytosis. The rapid release of CXCL-8, a preformed chemokine contained in Weibel–Palade bodies, confirmed that haemozoin induces a perturbation of the intracellular endothelial trafficking, including the exocytosis of MTT-formazan containing vesicles. The haem moiety of haemozoin is responsible for the observed effect. Moreover, this work underlines that MTT assay should not be used to measure cytotoxicity induced by haemozoin and other methods should be preferred.

Introduction

Haemozoin (HZ) or malaria pigment is a crystal made of ferriprotoporphylin IX dimers derived from the catabolism of haemoglobin by intraerythrocytic malaria parasites (Pagola *et al.*, 2000; Egan, 2008). HZ is released in the circulation during the intraerythrocytic cycle of parasites and can modulate the functions of different host cell types (Tyberghin *et al.*, 2014; Deroost *et al.*, 2016). On endothelial cells, HZ modulates the expression of adhesion molecules and the production of inflammatory mediators such as cytokines, chemokines and metalloproteinases (Taramelli *et al.*, 1998; Basilico *et al.*, 2010, 2017). In complicated *Plasmodium falciparum* malaria, such as cerebral and placental malaria, activation of endothelial cells contributes to the cytoadherence of infected red blood cells to the microvascular endothelium, leading to microcirculatory obstruction and tissue hypoxia (Beeson *et al.*, 2001; Brown *et al.*, 2001; Andrews and Lanzer, 2002; Pongponratn *et al.*, 2003; Khaw *et al.*, 2013; Guiguemde *et al.*, 2014).

Free haem is able to induce brain endothelial cells apoptosis contributing to the destruction of endothelial integrity in cerebral malaria (Liu *et al.*, 2016). No data on the cytotoxic effect of haemozoin on endothelial cells are available.

The MTT [3-(4,5-Dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide] assay is a commonly used test for *in vitro* cytotoxicity evaluation (Mosmann, 1983; Berridge *et al.*, 2005). The MTT tetrazolium salt is taken up by viable cells by endocytosis and it is reduced mostly in the mitochondria forming the blue formazan that reflects viable cell number. MTT is also internalized and reduced to formazan inside endosomes (Liu *et al.*, 1997). Formazan is then transported to the cell surface forming needle-like MTT-formazan crystals. It has been reported that β -amyloid peptides, chloroquine, cholesterol and silica nanoparticles increase the exocytosis of formazan crystals (Abe and Saito, 1998; Claus *et al.*, 1998; Liu and Schubert, 1998; Isobe *et al.*, 1999; Fisichella *et al.*, 2009).

The aim of the present work was to study the effect of malaria pigment, HZ, on the process of exocytosis in human endothelial cells. This analysis was stimulated by a preliminary observation that the reduction of MTT by endothelial cells treated with HZ did not correspond to the actual viability of the cells and that needle-like structures were seen outside endothelial cells monolayers early after the addition of MTT.

Materials and methods

Materials

All materials were from Sigma-Aldrich, unless otherwise stated. The MCDB-131 medium was from Life Technologies; RPMI 1640 medium was from EuroClone; fetal bovine serum was from HyClone; epidermal growth factor was from Cell Signalling Technology.

Cells and *Plasmodium falciparum* cultures

A long-term cell line of dermal microvascular endothelial cells (HMEC-1) immortalized by SV 40 large T antigen (Ades *et al.*, 1992) was kindly provided by the Center for Disease Control (Atlanta, GA). Cells were maintained in MCDB-131 medium supplemented with 10% fetal bovine serum, 10 ng mL⁻¹ of epidermal growth factor, 1 µg mL⁻¹ of hydrocortisone, 2 mM L-glutamine, 100 units mL⁻¹ of penicillin, 100 µg mL⁻¹ of streptomycin (EuroClone) and 20 mM Hepes buffer (EuroClone), pH 7.4. *Plasmodium falciparum* parasites (D10 and W2 strain; mycoplasma free) were kept in culture as described (D'Alessandro *et al.*, 2015) at 5% hematocrit (human type A+ RBCs) at 37 °C in RPMI 1640 medium supplemented with 10% heat-inactivated A+ human plasma, 20 mM Hepes buffer, pH 7.4, in a standard gas mixture consisting of 1% O₂, 5% CO₂, 94% N₂.

Preparation of HZ

To isolate HZ, *Plasmodium*-infected erythrocytes (4–8% parasitemia) were washed twice with serum-free culture medium, resuspended to 25% hematocrit and fractionated on a discontinuous Percoll/4% sorbitol (wt vol⁻¹) gradient (0, 40, 80%) (Omodeo-Sale *et al.*, 2003). After centrifugation at 1075g, HZ was collected at the top of the 0–40% gradient interphase, washed three times with PBS and stored at –20 °C. The haem content of a weighed amount of HZ dissolved in 0.1 M NaOH was determined by reading the absorbance at 405 nm and by using a standard curve of hemin dissolved in 0.1 M NaOH.

HMEC-1 treatment and CXCL8 determination

HMEC-1 cells were seeded in complete medium at 10⁵ cells well⁻¹ in 24-well flat bottom tissue culture plates or at 10⁴ cells well⁻¹ in 96-well plates. After overnight incubation to allow the cells to adhere, monolayers were exposed to HZ (5–20 µg mL⁻¹) or Lipopolysaccharide (LPS, 100 ng mL⁻¹) in a humidified CO₂/air-incubator at 37 °C for 2 or 24 h. In some experiments, after 24 h of HZ treatment, cells were incubated with genistein, 50 µM for 1 h and the MTT assay was performed afterward. In other experiments, cells were incubated 24 h with HZ 10 µg mL⁻¹ in the presence of the mitochondrial protectors polydatin 25 µM or cyclosporine A 5 µM. All the experiments were performed in serum-free medium. At the end of treatment, supernatants were collected and CXCL8 levels were measured by DuoSet ELISA Kit (R&D System), following the manufacturer's instructions.

MTT assay

Cell viability was evaluated using the 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide (MTT) assay. After HMEC-1 treatment, 20 µL of a 5 mg mL⁻¹ solution of MTT in PBS were added to the cells for 3 additional hours at 37 °C in the dark. The supernatants were then discarded and the dark blue formazan crystals dissolved using 100 µL of lysis buffer containing 20% (wt vol⁻¹) sodium dodecylsulfate, 40% N,N-dimethylformamide (pH 4.7 in 80% acetic acid). The plates were then read on a Synergy 4 (Biotek®) microplate reader at a test wavelength of 550 nm and at a reference wavelength of 650 nm. In some experiments, MTT reaction was stopped after 15, 30, 60, 120 min of incubation. To check the presence of needle-like crystals on cells surface, pictures were taken following incubation with MTT using a Nikon Ti-S microscope and a Nikon DS-F11C COOLED camera. The percentage of cells exocytosing MTT formazan was established by counting 200–250 cells by light microscopy at 400× magnification, in

multiple fields after the different time of incubation with MTT, as described (Liu *et al.*, 1998).

LDH assay

The potential cytotoxic effect of HZ was measured as the release of lactate dehydrogenase (LDH) from HMEC-1 into the extracellular medium using the LDH Cytotoxicity Assay kit following the manufacturer's instructions. LDH was measured both in the extracellular medium and in the cells pellet. Briefly, cells were incubated for 24 h with HZ (10 µg mL⁻¹) in a humidified CO₂/air-incubator at 37 °C. Then, cell supernatants were collected and centrifuged at 13 000g for 2 min. Cells were washed with PBS and 0.5 mL of Triton × 100 (2% final concentration) were added to lyse the cells. One hundred microlitres of this solution or 100 µL of supernatant were mixed with 100 µL of LDH reaction mix, containing the LDH substrate, and incubated for 10 min at room temperature in the dark. Absorbance was then read at 490 nm with a reference wavelength of 650 nm using Synergy 4 (Biotek®) microplate reader. Percent cytotoxicity was calculated following the manufacturer's instructions.

Cell count by trypan blue

After HZ treatment for 24–48–72 h, cells were detached by trypsin treatment and counted in a Neubauer chamber by trypan blue exclusion. Data are expressed as a number of live cells per well. The number of dead cells never exceeded 5% of the total in all the conditions.

Apoptosis determination

HMEC-1 cells were treated for 24 h with HZ (10 µg mL⁻¹) or exposed to UV-B irradiation for 30 min and then incubated for 24 h (positive control of apoptosis). The percentage of apoptotic cells was evaluated using FITC-conjugated annexin V and propidium iodide (PI, BD Biosciences, Franklin Lakes, USA) staining, according to the manufacturer's protocol. FITC Annexin V staining precedes the loss of membrane integrity of the latest stages of cell death, thus early apoptotic cells are PI negative and FITC Annexin V positive, whereas cells in late apoptosis or already dead (thus permeable to PI because of the loss of membrane integrity) are both FITC Annexin V and PI positive. Viable cells are FITC Annexin V and PI negative. The analyses were performed using a FACSCalibur and CellQuest software (BD Biosciences, Franklin Lakes, USA).

Cell cycle analysis

HMEC-1 cells were treated with HZ (10 µg mL⁻¹) for 24 h. Cells were harvested, washed with cold PBS and fixed with 70% ethanol at 4 °C for 30 min. The fixed cells were centrifuged and washed twice with cold PBS containing 5% fetal bovine serum. Cells were then resuspended in 0.5 mL of PBS containing 1 mg mL⁻¹ RNase A and 1% NP-40 at 37 °C for 30 min and stained with 1 mg mL⁻¹ propidium iodide (BD Biosciences) at 4 °C overnight. The cellular DNA content was measured using FACSCalibur and CellQuest software (BD Biosciences).

Acridine orange exocytosis assay

HMEC-1 cells were seeded at 10⁴ cells well⁻¹ in 96-well flat bottom tissue culture plates. After an overnight incubation at 37 °C for adhesion, cells were treated for 24 h with HZ (10 µg mL⁻¹). Cells were then treated with Acridine Orange (AO) 40 µM in PBS for 15 min at 37 °C to allow AO to enter the cells.

Fluorescence was evaluated in the supernatants and on cells monolayer by using a Synergy 4 (Biotek®) microplate reader with an excitation wavelength of 535 nm and an emission wavelength of 590 nm. After incubation of cells with AO, the supernatants were discarded. The cell monolayer was read at time 0 (T0), then 100 μL PBS were added to the cells and the plate incubated for 30 min at 37 °C. The supernatants were transferred in a new plate and read. The cell monolayer was read again (T30). The release of AO was expressed as fluorescence units (FU) in the supernatants and as the difference between the fluorescence of cells monolayer before and after PBS incubation (T0–T30).

Statistical analysis

All data were obtained from at least three independent experiments. Results are shown as means \pm standard deviation. Differences between groups were analysed by two-tailed Student's *t*-test or by one-way or two-way ANOVA analysis and posthoc multiple comparisons tests (Sidak, Tukey), using the software GraphPad Prism6.

Results

Effect of HZ on MTT reduction by HMEC-1

HMEC-1 were treated with different doses of HZ for 24 h and viability was measured by LDH and MTT assays. A direct toxicity of HZ on HMEC-1 was not observed since the release of LDH was not significantly different in control HMEC-1 or in HZ-treated cells (Fig. 1A). However, a decrease in MTT reduction, a measure of cell respiratory capacity and viability, was observed in HZ treated cells, at the end of 24 h experiments. (Fig. 1B). A high dose (50 $\mu\text{g mL}^{-1}$) of the antimalarial drug, dihydroartemisinin (DHA) was used as a control for cytotoxicity.

Compared with control, MTT reduction decreased to 80.3, 59.6 and 47.8% by 2.5, 5 or 10 $\mu\text{g mL}^{-1}$ of HZ, respectively. To solve the discrepancy between the results obtained by the MTT and the LDH assays, other methods for measuring cell viability were used. The actual number of cells after 24–48–72 h of HZ treatment, was counted by the trypan blue exclusion method. At the end of the 72 h incubation, the cell number doubled in both control and HZ treated cells, with less than 5% mortality. On the contrary, DHA used as a control, induced 50% decrease in viability and cell number. (Fig. 2A).

Both early apoptosis (FITC Annexin V positive and PI negative cells) and late apoptosis (FITC Annexin V and PI positive cells) were investigated. UV irradiation was used as positive control. As shown in Fig. 2B, differently from UV irradiation, HZ did not induce signs of early or late apoptosis in endothelial cells.

Since resting cells, in contrast to proliferating cells, can be metabolically quiescent and reduce low amounts of MTT, the cell cycle was analysed to establish if the inhibition of MTT reduction was dependent on a shift from the proliferative to the resting status. As shown in Fig. 2C, the percentage of cells in the different phases of the cell cycle was the same in control and in HZ-treated cells indicating that HZ did not induce an alteration in the cell cycle.

To verify if the inhibition of MTT reduction by HZ on endothelial cells was mediated by suppression of mitochondrial activity, the mitochondrial protectors polydatin and cyclosporine A were also used. None of these compounds recovered the MTT metabolism (Fig. 2D).

HZ enhanced the formation of needle-like MTT formazan crystals at cell surface

Needle-like formazan crystals were observed by microscopy after 30 min of MTT addition in HZ treated cells and increased over




Fig. 1. HZ treatment does not induce the release of LDH, but alters MTT metabolism. HMEC-1 were treated with different doses of HZ (2.5, 5, 10 $\mu\text{g mL}^{-1}$), for 24 h. At the end of incubation, the percentage of cytotoxicity was measured by the release of LDH (A), whereas the percentage of cell viability was measured by MTT assay (B). DHA at 50 $\mu\text{g mL}^{-1}$ was used as positive control. The results are the mean and standard deviation of three independent experiments. Ordinary one-way ANOVA, Tukey's multiple comparisons test: * $P < 0.05$ *** $P < 0.01$ **** $P < 0.001$ vs. control.

time (Fig. 3, arrows). In control cells, the majority of formazan crystals remained localized in intracellular granules and only after 180 min, some crystals appeared on the surface of control cells. At the same time points, the MTT decreased, as well (Fig. 3, histogram). The percentage of cells exocytosing MTT formazan is shown in Table 1. A significant increase above control was seen already after 30 min (69.2%) and reached a plateau after 120 min (82.7%). In the control, the percentage of cells exocytosing MTT reached 3.2% and 5.1% after 120 or 180 min, respectively. These results suggested an accelerated exocytosis of formazan crystals induced by HZ. To verify that HZ alone was unable to directly induce needle-like formazan crystals, HZ was incubated with MTT in a cell-free system. HZ did not induce crystals formation, indicating that the phenomenon depends on metabolically active cells.

MTT-formazan exocytosis in genistein-treated cells

Genistein, a compound known to inhibit MTT-formazan exocytosis, was used with the aim to counterbalance HZ effects (Liu *et al.*, 1997; Fisichella *et al.*, 2009). Genistein added to the cells after incubation with different concentrations of HZ, was able to inhibit the enhanced MTT exocytosis induced by HZ and a recovery of MTT reduction was observed (Fig. 4A). Similarly, the percentage of cells exocytosing MTT formazan decreased from 87.6% \pm 10.6 of HZ treated HMEC-1 to 28.4% \pm 7.9 in the presence of Genistein, as shown in the microscopic pictures (Fig. 4, panel B). Chloroquine at 50 $\mu\text{g mL}^{-1}$ was used as positive control since it has been reported that it enhances MTT-formazan




Fig. 2. HZ treatment does not induce HMEC-1 cell death and does not alter mitochondrial activity. HMEC-1 cells were treated with HZ and cells number (A), apoptosis (B) or cell cycle (C) were measured. (A) Viable cells were counted by trypan blue exclusion method. DHA at $50 \mu\text{g mL}^{-1}$ was used as positive control. (B) Apoptosis was determined by FACS analysis. Viable cells are both Annexin V and PI negative; cells in early apoptosis are Annexin V positive and PI negative; cells in late apoptosis or already dead are both Annexin V and PI positive. (C) cell cycle was measured by FACS analysis and PI staining. (D) HMEC-1 cells were treated with HZ in the presence of polydatin or cyclosporin A and MTT assay was performed. The results are the mean and standard deviation of three independent experiments. Two-way ANOVA, Tukey's multiple comparisons test * $P < 0.05$ ** $P < 0.001$, vs. control.

exocytosis in rat astrocytes cells (Isobe *et al.*, 1999). A significant decrease in MTT reduction was also observed in CQ-treated cells compared with controls and genistein partially restored the decrease of MTT induced by CQ. (Fig. 4A).

Accelerated release of preformed CXCL8 and AO exocytosis by HMEC treated with HZ

HMEC-1 were treated with HZ or LPS for 2 and 24 h and CXCL8 levels were measured in cells supernatants. As shown in Fig. 5A, 2 h of HZ, but not of LPS treatment induced significant release of CXCL8 indicating that HMEC-1 respond to HZ inducing exocytosis with a rapid release of preformed CXCL8. After 24 h of treatment, both HZ and LPS induced much stronger activation of HMEC-1 as demonstrated by the significant increase in the release of CXCL8 suggesting a transcriptional effect and the novo chemokine synthesis.

Since CXCL8 can be exocytosed by Weibel–Palade bodies (WPBs), which are lysosome-related secretory organelles of endothelial cells, AO, a fluorescent dye that once protonated, is nonspecifically trapped into acidic vesicles (Traganos and Darzynkiewicz, 1994) was used to evaluate acidic vesicle exocytosis. A slight increase, although not significant, in the release of AO was observed both in the supernatants of HZ treated cells compared with control and also by evaluating the loss of cell-associated AO at different times ($T_0 - T_{30}$) (Fig. 5B).

Beta-Hematin, but not other HZ components increased MTT exocytosis

In order to evaluate which HZ component contributed to the decrease in MTT reduction and accelerated crystal exocytosis, synthetic HZ (beta-haematin, BH, constituted by the sole haem backbone), 4-HNE or 15-HETE (two liperoxidation derivatives

generated by HZ from arachidonic acid), host fibrinogen (which is always associated with HZ) were also used to treat HMEC-1. The doses of each HZ component used were comparable with those able to induce biological activities in other *in vitro* models, as reported (Giribaldi *et al.*, 2004; Barrera *et al.*, 2011; Polimeni *et al.*, 2013; Basilio *et al.*, 2017). The results shown in Fig. 6, confirmed a significant decrease in MTT metabolism induced by BH, whereas 4HNE, 15-HETE and fibrinogen did not significantly interfere.

Discussion

Endothelial activation is a common feature observed in severe malaria. HZ is present at the site of parasite sequestration where can contribute to the endothelial activation. *In vitro* experiments have shown that HZ is able to induce the production of metalloproteinases and chemokines by endothelial cells (Prato *et al.*, 2011; Basilio *et al.*, 2017) and, when phagocytized by monocytes/macrophages, can induce both inhibitory and stimulatory effects (Basilico *et al.*, 2003; Khadjavi *et al.*, 2010; Prato and Giribaldi, 2011).

Here, we show that HZ was not cytotoxic on human endothelial cells, but was able to affect MTT metabolism accelerating formazan exocytosis.

Firstly, our results indicate that the MTT assay cannot be used for measuring cell viability in the presence of HZ. Alternatively, an accurate selection of the incubation time in the presence of MTT has to be done, since up to 30 min the difference between control and treated cells was not significant. After 1 or 3 h of incubation with MTT, despite an apparent reduction in MTT metabolism, cells were alive, as demonstrated by the release of LDH in the supernatants, a measure of cell cytotoxicity, or by cell count with the vital dye trypan blue. In parallel, a significant increase in the percent of cells exocytosing MTT formazan




Fig. 3. Accelerated formation of MTT-formazan needle-like crystals in HMEC-1 cells treated with HZ ($10 \mu\text{g mL}^{-1}$) for 24 h. After HZ treatment, MTT was added to the cells and light microscopy pictures of the cells (magnification $400\times$) were taken after 30 min, 1 h and 3 h of incubation at 37°C . Arrows show needle-like crystals. The histogram shows the decrease of MTT reduction over time (the data are expressed as % of untreated control).

crystals was observed in HZ treated cells, suggesting that the decrease in MTT metabolism was not due to cell death but most likely to modification of the exocytic pathways. The ability to accelerate formazan-MTT exocytosis is not specific for HZ, but it was described for different agents, such as β -amyloid peptides, chloroquine, cholesterol and silica nanoparticles (Abe and

Saito, 1998; Claus *et al.*, 1998; Liu and Schubert, 1998; Isobe *et al.*, 1999; Fisichella *et al.*, 2009). Moreover, the kinetic of formazan exocytosis and needle-like crystals formation varies depending on the cell type (Molinari *et al.*, 2005). These observations suggest to carefully define, for each cell type, the time of incubation with MTT assay. These results also suggest that when new compounds are tested for cytotoxicity using the MTT assay, it is important to exclude that the reduction of MTT is due to an accelerated formazan exocytosis to avoid false positive results.

It has been described that HZ can induce apoptosis in type II pneumocytes and erythroid precursors and can interfere with cell cycle progression of erythroid cells (Lamikanra *et al.*, 2009; Skorokhod *et al.*, 2010; Maknitikul *et al.*, 2018). However, here we show that HZ did not induce apoptosis or modification of the cell cycle in endothelial cells, indicating that HZ can induce different effects depending on the cell type. Moreover, in our experiments HMEC-1 cells were stimulated with a concentration of HZ relevant to physiological conditions, using a single treatment for 24 h. We cannot exclude that higher doses, longer time of incubation or repeated treatments with HZ could be toxic for endothelial cells, as well.

In the MTT assay, the reduction of MTT, a yellow tetrazolium dye, to purple formazan crystals, has been mainly attributed to the mitochondrial dehydrogenases (Mosmann, 1983). Thus, it is commonly accepted that the MTT assay is a suitable indicator of mitochondrial activity (Berridge and Tan, 1993; Berridge *et al.*, 1996). However, the hypothesis that HZ could interfere with MTT metabolism in HMEC-1 by suppression of mitochondrial activity was excluded, since mitochondrial protectors, such as polydatin and cyclosporine A, were unable to recover MTT metabolism. Moreover, the presence of needle-like crystals on the surface of the cells after only 30 min of incubation with MTT supported the fact that an acceleration of MTT-formazan exocytosis occurred in HZ treated cells.

Exocytosis is a process in which intracellular vesicles fuse to the plasma membrane and release different molecules. WPBs are endothelial specific secretory organelles containing bioactive molecules involved in inflammation and hemostasis (Metcalf *et al.*, 2008). Their membranes express the late-endosome/lysosome marker CD63 (Vischer and Wagner, 1993). Activation of endothelial cells by different stimuli results in WPBs exocytosis and in the release of peptides such as von Willebrand Factor, Angiopoietin 2 and Interleukin 8 (CXCL8) (Utgaard *et al.*, 1998). It has been reported that in *P. falciparum* malaria, plasma levels of von Willebrand factor, Angiopoietin 2 and CXCL8 are higher in patients with severe vs non-severe disease (Hollestelle *et al.*, 2006; Yeo *et al.*, 2008; Ayimba *et al.*, 2011). In pediatric cerebral malaria, high levels of CXCL8 have also been observed in the cerebrospinal fluid (Armah *et al.*, 2007). Here, we showed that HZ rapidly induced the release of CXCL8 from HMEC-1 suggesting exocytosis of the chemokine rather than *de novo* synthesis. On the contrary, LPS, a potent microbial stimulus did not induce the release of CXCL8 after 2 h, but only after 24 h of incubation. This is consistent with the reported observation that LPS

Table 1. Effect of haemozoin on MTT formazan exocytosis by HMEC-1 cells

Treatment	% of cells exocytosing MTT formazan ^a			
	30'	60'	120'	180'
Control	2.4 ± 1.9	2.4 ± 2.1	3.2 ± 1.8	5.1 ± 2.7
Haemozoin $10 \mu\text{g mL}^{-1}$	69.2* ± 8.6	72.8* ± 8.0	82.7* ± 10.2	87.6* ± 10.6

The percentage of cells exocytosing MTT formazan was determined at different times by counting 200 cells in multiple fields under a light microscope. Data are mean ± s.d. of at least three different experiments.

*Significantly different from control ($P < 0.05$) by two-tailed Student's *t*-test.

^aHMEC-1 cells were treated with $10 \mu\text{g mL}^{-1}$ of haemozoin for 24 h before the addition of MTT.

Fig. 4. Inhibition of MTT-formazan exocytosis by genistein. (A) After 24 h treatment with different doses of HZ (2.5, 5, 10 $\mu\text{g mL}^{-1}$) or CQ (50 $\mu\text{g mL}^{-1}$), HMEC-1 cells were incubated with genistein (50 μM) for 1 h and MTT assay was performed. Data are expressed as percent of control untreated cells. (B) Pictures were taken after 180 min of incubation with MTT of HMEC-1 cells treated with 10 $\mu\text{g mL}^{-1}$ of HZ alone or with genistein. The arrows show intracellular formazan crystals. The results are the mean and standard deviation of at least three independent experiments. Two-way ANOVA, Tukey's multiple comparisons test $**P < 0.001$ vs. control.




Fig. 5. CXCL8 and acridine orange release by HMEC-1 treated with HZ. (A) HMEC-1 were treated for 2 and 24 h with HZ (10 $\mu\text{g mL}^{-1}$) or LPS (100 ng mL^{-1}). At the end of the treatment cell supernatants were collected and assayed for the presence of CXCL8. (B) HMEC-1 cells were incubated with HZ 10 $\mu\text{g mL}^{-1}$ for 24 h and subsequently loaded with AO 40 μM for 15 min (T0). AO release was expressed as fluorescence units (FU $\text{ex } \lambda = 535 \text{ nm}$; $\text{em } \lambda = 590 \text{ nm}$) and evaluated both in the supernatants and in the cells monolayer at T0 (immediately after AO load) and at T30 (after 30 min of incubation to allow AO exocytosis), and calculating the difference in FU between T0 and T30. The results are expressed as means \pm s.d. of three independent experiments. Two-way ANOVA, Tukey's multiple comparisons test $*P < 0.05$, $***P < 0.0001$ vs. control.





Fig. 6. BH is the HZ component involved in the increase of MTT - formazan exocytosis. HMEC-1 cells were treated for 24 h with different HZ components: BH at 10 $\mu\text{g mL}^{-1}$, 4-hydroxynonenal (HNE) at 100 nM , 15-HETEs at 10 μM , fibrinogen (FG) at 200 $\mu\text{g mL}^{-1}$. The MTT assay was performed at the end of treatment and the data are expressed as percent of control. The results are the mean and standard deviation of three independent experiments. Ordinary one-way ANOVA, Tukey's multiple comparisons test: $**P < 0.01$ vs. control.

is unable to induce Weibel-Palade exocytosis by human aortic endothelial cells (Into *et al.*, 2007).

WPBs can be classified as lysosome-related organelles, thus AO exocytosis was evaluated in HZ treated cells. In our experiments, HZ induced only a slight, not significant enhancement of AO exocytosis indicating that even if the WPBs could be involved in exocytosis induced by HZ, trafficking of other acidic compartments, such as endothelial lysosomes, seems not to be altered. Moreover, CXCL8 is stored in WPBs, while the precise localization of the reduced formazan product inside the cells is

still unclear. It has been reported that MTT formazan granules co-localize with lysosomes/endosomes (Liu *et al.*, 1997; Liu and Schubert, 1997), mitochondria or, more recently, in lipid droplets (Diaz *et al.*, 2007; Stockert *et al.*, 2012). The result on AO also suggests that: (i) formazan crystals do not completely co-localize with lysosomes; (ii) HZ does not, or only partially, perturb intracellular vesicles trafficking of acidic vesicles; (iii) further studies are needed to better understand which compartments are involved in MTT-formazan or CXCL8 exocytosis by HZ.

HZ activates endothelial cells, but little is known on the mechanisms involved in this process. The accelerated MTT formazan exocytosis suggests that HZ, perturbing the intracellular trafficking, could stimulate also the exocytosis of WPBs, contributing to the release of different vasoactive molecules, as demonstrated with CXCL8. Genistein, used as an inhibitor of exocytosis (Liu and Schubert, 1997; Fisichella *et al.*, 2009), was able to decrease the MTT-formazan exocytosis induced by HZ. Interestingly, it has been reported that genistein is also able to block the VEGF-induced von Willebrand factor exocytosis by human aortic endothelial cells (Matsushita *et al.*, 2005).

HZ is a ferriprotoporphyrin-IX crystal bound to lipids, DNA, and plasma proteins such as fibrinogen and it can generate lipo-peroxidation products (15-HETE and 4-HNE) (Goldie *et al.*, 1990; Schwarzer *et al.*, 2003; Barrera *et al.*, 2011). All the different components of HZ are responsible for endothelial cells and monocytes/macrophages activation. Some effects of HZ are indeed mediated by the ferriprotoporphyrin-IX moiety, others by 15-HETE and 4-HNE or by parasite DNA or host fibrinogen associated with HZ (Omodeo-Salè *et al.*, 1998; Taramelli *et al.*, 2000; Schwarzer *et al.*, 2003; Jaramillo *et al.*, 2005; Parroche *et al.*, 2007; Prato *et al.*, 2008; Shio *et al.*, 2010; Skorokhod

et al., 2010; Barrera *et al.*, 2011). This study indicates that accelerated MTT-formazan exocytosis in endothelial cells is mediated by the Fe(III)protoporphyrin IX moiety and not by other HZ components. We found indeed that both the native HZ and the synthetic HZ enhance MTT-formazan exocytosis, as in other studies where both the native and synthetic HZ showed similar activity (Jaramillo *et al.*, 2005).

In conclusion, we demonstrated that HZ was not cytotoxic on human microdermal endothelial cells, but accelerated the MTT-formazan exocytosis through the action of its haem moiety. Thus, MTT assay is not suitable to measure cell viability in the presence of compounds that alter MTT formazan exocytosis. Moreover, HZ, by inducing perturbation of intracellular vesicles trafficking, could contribute to the pathogenesis of severe malaria through the release of bioactive molecules in the circulation.

Acknowledgements. Thanks are due to Laura Galastri, Tiziana Bianchi and Paola Verducci from Associazione Volontari Italiani Sangue (AVIS Comunale Milano) for providing fresh blood to culture *Plasmodium falciparum*.

Financial support. This work was supported by the Ministero dell'Istruzione, dell'Università e della Ricerca (PRIN 2015.4JRP004) and by the Università degli Studi di Milano (Piano Sviluppo linea B 2016).

Conflict of interest. None

Ethical standards. Not applicable

References

- Abe K and Saito H (1998) Amyloid beta protein inhibits cellular MTT reduction not by suppression of mitochondrial succinate dehydrogenase but by acceleration of MTT formazan exocytosis in cultured rat cortical astrocytes. *Neuroscience Research* **31**, 295–305.
- Ades EW, Candal FJ, Swerlick RA, George VG, Summers S, Bosse DC and Lawley TJ (1992) HMEC-1: establishment of an immortalized human microvascular endothelial cell line. *Journal of Investigative Dermatology* **99**, 683–690.
- Andrews KT and Lanzer M (2002) Maternal malaria: *Plasmodium falciparum* sequestration in the placenta. *Parasitology Research* **88**, 715–723.
- Armah HB, Wilson NO, Sarfo BY, Powell MD, Bond VC, Anderson W, Adjei AA, Gyasi RK, Tettey Y, Wiredu EK, Tongren JE, Udhayakumar V and Stiles JK (2007) Cerebrospinal fluid and serum biomarkers of cerebral malaria mortality in Ghanaian children. *Malaria Journal* **6**, 147.
- Ayimba E, Hegewald J, Ségbéna AY, Gantin RG, Lechner CJ, Agossou A, Banla M and Soboslay PT (2011) Proinflammatory and regulatory cytokines and chemokines in infants with uncomplicated and severe *Plasmodium falciparum* malaria. *Clinical and Experimental Immunology* **166**, 218–226.
- Barrera V, Skorokhod OA, Baci D, Gremo G, Arese P and Schwarzer E (2011) Host fibrinogen stably bound to hemozoin rapidly activates monocytes via TLR-4 and CD11b/CD18-integrin: a new paradigm of hemozoin action. *Blood* **117**, 5674–5682.
- Basilico N, Tognazioli C, Picot S, Ravagnani F and Taramelli D (2003) Synergistic and antagonistic interactions between haemozoin and bacterial endotoxin on human and mouse macrophages. *Parassitologia* **45**, 135–140.
- Basilico N, Parapini S, Sisto F, Omodeo-Salè F, Coghi P, Ravagnani F, Oliaro P and Taramelli D (2010) The lipid moiety of haemozoin (Malaria Pigment) and *P. falciparum* parasitised red blood cells bind synthetic and native endothelin-1. *Journal of Biomedicine & Biotechnology* **2010**, 854927.
- Basilico N, Corbett Y, D' Alessandro S, Parapini S, Prato M, Girelli D, Misiano P, Oliaro P and Taramelli D (2017) Malaria pigment stimulates chemokine production by human microvascular endothelium. *Acta Tropica* **172**, 125–131.
- Beeson JG, Reeder JC, Rogerson SJ and Brown GV (2001) Parasite adhesion and immune evasion in placental malaria. *Trends in Parasitology* **17**, 331–337.
- Berridge MV and Tan AS (1993) Characterization of the cellular reduction of 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide (MTT): subcellular localization, substrate dependence, and involvement of mitochondrial electron transport in MTT reduction. *Archives of Biochemistry and Biophysics* **303**, 474–482.
- Berridge M, Tan A, Mccoy K and Wang R (1996) The biochemical and cellular basis of cell proliferation assays that use tetrazolium salts. *Biochemica* **4**, 14–16.
- Berridge MV, Herst PM and Tan AS (2005) Tetrazolium dyes as tools in cell biology: new insights into their cellular reduction. *Biotechnology Annual Review* **11**, 127–152.
- Brown H, Rogerson S, Taylor T, Tembo M, Mwenechanya J, Molyneux M and Turner G (2001) Blood-brain barrier function in cerebral malaria in Malawian children. *American Journal of Tropical Medicine and Hygiene* **64**, 207–213.
- Claus V, Jahraus A, Tjelle T, Berg T, Kirschke H, Faulstich H and Griffiths G (1998) Lysosomal enzyme trafficking between phagosomes, endosomes, and lysosomes in J774 macrophages. Enrichment of cathepsin H in early endosomes. *Journal of Biological Chemistry* **273**, 9842–9851.
- D'Alessandro S, Corbett Y, Iboudo DP, Misiano P, Dahiya N, Abay SM, Habluetzel A, Grande R, Gismondo MR, Decherig KJ, Koolen KM, Sauerwein RW, Taramelli D, Basilico N and Parapini S (2015) Salinomycin and other ionophores as a new class of antimalarial drugs with transmission-blocking activity. *Antimicrobial Agents and Chemotherapy* **59**, 5135–5144.
- Deroost K, Pham TT, Opdenakker G and Van den Steen PE (2016) The immunological balance between host and parasite in malaria. *FEMS Microbiology Reviews* **40**, 208–257.
- Diaz G, Melis M, Musin A, Piludu M, Piras M and Falchi AM (2007) Localization of MTT formazan in lipid droplets. An alternative hypothesis about the nature of formazan granules and aggregates. *European Journal of Histochemistry* **51**, 213–218.
- Egan TJ (2008) Haemozoin formation. *Molecular and Biochemical Parasitology* **157**, 127–136.
- Fischella M, Dabboue H, Bhattacharyya S, Saboungi ML, Salvetat JP, Hevor T and Guerin M (2009) Mesoporous silica nanoparticles enhance MTT formazan exocytosis in HeLa cells and astrocytes. *Toxicology in Vitro* **23**, 697–703.
- Giribaldi G, Ulliers D, Schwarzer E, Roberts I, Piacibello W and Arese P (2004) Hemozoin- and 4-hydroxynonenal-mediated inhibition of erythropoiesis. Possible role in malarial dyserythropoiesis and anemia. *Haematologica* **89**, 492–493.
- Goldie P, Roth EF, Oppenheim J and Vanderberg JP (1990) Biochemical characterization of *Plasmodium falciparum* hemozoin. *American Journal of Tropical Medicine and Hygiene* **43**, 584–596.
- Guiguemde WA, Hunt NH, Guo J, Marciano A, Haynes RK, Clark J, Guy RK and Golenser J (2014) Treatment of murine cerebral malaria by artemisone in combination with conventional antimalarial drugs: antiplasmodial effects and immune responses. *Antimicrobial Agents and Chemotherapy* **58**, 4745–4754.
- Hollestelle MJ, Donkor C, Mantey EA, Chakravorty SJ, Craig A, Akoto AO, O'Donnell J, van Mourik JA and Bunn J (2006) von Willebrand factor propeptide in malaria: evidence of acute endothelial cell activation. *British Journal of Haematology* **133**, 562–569.
- Into T, Kanno Y, Dohkan J, Nakashima M, Inomata M, Shibata K, Lowenstein CJ and Matsushita K (2007) Pathogen recognition by Toll-like receptor 2 activates Weibel-Palade body exocytosis in human aortic endothelial cells. *Journal of Biological Chemistry* **282**, 8134–8141.
- Isobe I, Michikawa M and Yanagisawa K (1999) Enhancement of MTT, a tetrazolium salt, exocytosis by amyloid beta-protein and chloroquine in cultured rat astrocytes. *Neuroscience Letters* **266**, 129–132.
- Jaramillo M, Godbout M and Olivier M (2005) Hemozoin induces macrophage chemokine expression through oxidative stress-dependent and -independent mechanisms. *Journal of Immunology* **174**, 475–484.
- Khadjavi A, Giribaldi G and Prato M (2010) From control to eradication of malaria: the end of being stuck in second gear? *Asian Pacific Journal of Tropical Medicine* **3**, 412–420.
- Khaw LT, Ball HJ, Golenser J, Combes V, Grau GE, Wheway J, Mitchell AJ and Hunt NH (2013) Endothelial cells potentiate interferon- γ production in a novel tripartite culture model of human cerebral malaria. *PLoS ONE* **8**, e69521.
- Lamkanra AA, Theron M, Kooij TW and Roberts DJ (2009) Hemozoin (malarial pigment) directly promotes apoptosis of erythroid precursors. *PLoS ONE* **4**, e8446.
- Liu Y and Schubert D (1997) Cytotoxic amyloid peptides inhibit cellular 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide (MTT) reduction

- by enhancing MTT formazan exocytosis. *Journal of Neurochemistry* **69**, 2285–2293.
- Liu Y and Schubert D** (1998) Steroid hormones block amyloid fibril-induced 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide (MTT) formazan exocytosis: relationship to neurotoxicity. *Journal of Neurochemistry* **71**, 2322–2329.
- Liu Y, Peterson DA, Kimura H and Schubert D** (1997) Mechanism of cellular 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide (MTT) reduction. *Journal of Neurochemistry* **69**, 581–593.
- Liu Y, Peterson DA and Schubert D** (1998) Amyloid beta peptide alters intracellular vesicle trafficking and cholesterol homeostasis. *Proceedings of the National Academy of Sciences of the United States of America* **95**, 13266–13271.
- Liu M, Dickinson-Copeland C, Hassana S and Stiles JK** (2016) Plasmodium-infected erythrocytes (pRBC) induce endothelial cell apoptosis via a heme-mediated signaling pathway. *Drug Design, Development and Therapy* **10**, 1009–1018.
- Maknitikul S, Luplertlop N, Chaisri U, Maneerat Y and Ampawong S** (2018) Featured article: immunomodulatory effect of hemozoin on pneumocyte apoptosis via CARD9 pathway, a possibly retarding pulmonary resolution. *Experimental Biology and Medicine (Maywood)* **243**, 395–407.
- Matsushita K, Yamakuchi M, Morrell CN, Ozaki M, O'Rourke B, Irani K and Lowenstein CJ** (2005) Vascular endothelial growth factor regulation of Weibel-Palade-body exocytosis. *Blood* **105**, 207–214.
- Metcalf DJ, Nightingale TD, Zenner HL, Lui-Roberts WW and Cutler DF** (2008) Formation and function of Weibel-Palade bodies. *Journal of Cell Science* **121**, 19–27.
- Molinari BL, Tasat DR, Palmieri MA and Cabrini RL** (2005) Kinetics of MTT-formazan exocytosis in phagocytic and non-phagocytic cells. *Micron* **36**, 177–183.
- Mosmann T** (1983) Rapid colorimetric assay for cellular growth and survival: application to proliferation and cytotoxicity assays. *Journal of Immunological Methods* **65**, 55–63.
- Omodeo-Salè F, Basilico N, Folini M, Olliaro P and Taramelli D** (1998) Macrophage populations of different origins have distinct susceptibilities to lipid peroxidation induced by beta-haematin (malaria pigment). *FEBS Letters* **433**, 215–218.
- Omodeo-Sale F, Motti A, Basilico N, Parapini S, Olliaro P and Taramelli D** (2003) Accelerated senescence of human erythrocytes cultured with *Plasmodium falciparum*. *Blood* **102**, 705–711.
- Pagola S, Stephens PW, Bohle DS, Kosar AD and Madsen SK** (2000) The structure of malaria pigment beta-haematin. *Nature* **404**, 307–310.
- Parroche P, Lauw FN, Goutagny N, Latz E, Monks BG, Visintin A, Halmen KA, Lamphier M, Olivier M, Bartholomeu DC, Gazzinelli RT and Golenbock DT** (2007) Malaria hemozoin is immunologically inert but radically enhances innate responses by presenting malaria DNA to Toll-like receptor 9. *Proceedings of the National Academy of Sciences of the United States of America* **104**, 1919–1924.
- Polimeni M, Valente E, Aldieri E, Khadjavi A, Giribaldi G and Prato M** (2013) Role of 15-hydroxyeicosatetraenoic acid in hemozoin-induced lysosome release from human adherent monocytes. *Biofactors* **39**, 304–314.
- Pongponratn E, Turner GD, Day NP, Phu NH, Simpson JA, Stepniewska K, Mai NT, Viriyavejakul P, Looreesuwan S, Hien TT, Ferguson DJ and White NJ** (2003) An ultrastructural study of the brain in fatal *Plasmodium falciparum* malaria. *American Journal of Tropical Medicine and Hygiene* **69**, 345–359.
- Prato M and Giribaldi G** (2011) Matrix metalloproteinase-9 and haemozoin: wedding rings for human host and *Plasmodium falciparum* parasite in complicated malaria. *Journal of Tropical Medicine* **2011**, ID 628435.
- Prato M, Gallo V, Giribaldi G and Arese P** (2008) Phagocytosis of haemozoin (malarial pigment) enhances metalloproteinase-9 activity in human adherent monocytes: role of IL-1beta and 15-HETE. *Malaria Journal* **7**, 157.
- Prato M, D'Alessandro S, Van den Steen PE, Opdenakker G, Aresel P, Taramelli D and Basilico N** (2011) Modulation of Matrix Metalloproteinases and their tissue inhibitors in haemozoin-treated human microvascular endothelial cells. *Febs Journal* **278**, 310–310.
- Schwarzer E, Kuhn H, Valente E and Arese P** (2003) Malaria-parasitized erythrocytes and hemozoin nonenzymatically generate large amounts of hydroxy fatty acids that inhibit monocyte functions. *Blood* **101**, 722–728.
- Shio MT, Kassa FA, Bellemare MJ and Olivier M** (2010) Innate inflammatory response to the malarial pigment hemozoin. *Microbes and Infection* **12**, 889–899.
- Skorokhod OA, Caione L, Marrocco T, Migliardi G, Barrera V, Arese P, Piacibello W and Schwarzer E** (2010) Inhibition of erythropoiesis in malaria anemia: role of hemozoin and hemozoin-generated 4-hydroxynonenal. *Blood* **116**, 4328–4337.
- Stockert JC, Blázquez-Castro A, Cañete M, Horobin RW and Villanueva A** (2012) MTT assay for cell viability: intracellular localization of the formazan product is in lipid droplets. *Acta Histochemica* **114**, 785–796.
- Taramelli D, Basilico N, De Palma AM, Saresella M, Ferrante P, Mussoni L and Olliaro P** (1998) The effect of synthetic malaria pigment (beta-haematin) on adhesion molecule expression and interleukin-6 production by human endothelial cells. *Transactions of the Royal Society of Tropical Medicine and Hygiene* **92**, 57–62.
- Taramelli D, Recalcati S, Basilico N, Olliaro P and Cairo G** (2000) Macrophage preconditioning with synthetic malaria pigment reduces cytokine production via heme iron-dependent oxidative stress. *Laboratory Investigation* **80**, 1781–1788.
- Traganos F and Darzynkiewicz Z** (1994) Lysosomal proton pump activity: supravital cell staining with acridine orange differentiates leukocyte subpopulations. *Methods in Cell Biology* **41**, 185–194.
- Tyberghein A, Deroost K, Schwarzer E, Arese P and Van den Steen PE** (2014) Immunopathological effects of malaria pigment or hemozoin and other crystals. *Biofactors* **40**, 59–78.
- Utgaard JO, Jahnsen FL, Bakka A, Brandtzaeg P and Haraldsen G** (1998) Rapid secretion of prestored interleukin 8 from Weibel-Palade bodies of microvascular endothelial cells. *Journal of Experimental Medicine* **188**, 1751–1756.
- Vischer UM and Wagner DD** (1993) CD63 is a component of Weibel-Palade bodies of human endothelial cells. *Blood* **82**, 1184–1191.
- Yeo TW, Lampah DA, Gitawati R, Tjitra E, Kenangalem E, Piera K, Price RN, Duffull SB, Celermajer DS and Anstey NM** (2008) Angiotensin-2 is associated with decreased endothelial nitric oxide and poor clinical outcome in severe falciparum malaria. *Proceedings of the National Academy of Sciences of the United States of America* **105**, 17097–17102.