

# *Crithidia bombi* can infect two solitary bee species while host survivorship depends on diet

Laura L. Figueroa<sup>1</sup> , Cali Grincavitch<sup>1,2</sup> and Scott H. McArt<sup>1</sup><sup>1</sup>Department of Entomology, Cornell University, Ithaca, NY 14853, USA and <sup>2</sup>Department of Integrative Biology at Harvard, Harvard University, Cambridge, MA 02138, USA

## Research Article

**Cite this article:** Figueroa LL, Grincavitch C, McArt SH (2021). *Crithidia bombi* can infect two solitary bee species while host survivorship depends on diet. *Parasitology* **148**, 435–442. <https://doi.org/10.1017/S0031182020002218>

Received: 10 July 2020

Revised: 30 October 2020

Accepted: 19 November 2020

First published online: 1 December 2020

### Key words:

Alfalfa leafcutting bee; blue orchard bee; mortality; nutrition; orchard mason bee; pollinator health; solitary bees; survivorship curves; trypanosomes

### Author for correspondence:

Laura L. Figueroa, E-mail: [llf44@cornell.edu](mailto:llf44@cornell.edu)

## Abstract

Pathogens and lack of floral resources interactively impair global pollinator health. However, epidemiological and nutritional studies aimed at understanding bee declines have historically focused on social species, with limited evaluations of solitary bees. Here, we asked whether *Crithidia bombi*, a trypanosomatid gut pathogen known to infect bumble bees, could infect the solitary bees *Osmia lignaria* (females) and *Megachile rotundata* (males), and whether nutritional stress influenced infection patterns and bee survival. We found that *C. bombi* was able to infect both solitary bee species, with 59% of *O. lignaria* and 29% of *M. rotundata* bees experiencing pathogen replication 5–11 days following inoculation. Moreover, access to pollen resulted in *O. lignaria* living longer, although it did not influence *M. rotundata* survival. Access to pollen did not affect infection probability or resulting pathogen load in either species. Similarly, inoculating with the pathogen did not drive survival patterns in either species during the 5–11-day laboratory assays. Our results demonstrate that solitary bees can be hosts of a known bumble bee pathogen, and that access to pollen is an important contributing factor for bee survival, thus expanding our understanding of factors contributing to solitary bee health.

## Introduction

Pathogen pressure, inadequate nutrition and the interaction between these stressors can drastically impair pollinator health (Goulson *et al.*, 2015). The relationship between nutrition and bee disease dynamics is complex, as diet quality and diversity can support immunocompetence and reduce infections in some instances, while increasing disease burdens in other cases (Koch *et al.*, 2017; Dolezal and Toth, 2018). A bee's diet is made up of nectar and pollen, both of which have primary metabolites such as sugars, amino acids and lipids, as well as secondary compounds, such as flavonoids, terpenoids and alkaloids, with the potential to decrease (Richardson *et al.*, 2015) or increase (Palmer-Young and Thursfield, 2017) infections. In a world dominated by agricultural landscapes, with low floral diversity and frequent movement of commercial bee colonies that can potentially introduce pathogens into wild solitary bee populations (Kremen *et al.*, 2002; Otterstatter and Thomson, 2008; Furst *et al.*, 2014; Alger *et al.*, 2019), understanding the interplay between nutrition and pathogen burdens is important for protecting pollinator health.

Immune defence against pathogens is energetically costly, yet the resulting physiological trade-offs between immunity and other fitness components can be compensated by changes in diet (Moret and Schmid-Hempel, 2000). When *Bombus terrestris* bees are starved of pollen, infection with *Crithidia bombi*, a trypanosomatid gut pathogen, markedly increases host mortality compared to bees with access to pollen (Brown *et al.*, 2000). However, when both *B. terrestris* and *B. impatiens* have access to pollen, they can present higher *C. bombi* loads than pollen-starved counterparts (Logan *et al.*, 2005; Conroy *et al.*, 2016). Similarly, access to pollen can increase honey bee survival, while simultaneously increasing pathogen loads of the microsporidians *Nosema apis* (Rinderer and Dell Elliott, 1977) and *Nosema ceranae* (Zheng *et al.*, 2014; Jack *et al.*, 2016). As such, disentangling the effects of nutritional stress and pathogen infection on bee survival is important for understanding pollinator health, especially for solitary bees, which contribute substantially to pollination services worldwide yet have been historically understudied (Danforth *et al.*, 2019).

Honey bees and bumble bees have been the predominant model systems for addressing questions regarding bee health, especially for nutritional and epidemiological evaluations (Schmid-Hempel, 1998). Recent advances in molecular surveillance have revealed widespread pathogen prevalence across solitary bee taxa from most bee families (Andrenidae, Apidae, Colletidae, Halictidae and Megachilidae). This includes pathogens known to infect honey bees and bumble bees (Apidae), including *Apicystis bombi*, *Ascospaera* spp., *C. bombi*, *C. mellifica*, *N. ceranae* and numerous viruses (Singh *et al.*, 2010; Evison *et al.*, 2012; Ravoet *et al.*, 2014; Schoonvaere *et al.*, 2018; Figueroa *et al.*, 2020). However, except for single-stranded RNA viruses which allow for strand-specific PCR assays to detect viral replication, we currently cannot distinguish between transient passage through the bee gut and active infections, nor do we know if there are negative consequences for the host based solely on molecular screenings

(Bramke et al., 2019). Some existing studies that have experimentally evaluated the impacts on solitary bee health have shown increased mortality associated with infections [e.g. *Megachile rotundata* larvae infected with the fungus *Ascospheera aggregata* (James, 2005) and *Osmia bicornis* infected with the neogregarine *A. bombi* and the microsporidian *N. ceranae* (Tian et al., 2018; Bramke et al., 2019)], further highlighting the need to address the host range of bee pathogens and negative consequences for these understudied bee species.

Pathogens are spread between bee species via shared use of floral resources (Graystock et al., 2015). Despite ample possible routes of indirect transmission via flowers at the community level (Figueroa et al., 2020), the host range of many bee pathogens remains currently unknown. Infection with the trypanosomatid *C. bombi* can affect bumble bee foraging behaviour, cognitive function (Gegear et al., 2005, 2006) and reproduction (Goulson et al., 2018). While *C. bombi* is known to infect multiple bumble bee species (Colla et al., 2006; Cordes et al., 2012; Ruiz-González et al., 2012), honey bees are not a known host (Ruiz-González and Brown, 2006; Graystock et al., 2015), even though both groups belong to the same family. It is largely unknown whether solitary bee species, which frequently test positive for *C. bombi* via PCR-based screenings (Figueroa et al., 2020; Graystock et al., 2020), are actually infected by this pathogen (Ravoet et al., 2014). The solitary bees *Osmia lignaria* and *M. rotundata* are cavity-nesting species that provide important pollination services for fruits and vegetables in North America and Europe (Velthuis and van Doorn, 2006; Pitts-Singer and Cane, 2011; Brittain et al., 2013) and can serve as model organisms for experimentally evaluating epidemiological questions due to their commercial availability. Recent experimental work has shown that *C. bombi* collected from *B. impatiens* can infect these species (Ngor et al., 2020), further highlighting the need to understand the biotic and abiotic factors, such as diet, that influence disease dynamics in species beyond honey bees and bumble bees.

To fill this critical knowledge gap, we conducted a study to understand the influence of nutritional stress on the susceptibility of solitary bees to pathogens. Specifically, we asked: (1) how frequently after exposure to *C. bombi* do the solitary bee species *O. lignaria* and *M. rotundata* become infected, (2) does pollen access influence the likelihood of infection and/or subsequent load of *C. bombi* in *O. lignaria* and *M. rotundata*, and (3) does pollen access and/or *C. bombi* exposure influence *O. lignaria* and *M. rotundata* survival?

## Materials and methods

### Study system

For our experiments, we used the trypanosomatid pathogen *C. bombi* (Kinetoplastea, Trypanosomatida) and two solitary bee species: *M. rotundata* (Hymenoptera, Megachilidae; Crown Bees, Woodinville, WA, USA) and *O. lignaria* (Hymenoptera, Megachilidae; Watts bees, Bothell, WA, USA). The bees were obtained as cocoons and allowed to eclose in individually marked housing units: inverted 59 mL salsa cups (Fabi-Kal® Greenware, Lancaster, PA, USA) lined with filter paper (Whatman®, Marlborough, PA, USA) that provided access to 30% sucrose solution through a small opening at the tip of a 1.5 mL microcentrifuge tube (VWR™, Radnor, PA, USA) (Fig. S1). The sex of these individuals can be easily determined visually upon eclosion. As such, only females were maintained for the experiment with *O. lignaria*. Females represented <10% of the eclosed *M. rotundata*, and we therefore conducted the experiments exclusively with males for this species in order to have sufficient sample sizes. As such, we do not make any formal species or sex comparisons in our analyses.

The *C. bombi* we used in our trials was collected from wild *Bombus impatiens* (Hymenoptera, Apidae) workers from Massachusetts, USA (GPS: 42822°17.5300N, 72835°13.5200W). The strain was sustained in laboratory bumble bee colonies (Biobest, Leamington, Ontario, Canada). The two solitary bee species are commonly used in commercial agriculture in the USA, though *O. lignaria* is native to North America while *M. rotundata* is European (Velthuis and van Doorn, 2006).

### Experimental design

To investigate *C. bombi* replication and its effect on bee survival, we conducted a 2 × 2 factorial experiment for each bee species, contrasting nutritional stress (presence/absence of pollen) and infection status (inoculated/sham-inoculated with *C. bombi*). We prepared *C. bombi* inoculum fresh for each inoculation day by dissecting the gut of infected *B. impatiens* bees from the laboratory source colony. We homogenized the bee guts in distilled water and quantified *C. bombi* cells using a haemocytometer. We diluted the mixture to 1200 *C. bombi* cells  $\mu\text{L}^{-1}$ , which we then combined 1:1 with 30% sucrose solution for an inoculum of 600 cells  $\mu\text{L}^{-1}$ , a standard inoculum concentration for infecting bumble bees with *C. bombi* (Richardson et al., 2015; Figueroa et al., 2019). We used 30% sucrose without *C. bombi* as a control (sham) inoculum. The sucrose solution was coloured with blue food colouring (McCormick & Company, Baltimore, MD, USA) to facilitate confirmation of consumption and availability.

For each trial, half of the bees were inoculated with *C. bombi* and half received the sham inoculation. Each bee species was inoculated on separate dates: *O. lignaria* on 27 June 2019 ( $n = 97$ ), 22 July 2019 ( $n = 48$ ) and 23 July 2019 ( $n = 52$ ), while *M. rotundata* was inoculated on 7 July 2019 ( $n = 100$ ). The unit of replication for this study was individual bees. The lower sample size for *M. rotundata* was due to lower availability of bees. The bees had eclosed 1–2 days before being inoculated in their individual housing units. Each treatment was evenly represented on each inoculation date. Standard *C. bombi* inoculations in bumble bees are conducted by eliciting proboscis extension from the sugar in the inoculum and directly feeding the bees 10  $\mu\text{L}$  of inoculum (Richardson et al., 2015; Figueroa et al., 2019). However, in a pilot study, we found that *O. lignaria* and *M. rotundata* bees were uncooperative and would not consume the inoculum droplets using the standardized bumble bee protocol. When we subdued the bees using CO<sub>2</sub>, we were able to place the inoculum droplet directly on their extended proboscis, finding that this method was effective at infecting the solitary bees.

To administer the inoculum for this experiment, we exposed individual bees to CO<sub>2</sub> gas for 45 s, during which time most bees extended their proboscis. We then placed 5  $\mu\text{L}$  of inoculum on their proboscis and on the pollen, if present, or on the side of the feeding tube for pollen-free treatments, to maximize exposure to the pathogen (10  $\mu\text{L}$  on the first day of trial), *O. lignaria* received an additional 5  $\mu\text{L}$  dose on the pollen/feeding tube for two consecutive days for a total of 20  $\mu\text{L}$  (12 000 *C. bombi* cells) administered compared to the 10  $\mu\text{L}$  (6000 *C. bombi* cells) for *M. rotundata*. The additional doses were administered to increase the likelihood of exposure to the pathogen in the larger of the two species (*O. lignaria*). Trays containing the bees were separated by treatment to avoid cross-contamination, but were all maintained on the same laboratory bench for the duration of the assay.

Half of the bees in both the inoculated and sham-inoculated treatments were provided ~36 mg balls of pollen, made from a mixture of honey bee collected poly-floral pollen (Bee Pollen Granules, CC Pollen High Desert, Phoenix AZ, USA), and all bees were provided 30% sucrose solution. The sucrose solution was replaced every 3 days and pollen balls were given to the pollen

treatment bees every other day. We verified that the pollen had little to no pesticides by screening a pollen sample for 267 pesticides using liquid chromatography/mass spectrometry (Urbanowicz *et al.*, 2019). We detected only one pesticide, the acaricide coumaphos, at a level below the limit of quantification ( $<0.525 \text{ ng g}^{-1}$ ), and thus concluded that pesticide exposure would not be a primary driver of any pattern found with pollen access. The pollen was not a source of *C. bombi* as can be verified by the absence of *C. bombi* in our sham-inoculated bees fed pollen (Fig. 1). While *O. lignaria* larvae tend to develop more quickly and larger body sizes on pollen collected by members of its own species, they will nonetheless develop on honey bee-collected pollen (Levin and Haydak, 1957). Whether this translates into differences in pollen feeding by adults (both female *O. lignaria* and male *M. rotundata*) has not been tested. Though we did not quantify pollen consumption, we observed frass with pollen residues for some of the bees in the pollen-access treatments, indicating consumption. The bees were maintained in laboratory conditions at an average temperature of approximately 20°C in constant dark.

We checked bee survival daily for the duration of the trial (terminated after 11 days for *M. rotundata*, and 8, 6 and 5 days for the three *O. lignaria* trials, respectively), and recorded daily mortality for each bee. Trial lengths varied due to differences in mortality; while *M. rotundata* had low overall mortality (18% died by the end of the 11-day trial), mortality for *O. lignaria* was overall higher and varied greatly (ranged from 14 to 54% depending on the trial). We dissected any bee that had died within 24 h to screen for *C. bombi* (checked daily), as well as all bees that survived until the end of the trial. Given that our pathogen counts are based on motile cells, we expected the most accurate counts from bees that were alive. As such, we shortened the trial times for *O. lignaria* in order to have enough live bees to accurately quantify infection because observing live *C. bombi* in dead bees, while possible, is likely less accurate. For example, only 8% of recorded *C. bombi*-positive *O. lignaria* bees had died before the end of the trial and had much lower corresponding pathogen loads (mean of 71 active *C. bombi* cells  $\mu\text{L}^{-1}$  compared to 171 cells  $\mu\text{L}^{-1}$  in bees that survived the length of the trial). We cannot determine whether the lower likelihood of detection and lower counts were products of insufficient time for the pathogen to replicate or whether the pathogen had died within the host before dissection and could not be visualized. As such, infection analyses were conducted only on bees that survived until the end of the trial. Nonetheless, the shorter trial times for *O. lignaria* are within time frames relevant for *C. bombi* replication in bumble bees (Schmid-Hempel and Schmid-Hempel, 1993). We homogenized the dissected bee guts in 200  $\mu\text{L}$  of distilled water, incubated the mixture for 4 h at room temperature and finally quantified motile *C. bombi* using a haemocytometer (Richardson *et al.*, 2015; Figueroa *et al.*, 2019).

For the bees that had motile *C. bombi*, we evaluated whether the values indicated active replication by the pathogen. To do this, we compared estimated whole gut counts to the value in the entirety of the inoculum provided to the bees. The *C. bombi*  $\mu\text{L}^{-1}$  observed for each bee was multiplied by 200  $\mu\text{L}$ , the volume of water in which the gut was incubated, indicating the total number of *C. bombi* cells estimated to be in the bee gut. This is a conservative estimate as we are not including the volume of the gut itself. Values above 12 000 for *O. lignaria* and 6000 for *M. rotundata* indicate active pathogen replication in the bees as these values correspond to the maximum possible cells consumed in the inoculum. We report these numbers as 'whole gut *C. bombi* estimates' in the results.

At the end of the test periods, we measured the inter-tegular distance (ITD) of all of the bees using an Olympus SZX10 microscope and cellSens Standard software (Olympus Corporation of the Americas, Scientific Solutions Group, Waltham,

Massachusetts, USA). The ITD is commonly used as a proxy for bee body size (Greenleaf *et al.*, 2007), which is a factor related to immunity in bumble bees (Otterstatter and Thomson, 2006).

### Statistical analyses

Data analyses were conducted using R version 3.5.1 (R Development Core Team, 2008) with the lme4, glmmTMB and coxme packages (Bates *et al.*, 2015; Therneau and Therneau, 2015; Brooks *et al.*, 2017). We conducted analyses on *M. rotundata* and *O. lignaria* separately.


To determine the role of pollen access on *C. bombi* infection patterns in the solitary bees, we employed a manual two-step hurdle model due to zero-inflation and overdispersion in the data (Zuur *et al.*, 2009). As mentioned above, these analyses were only conducted on inoculated bees that survived until the last day of the trial to increase the accuracy of *C. bombi* counts (11 days for *M. rotundata* and 5, 6 or 8 days for the three *O. lignaria* trials). We first evaluated whether pollen access predicted the presence or absence of *C. bombi* in the gut of inoculated bees, followed by an evaluation of the *C. bombi* counts for those that were infected. For *O. lignaria*, we first constructed a generalized linear mixed model (GLMM) to evaluate whether the solitary bee species harboured *C. bombi* in their gut as the response (yes/no), predicted by access to pollen (yes/no) and bee size ( $n = 59$ ). The model included inoculation date as the random effect and a binomial distribution (logit link). To determine significance, we employed a likelihood ratio test and compared the model to one with the same response, distribution and random-effect structure, but which excluded pollen as a predictor. We then developed a GLMM that included *C. bombi* count in *C. bombi*-positive bees as the response variable, predicted by access to pollen (presence/absence) and bee size ( $n = 49$ ). The model included inoculation date as the random effect and fit a truncated negative binomial distribution, which is suitable for count data with overdispersion (Brooks *et al.*, 2017). Significance was similarly determined using a likelihood ratio test. We verified model assumptions using the DHARMA package (Hartig, 2017). The statistical analyses for *M. rotundata* were likewise evaluated, with the sole difference that the manual hurdle model was conducted using a Generalized Linear Model instead of a GLMM because the species was inoculated in a single day and thus did not require inoculation date as a random effect ( $n = 35$  for binomial response and  $n = 10$  for *C. bombi* count as a response).

To evaluate whether exposure to *C. bombi*, access to pollen and their interactions influenced solitary bee survival, we conducted survival analyses using Cox proportional hazards models ( $n = 179$  and  $n = 93$  for *O. lignaria* and *M. rotundata*, respectively). For *O. lignaria*, the survival analysis evaluated bee survival (death/days elapsed) as the response, *C. bombi* inoculation and access to pollen as explanatory variables, and inoculation date as the random effect. To determine the significance of the treatments (pollen and inoculation), we conducted a likelihood ratio test comparing the full model with a model that included the same random-effect structure but excluded either explanatory variable or included an additive relationship instead of an interaction. The analyses for *M. rotundata* were conducted using the *coxph* function (no random effect since all bees were inoculated on the same day) while the *coxme* function was used for *O. lignaria* (inoculation date as a random effect) (Therneau and Therneau, 2015).

### Results

We found that both *O. lignaria* females and *M. rotundata* males became infected with *C. bombi* sourced from the common eastern






**Fig. 1.** Effect of pollen access on motile *C. bombi* presence in solitary bees that survived the length of the trial. Pollen access did not affect the likelihood of *C. bombi* presence in (A) *Osmia lignaria* females or (B) *Megachile rotundata* males. Differences in sample size are a product of mortality. (n.s.) indicates  $P > 0.05$ .

bumble bee (*B. impatiens*). Specifically, 76% of inoculated *O. lignaria* and 29% of *M. rotundata* harboured active *C. bombi* in their gut 5–11 days post inoculation; we did not detect *C. bombi* in any sham-inoculated bees (Fig. 1). The median *C. bombi* load at the end of the trial for inoculated *O. lignaria* females was 100 moving cells  $\mu\text{L}^{-1}$  of bee gut (range: 0–1600 cells), while for *M. rotundata* males, the median was 0 cells  $\mu\text{L}^{-1}$  (range: 0–300 cells). The *O. lignaria* females were inoculated with 12 000 *C. bombi* cells and final counts averaged 20 000 cells (range: 0–320 000 whole gut cell estimates), indicating active replication of *C. bombi* in *O. lignaria* (Fig. 2). While the median *C. bombi* whole gut count was zero in *M. rotundata* males, the bees with the highest gut counts in this species (60 000) had more than the administered 6000 *C. bombi* cells, indicating replication of *C. bombi* in *M. rotundata* was also possible (Fig. 2). Specifically, 56% of inoculated *O. lignaria* females and 29% of *M. rotundata* males had estimated whole gut *C. bombi* counts above the inoculation concentration indicating active pathogen replication (Fig. 2).

Pollen access did not influence the likelihood of becoming infected with *C. bombi* for *O. lignaria* females or *M. rotundata* males ( $\chi^2_1 = 1.86$ ,  $P = 0.173$  and  $z = 0.05$ ,  $P = 0.962$ , respectively; Fig. 1) nor subsequent load for infected bees ( $\chi^2_1 = 0.05$ ,  $P = 0.830$  and  $z = 1.78$ ,  $P = 0.076$ , respectively for *O. lignaria* females and *M. rotundata* males; Fig. 3). Bee size did not explain the likelihood of infection with *C. bombi* nor subsequent load for the infected bees in *O. lignaria* females or *M. rotundata* males ( $z < 0.84$  and  $P > 0.402$  for all).

Access to pollen increased *O. lignaria* female bee survival rates ( $\chi^2_1 = 4.13$ ,  $P = 0.042$ ; Fig. 4A), though not for *M. rotundata* males ( $\chi^2_1 = 1.19$ ,  $P = 0.275$ ; Fig. 4C). Inoculation with *C. bombi* did not influence survival in *O. lignaria* females or *M. rotundata* males ( $\chi^2_1 = 1.56$ ,  $P = 0.212$  and  $\chi^2_1 = 2.02$ ,  $P = 0.155$ , respectively;




**Fig. 2.** Estimated whole gut counts of *C. bombi* in solitary bees 5–11 days post-inoculation. Blue circles indicate estimated whole gut counts in inoculated *M. rotundata* males and *O. lignaria* females (see Methods). The dashed red line indicates the total number of *C. bombi* cells in inoculum provided to *M. rotundata* males and the solid red line indicates the total number of *C. bombi* cells in inoculum provided to *O. lignaria* females. Values above these thresholds indicate active pathogen replication in the host bees.

Fig. 4B and D), nor was there pollen access by inoculation interaction for survival in *O. lignaria* females ( $\chi^2_1 = 2.26$ ,  $P = 0.133$ ) or *M. rotundata* males ( $\chi^2_1 = 1.73$ ,  $P = 0.189$ ).


## Discussion

In this study, we found that the solitary bee species *O. lignaria* and *M. rotundata* could become infected with *C. bombi* collected from bumble bees. We found evidence for *C. bombi* replication in 56% of inoculated *O. lignaria* females and 29% of inoculated *M. rotundata* males. Access to pollen did not affect *C. bombi* infection in *O. lignaria* females or *M. rotundata* males. However, pollen-fed *O. lignaria* females survived longer than their pollen-starved counterparts. Inoculation with *C. bombi* did not affect survival in either solitary bee species during the 5–11-day laboratory assay. Overall, our study illustrates that solitary bees can be hosts for a pathogen known to infect bumble bees and that diet can play a key role in the health of these important but historically understudied solitary bee taxa.

Most bee species within a community can be exposed to numerous pathogens when foraging at flowers, including *C. bombi* (Figueroa et al., 2020; Graystock et al., 2020). Our results support a growing body of literature indicating the need to assess the host range of bee pathogens, including assessments of replication and impacts on survival (Bramke et al., 2019; Müller et al., 2019). The presence of *C. bombi* in bee feces has been used to identify active infections in bumble bees (Imhoof and Schmid-Hempel, 1999), where it can be detected in as quickly as 5 days post-inoculation (Logan et al., 2005). We detected active *C. bombi* in solitary bee guts at the end of the trials, 5–11 days post-inoculation, for a total of 76% of inoculated *O. lignaria* females and 29% of inoculated *M. rotundata* males harbouring motile cells. Moreover, 56% of the inoculated *O. lignaria* females and 29% of the inoculated *M. rotundata* males had estimated whole gut counts above the concentration of inoculum provided,



**Fig. 3.** Effect of pollen access on *C. bombi* load in solitary bees. Pollen access did not affect *C. bombi* load (cells  $\mu\text{L}^{-1}$ ) in (A) *Osmia lignaria* females or (B) *Megachile rotundata* males. (n.s.) indicates  $P > 0.05$ . This figure represents subsequent load for inoculated bees that presented motile *C. bombi*.



**Fig. 4.** Solitary bee survival across pollen access and *C. bombi* inoculation treatments. (A) *Osmia lignaria* females with access to pollen survived longer than those without access to pollen. (B). Inoculation with *C. bombi* did not influence *O. lignaria* survivorship. (C) Pollen access did not affect *Megachile rotundata* male survivorship, (D) nor did inoculation with *C. bombi*. Trial lengths varied according to general mortality, which was greater in *M. rotundata* than for *O. lignaria* (see Methods). Shaded error bands indicate 95% confidence intervals. (\*) denotes  $P < 0.05$  while (n.s.) indicates  $P > 0.05$ .

indicating active pathogen replication. These values are comparable to infection probability in bumble bees [e.g. 57% of inoculated *B. terrestris* bees presented active infections after a 7-day trial (Näpflin and Schmid-Hempel, 2018)]. Furthermore, our estimates for the percentage of bees experiencing *C. bombi* replication are conservative (i.e. low) since we assume the bee gut did not contribute to the volume of the extraction and that the bees consumed the entirety of the inoculum droplet, further highlighting the extent to which solitary bees can develop active *C. bombi* infections.

While phylogenetic distance often predicts the ability of hosts to be infected by a given pathogen (Gilbert and Webb, 2007; Streicker *et al.*, 2010), this may not be the case with *C. bombi* and its known hosts. Specifically, one recent study (Ngor *et al.*, 2020) has found that *C. bombi* does not replicate in honey bees (*Apis mellifera*), but this pathogen does replicate in bumble bees (*Bombus* spp.). Both honey bees and bumble bees are in the Apidae family, while the two evaluated solitary bees evaluated in the present study are in the Megachilidae family. The factors that enable honey bees to avoid chronic *C. bombi* infections, despite being able to transmit the pathogen *via* flowers (Ruiz-González and Brown, 2006; Graystock *et al.*, 2015) and

being more phylogenetically similar to bumble bees than *O. lignaria* and *M. rotundata*, remain unknown. Future evaluations of *C. bombi* infection and health impacts on solitary and social species across additional bee families are clearly warranted.

Host nutrition can increase infection intensity by providing resources to pathogens, but may also support host tolerance of infection. Specifically, increased access to food could increase infections by (1) directly providing nutrients to the pathogen; or (2) improving host quality for the pathogen. Conversely, food availability could suppress pathogens by (3) enabling physiologically costly immune responses; and/or (4) providing antimicrobial compounds (Conroy *et al.*, 2016; Palmer-Young and Thursfield, 2017). While we did not find that pollen access directly influenced the likelihood or severity of *C. bombi* infections (Figs 1 and 3), others have found that pollen-deprivation in bumble bees can reduce *C. bombi* infections (Logan *et al.*, 2005; Conroy *et al.*, 2016). We found that pollen access increased *O. lignaria* female survival, independent of *C. bombi* infection. While we did not find an effect of pollen on *M. rotundata*, we cannot determine whether this was due to low mortality during the 11-day trial and whether patterns would differ in longer time scales. Similarly to *O. lignaria*, access to pollen can increase

honey bee survival rates despite greater pathogen loads [e.g. *N. apis*, *N. ceranae* and numerous RNA viruses (Rinderer and Dell Elliott, 1977; Zheng *et al.*, 2014; Jack *et al.*, 2016; Zhang *et al.*, 2020)], further illustrating the importance of diet in pollinator health. Access to pollen in provisioning female solitary bees has been shown to increase the number of brood cells and proportion of offspring surviving to adulthood [e.g. in *O. bicornis* and *O. californica* (Cane, 2016; Bukovinszky *et al.*, 2017)], suggesting that differences in physiological demands between sexes could result in differential effects of pollen access. Furthermore, a recent study that experimentally inoculated *O. cornuta* with *C. mellificae*, a trypanosomatid known to infect honey bees, found that while both sexes had higher pathogen loads over time, only male survival was significantly reduced by infection (Strobl *et al.*, 2019). Future evaluations of the impact of *C. bombi* on physiology and survival in solitary bees across sexes are well justified, especially in the field, and could incorporate assessments of resource availability and bee immune responses.

Bumble bees infected with *C. bombi* (Richardson *et al.*, 2015) and honey bees infected with *N. ceranae* (Mayack and Naug, 2009) often consume more pollen, likely to compensate for the energetic cost of mounting an immune response (Moret and Schmid-Hempel, 2000). Interestingly, the role of pollen on *C. bombi* growth varies depending on whether the assessments are conducted *in vitro* or *in vivo*; for example, sunflower pollen *in vitro* increases pathogen growth (Palmer-Young and Thursfield, 2017), while *in vivo* this pollen type has a strong inhibitory effect on the pathogen when consumed by bumble bees (Giacomini *et al.*, 2018; LoCascio *et al.*, 2019). When inhibitory, secondary compounds appear to facilitate these anti-*C. bombi* interactions in bumble bees (Richardson *et al.*, 2015; Koch *et al.*, 2017). These differences suggest that pollen may interact in important ways within the host, leading to varying effects on *C. bombi* growth. Future evaluations of the role of pollen quality, quantity and diversity on susceptibility to pathogens mediated by interactions between immune responses, gut physiology and microbiota are important avenues for pollinator research (Alaux *et al.*, 2010; Cotter *et al.*, 2011; Dolezal and Toth, 2018), especially focusing on solitary bees and other understudied pollinators.

Life-history and functional traits are important mediators of disease transmission and dynamics within a host. While we did not find a role of pollen in *M. rotundata* male infection patterns, recent work has shown that adult male solitary bees (*Andrena* spp.) frequently harbour pollen in their digestive tract (Urban-Mead *et al.*, unpublished results). The importance of males in dispersing pathogens in plant–pollinator networks is not well established, despite male and female solitary bees having comparable infection rates (Müller *et al.*, 2019; Strobl *et al.*, 2019; Ngor *et al.*, 2020) and marked differences in floral preference (Roswell *et al.*, 2019). Moreover, natural rates of pathogen acquisition or deposition by solitary bees while foraging is not known (Figueroa *et al.*, 2019), including the threshold of pathogen cells necessary for solitary bees to develop active infections, all of which could strengthen disease spread models. While we report high rates of *C. bombi* infection for the two solitary species evaluated, especially for *O. lignaria* females, we did not find that experimentally exposing the solitary bees to *C. bombi* influenced host survival in the 5–11-day laboratory assay. Characterizing naturally occurring infection rates as well as the corresponding impacts on mortality and reproduction in the field, especially alongside numerous co-occurring stressors (e.g. inadequate diet, pesticide exposure and co-infections), is an important future direction. Differences in life histories across solitary bees may influence these dynamics. For example, *O. lignaria* overwinter as adults, similar to bumble bee queens [a life stage highly vulnerable to *C. bombi* (Brown *et al.*, 2003)], while *M. rotundata* overwinter

as prepupae (Kemp *et al.*, 2004). Similarly, given that we know infection with *C. bombi* alters resource allocation patterns in *B. terrestris*, whereby infected bees invest more in their fat body and less in reproduction than non-infected bees (Brown *et al.*, 2000), assessing differences in resource allocation, foraging behaviour and pollen-provisioning abilities for infected solitary bees compared to uninfected counterparts could advance our understanding of pollinator health in wild bee communities.

Overall, our work supports the importance of improving knowledge concerning solitary bee species in pollinator epidemiology, including their assessment as hosts for what have traditionally been considered social bee pathogens, and considering resource availability when evaluating host–pathogen interactions. Future evaluations of the likelihood of transmission between solitary and social species on flowers would benefit from this knowledge, as transmission probability is already known to vary between bumble bee species (Ruiz-González *et al.*, 2012). Moreover, the activity period of *Osmia* spp. and *Megachile* spp. often greatly overlaps with bumble bees, further highlighting the need to understand interspecies disease transmission networks (Figueroa *et al.*, 2020). In addition, pathogen transmission between social and solitary bees could occur *via* spillover by introduced commercial colonies (Colla *et al.*, 2006; Otterstatter and Thomson, 2008; Graystock *et al.*, 2013), and could occur in various directions (Graystock *et al.*, 2016), further highlighting the need to understand host ranges and impacts of disease on pollinator communities. In order to respond to the growing dependence of pollinators for food security (Aizen *et al.*, 2019), which is in large part contributed by wild bees (Garibaldi *et al.*, 2013; Winfree *et al.*, 2018), we must expand our understanding of the role of pathogens, nutrition and other stressors on solitary bee health.

**Supplementary material.** The supplementary material for this article can be found at <https://doi.org/10.1017/S0031182020002218>

**Acknowledgements.** We thank Abby Davis, Julie Davis, Mesly Mata Loya and Alondra Torres Gonzalez for assistance with the experimental set-up and trials, Laura Harrington, Quinn McFrederick, Evan Palmer-Young, Todd Ugine and members of the Cornell Pollinator Reading Group for reading a previous version of the manuscript, and Wee Hao Ng for assistance with statistical analyses. We would also like to thank Jonathan Giacomini, Bryanna Joyce, Simon Pinilla-Gallego and Jenny Van Wyk for assistance in obtaining source bees for making the inoculum used in the trials.

**Financial support.** This work was supported by the National Science Foundation Graduate Research Fellowship (grant number DGE-1650441), the National Science Foundation Postdoctoral Research Fellowships in Biology Program (grant number NSF-2010615), and the National Institute of General Medical Sciences of the National Institutes of Health (grant number R01GM122062). The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health or the National Science Foundation. This material is based upon work supported by the NSF Postdoctoral Research Fellowships in Biology Program under Grant No. 2010615.

**Conflict of interest.** None.

**Ethical standards.** Not applicable.

## References

- Aizen MA, Aguiar S, Biesmeijer JC, Garibaldi LA, Inouye DW, Jung C, Martins DJ, Medel R, Morales CL, Ngo H, Pauw A, Paxton RJ, Sáez A and Seymour CL (2019) Global agricultural productivity is threatened by increasing pollinator dependence without a parallel increase in crop diversification. *Global Change Biology* 25, 3516–3527.
- Alaux C, Ducloz F, Crauser D and Conte YL (2010) Diet effects on honeybee immunocompetence. *Biology Letters* 6, 562–565.
- Alger SA, Burnham PA, Boncristiani HF and Brody AK (2019) RNA virus spillover from managed honeybees (*Apis mellifera*) to wild bumblebees (*Bombus* spp.). *PLoS ONE* 14, e0217822.



- Bates D, Mächler M, Bolker B and Walker S (2015) Fitting linear mixed-effects models using lme4. *Journal of Statistical Software* **67**, 1–48.
- Bramke K, Müller U, McMahon DP and Rolff J (2019) Exposure of larvae of the solitary bee *Osmia bicornis* to the honey bee pathogen *Nosema ceranae* affects life history. *Insects* **10**, 380.
- Brittain C, Williams N, Kremen C and Klein A-M (2013) Synergistic effects of non-Apis bees and honey bees for pollination services. *Proceedings of the Royal Society B-Biological Sciences* **280**, 20122767.
- Brooks ME, Kristensen K, van Benthem KJ, Magnusson A, Berg CW, Nielsen A, Skaug HJ, Machler M and Bolker BM (2017) glmmTMB balances speed and flexibility among packages for zero-inflated generalized linear mixed modeling. *The R Journal* **9**, 378–400.
- Brown M, Loosli R and Schmid-Hempel P (2000) Condition-dependent expression of virulence in a trypanosome infecting bumblebees. *Oikos* **91**, 421–427.
- Brown MJ, Schmid-Hempel R and Schmid-Hempel P (2003) Strong context-dependent virulence in a host–parasite system: reconciling genetic evidence with theory. *Journal of Animal Ecology* **72**, 994–1002.
- Bukovinsky T, Rikken I, Evers S, Wäckers FL, Biesmeijer JC, Prins HH and Kleijn D (2017) Effects of pollen species composition on the foraging behaviour and offspring performance of the mason bee *Osmia bicornis* (L.). *Basic and Applied Ecology* **18**, 21–30.
- Cane JH (2016) Adult pollen diet essential for egg maturation by a solitary *Osmia* bee. *Journal of Insect Physiology* **95**, 105–109.
- Colla SR, Otterstatter MC, Gegear RJ and Thomson JD (2006) Plight of the bumble bee: pathogen spillover from commercial to wild populations. *Biological Conservation* **129**, 461–467.
- Conroy TJ, Palmer-Young EC, Irwin RE and Adler LS (2016) Food limitation affects parasite load and survival of *Bombus impatiens* (Hymenoptera: Apidae) infected with *Crithidia* (Trypanosomatida: Trypanosomatidae). *Environmental Entomology* **45**, 1212–1219.
- Cordes N, Huang WF, Strange JP, Cameron SA, Griswold TL, Lozier JD and Solter LF (2012) Interspecific geographic distribution and variation of the pathogens *Nosema bombi* and *Crithidia* species in United States bumble bee populations. *Journal of Invertebrate Pathology* **109**, 209–216.
- Cotter SC, Simpson SJ, Raubenheimer D and Wilson K (2011) Macronutrient balance mediates trade-offs between immune function and life history traits. *Functional Ecology* **25**, 186–198.
- Danforth BN, Minckley RL, Neff JL and Fawcett F (2019) *The Solitary Bees: Biology, Evolution, Conservation*. Princeton, New Jersey: Princeton University Press.
- Dolezal AG and Toth AL (2018) Feedbacks between nutrition and disease in honey bee health. *Current Opinion in Insect Science* **26**, 114–119.
- Evison SEF, Roberts KE, Laurenson L, Pietravalle S, Hui J, Biesmeijer JC, Smith JE, Budge G and Hughes WOH (2012) Pervasiveness of parasites in pollinators. *PLoS ONE* **7**, e30641.
- Figuerola LL, Blinder M, Grincavitch C, Jelinek A, Mann EK, Merva LA, Metz LE, Zhao AY, Irwin RE and McArt SH (2019) Bee pathogen transmission dynamics: deposition, persistence and acquisition on flowers. *Proceedings of the Royal Society B-Biological Sciences* **286**, 20190603.
- Figuerola LL, Grab H, Ng WH, Myers CR, Graystock P, McFrederick QS and McArt SH (2020) Landscape simplification shapes pathogen prevalence in plant-pollinator networks. *Ecology Letters* **23**, 1212–1222.
- Furst MA, McMahon DP, Osborne JL, Paxton RJ and Brown MJF (2014) Disease associations between honeybees and bumblebees as a threat to wild pollinators. *Nature* **506**, 364–366.
- Garibaldi LA, Steffan-Dewenter I, Winfree R, Aizen MA, Bommarco R, Cunningham SA, Kremen C, Carvalheiro LG, Harder LD, Afik O, Bartomeus I, Benjamin F, Boreux V, Cariveau D, Chacoff NP, Dudenhoffer JH, Freitas BM, Ghazoul J, Greenleaf S, Hipolito J, Holzschuh A, Howlett B, Isaacs R, Javorek SK, Kennedy CM, Krewenka KM, Krishnan S, Mandelik Y, Mayfield MM, Motzke I, Munyuli T, Nault BA, Otieno M, Petersen J, Pisanty G, Potts SG, Rader R, Ricketts TH, Rundlof M, Seymour CL, Schuëpp C, Szentgyorgyi H, Taki H, Tscharntke T, Vergara CH, Viana BF, Wanger TC, Westphal C, Williams N and Klein AM (2013) Wild pollinators enhance fruit set of crops regardless of honey bee abundance. *Science (New York, NY)* **339**, 1608–1611.
- Gegear RJ, Otterstatter MC and Thomson JD (2005) Does parasitic infection impair the ability of bumblebees to learn flower-handling techniques? *Animal Behaviour* **70**, 209–215.
- Gegear RJ, Otterstatter MC and Thomson JD (2006) Bumble-bee foragers infected by a gut parasite have an impaired ability to utilize floral information. *Proceedings of the Royal Society B-Biological Sciences* **273**, 1073–1078.
- Giacomini JJ, Leslie J, Tarpay DR, Palmer-Young EC, Irwin RE and Adler LS (2018) Medicinal value of sunflower pollen against bee pathogens. *Scientific Reports* **8**, 14394.
- Gilbert GS and Webb CO (2007) Phylogenetic signal in plant pathogen-host range. *Proceedings of the National Academy of Sciences of the USA* **104**, 4979–4983.
- Goulson D, Nicholls E, Botías C and Rotheray EL (2015) Bee declines driven by combined stress from parasites, pesticides, and lack of flowers. *Science (New York, NY)* **347**, 1255957.
- Goulson D, O'Connor S and Park KJ (2018) The impacts of predators and parasites on wild bumblebee colonies. *Ecological Entomology* **43**, 168–181.
- Graystock P, Yates K, Evison SEF, Darvill B, Goulson D and Hughes WOH (2013) The Trojan hives: pollinator pathogens, imported and distributed in bumblebee colonies. *Journal of Applied Ecology* **50**, 1207–1215.
- Graystock P, Goulson D and Hughes WOH (2015) Parasites in bloom: flowers aid dispersal and transmission of pollinator parasites within and between bee species. *Proceedings of the Royal Society B-Biological Sciences* **282**, 20151371.
- Graystock P, Blane EJ, McFrederick QS, Goulson D and Hughes WOH (2016) Do managed bees drive parasite spread and emergence in wild bees? *International Journal for Parasitology: Parasites and Wildlife* **5**, 64–75.
- Graystock P, Ng W, Parks K, Tripodi A, Muñoz P, Fersch A, Myers C, McFrederick Q and McArt S (2020) Dominant bee species and floral abundance drive parasite temporal dynamics in plant-pollinator communities. *Nature Ecology and Evolution* **4**, 1358–1367.
- Greenleaf SS, Williams NM, Winfree R and Kremen C (2007) Bee foraging ranges and their relationship to body size. *Oecologia* **153**, 589–596.
- Hartig F (2017) DHARMa: residual diagnostics for hierarchical (multi-level/mixed) regression models. R package version 0.1.5.
- Imhoof B and Schmid-Hempel P (1999) Colony success of the bumble bee, *Bombus terrestris*, in relation to infections by two protozoan parasites, *Crithidia bombi* and *Nosema bombi*. *Insectes Sociaux* **46**, 233–238.
- Jack CJ, Uppala SS, Lucas HM and Sagili RR (2016) Effects of pollen dilution on infection of *Nosema ceranae* in honey bees. *Journal of Insect Physiology* **87**, 12–19.
- James RR (2005) Temperature and chalkbrood development in the alfalfa leaf-cutting bee, *Megachile rotundata*. *Apidologie* **36**, 15–23.
- Kemp W, Bosch J and Dennis B (2004) Oxygen consumption during the life cycles of the prepupa-wintering bee *Megachile rotundata* and the adult-wintering bee *Osmia lignaria* (Hymenoptera: Megachilidae). *Annals of the Entomological Society of America* **97**, 161–170.
- Koch H, Brown MJF and Stevenson PC (2017) The role of disease in bee foraging ecology. *Current Opinion in Insect Science* **21**, 60–67.
- Kremen C, Williams NM and Thorp RW (2002) Crop pollination from native bees at risk from agricultural intensification. *Proceedings of the National Academy of Sciences of the USA* **99**, 16812–16816.
- Levin MD and Haydak MH (1957) Comparative value of different pollens in the nutrition of *Osmia lignaria*. *Bee World* **38**, 221–226.
- LoCascio GM, Aguirre L, Irwin RE and Adler LS (2019) Pollen from multiple sunflower cultivars and species reduces a common bumblebee gut pathogen. *Royal Society Open Science* **6**, 190279.
- Logan A, Ruiz-González M and Brown M (2005) The impact of host starvation on parasite development and population dynamics in an intestinal trypanosome parasite of bumble bees. *Parasitology* **130**, 637–642.
- Mayack C and Naug D (2009) Energetic stress in the honeybee *Apis mellifera* from *Nosema ceranae* infection. *Journal of Invertebrate Pathology* **100**, 185–188.
- Moret Y and Schmid-Hempel P (2000) Survival for immunity: the price of immune system activation for bumblebee workers. *Science (New York, NY)* **290**, 1166–1168.
- Müller U, McMahon DP and Rolff J (2019) Exposure of the wild bee *Osmia bicornis* to the honey bee pathogen *Nosema ceranae*. *Agricultural and Forest Entomology* **65**, 191–199.
- Näpflin K and Schmid-Hempel P (2018) High gut microbiota diversity provides lower resistance against infection by an intestinal parasite in bumblebees. *The American Naturalist* **192**, 131–141.
- Ngor L, Palmer-Young EC, Nevarez RB, Russell KA, Leger L, Giacomini SJ, Pinilla-Gallego MS, Irwin RE and McFrederick QS (2020) Cross-infectivity of honey and bumble bee-associated parasites across three bee families. *Parasitology* **147**, 1290–1304.

- Otterstatter MC and Thomson JD (2006) Within-host dynamics of an intestinal pathogen of bumble bees. *Parasitology* **133**, 749–761.
- Otterstatter MC and Thomson JD (2008) Does pathogen spillover from commercially reared bumble bees threaten wild pollinators? *PLoS ONE* **3**, 9.
- Palmer-Young EC and Thursfield L (2017) Pollen extracts and constituent sugars increase growth of a trypanosomatid parasite of bumble bees. *PeerJ* **5**, e3297.
- Pitts-Singer TL and Cane JH (2011) The alfalfa leafcutting bee, *Megachile rotundata*: the world's most intensively managed solitary bee. In Berenbaum MR, Carde RT and Robinson GE (eds), *Annual Review of Entomology*. Palo Alto, CA, USA: Annual Reviews, vol. **56**. pp. 221–237.
- Ravoet J, De Smet L, Meeus I, Smaghe G, Wenseleers T and de Graaf DC (2014) Widespread occurrence of honey bee pathogens in solitary bees. *Journal of Invertebrate Pathology* **122**, 55–58.
- R Development Core Team (2008) *R: A Language and Environment for Statistical Computing*. Vienna, Austria: R Foundation for Statistical Computing.
- Richardson LL, Adler LS, Leonard AS, Andicoechea J, Regan KH, Anthony WE, Manson JS and Irwin RE (2015) Secondary metabolites in floral nectar reduce parasite infections in bumblebees. *Proceedings of the Royal Society B-Biological Sciences* **282**, 8.
- Rinderer TE and Dell Elliott K (1977) Worker honey bee response to infection with *Nosema apis*: influence of diet. *Journal of Economic Entomology* **70**, 431–433.
- Roswell M, Dushoff J and Winfree R (2019) Male and female bees show large differences in floral preference. *PLoS ONE* **14**, e0214909.
- Ruiz-González MX and Brown MJ (2006) Honey bee and bumblebee trypanosomatids: specificity and potential for transmission. *Ecological Entomology* **31**, 616–622.
- Ruiz-González MX, Bryden J, Moret Y, Reber-Funk C, Schmid-Hempel P and Brown MJF (2012) Dynamic transmission, host quality, and population structure in a multi-host parasite of bumblebees. *Evolution* **66**, 3053–3066.
- Schmid-Hempel P (1998) *Parasites in Social Insects*. Princeton, New Jersey: Princeton University Press.
- Schmid-Hempel P and Schmid-Hempel R (1993) Transmission of a pathogen in *Bombus terrestris*, with a note on division of labour in social insects. *Behavioral Ecology and Sociobiology* **33**, 319–327.
- Schoonvaere K, Smaghe G, Francis F and de Graaf DC (2018) Study of the metatranscriptome of eight social and solitary wild bee species reveals novel viruses and bee parasites. *Frontiers in Microbiology* **9**, 177.
- Singh R, Levitt AL, Rajotte EG, Holmes EC, Ostiguy N, vanEngelsdorp D, Lipkin WI, dePamphilis CW, Toth AL and Cox-Foster DL (2010) RNA viruses in Hymenopteran pollinators: evidence of inter-taxa virus transmission via pollen and potential impact on non-*Apis* Hymenopteran species. *PLoS ONE* **5**, e14357.
- Streicker DG, Turmelle AS, Vonhof MJ, Kuzmin IV, McCracken GF and Rupprecht CE (2010) Host phylogeny constrains cross-species emergence and establishment of rabies virus in bats. *Science (New York, NY)* **329**, 676–679.
- Strobl V, Yañez O, Straub L, Albrecht M and Neumann P (2019) Trypanosomatid parasites infecting managed honeybees and wild solitary bees. *International Journal for Parasitology* **49**, 605–613.
- Therneau TM and Therneau MTM (2015) Package 'coxme'. *Mixed Effects Cox Models. R package version 2.2-16*.
- Tian T, Piot N, Meeus I and Smaghe G (2018) Infection with the multi-host micro-parasite *Apicystis bombi* (Apicomplexa: Neogregarinorida) decreases survival of the solitary bee *Osmia bicornis*. *Journal of Invertebrate Pathology* **158**, 43–45.
- Urbanowicz C, Baert N, Bluhner SE, Böröczky K, Ramos M and McArt SH (2019) Low maize pollen collection and low pesticide risk to honey bees in heterogeneous agricultural landscapes. *Apidologie* **50**, 379–390.
- Velthuis HHW and van Doorn A (2006) A century of advances in bumblebee domestication and the economic and environmental aspects of its commercialization for pollination. *Apidologie* **37**, 421–451.
- Winfree R, Reilly JR, Bartomeus I, Cariveau DP, Williams NM and Gibbs J (2018) Species turnover promotes the importance of bee diversity for crop pollination at regional scales. *Science (New York, NY)* **359**, 791–793.
- Zhang G, St. Clair AL, Dolezal A, Toth AL and O'Neal M (2020) Honey bee (Hymenoptera: Apidae) pollen forage in a highly cultivated agroecosystem: limited diet diversity and its relationship to virus resistance. *Journal of Economic Entomology* **113**, 1062–1072.
- Zheng H-Q, Lin Z-G, Huang S-K, Sohr A, Wu L and Chen YP (2014) Spore loads may not be used alone as a direct indicator of the severity of *Nosema ceranae* infection in honey bees *Apis mellifera* (Hymenoptera: Apidae). *Journal of Economic Entomology* **107**, 2037–2044.
- Zuur AF, Ieno EN, Walker NJ, Saveliev AA and Smith GM (2009) Zero-truncated and zero-inflated models for count data. In Gail M, Krickeberg K, Samet JM, Tsiatis A and Wong W (eds), *Mixed Effects Models and Extensions in Ecology with R*. New York, New York: Springer, pp. 261–293.