

Characterization of *Dermotheca* sp. Infection in a midwestern state-endangered salamander (*Ambystoma platineum*) and a co-occurring common species (*Ambystoma texanum*)

Research Article

Cite this article: Adamovicz L, Woodburn DB, Virrueta Herrera S, Low K, Phillips CA, Kuhns AR, Crawford JA, Allender MC (2020). Characterization of *Dermotheca* sp. Infection in a midwestern state-endangered salamander (*Ambystoma platineum*) and a co-occurring common species (*Ambystoma texanum*). *Parasitology* **147**, 360–370. <https://doi.org/10.1017/S0031182019001677>

Received: 28 July 2019

Revised: 11 November 2019

Accepted: 14 November 2019


First published online: 8 January 2020

Key words:

Ambystoma jeffersonianum complex; amphibian; emerging infectious disease; Ichthyosporea; Mesomycetozoea.

Author for correspondence:

Laura Adamovicz,
E-mail: adamovi2@illinois.edu

Laura Adamovicz¹ , Daniel B. Woodburn², Stephany Virrueta Herrera^{3,4}, Kelsey Low^{3,4}, Christopher A. Phillips⁴, Andrew R. Kuhns⁴, John A. Crawford⁵ and Matthew C. Allender¹

¹Wildlife Epidemiology Lab, University of Illinois, 2001 S Lincoln Ave, Urbana, IL 61802, USA; ²Zoological Pathology Program, College of Veterinary Medicine, University of Illinois, 3300 Golf Road, Brookfield, Illinois, USA; ³Program in Ecology, Evolution, and Conservation Biology, University of Illinois, 505 South Goodwin Ave, Urbana, IL 61801, USA; ⁴Illinois Natural History Survey, Prairie Research Institute, 615 E Peabody Dr, Champaign, IL 61820, USA and ⁵National Great Rivers Research and Education Center, One Confluence Way, East Alton, IL 62024, USA

Abstract

Ichthyosporean parasites (order Dermocystida) can cause morbidity and mortality in amphibians, but their ecology and epidemiology remain understudied. We investigated the prevalence, gross and histologic appearance, and molecular phylogeny of a novel dermocystid in the state-endangered silvery salamander (*Ambystoma platineum*) and the co-occurring, non-threatened small-mouthed salamander (*Ambystoma texanum*) from Illinois. Silvery salamanders ($N = 610$) were sampled at six ephemeral wetlands from 2016 to 2018. Beginning in 2017, 1–3 mm raised, white skin nodules were identified in 24 silvery salamanders and two small-mouthed salamanders from five wetlands (prevalence = 0–11.1%). Skin biopsy histology ($N = 4$) was consistent with dermocystid sporangia, and necropsies ($N = 3$) identified infrequent hepatic sporangia. Parasitic 18S rRNA sequences ($N = 5$) from both salamander species were identical, and phylogenetic analysis revealed a close relationship to *Dermotheca viridescens*. Dermocystids were not identified in museum specimens from the same wetlands ($N = 125$) dating back to 1973. This is the first report of *Dermotheca* sp. affecting caudates in the Midwestern United States. Future research is needed to determine the effects of this pathogen on individual and population health, and to assess whether this organism poses a threat to the conservation of ambystomatid salamanders.

Introduction

Global amphibian declines have been attributed to multiple factors including habitat loss and fragmentation, invasive species, pollution, climate change and disease (Kiesecker *et al.*, 2001; Daszak *et al.*, 2003; Blaustein *et al.*, 2012). Pathogens such as ranavirus and *Batrachochytrium dendrobatidis*, which have wide spatial distributions and contribute to mortality events that can affect population stability, are obvious targets for research (Skerratt *et al.*, 2007; Duffus and Cunningham, 2010; Price *et al.*, 2014). However, many other pathogens affect free-living amphibians and may have population-level effects which are poorly understood or masked by co-infection with more virulent agents (Pessier 2008). One example is the Ichthyosporea, a class of protists in the Opisthokonta. Within the Ichthyosporea (junior synonym Mesomycetozoea), the order Dermocystida consists entirely of pathogens affecting mammals, birds, fish and amphibians (Cavalier-Smith, 1998; Mendoza *et al.*, 2002).

Until recently, Dermocystid parasites infecting amphibians were classified into two genera, *Amphibiocystidium* and *Amphibiothecum* (Pascolini *et al.*, 2003; Feldman *et al.*, 2005; Pereira *et al.*, 2005). Currently, amphibian dermocystids are assigned to six genera (*Amphibiocystidium*, *Dermocystidium*, *Dermosporidium*, *Dermotheca*, *Sphaerothecum* and *Valentines*) to ensure monophyly within each genus and reflect the discovery of *Sphaerothecum caipira* infecting frogs (Borteiro *et al.*, 2018a). Representatives of these genera have been documented most frequently in European anurans and urodeles including common frogs (*Rana temporaria*), water frogs (*Pelophylax esculentus* complex), Italian stream frogs (*Rana italica*), common midwife toads (*Alytes obstetricans*), palmate newts (*Lissotriton helveticus*), common newts (*Lissotriton vulgaris*), marbled newts (*Triturus marmoratus*) and crested newts (*Triturus cristatus*) from Italy, France, Germany, Switzerland, Czechoslovakia, Scotland and the Netherlands (Pérez, 1907; Moral, 1913; Pérez, 1913; Granata, 1919; Guyénot and Naville, 1922; Remy, 1931; Poisson, 1937; Broz and Privora, 1952; Broz and Kulda, 1954; Pascolini *et al.*, 2003; Pereira *et al.*, 2005; Duffus and Cunningham, 2010; González-Hernández *et al.*, 2010; Courtois *et al.*, 2013; Stark and Guex, 2014; Federici *et al.*, 2015; Fiegna *et al.*, 2017; Fagotti *et al.*, 2019). North American dermocystids include *Dermocystidium penneri*, which affects American toads (*Anaxyrus americanus*), Fowler's toads (*Anaxyrus fowleri*), and Yosemite toads (*Anaxyrus*

canorus) (Jay and Pohley, 1981; Green and Sherman, 2001; Green *et al.*, 2002; Converse and Green, 2005), and *Dermotheca viridescens*, which affects eastern newts (*Notophthalmus viridescens*) from Pennsylvania, Massachusetts and West Virginia (Raffel *et al.*, 2008).

Amphibians infected with dermocyctids develop single or multiple cutaneous lesions, typically appearing as either 1–5 mm round, c-shaped, u-shaped, or dumbbell-shaped white nodules or 1–7 mm vesicular structures which may coalesce and ulcerate. Lesions have also been observed infrequently within gastrointestinal mucosa and internal organs (Raffel *et al.*, 2008; Fiegna *et al.*, 2017). Histologically, nodular foci consist of sporangia containing microscopic endospores (Mendoza *et al.*, 2002). Host response to the sporangia is typically minimal until they rupture, at which point a localized immune response can cause lesions to appear inflamed (Pérez, 1913; Broz and Kulda, 1954; Pascolini *et al.*, 2003; Raffel *et al.*, 2008; Fiegna *et al.*, 2017). Information on recovery period is limited, though anecdotal evidence suggests that the complete resolution of these lesions in mildly affected individuals is possible within a few weeks (Raffel *et al.*, 2008).

Infection with dermocyctid pathogens was initially believed to be fairly benign; however, heavily infected newts from France, Scotland, Germany and the USA have demonstrated more severe clinical signs and pathologic changes, including regional oedema, haemorrhages, extensive cutaneous ulceration, development of secondary bacterial infections and death (Moral, 1913; Gambier, 1924; González-Hernández *et al.*, 2010; Raffel *et al.*, 2008; Fiegna *et al.*, 2017). Amphibian skin plays a major role in osmoregulation, electrolyte balance and defence against pathogen invasion (Woodhams *et al.*, 2007; Rollins-Smith, 2009). Disruption of these vital functions in animals with heavy dermocyctid burdens may result in death due to fatal electrolyte disturbances, similar to chytridiomycosis (Campbell *et al.*, 2012). Indeed, an increased mortality rate was reported for captive *N. viridescens* infected with *D. viridescens*, compared to newts without visible lesions (Raffel *et al.*, 2008). Alternatively, development of opportunistic secondary infections may contribute to morbidity and mortality. Amphibian dermocyctid infections have also been linked to cutaneous microbiome changes favouring fungal colonization and dermal granular gland shrinkage, which may indicate the reduction of antimicrobial peptide secretion and subsequent immune dysfunction (Federici *et al.*, 2015; Fiegna *et al.*, 2017). Further study of dermocyctid parasites is needed to understand the threat that these organisms may pose to vulnerable amphibian populations. This study seeks to characterize dermocyctid infection in a North American salamander species of conservation concern.

The triploid, unisexual silvery salamander (*Ambystoma platineum*) occurs in the Northeastern and Midwestern United States and originated through hybridization of blue-spotted salamanders (*A. laterale*) and Jefferson salamanders (*A. jeffersonianum*), resulting in retention of two sets of *A. jeffersonianum* chromosomes and one set of *A. laterale* chromosomes (JLL; Uzzell, 1964; Uzzell and Goldblatt, 1967). Silvery salamanders typically coexist in habitats with one of their parental species and act as sexual parasites on the males of that species. However, some populations, including those in Illinois, do not occur with either parental species. In these populations, silvery salamanders utilize sperm from small-mouthed salamanders (*A. texanum*) for kleptogenic reproduction (Uzzell and Goldblatt, 1967; Morris and Brandon, 1984; Spolsky *et al.*, 1992; Phillips *et al.*, 1997; Teltser and Greenwald, 2015).

Silvery salamanders are found in only two counties in Illinois and are considered state-endangered (Illinois Endangered Species Protection Board, 2015). Threats to the persistence of silvery salamanders in Illinois are largely unknown, but an ongoing demographic and health assessment study has been initiated to

examine conservation challenges for this species. The primary objective of this study is to provide a morphologic, histologic and molecular description of dermocyctid infection in free-living silvery salamanders in Illinois, and secondarily, to describe the incidental detection of dermocyctids in sympatric small-mouthed salamanders.

Methods

Demographic survey

Silvery salamanders were evaluated at six ephemeral wetlands distributed among three study sites including Collison (wetland 71), Middle Fork (MF, wetlands 76 & 285) and Kickapoo State Park (KSP, wetlands 67, 282, and 283) in Vermilion County, Illinois. Exact wetland locations are not provided to protect the animals but are available upon request. In 2016, salamanders were captured using minnow traps and dip-nets with a standardized number of traps and dip-net sweeps used based on wetland size. In 2017 and 2018, study wetlands were encircled with drift-fences and associated pitfall traps to monitor salamander ingress and egress during breeding migrations. Traps were checked daily from February 1st to April 15th 2017 and 2018 and salamanders were toe-clipped using a wetland specific numbering system to estimate fence trespass rates.

Health assessments

Salamander health assessments were performed for 3–4 days each year during breeding migrations. Animals were measured (snout-vent length and total length; mm) and weighed (g) and a complete physical examination was performed by a veterinarian (LA) noting appearance of the eyes, nose, oral cavity, integument, musculoskeletal system and cloaca. During health assessments in 2017 white cutaneous nodules consistent with dermocyctid infection were observed on multiple animals. Digits with nodules from two different salamanders were opportunistically excised during routine toe-clipping and immediately placed in 100% ethanol. A single male small-mouthed salamander from wetland 67 was found with white cutaneous nodules on the trunk. The skin overlying the nodules was swabbed and the animal was released. In 2018 one heavily affected silvery salamander (21 nodules) and one heavily affected small-mouthed salamander (>50 nodules) from pond 283 were euthanized using an overdose of buffered MS-222 (3-Aminobenzoic Acid Ethyl Ester, Sigma Chemical Co. St. Louis, MO 61378). An additional heavily affected silvery salamander (>50 nodules) was found dead within a bucket trap at the same site. Two skin nodules from each animal were collected using a sterile scalpel blade and dry-frozen at –20°C. The bodies were then preserved in 10% buffered formalin and submitted for histopathology.

Pathologic evaluation

Fixed tissues collected during health assessments were processed routinely for histopathology and stained with hematoxylin and eosin. Slides were evaluated *via* light microscopy by an anatomic pathologist. Images were taken using an Olympus DP71 digital microscope camera.

DNA extraction, polymerase chain reaction (PCR) and sequencing

DNA was extracted from the skin swab and frozen and ethanol-preserved tissue samples using a commercial kit (DNeasy Blood & Tissue Kit, Qiagen, Valencia, CA, USA). Quality and purity

of resulting DNA samples was evaluated spectrophotometrically (NanoDrop 1000, Thermo Fisher Scientific, 260 Waltham, MA, USA). Two sets of existing consensus PCR primers targeting the 18S rRNA gene were used to screen tissue samples for the presence of dermocyctids (González-Hernández *et al.*, 2010; Fiegna *et al.*, 2017). Each reaction contained 1 × PCR buffer, 2.5 mM magnesium chloride, 0.2 mM dNTPs, 2.5 U Platinum Taq polymerase, 0.2 μM of each primer, 1 μL template DNA (<1 μg) and sterile deionized water to bring the final volume to 50 μL (Platinum Taq DNA Polymerase, Invitrogen, Carlsbad, CA 92008). Thermocycler settings consisted of a 5-minute initial denaturation at 95°C followed by 40 cycles of 95°C for 30 s, 57.7°C for 30 s and 72°C for 120 s, then a final elongation step of 72°C for 7 minutes. The optimal annealing temperature of 57.7°C was determined using a thermal gradient testing 10 temperatures from 53 to 65°C (data not shown). A positive control consisting of sequence-confirmed *D. penneri* from a Fowler's toad (*Anaxyrus fowleri*) and a non-template control (sterile water) were included in each run. Products were resolved on a 1% agarose gel and samples with appropriately sized bands (1400 bp) were treated with ExoSAP-IT (USB Corporation, 26111 miles Road, Cleveland, OH, USA) and commercially sequenced in both directions. Sequences were trimmed of primers and low-quality base-pair calls in Geneious v. 11.1.3 (Kearse *et al.*, 2012) and compared to known sequences in the NCBI GenBank database using BLASTN (Benson *et al.*, 2008).

Phylogenetic analysis

18s rRNA sequences that were 92–99% similar to the silvery and small-mouthed salamander dermocyctids were recovered from GenBank using BLASTN. Sequences were aligned using MAFFT v. 7.388 (Kato and Standley, 2013) with default parameters as implemented in Geneious. The ends of the resulting alignment file were manually adjusted and trimmed in Geneious. In order to estimate the best substitution model for phylogenetic analysis, PartitionFinder v. 2.1.1 (Lanfear *et al.*, 2017) was used to search through all 195 nucleotide models using the corrected Akaike Information Criterion (AICc; Sugiura 1978). Maximum likelihood phylogenetic reconstruction was performed in RAxML v.8.1.3 using a GTR GAMMA model with 100 bootstrap replicates (Stamatakis, 2014). Bootstrap support (BS) was then summarized on the best tree. Bayesian phylogenetic reconstruction was performed using MrBayes v. 3.2.6 (Ronquist *et al.*, 2012) with a GTR + I + G substitution model and 20 million generations of Markov Chain Monte Carlo (MCMC) models for three runs of four chains each, sampling every 1000 trees. Trace files were viewed in Tracer v. 1.6 (Rambaut and Drummond, 2007) to assess parameter convergence. To assess topological convergence, .t files were assessed using the R package RWTY v. 1.0.1 (Warren *et al.*, 2017). Based on these assessments, the first 10% (2000 trees) were discarded as burn-in.

Museum specimen survey

A total of 125 *A. platineum* specimens collected from Vermilion County, Illinois and maintained in the Illinois Natural History Survey (INHS) and University of Illinois Museum of Natural History (UIMNH) collections were visually examined for the presence of white skin nodules consistent with dermocyctid infection.

Statistical analysis

All statistical evaluations were performed in R v 3.5.1 (R Core Team 2018). Descriptive statistics for skin nodule prevalence

were tabulated by year and wetland. Ninety-five per cent confidence intervals for binomial probabilities were constructed around prevalence estimates using the binconf function in the Hmisc package. The effect of wetland ID and study site on the probability of dermocyctid detection was assessed in each year using generalized linear models, package brglm (Kosmidis 2017). Post-hoc between group differences were evaluated using the contrasts function in the lsmeans package (Lenth 2016).

Sample size calculations were performed to determine the number of salamander examinations needed to detect varying levels of dermocyctid prevalence (1%, 2%, 3%, 4%, 5%, 10%, 20%), at different salamander population sizes (50, 100, 200, 300, 500, 1000 and 5000). The specificity for visual detection of dermocyctid skin nodules was assumed to be one, while a conservative value of 0.8 was chosen for sensitivity, based on previous studies (Raffel *et al.*, 2008, González-Hernández *et al.*, 2010; Fiegna *et al.*, 2017). Herd sensitivity, or the probability of infected salamanders being detected at a wetland where dermocyctid infection is actually present, was set at 0.95. These calculations were performed using an online tool provided at epitools.ausvet.com.au, based on an approximation of the hypergeometric distribution modified to account for imperfect diagnostic test sensitivity (MacDiarmid 1988).

Results

Sample population

In total 36 silvery salamanders were examined on March 3–4th 2016, 390 individuals were assessed on February 24th, February 28th and March 7th, 2017 and 220 were examined on February 19–21st and 25th, 2018. The proportion of salamanders examined compared to the total number of unique captures per wetland each year ranged from 12 to 83%, reflecting somewhat uneven sampling effort for the health assessment component of this study (Table 1).

White cutaneous nodules consistent with dermocyctid infection were not observed in 2016. In 2017, the prevalence of nodules was higher at KSP than MF (OR = 6.65, 95% CI = 3.59–36.18, $P = 0.028$) and Collision (OR = 20, 95% CI = 1.2–340, $P = 0.039$). Prevalence was also higher at wetland 283 than wetlands 76 (OR = 6.95, 95% CI = 1.23–39.19, $P = 0.028$) and 71 (OR = 25.8, 95% CI = 1.45–448.8, $P = 0.026$). In 2018, the prevalence of nodules was not significantly different between sites or wetlands. Skin nodules consistent with dermocyctid infection were not detected in Collision (wetland 71) at any point during the study.

Sample size calculations indicate that the number of salamanders examined per wetland prior to 2017 was inadequate to detect dermocyctids at a prevalence of 20% or lower (Table 2). Using a total number of unique captures as a (limited) proxy for population size, sampling effort was also likely inadequate to detect dermocyctids at a prevalence of 20% or lower at wetland 285 in 2017 and wetlands 67 and 282 in 2018.

Gross and histologic lesion description

Skin nodules consistent with dermocyctid infection in silvery salamanders consisted of 1–3 mm white, raised, slightly firm subcutaneous nodules concentrated on the trunk (Figs 1 and 2). Individual nodules were round to dumbbell-shaped and occasionally curved or C-shaped. Most salamanders had fewer than 10 lesions. Hepatic nodules were identified in the most heavily affected silvery salamander at necropsy, but internal lesions were not identified in the other two necropsied animals.

On histologic examination, grossly apparent skin and hepatic nodules were consistent with ichthyosporean sporangia (Fig. 3).

Table 1. Sample size and *Dermostheca* sp. prevalence in adult silvery salamanders (*Ambystoma platineum*) surveyed in Vermilion County, Illinois in 2017 and 2018.

Wetland	Site	Year	Salamanders examined	Total unique captures	Proportion of captures examined (%)	<i>Dermostheca</i> sp. prevalence (%) (95% CI)
282	KSP	2017	30	36	83	3.3 (0.17–16.7)
		2018	19	68	28	0
283	KSP	2017	108	742	15	11.1 (6.5–18.4)
		2018	70	395	18	7.1 (3.1–15.7)
67	KSP	2017	53	75	71	7.5 (3–17.9)
		2018	15	127	12	0
71	Collison	2017	99	310	32	0
		2018	57	223	26	0
76	MF	2017	81	129	63	1.3 (0.06–6.7)
		2018	39	58	67	0
285	MF	2017	19	36	53	0
		2018	20	37	54	5 (0.26–23.6)

Table 2. Number of silvery salamander (*Ambystoma platineum*) examinations needed to detect *Dermostheca* sp. at different levels of prevalence and different salamander population sizes

Population Size	<i>Dermostheca</i> sp. Prevalence						
	1%	2%	3%	4%	5%	10%	20%
50	50	50	49	49	40	29	17
100	100	98	79	66	57	33	18
200	195	132	99	79	65	35	19
300	237	148	107	83	68	36	19
500	282	162	114	87	71	37	19
1000	324	174	119	91	73	37	19
5000	364	185	124	93	75	38	19

Calculations were performed using an approximation of the hypergeometric distribution modified to account for imperfect diagnostic test sensitivity. Sensitivity = 0.8, specificity = 1, herd sensitivity = 0.95.

**Fig. 1.** Gross appearance of skin lesions associated with *Dermostheca* sp. infection in free-living silvery salamanders (*Ambystoma platineum*) from Vermilion County, IL 2017–2018.

Sporangia were located within the superficial dermis and subcapsular hepatic parenchyma and mildly compressed surrounding tissue. Each sporangium ranged from 300 to 1000 μm in diameter and had a thin (5–10 μm) eosinophilic, hyalinized wall. Sporangia contained numerous 6–10 μm round endospores with 5 μm round eosinophilic cytoplasmic bodies, colorless cytoplasm and small round peripheral nuclei. Only minimal tissue reaction was observed in association with these sporangia and consisted primarily of mild hyperplasia (up to seven cell layers) in the overlying epidermis. Rare sporangia were segmentally ruptured with low numbers of free endospores and minimal heterophilic inflammation in the immediately adjacent dermis or hepatic parenchyma. Most observed sporangia were intact with no associated inflammation, though a few larger sporangia contained low numbers of central viable and necrotic heterophils. The sporangia capsules stained strongly with Gomori Methenamine-Silver Nitrate Stain (GMS), and the endospores had diffusely strong reactions with Periodic acid–Schiff (PAS) stain.

Molecular & phylogenetic analysis

An approximately 1350 bp product was amplified from the skin swab and two digits collected in 2017 and the euthanized/




Fig. 2. Intensity (A) and distribution (B) of *Dermotheca* sp. lesions in 24 affected adult silvery salamanders (*Ambystoma platineum*) from Vermilion County, IL 2017–2018.




Fig. 3. Skin from *Ambystoma platineum*. Within the superficial dermis is a single ichthyosporan sporangium that elevates the epidermis and slightly compresses surrounding tissues (A, hematoxylin and eosin stain, 100X). The sporangium has a thin eosinophilic hyalinized wall and contains numerous 6–10 μ m round endospores (B, hematoxylin and eosin stain, 600X).

deceased silvery and small-mouthed salamanders collected in 2018 using both sets of primers. The aligned portions of all sequences were identical, and representative data were deposited in GenBank (Accession numbers MK330881.1, MK757499.1, MK757500.1, MK757501.1). Sequences from the present study were 99% similar to *D. viridescens* and a *Dermotheca* sp. infecting Italian stream frogs in Europe (Accession # EU650666).

After the 10% burn-in, all parameters and topologies from the Bayesian phylogenetic analysis had ESS values $\gg 200$ or average standard deviation of split frequencies < 0.01 , indicating the MCMC runs had converged to stationarity. BS values in the maximum likelihood analysis averaged 67%, and posterior probability (PP) support averaged 97% in the Bayesian analysis. The maximum likelihood and Bayesian phylogenetic analyses produced similar trees, recovering *Sphaerothecum* as sister to *Rhinosporidium*, *Valentines*, *Dermocystidium*, *Amphibiocystidium* and *Dermotheca* with $> 66\%$ BS and $> 97\%$ PP (Figs 4 and 5). The silvery and small-mouthed salamander sequences were recovered in a monophyletic group with *Dermotheca* spp. (70 BS/0.99 PP).

Historic occurrence

A total of 125 *A. platineum* specimens collected in Vermilion county, Illinois from 1973 to 2018 were visually inspected for the presence of white nodules consistent with dermocystid infection. Sampling effort during this time period was haphazard, with the most specimens collected in the early 1980s (18 salamanders in 1980, 32 in 1981). Following this peak, sample sizes only exceeded 10 individuals per year in 2000 and 2018. Cutaneous nodules were not identified in any specimen. Tissue sampling for molecular screening was not pursued due to initial fixation of specimens in formalin prior to long-term storage in ethanol.

Discussion

This is the first report of *Dermotheca* sp. in the Midwestern United States and the first detection of this organism in

ambystomatid salamanders. The gross appearance of *Dermotheca* sp. is consistent with *D. viridescens* and slightly different from all other dermocystid species (Raffel et al., 2008). Lesions in *Amphibiocystidium ranae* infections are similar in size and color, but mature into more dramatic U-shaped sporangia (Guyénot and Naville, 1922; Remy, 1931; Broz and Privora, 1952; Broz and Kulda, 1954; Pascolini et al., 2003; Pereira et al., 2005). *Dermocystidium pusula* lesions start out round and similarly sized (1–3 mm), but have varying coloration and can progress to much larger (5–8 mm) cysts with 1–4 openings prior to ulceration. These lesions are also occasionally hemorrhagic and may induce localized or systemic oedema, which has not been described for other dermocystid infections (Pérez, 1907, 1913; Moral, 1913; Granata, 1919; González-Hernández et al., 2010; Fiegna et al., 2017). *Dermocystidium penneri* lesions are large (4–6 mm) and range in shape from round to oval to serpiginous tracts of divided chambers (Jay and Pohley, 1981; Feldman et al., 2005). *Sphaerothecum caipira*, *Dermosporidium hylarum* and *Valentines rwandae* all produce variably colored, 0.5–1 mm round lesions with or without ulceration (Carini, 1940; Borteiro et al., 2014; Scheid et al., 2015; Borteiro et al., 2018a). Elongated, curved and dumbbell-shaped lesions have not been reported for these three dermocystid species.

Cutaneous *Dermotheca* sp. nodules in silvery salamanders are distributed across the body in a similar way to most other dermocystids. The dorsal trunk is most commonly affected by *S. caipira*, *V. rwandae*, and *D. pusula*, and the ventral trunk by *A. ranae*, *D. viridescens* and *D. penneri* infections (Moral, 1913; Broz and Privora, 1952; Jay and Pohley, 1981; Pascolini et al., 2003; Feldman et al., 2005; Pereira et al., 2005; Raffel et al., 2008; González-Hernández et al., 2010; Scheid et al., 2015; Fiegna et al., 2017; Borteiro et al., 2018a). The head is a common site for lesion development in *S. caipira* and *V. rwandae* infections, the limbs are frequently affected by *A. ranae* and *D. pusula*, and the tail often hosts nodules in *D. viridescens* and *D. pusula* infections (Moral 1913; Guyénot and Naville, 1922; Raffel et al., 2008; González-Hernández et al., 2010; Fiegna et al., 2017;




Fig. 4. Phylogenetic tree for ichthyosporean organisms produced by maximum-likelihood analytic methods.




Fig. 5. Phylogenetic tree for ichthyosporean organisms produced by Bayesian analytic methods.

Borteiro et al., 2018a). Internal dermocystid infections have also been described, with *D. viridescens* sporangia identified in the liver, *V. rwandae* sporangia in cardiac muscle, *Dermostheca* sp. sporangia in palpebral conjunctivae and oral mucosa, and *D. pusula* sporangia in muscle tissue, oral mucosa, liver and cloacal stroma (Pérez 1907; Gambier, 1924; Raffel et al., 2008; Fiegna et al., 2017; Borteiro et al., 2018a; Fagotti et al., 2019). Systemic dissemination of *D. penneri* spores to the liver, kidneys and fat pads has also been identified in a Yosemite toad (Green and Sherman, 2001). While hepatic sporangia were detected in a single silvery salamander with heavy cutaneous dermocystid infection, the frequency and intensity of internal and disseminated infection in ambystomatid salamanders cannot be assessed without further necropsy surveillance.

Dermostheca sp. sporangia from the present study were similar in size and structure compared to other dermocystid species, with the possible exception of *A. ranae*, *D. pusula*, *D. penneri* and (in a single instance) *D. viridescens*. Sporangia in these species contain internal septations dividing endospores into chambers – apparently based on spore developmental stage; this was not observed in the *Dermostheca* sp. from silvery salamanders (Granata 1919; Guyénot and Naville, 1922; Broz and Privora, 1952; Broz and Kulda, 1954; Jay and Pohley, 1981; Raffel et al., 2008; González-Hernández et al., 2010; Fiegna et al., 2017; Fagotti et al., 2019). *Dermostheca* sp. endospores share similar histologic features with other dermocystids except the endospores of *D. viridescens*, *D. hylarum*, *A. ranae*, *D. penneri* and *D. pusula*, which contain variable numbers of cytoplasmic granules in addition to the larger cytoplasmic inclusion body (Carini, 1940; Broz and Privora, 1952; Broz and Kulda, 1954; Jay and Pohley, 1981; Green and Sherman, 2001; Pascolini et al., 2003; Feldman et al., 2005; Raffel et al., 2008; González-Hernández et al., 2010; Fiegna et al., 2017). Sporangia in all dermocystid species, including the *Dermostheca* sp. of the present study, are associated with mild epidermal hyperplasia. Most sporangia induced a minimal host inflammatory response that became more intense around ruptured or degenerating structures, similar to previous reports (Granata, 1919; Pérez, 1913; Pascolini et al., 2003; Pereira et al., 2005; Fiegna et al., 2017; Fagotti et al., 2019).

The gross appearance, lesion distribution and histologic characteristics of *Dermostheca* sp. infection in ambystomatid salamanders from Illinois are generally consistent with those of other dermocystid parasites. However, these characteristics alone are inadequate for species identification within this group, and more recent classifications have relied upon molecular phylogenetics instead (Pascolini et al., 2003; Feldman et al., 2005; Pereira et al., 2005; Raffel et al., 2008; González-Hernández et al., 2010; Scheid et al., 2015; Fiegna et al., 2017; Borteiro et al., 2018a). These analyses have resulted in multiple species reclassifications and their placement within several new genera, with the most recent report proposing a major taxonomic restructuring to ensure monophyly within each genus (Borteiro et al., 2018a). While molecular phylogenetics has improved dermocystid taxonomic resolution compared to morphology alone, limitations persist due to the restricted number and size of available sequences and the reliance upon one gene target (18S) for phylogenetic assessment (Feldman et al., 2005). These obstacles may be overcome with the publication of more 18S sequences and/or the analysis of sequence data from additional gene targets. However, until additional data is available, interpretation of dermocystid taxonomy should be performed cautiously.

With these caveats in mind, the phylogenetic analysis conducted in the present study recovered similar dendrogram topology to that of Borteiro et al. (2018a). Clades including amphibian *Dermostheca* spp., *Amphibiocystidium ranae*, *Rhinosporidium* spp., *Valentines rwandae* and *Sphaerothecum*

spp. were reliably recovered with high support (>85% BS, >96% PP), however, monophyly was not fully consistent between analyses. Unlike the analysis of Borteiro et al. (2018a), the present study recovered *Dermocystidium pusula* and *Dermocystidium penneri* in separate, well-supported clades (100% BS, 100% PP) in every dendrogram. This may indicate that these parasites belong in different genera; however, sequencing of additional gene targets would likely be necessary to improve phylogenetic resolution and confidently settle this issue. Similarly, while branch support was high for amphibian *Dermostheca* spp. and *Dermostheca percae*, resolution was poorer for other *Dermostheca* species infecting fish, including *D. fenicum*, *D. salmonis* and *D. borealis*. This illustrates the phylogenetic uncertainty within the order Dermocystida and highlights the need for additional evolutionary history research within this taxon.

The dermocystids infecting silvery salamanders and small-mouthed salamanders from Illinois were recovered in a monophyletic clade with *D. viridescens* and a currently un-named *Dermostheca* sp. from Italy with high BS and PP. The sequence variation within amphibian *Dermostheca* spp. is similar to or lower than that of *R. seeberi* and *D. percae*, potentially indicating that all amphibian *Dermostheca* spp. should be classified as *D. viridescens*. We did not assign our organism a specific epithet due to the aforementioned phylogenetic uncertainty associated with single-gene analyses, however, future multi-gene sequencing may confirm that the ambystomatid *Dermostheca* sp. from Illinois is actually *D. viridescens*. *Dermostheca viridescens* was originally identified in eastern newts from Pennsylvania, Massachusetts and West Virginia (Raffel et al., 2008). Eastern newts occur in northern Illinois but are not present at the wetlands sampled in the present study. In fact, the closest known eastern newt population is approximately 64 km away in Montgomery County, Indiana (Minton 2001). Movement of this parasite across large distances may have been facilitated by human transport of infected amphibians, as has been previously documented for ranavirus and chytridiomycosis (Jancovich et al., 2005; Picco et al., 2007; Picco and Collins, 2008; Farrer et al., 2011; Schloegel et al., 2006). Alternatively, infection with *Dermostheca* sp. may be more widespread than currently recognized and could be naturally present in understudied amphibian populations linking affected areas. Broader surveillance efforts will be necessary to define the current spatial distribution of *Dermostheca* sp. and to characterize the host range of this poorly understood parasite.

The origin of *Dermostheca* sp. in ambystomatid salamanders from Illinois is unclear, and so is the length of time this parasite has been present in affected populations. Sampling effort for focused health assessments prior to 2017 was insufficient to detect *Dermostheca* sp. at a wetland prevalence of 10% or lower. Observed *Dermostheca* sp. prevalence in 2017 and 2018 ranged from 1 to 11%, indicating that previous health surveys could easily have missed the parasite if it was present at similarly low levels. Salamander ecology studies at the same wetlands occasionally had large enough sample sizes to detect *Dermostheca* sp., but the primary focus of these studies was not physical examination and cutaneous nodules may have been present but not detected (Phillips et al., 1997; Phillips et al., 2002; McAllister et al., 2010). While it may be tempting to assume that *Dermostheca* sp. was suddenly introduced into affected salamander populations in 2017, it appears equally possible that this is an endemic parasite normally occurring at a low prevalence in ambystomatids. Ultimately, freedom from *Dermostheca* sp. infection prior to 2017 cannot be confidently assumed due to a lack of historic parasite surveillance data.

This limitation is not confined to the present study. Comprehensive dermocystid surveillance is rarely undertaken in

wild amphibian populations and many existing reports depend on opportunistic observations of infected animals. While robust epidemiologic data for amphibian dermocyctids is limited, a few consistent trends emerge when all available literature is considered together. Several studies demonstrate that dermocyctid prevalence is highest in cooler seasons including late fall, winter and early spring (Pérez, 1907; Pérez, 1913; Guyénot and Naville, 1922; Remy, 1931; Poisson, 1937; Broz and Privora, 1952; Jay and Pohley, 1981; Green *et al.*, 2002; Raffel *et al.*, 2008; Borteiro *et al.*, 2018b). This could be due to temperature-dependent alterations in host immunity, seasonal behavioral changes facilitating infection and/or cooler temperatures promoting parasite replication, as documented for *B. dendrobatidis* (Berger 2004; Kriger and Hero, 2007; Rowley and Alford, 2007). Salamanders were only sampled during February and March in the present study, potentially pairing search effort with periods of maximal dermocyctid prevalence and increasing the likelihood of detecting infections. Dermocyctid prevalence was not investigated during other time points because ambystomatids are fossorial and impractical to locate in large numbers outside of the breeding season. While this will likely remain a significant limitation to the study of dermocyctids in ambystomatids, more thorough characterization of seasonal prevalence in other species may provide insights into the factors driving lesion development in amphibians.

Dermocyctid prevalence can vary between wetlands and years, indicating that host and environmental factors likely influence lesion development (Broz and Privora, 1952; Raffel *et al.*, 2008; Courtois *et al.*, 2013; Fiegna *et al.*, 2017; Borteiro *et al.*, 2018b; Fagotti *et al.*, 2019). Furthermore, dermocyctid prevalence can vary between sympatric species examined within a single sampling period. Scheid *et al.* (2015) examined 16 anuran species within the same swamp and detected *V. rwandae* in only two species, one of which (*Hyperolius lateralis*) had a much higher dermocyctid prevalence (18.2%) than the other (*H. viridiflavus*, 0.7%). Prevalence of *A. ranae* was significantly higher in a population of declining *R. lessonae* (45.5–52.4%) than a stable population of *R. esculenta* (14.3–17.2%) from the same wetland complex in Italy (Pascolini *et al.*, 2003). Borteiro *et al.* (2018b) identified *D. hylarum* in *Scinax fuscovarius*, but not in any other amphibian species examined; Jay and Pohley (1981) found *D. penneri* in *A. americanus*, but not in six other species from the same study site; and Broz and Privora (1952) identified *A. ranae* in *R. temporaria*, but not *R. esculenta* or any other species occupying the same habitat. The present study documented *Dermostheca* sp. infection in silvery salamanders and small-mouthed salamanders, but the infection was not observed in several other sympatric amphibian species including two ambystomatid salamanders (*Ambystoma maculatum* and *Ambystoma opacum*) and several anurans. This may suggest a difference in exposure or susceptibility between species, underscoring the need for additional research documenting dermocyctid host range, life history characteristics, modes of transmission and pathogenesis.

Common methods for diagnosing amphibian dermocyctid infections include visualization of grossly consistent cutaneous nodules, microscopic examination of tissue sections *via* histology or transmission electron microscopy and PCR performed on tissue samples. Many of these methods require biopsy or sacrifice of affected animals, making visual examination an attractive option. Several studies rely on visual detection of dermocyctids exclusively, however, few have critically evaluated the sensitivity and specificity of this method (Courtois *et al.*, 2013; Diego-Rasilla 2017; Galán and Dopereiro, 2017; Borteiro *et al.*, 2018b). Raffel *et al.* (2008) dissected 106 eastern newts and found that 12 newts with visible cutaneous *D. viridescens* lesions had consistent sporangia in the skin on histologic examination. Four of these

animals also had sporangia in the liver, while three newts had hepatic sporangia with no evidence of cutaneous disease. Overall sensitivity for visual examination in this report was 0.8 (95% CI: 0.51–0.95) and specificity was 1 (95% CI: 0.95–1). González-Hernández *et al.* (2010) dissected 21 palmate newts and detected *D. pusula* DNA in skin biopsies from all animals with visibly apparent cutaneous nodules ($N=10$) and one newt which had histologic skin changes consistent with recent dermocyctid infection, but no cutaneous nodules. In this case, sensitivity of visual examination was 0.91 (95% CI: 0.57–1) and specificity was 1 (95% CI: 0.66–1). Fiegna *et al.* (2017) dissected 30 palmate newts and found *D. pusula* sporangia in 18 animals with and two animals without visually apparent cutaneous lesions. Sensitivity for visual examination in this report was 0.9 (95% CI: 0.67–0.98) and specificity was 1 (95% CI: 0.66–1). Amphibian dermocyctid infections produce very characteristic cutaneous lesions, and visual identification of these lesions has high specificity for dermocyctid detection. However, animals with internal dermocyctid infections and those with early or resolving lesions may be misclassified as dermocyctid negative using visual examination alone. The prevalence estimates obtained in the present study relied heavily upon visual examination and may be underestimates of true *Dermostheca* sp. prevalence in silvery salamanders.

PCR screening of skin swabs is one non-invasive diagnostic option which may reduce false-negative rates in dermocyctid surveys. González-Hernández *et al.* (2010) found that skin swabs from 24 newts with cutaneous dermocyctid lesions were PCR positive for *D. pusula*, while swabs from 16 animals with no gross lesions were PCR negative. Additional studies are needed to determine the sensitivity and specificity of this technique, however, this may provide a practical means of dermocyctid detection to augment existing methodologies. Non-invasive antemortem diagnostics are especially useful for threatened and endangered species, where invasive testing is not feasible or ethical in large numbers of animals.

From a disease ecology perspective, ichthyosporeans share a suite of biological characteristics with parasites capable of causing host extinction, including low host specificity, the potential for high virulence, presence of a free-living infectious stage and the ability to survive prolonged adverse environmental conditions (Mendoza *et al.*, 2002; Fisher *et al.*, 2012; Rowley *et al.*, 2013). Unfortunately, studies documenting the population level impacts of dermocyctid infections are extremely limited (González-Hernández *et al.*, 2010; Raffel *et al.*, 2008; Fiegna *et al.*, 2017). As a result, the real-world importance of these parasites for amphibian conservation is hard to determine, and likely context-dependent. Long-term research on the *Pelophylax esculentus* complex in central Italy documented steep population declines in *P. lessonae* with severe *A. ranae* infections, while sympatric *P. esculentus* with lower dermocyctid burdens remained stable (Simoncelli *et al.*, 2005; Fagotti *et al.*, 2005; Di Rosa *et al.*, 2007). Extreme weather events, elevated tissue organochlorine levels and chytridiomycosis were also identified during these declines, and it is difficult to disentangle the complex interaction between disease and other anthropogenic and climactic stressors in this case (Simoncelli *et al.*, 2005; Fagotti *et al.*, 2005; Di Rosa *et al.*, 2007). If *A. ranae* was not the proximate cause of the *P. lessonae* declines, then monitoring dermocyctid prevalence may provide information on the overall wellness of infected populations, rising as health status deteriorates secondary to multiple stressors. However, a 7-year study in Italian stream frogs failed to demonstrate differences in weight, survival probability, or capture probability between *Dermostheca* sp. infected and uninfected frogs, despite high parasite prevalence (>50% infected) (Fagotti *et al.*, 2019). This outcome disparity demonstrates the importance of characterizing natural dermocyctid infection dynamics and

identifying pathogen, host and environmental factors promoting severe disease sequelae.

In the present study, salamander health status did not differ based on the presence of *Dermostheca* sp. infection, however, all animals were released immediately after physical examination and long-term survival and fitness consequences are unknown. Furthermore, only animals healthy enough to attempt breeding were evaluated, and it is possible that more heavily affected salamanders remained in burrows and were not examined. The cross-sectional nature of this study and the possibility of sampling bias ultimately makes it difficult to infer the effects of *Dermostheca* sp. on salamander health. The importance of other amphibian species for the maintenance of *Dermostheca* sp. in Vermilion County wetlands is also unknown. Small-mouthed salamanders are common at all study sites and could potentially serve as a reservoir for *Dermostheca* sp. infection in silvery salamanders. This is potentially concerning, as the presence of an abundant reservoir host can promote high levels of infection in rarer species, releasing pathogens from density dependence and contributing to differentially large disease impacts (McCallum and Dobson, 1995). This has been demonstrated for dermocystids infecting fish (Gozlan et al., 2005; Hershberger et al., 2010; Andreou et al., 2012). Future research should incorporate dermocystid surveillance in sympatric species to determine relative host susceptibility and assess the potential for reservoir host development. Longitudinal evaluation of infected salamanders in laboratory or mesocosm settings will be necessary to determine the impacts of *Dermostheca* sp. on individual health and survival. Continued population monitoring is needed to evaluate whether dermocystid parasites threaten silvery salamander conservation efforts.

Emerging infectious diseases can pose a significant threat to amphibians, especially species of conservation concern. Dermocystid parasites can affect individual survival and have the potential to impact population stability alone or in conjunction with other stressors. The identification of *Dermostheca* sp. in a state-endangered salamander is concerning and further robust studies are needed to determine its effects on individual and population health. The present study provides useful baseline information on morphology, pathology, phylogeny and prevalence that can be used to direct future research efforts and to monitor changes in the affected populations over time. Understanding the health of free-living amphibians may promote more effective management strategies and ultimately improve conservation outcomes.

Acknowledgements. We thank Andrew D. Sweet, PhD for guidance in phylogenetic analyses. We thank Connor T. Dempsey, Jaclyn M. Adams, Rachel Neubauer and Victoria Alessa Laserna for their assistance with collecting and sampling salamanders. We thank Jason P. Ross, John Warner, Allison Sieia, Eleanor Moen and Lindsay Hsieh for constructing the drift fences around each wetland.

Financial support. This work was supported by Illinois State Wildlife Grants (grant numbers T-104-R-1 and T-108-R-1).

Ethical standards. All animal handling and procedures were approved by the University of Illinois Institutional Animal Care and Use Committee (protocol #15180) and Illinois Endangered and Threatened Species Permits 05-11S and 1042.

Conflicts of interest. None

References

- Andreou D, Arkush KD, Guégan J-F and Gozlan RE (2012) Introduced pathogens and native freshwater biodiversity: a case study of *Sphaerothecum destruens*. *PLoS ONE* 7, e36998.
- Benson DA, Karsch-Mizrachi I, Lipman DJ, Ostell J and Wheeler DL (2008) Genbank. *Nucleic Acids Research* 36, D25–D30.

- Berger L, Speare R, Hines HB, Marantelli G, Hyatt AD, McDonald KR, Skerratt LF, Olsen V, Clarke JM, Gillespie G, Mahony M, Sheppard N, Williams C and Tyler MJ (2004) Effect of season and temperature on mortality in amphibians due to chytridiomycosis. *Australian Veterinary Journal* 82, 434–439.
- Blaustein AR, Gervasi SS, Johnson PTJ, Hoverman JT, Belden LK, Bradley PW and Xie GY (2012) Ecophysiology meets conservation: understanding the role of disease in amphibian population declines. *Philosophical Transactions of the Royal Society B: Biological Sciences* 367, 1688–1707.
- Borteiro C, Cruz JC, Kolenc F, Verdes JM, Moraña A, Martínez Debat C, Kun A, Ubilla M and Okada K (2014) Dermocystid-chytrid coinfection in the Neotropical frog *Hypsiboas pulchellus* (Anura: Hylidae). *Journal of Wildlife Diseases* 50, 150–153.
- Borteiro C, Baldo D, Maronna MM, BaÊta D, Sabbag AF, Kolenc F, Debat CM, Haddad CFB, Cruz JC, Verdes JM and Ubilla M (2018a) Amphibian parasites of the order *Dermocystida* (Ichthyosporae): current knowledge, taxonomic review and new records from Brazil. *Zootaxa* 4461, 499–518.
- Borteiro C, Gobel N, Kolenc F, Laufer G, Martínez Debat C and Ubilla M (2018b) Skin-mates or neighbors? A seasonal study of amphibian chytrid and dermocystid infection in *Boana pulchella* (Anura: Hylidae). *Cuadernos de Herpetología* 32, 101–108.
- Broz O and Kulda J (1954) *Dermosporidium Multigranulare* n.sp parasite z kuze *Rana esculenta*. *Acta Societatis Zoologicae Bohemicae* 18, 91–97.
- Broz O and Privora M (1952) Two skin parasites of *Rana Temporaria*: *Dermocystidium ranae* Guénot et Naville and *Dermosporidium granulatum* n. sp. *Parasitology* 42, 65–69.
- Campbell CR, Voyles J, Cook DI and Dinudom A (2012) Frog skin epithelium: electrolyte transport and chytridiomycosis. *International Journal of Biochemistry and Cell Biology* 44, 431–434.
- Carini A (1940) Sobre um parasito semelhante a um “*Rhinosporidium*”, encontrado em quistos da pele de uma “Hyla”. *Arquivos do Instituto Biológico (São Paulo)* 11, 93–98.
- Cavalier-Smith T (1998) Neomonada and the origin of animals and fungi. In Coombs GH, Vickerman K, Sleigh MA and Warren A (eds), *Evolutionary Relationships Among Protozoa*. Norwell, Massachusetts, USA: Kluwer Academic Publishers, pp. 375–407.
- Converse KA and Green DE (2005) Diseases of frogs and toads. In Majumdar SK, Huffman JE, Brenner FJ and Panah AI (eds), *Wildlife Diseases: Landscape Epidemiology, Spatial Distribution and Utilization of Remote Sensing Technology*. Easton, PA: The Pennsylvania Academy of Science, pp. 89–117.
- Courtois EA, Cornuau JH, Loyau A and Schmeller DS (2013) Distribution of *Amphibiocystidium* sp. in palmate newts (*Lissotriton helveticus*) in Ariège, France. *Herpetology Notes* 6, 539–543.
- Daszak P, Cunningham AA and Hyatt AD (2003) Infectious disease and amphibian population declines. *Diversity and Distributions* 9, 141–150.
- Diego-Rasilla FJ (2017) Parasitic dermic cyst in *Mesotriton alpestris* (caudata: salamandridae). *Boletín de la Asociación Herpetológica Española* 28, 56–57.
- Di Rosa I, Simoncelli F, Fagotti A and Pascolini R (2007) Ecology: the proximate cause of frog declines? *Nature* 447, E4–E5.
- Duffus ALJ and Cunningham AA (2010) Major disease threats to European amphibians. *Herpetological Journal* 20, 117–127.
- Fagotti A, Morosi L, Di Rosa I, Clarioni R, Simoncelli F, Pascolini R, Pellegrino R, Guex GD and Hotz H (2005) Bioaccumulation of organochlorine pesticides in frogs of the *Rana esculenta* complex in central Italy. *Amphibia-Reptilia* 26, 93–104.
- Fagotti A, Rossi R, Canestrelli D, La Porta G, Paracucchi R, Lucentini L, Simoncelli F and Di Rosa I (2019) Longitudinal study of *Amphibiocystidium* sp. Infection in a natural population of the Italian stream frog (*Rana italica*). *Parasitology* 28, 1–8.
- Farrer RA, Weinert LA, Bielby J, Garner TW, Balloux F, Clare F, Bosch J, Cunningham AA, Weldon C, du Preez LH, Anderson L, Pond SL, Shahar-Golan R, Henk DA and Fisher MC (2011) Multiple emergences of genetically diverse amphibian-infecting chytrids include a globalized hypervirulent recombinant lineage. *Proceedings of the National Academy of Sciences USA* 108, 18732–18736.
- Federici E, Rossi R, Fidati L, Paracucchi R, Scargetta S, Montalbani E, Franzetti A, La Porta G, Fagotti A, Simoncelli F, Cenci G and Di Rosa I (2015) Characterization of the skin microbiota in Italian stream frogs (*Rana italica*) infected and uninfected by a cutaneous parasitic disease. *Microbes and Environments* 30, 262–269.

- Feldman SH, Wimsatt JH and Green DE (2005) Phylogenetic classification of the frog pathogen *Amphibiothecum* (Dermosporidium) penneri based on small ribosomal subunit sequencing. *Journal of Wildlife Disease* **41**, 701–706.
- Fiegna C, Clarke CL, Shaw DJ, Bailly JL, Clare FC, Gray A, Garner TW and Meredith AL (2017) Pathological and phylogenetic characterization of *Amphibiothecum* sp. Infection in an isolated amphibian (*Lissotriton helveticus*) population on the island of Rum (Scotland). *Parasitology* **144**, 484–496.
- Fisher MC, Henk DA, Briggs CJ, Brownstein JS, Madoff LC, McCraw SL and Gurr SJ (2012) Emerging fungal threats to animal, plant and ecosystem health. *Nature* **484**, 186–194.
- Galán P and Dopereiro D (2017) Infección por dermocistidios (*Dermocystida*) en una población de *Lissotriton helveticus* de A Limia (Ourense, Galicia). *Boletín de la Asociación Herpetológica Española* **28**, 74–77.
- Gambier H (1924) Sur un Protiste parasite et pathogene des Tritons: *Hepatosphera molgarum* n. g., n. sp. *Comptes rendus des séances de la Société de biologie et de ses filiales* **90**, 439–441.
- González-Hernández M, Denoël M, Duffus AJL, Garner TWJ, Cunningham AA and Acevedo-Whitehouse K (2010) Dermocystid infection and associated skin lesions in free-living palmate newts (*Lissotriton helveticus*) from southern France. *Parasitology International* **59**, 44–350.
- Gozlan RE, St-Hilaire S, Feist SW, Martin P and Kent ML (2005) Biodiversity: disease threat to European fish. *Nature* **435**, 1046.
- Granata L (1919) *Dermomycooides beccarii* n. g. n. sp. nuovo enigmatico parassita di *Molge vulgaris*. *Monitore zoologico italiano* **20**, 153–160.
- Green DE and Sherman CK (2001) Diagnostic histological findings in Yosemite toads (*Bufo canorus*) from a die-off in the 1970s. *Journal of Herpetology* **35**, 92–103.
- Green DE, Converse KA and Schrader AK (2002) Epizootiology of sixty-four amphibian morbidity and mortality events in the USA, 1996–2001. *Annals of the New York Academy of Sciences* **969**, 323–339.
- Guyénot E and Naville A (1922) Un nouveau protiste du genre *Dermocystidium* parasite de la Grenouille *Dermocystidium ranae* nov. spec. *Revue Suisse de Zoologie* **29**, 133–145.
- Hershberger PK, van der Leeuw BK, Gregg JL, Grady CA, Lujan KM, Gutenberger SK, Purcell MK, Woodson JC, Winton JR and Parsley MJ (2010) Amplification and transport of an endemic fish disease by an introduced species. *Biological Invasions* **12**, 3665–3675.
- Illinois Endangered Species Protection Board (2015) Checklist of Endangered and Threatened Animals and Plants of Illinois. Retrieved from the Illinois Department of Natural Resources website: https://www.dnr.illinois.gov/ESPB/Documents/2015_ChecklistFINAL_for_webpage_051915.pdf (Accessed 15 September 2018).
- Jancovich JK, Davidson EW, Parameswaran N, Mao J, Chinchar VG, Collins JP, Jacobs BL and Storfer A (2005) Evidence for emergence of an amphibian iridoviral disease because of human-enhanced spread. *Molecular Ecology* **14**, 213–224.
- Jay JM and Pohley WJ (1981) *Dermosporidium penneri* sp n from the skin of the American toad, *Bufo americanus* (Amphibia, Bufonidae). *Journal of Parasitology* **67**, 108–110.
- Katoh K and Standley DM (2013) MAFFT Multiple sequence alignment software version 7: improvements in performance and usability. *Molecular Biology and Evolution* **30**, 772–780.
- Kearse M, Moir R, Wilson A, Stones-Havas S, Cheung M, Sturrock S, Buxton S, Cooper A, Markowitz S, Duran C, Thierer T, Ashton B, Mentjies P and Drummond A (2012) Geneious basic: an integrated and extendable desktop software platform for the organization and analysis of sequence data. *Bioinformatics (Oxford, England)* **28**, 1647–1649.
- Kiesecker JM, Blaustein AR and Belden LK (2001) Complex causes of amphibian population declines. *Nature* **410**, 681–684.
- Kosmidis I (2017) brglm: Bias reduction in binary-response generalized linear models. R package version 0.6.1, <http://www.ucl.ac.uk/~ucakiko/software.html>.
- Kruger KM and Hero J-M (2007) Large-scale seasonal variation in the prevalence and severity of chytridiomycosis. *Journal of Zoology* **271**, 352–359.
- Lanfear R, Senfeld T, Frandsen P, Wright A and Calcott B (2017) Partitionfinder 2: new methods for selecting partitioned models of evolution for molecular and morphological phylogenetic analyses. *Molecular Biology and Evolution* **34**, 772–773.
- Lenth R (2016) Least-squares means: the R package lsmeans. *Journal of Statistical Software* **69**, 1–33.
- MacDiarmid SC (1988) Future options for brucellosis surveillance in New Zealand beef herds. *New Zealand Veterinary Journal* **36**, 39–42.
- McAllister CT, Bursey CR, Crawford JA, Kuhns AR, Shaffer C and Trauth SE (2010) Metacercariae of *Clinostomum* (Trematoda: Digenea) from three species of *Ambystoma* (Caudata: Ambystomatidae) from Arkansas and Illinois, U.S.A. *Comparative Parasitology* **77**, 25–30.
- McCallum H and Dobson A (1995) Detecting disease and parasite threats to endangered species and ecosystems. *Trends in Ecology and Evolution* **10**, 190–194.
- Mendoza L, Taylor JW and Ajello L (2002) The class Mesomycetozoea: a heterogeneous group of microorganisms at the animal-fungal boundary. *Annual Reviews of Microbiology* **56**, 315–344.
- Minton Jr SA . (2001) *Amphibians and Reptiles of Indiana*. Indianapolis, IN: Indiana Academy of Science.
- Moral H (1913) Über das Auftreten von *Dermocystidium Pusula* (Pérez), einem einzelligen parasiten der haut des molches bei Triton cristatus. *Archiv für mikroskopische Anatomie* **81**, 381–393.
- Morris MA and Brandon RA (1984) Gynogenesis and hybridization between *Ambystoma Platineum* and *Ambystoma texanum* in Illinois. *Copeia* **1984**, 324–337.
- Pascolini R, Daszak P, Cunningham AA, Tei S, Vagnetti D, Bucci S, Fagotti A and Di Rosa I (2003) Parasitism by *Dermocystidium Ranae* in a population of *Rana esculenta* complex in Central Italy and description of *Amphibiocystidium* n. gen. *Diseases of Aquatic Organisms* **56**, 65–74.
- Pereira CN, Di Rosa I, Fagotti A, Simoncelli F, Pascolini R and Mendoza L (2005) The pathogen of frogs *Amphibiocystidium Ranae* is a member of the order Dermocystida in the class Mesomycetozoea. *Journal of Clinical Microbiology* **43**, 192–198.
- Pérez C (1907) *Dermocystidium Pusula*, organismenouveau parasite de lapeau-destritons. *Comptesrendus de Seances de Societe de Biologie* **63**, 445–446.
- Pérez C (1913) *Dermocystidium Pusula*: parasite de la peau des tritons. *Archives de Zoologie Experimentale et Generale* **52**, 343–357.
- Pessier AP (2008) Management of disease as a threat to amphibian conservation. *International Zoo Yearbook* **42**, 30–39.
- Phillips CA, Uzzell T, Spolsky CM, Serb JM, Szafoni RE and Pollowy TR (1997) Persistent high levels of tetraploidy in salamanders of the *Ambystoma Jeffersonianum* complex. *Journal of Herpetology* **31**, 530–535.
- Phillips CA, Johnson JR, Dreslik MJ and Petzing JE (2002) Effects of hydroperiod on recruitment of mole salamanders (genus *Ambystoma*) at a temporary pond in Vermilion County, Illinois. *Transactions of the Illinois State Academy of Science* **95**, 131–139.
- Picco AM and Collins JP (2008) Amphibian commerce as a likely source of pathogen pollution. *Conservation Biology* **22**, 1582–1589.
- Picco AM, Brunner JL and Collins JP (2007) Susceptibility of the endangered California tiger salamander, *Ambystoma californiense*, to ranavirus infection. *Journal of Wildlife Diseases* **43**, 286–290.
- Poisson C (1937) Sur une nouvelle espèce du genre *Dermomycooides* Granata 1919: *Dermomycooides armoriacus* Poisson 1936 parasite cutane de *Triturus palmatus* (Schneider). Genèse et structure de la zoospore. *Bulletin Biologique de la France et de la Belgique* **71**, 91–116.
- Price SJ, Garner TWJ, Nichols RA, Balloux F, Ayres C, de Alba AMC and Bosch J (2014) Collapse of amphibian communities due to an introduced ranavirus. *Current Biology* **24**, 2586–2591.
- Raffel TR, Bommarito T, Barry DS, Witiak SM and Shackelton LA (2008) Widespread infection of the eastern redspotted newt (*Notophthalmus viridescens*) by a new species of *Amphibiocystidium*, a genus of fungus-like Mesomycetozoa parasites not previously reported in North America. *Parasitology* **135**, 203–215.
- Rambaut, A and Drummond, AJ (2007) Tracer, Vol. 1.6. Available at: <http://tree.bio.ed.ac.uk/software/tracer/>.
- R Core Team (2018) *R: A Language and Environment for Statistical Computing*. Vienna, Austria: R Foundation for Statistical Computing. URL <https://www.R-project.org/>.
- Remy P (1931) Presence de *Dermocystidium Ranae* (Guyénot et Naville) chez une *Rana esculenta* L. de Lorraine. *Annals of Parasitology* **9**, 1–3.
- Rollins-Smith LA (2009) The role of amphibian antimicrobial peptides in protection of amphibians from pathogens linked to global amphibian declines. *Biochimica et Biophysica Acta-Biomembranes* **1788**, 1593–1599.
- Ronquist F, Teslenko M, van der Mark P, Ayres DL, Darling A, Höhna S, Larget B, Liu L, Suchard MA and Huelsenbeck JP (2012) MrBayes 3.2: efficient Bayesian phylogenetic inference and model choice across a large model space. *Systematic Biology* **3**, 539–542.

- Rowley J and Alford RA** (2007) Behaviour of Australian rainforest stream frogs may affect the transmission of chytridiomycosis. *Diseases of Aquatic Organisms* **77**, 1–9.
- Rowley J, Gleason FH, Andreou D, Marshall W, Lilje O and Goslan R** (2013) Impacts of Mesomycetozoean parasites on amphibian and freshwater fish populations. *Fungal Biology Reviews* **27**, 100–111.
- Scheid P, Balczun C, Dehling JM, Ammon A and Sinsch U** (2015) Rhinosporidiosis in African reed frogs *Hyperolius* Spp. Caused by a new species of Rhinosporidium. *Diseases of Aquatic Organisms* **115**, 111–120.
- Schloegel LM, Toledo LF, Longcore JE, Greenspan SE, Vieira CA, Lee M, Zhao S, Wangen C, Ferreira CM, Hipolito M, Davies AJ, Cuomo CA, Daszak P and James TY** (2012) Novel, panzootic and hybrid genotypes of amphibian chytridiomycosis associated with the bullfrog trade. *Molecular Ecology* **21**, 5162–5177.
- Simoncelli F, Fagotti A, Dall’Olio R, Vagnetti D, Pascolini R and Di Rosa I** (2005) Evidence of *Batrachochytrium Dendrobatidis* infection in water frogs of the *Rana esculenta* complex in central Italy. *EcoHealth* **2**, 307–312.
- Skerratt LF, Berger L, Speare R, Cashins S, McDonald KR, Phillott A, Hines H and Kenyon N** (2007) Spread of chytridiomycosis has caused the rapid global decline and extinction of frogs. *EcoHealth* **4**, 125–134.
- Spolsky C, Phillips CA and Uzzell T** (1992) Gynogenetic reproduction in hybrid mole salamanders (Genus *Ambystoma*. *Evolution* **46**, 1935–1944.
- Stamatakis A** (2014) RAxML version 8: a tool for phylogenetic analysis and post-analysis of large phylogenies. *Bioinformatics (Oxford, England)* **30**, 1312–1313.
- Stark T and Guex G-D** (2014) Infecties bij amfi bieën in Nederland: *Amphibiocystidium*. *Ravon* **54**, 54–56.
- Sugiura N** (1978) Further analysis of the data by Akaike’s information criterion and the finite corrections. *Communications in Statistics* **7**, 13–26.
- Teltser C and Greenwald KR** (2015) Survivorship of ploidy-variable unisexual *Ambystoma* Salamanders across developmental stages. *Herpetologica* **71**, 81–87.
- Uzzell Jr TM** (1964) Relations of the diploid and triploid species of the *Ambystoma Jeffersonianum* complex (Amphibia, Caudata). *Copeia* **2**, 257–300.
- Uzzell Jr TM and Goldblatt SM** (1967) Serum proteins of salamanders of the *Ambystoma Jeffersonianum* complex, and the origin of the triploid species of this group. *Evolution* **21**, 345–354.
- Warren DL, Geneva AJ and Lanfear R** (2017) RWTY (R We There Yet): an R package for examining convergence of Bayesian phylogenetic analyses. *Molecular Biology and Evolution* **34**, 1016–1020.
- Woodhams DC, Rollins-Smith LA, Alford RA, Simon MA and Harris RN** (2007) Innate immune defenses of amphibian skin: antimicrobial peptides and more. *Animal Conservation* **10**, 425–428.