

Review

Photoprotection in lichens: adaptations of photobionts to high light

Richard Peter Beckett^{1,2} , Farida Minibayeva³ , Knut Asbjørn Solhaug⁴  and Thomas Roach⁵ 

¹School of Life Sciences, University of KwaZulu-Natal, Private Bag X01, Scottsville 3209, South Africa; ²Open Lab ‘Biomarker’, Kazan (Volga Region) Federal University, Kremlevskaya str. 18, Kazan 420008, Russia; ³Kazan Institute of Biochemistry and Biophysics, Federal Research Center ‘Kazan Scientific Center of RAS’, P.O. Box 261, Kazan 420111, Russia; ⁴Faculty of Environmental Sciences and Natural Resource Management, Norwegian University of Life Sciences, P.O. Box 5003, 1432 Ås, Norway and ⁵Department of Botany, University of Innsbruck, Sternwartestrasse 15, Innsbruck 6020, Austria

Abstract

Lichens often grow in microhabitats where they are exposed to severe abiotic stresses such as desiccation and temperature extremes. They are also often exposed to levels of light that are greater than lichen photobionts can use in carbon fixation. Unless regulated, excess energy absorbed by the photobionts can convert ground state oxygen to reactive oxygen species (ROS). These ROS can attack the photosynthetic apparatus, causing photoinhibition and photo-oxidative stress, reducing the ability of the photobionts to fix carbon. Here, we outline our current understanding of the effects of high light on lichens and the mechanisms they use to mitigate or tolerate this stress in hydrated and desiccated states. Tolerance to high light can be achieved first by lowering ROS formation, via synthesizing light screening pigments or by thermally dissipating the excess light energy absorbed; second, by scavenging ROS once formed; or third, by repairing ROS-induced damage. While the primary focus of this review is tolerance to high light in lichen photobionts, our knowledge is rather fragmentary, and therefore we also include recent findings in free-living relatives to stimulate new lines of research in the study of high light tolerance in lichens.

Key words: ascorbate glutathione pathway, high light stress, non-photochemical quenching, PSII repair cycle, secondary metabolites

(Accepted 28 October 2020)

Introduction

Lichens are the dominant life forms in c. 8% of the land surface of the earth (Ahmadjian 1995), mainly in polar regions and on mountain tops. Vegetation growing in these places experiences severe abiotic stresses such as desiccation, temperature extremes and high light intensities. Arguably, what makes lichens special, and what separates them from most other eukaryotic organisms, is their ability to tolerate extreme stresses. Lichens have been termed ‘extremophiles’, organisms that can thrive in conditions that do not permit other, less specialized organisms to survive. Understanding the physiological processes that lie behind stress injury, and how lichens tolerate environmental stress, is therefore of great importance in lichen biology. While desiccation tolerance has received a great deal of attention in recent years, given their natural habitats, many lichens also need to tolerate high light levels, not only when hydrated but also at times when metabolic activity is not possible. The aim of this review is to outline our current understanding of the effects of high light on lichens and mechanisms enabling its tolerance, with an emphasis on recent publications. We have not exhaustively reviewed all the existing literature on light stress in lichens. An additional aim is to suggest new lines of research for studying tolerance to high

light in lichens, often based on recent findings in free-living relatives of lichen photobionts.

This review is dedicated to Professor Peter Crittenden, who made, and continues to make, an immensely valuable contribution to lichenology. It was Peter that helped the first author more than 35 years ago by awarding him a postdoctoral grant to work in Finnish Lapland. RB learned much during the tenure of this grant which inspired a love of arctic and boreal lichens that continues to the present day.

Acclimation and adaptations of lichens to their light environment

The concept of high light stress is closely linked to acclimation or induced tolerance, and the ability of a lichen to cope with an unfavourable light environment. Generally, stress tolerance in lichens often increases as a result of exposure to prior stress (Beckett *et al.* 2008). A lichen is then said to be acclimated (or hardened) by means of phenotypic plasticity. Acclimation can be distinguished from adaptation, which usually refers to a genetically determined level of resistance acquired by a process of selection over many generations. As will become clear in this review, adaptation and acclimation to light stress result from changes that occur at all levels of organization, from the anatomical and morphological level to the cellular, biochemical and molecular level.

Lichens display considerable plasticity in their responses to high light stress, as evidenced in their ability to display seasonal variations in photosynthetic capacity, and the ability of lichens to show ‘sun’ or ‘shade’ forms. In this, they resemble other

Author for correspondence: Richard Peter Beckett. E-mail: rpbeckett@gmail.com

Cite this article: Beckett RP, Minibayeva F, Solhaug KA and Roach T (2021) Photoprotection in lichens: adaptations of photobionts to high light. *Lichenologist* 53, 21–33. <https://doi.org/10.1017/S0024282920000535>

photosynthetic organisms, which must be able to respond to rapidly changing environmental conditions to optimize their light usage, and deal with excess light (Osmond *et al.* 1997). Early work demonstrated that the way lichens respond to light varies throughout the year. For example, Stålfelt (1938) showed that *Evernia prunastri* and *Ramalina farinacea* have higher rates of photosynthesis in summer than in winter. Later studies, in particular the detailed gas exchange measurements made by Prof. Ken Kershaw's group, confirmed the ability of lichens to adapt to seasonal changes in their light environment. Furthermore, changes could be replicated by exposure to light under laboratory conditions (for review see Kershaw (1985)).

In addition to seasonal changes, there is abundant evidence that members of the same species of lichen can, as for higher plants, display sun and shade forms (Piccotto & Tretiach 2010). Lichens from shaded habitats tend to have lower light saturation and compensation points than those of sun-exposed habitats (Green *et al.* 1997), as observed in vascular plants, and shade forms also have less cortical pigments (Dietz *et al.* 2000). Interestingly, other parameters, such as photosynthetic capacity and chlorophyll content which clearly differ in the sun and shade leaves of higher plants, show less clear patterns in lichens. Possibly, pigments in the fungal upper cortex provide lichen photobionts with far stronger light screening than that given by cuticles to photosynthetic cells in plant leaves. Therefore, in heavily melanized sun-exposed thalli the photobiont could still be shade-adapted (Gauslaa & Goward 2020). Furthermore, although lichen thalli superficially resemble leaves, they are small 'organismic ecosystems'. Other than simply photosynthesizing, lichen thalli need to optimize water and nutrient absorption and carry out reproduction (Piccotto & Tretiach 2010). Therefore, the adaptations present in sun and shade thalli represent more than adaptations to light. For example, Piccotto & Tretiach (2010) showed that nitrogen supply can greatly affect the photosynthetic parameters of a particular species.

While the way lichen photobionts display significant plasticity in their response to light is notable, it is clear that genetic adaptation, even within a single species, is also important. It is not always easy to distinguish phenotypic changes from genetic adaptation. Williams *et al.* (2017) transplanted thalli of the widespread lichen *Psora decipiens* between climatic zones in Europe and showed that the photobionts were unable to adapt to transplantation, and tended to die. Gene sequencing indicated that the photobionts from different populations were genetically distinct (Williams *et al.* 2017). Such transplant experiments demonstrate that there are limits to photobiont plasticity, and the mycobiont 'switches' to a locally adapted photobiont when environmental challenges exceed a certain threshold.

Why is excess light absorption harmful to photosynthetic organisms?

Even in lichens adapted to high light conditions, when the maximum rate of photosynthesis (P_{\max}) has been reached, photosynthesis will no longer be able to use further light. In higher plants, even the hardiest individuals reach P_{\max} at less than full sunlight, and individual leaves on plants growing in full sun commonly experience excess light intensities (Pospíšil 2016). In fact, even under moderate light, energy transfer and electron transport in photosystem II (PSII) unavoidably lead to the production of various reactive oxygen species (ROS) (Pospíšil 2016). For example, the transfer of energy from triplet chlorophyll in the reaction

centre of a photosystem (e.g. PSII) to molecular oxygen produces the highly reactive singlet oxygen (1O_2). At the other end of the electron transport chain, at the acceptor side of photosystem I (PSI), electrons can be leaked to molecular oxygen, forming the superoxide anion radical (O_2^-). Superoxide dismutates into hydrogen peroxide (H_2O_2), which feeds into many redox signalling pathways but can also be reduced by Fe^{2+} or Cu^+ to form highly reactive hydroxyl radicals (HO^\cdot) (Foyer 2018). Reactive oxygen species production greatly increases when the light absorbed exceeds that which can be utilized for carbon fixation; this will occur particularly when a plant is stressed and fixation reduced (Liu *et al.* 2019). There are few studies that have directly demonstrated that ROS are induced by light stress in lichen photobionts. Carniel *et al.* (2015) used a histochemical technique based on dichlorofluorescein diacetate to show light increased ROS production in both cultured and symbiotic *Trebouxia* from the lichen *Parmotrema perlatum* recovering from desiccation stress. However, more detailed information is available from free-living algae. In *Chlamydomonas reinhardtii*, H_2O_2 production under saturating light is lowered when CO_2 availability is restricted (Roach *et al.* 2015), whereas symptoms of singlet oxygen (1O_2) stress are closely associated with excess light/photo-oxidative stress (Roach *et al.* 2017). These results are consistent with results obtained from higher plants (Triantaphylidès *et al.* 2008; Noctor *et al.* 2014). When the production of ROS, and particularly 1O_2 , exceeds the capacity of the plant's detoxification systems, they can react with thylakoid membranes, leading to lipid peroxidation or damage to the protein complexes of the photosynthetic apparatus, contributing to a reduction in photosynthesis in what is often termed 'photoinhibition' (Li *et al.* 2018). Strong evidence exists that photoinhibition regularly occurs in lichens in field situations. For example, Gauslaa & Solhaug (2000) and Jairus *et al.* (2009) both showed that lichens growing on trees with reduced canopy cover (e.g. as a result of the felling of surrounding trees) display sustained reductions in photosynthesis. Photoinhibition may occur on an annual basis in some environments. For example, Míguez *et al.* (2017a) showed that lichens from subalpine environments display an annual winter photoinhibitory response. However, even under normal mild temperate conditions, continuous field measurements of photosynthesis in *Lecanora muralis* over several days indicated that photoinhibition is a regular occurrence (Leisner *et al.* 1997).

Why are poikilohydric organisms particularly sensitive to light stress?

Poikilohydric organisms such as lichens may be particularly sensitive to high light stress for at least three reasons. First, during drying, carbon fixation often stops before photophosphorylation, increasing the 'leakage of electrons' to ground state oxygen and therefore stimulating ROS production (Challabathula *et al.* 2018). Second, even though lichens may rapidly dry out when exposed to high light, they can suffer from light stress even when desiccated (Kershaw & MacFarlane 1980). In bryophytes, desiccation does not stop the transfer of excitation energy from the light-harvesting pigments to the reaction centres (Heber *et al.* 2006). However, the highly quenched state of chlorophyll upon desiccation shows extremely efficient dissipation of photons, lowering 1O_2 formation. Third, even if light only causes the formation of tiny amounts of ROS in desiccated thalli, normal repair processes do not take place (Buffoni Hall *et al.* 2003). Enzyme reactions are severely restricted by 'rubbery' cytoplasmic states

that occur at the onset of desiccation, and are totally restricted in glassy cytoplasmic states (Fernandez-Marin *et al.* 2013), typically found in air-desiccated lichens during the day. Furthermore, recovery upon hydration is a key issue for poikilohydric organisms, with clear legacy effects of the duration of desiccation, which can be considered in some respects as 'ageing', in for example conversion rates of zeaxanthin back to violaxanthin and cellular redox states (Kranter *et al.* 2003). At the other end of the spectrum, the absence of a cuticle means that sometimes a lichen thallus can become oversaturated with water. Under these conditions CO₂ diffusion is very slow, reducing its fixation rate (Cowan *et al.* 1992). As saturated lichens will still be absorbing light, this excess energy needs to be dissipated. The study of Leisner *et al.* (1997), involving continuous measurements of photosynthesis in *Lecanora muralis*, showed that thalli regularly become saturated and display a strong depression of photosynthesis under field conditions, probably as a result of the combination of low CO₂ supply and photoinhibition.

Although not directly related to poikilohydry, as discussed in the Introduction, lichens often dominate in cold boreal and sub-polar regions, and on the tops of mountains. As carbon fixation is an enzymatic process, low temperatures will increase the chance that more light is harvested than can be used in metabolic activity, resulting in excess PSII excitation pressure (Huner *et al.* 1998; Öquist & Huner 2003). While lichens are clearly well adapted to cold environments (e.g. Cho *et al.* 2020), the specific adaptations of their photosynthetic apparatus to low temperatures remain unclear (Sahu *et al.* 2019). Further work is needed to elucidate precisely how lichens can protect themselves under contrasting environmental conditions.

Overview of tolerance mechanisms

Tolerance to high light levels in photosynthetic organisms has been reviewed many times (e.g. Derks *et al.* 2015; Liu *et al.* 2019). Tolerance mechanisms can be divided into three broad categories: 1) ROS formation can be reduced by synthesizing light screening pigments or by dissipating the excess energy absorbed radiationlessly as heat; 2) ROS can be scavenged once formed; 3) ROS-induced damage can be repaired. According to the classical model of stress resistance developed by Levitt (2012), screening, dissipation and scavenging would be classed as stress avoidance, while only repair would be classed as true tolerance. However, for simplicity, in this review any mechanism that reduces the potential for high light stress on lichens will be referred to as tolerance.

Avoidance by Light Screening

The algal partner in lichens is protected against high light by screening in the upper cortex of the lichen thallus. The amount of visible light transmitted by a moist cortex ranges from c. 90% for rainforest lichens to only 45% for lichens from high light exposed sites (Dietz *et al.* 2000). Transmittance is highly reduced in the dry state (Ertl 1951). Cortical screening results from secondary compounds that absorb radiation or from the optical properties of the fungal hyphae in the cortex. Increased screening in the dry state is important because, as discussed above, lichens have less opportunity to dissipate excess energy or repair damage from excess radiation when dry. Therefore, exposure to high or medium light levels of visible radiation for long periods

in the dry state may result in accumulated severe damage (Gauslaa & Solhaug 1996; Gauslaa *et al.* 2012; Mafole *et al.* 2019b).

Lichens often contain large amounts of fungal-produced secondary compounds. More than 1050 different secondary compounds have so far been isolated and characterized (Huneck & Yoshimura 1996; Molnar & Farkas 2010). Most of these secondary compounds, particularly those that occur in the cortex, absorb UV radiation and some also absorb visible radiation. The extinction coefficients (see Huneck & Yoshimura 1996) show that most lichen compounds absorb UV-B radiation very efficiently. The high concentration of secondary compounds that normally comprise several percent of the thallus dry mass indicates that these compounds effectively screen harmful UV-B. However, high extinction coefficients in organic solvents do not necessarily mean that the secondary compounds screen efficiently *in vivo*. Many secondary compounds occur as crystals outside fungal hyphae. Screening efficiency might then be less than predicted because light may pass between the crystals (McEvoy *et al.* 2007b; Solhaug & Gauslaa 2012). As most of the secondary compounds are almost insoluble in water, light transmission between crystals will also occur for lichen thalli in the moist state, and the screening efficiency *in vivo* might be less than what is indicated by absorbance spectra in an organic solvent. Although the absorbance spectra of secondary compounds show that they have high potential for screening radiation, other functions such as herbivore protection may be more important (reviewed by Solhaug & Gauslaa (2012)). In addition, there is a trade-off between visible light protection against photoinhibition and photosynthetic efficiency under low light. The blue light-absorbing compounds parietin in *Xanthoria* species and vulpinic acid in *Letharia vulpina* protect these lichens against photoinhibition, although the quantum yield of photosynthesis is reduced (Solhaug & Gauslaa 1996; Phinney *et al.* 2019).

Most secondary lichen compounds do not absorb visible light. Among the few coloured lichen compounds, the most widespread are yellow compounds that absorb blue light and can therefore protect against these wavelengths. Higher plants, algae and cyanobacteria are more photoinhibited by blue light than by red and green light (see Zavafer *et al.* 2015), which is probably a consequence of the action spectra of the two major processes responsible. The first process is damage to the oxygen-evolving complex of PSII, which has an action spectrum that increases steeply with decreasing wavelengths in the blue and UV region of the spectrum; the second process involves the destruction of PSII (Ohnishi *et al.* 2005). The second process has an action spectrum similar to the absorbance spectrum of chlorophyll. Strong UV and blue light photoinhibition can be explained by damage to the oxygen-evolving complex of PSII (Ohnishi *et al.* 2005; Zavafer *et al.* 2015). The dominance of yellow secondary compounds in lichens might therefore be explained by a greater need for screening against blue light-induced photoinhibition. Although not absorbing visible light, removal of atranorin from *Physcia aipolia* by acetone rinsing considerably reduced the reflectance of moist thalli, presumably because crystals either directly reflect light or raise reflection by preventing water from entering air spaces in the cortex (Solhaug *et al.* 2010).

Several lichens contain melanins (reviewed by Mafole *et al.* (2019a)). They may either be produced constitutively as melanins in the lower cortex such as in *Parmelia* species or throughout the outer cortex such as in fruticose lichens such as in dark *Bryoria* species; alternatively, they may be induced by UV-B radiation in the upper cortex as in *Lobaria pulmonaria* (Solhaug *et al.* 2003).

Interestingly, although melanins in the upper cortex are induced by UV-B radiation, they are not necessary for UV-B protection of the photobiont because non-melanized thalli tolerate extremely high UV-B levels (Gauslaa *et al.* 2017). However, for melanins there is also a trade-off between visible light screening and photosynthetic efficiency in *L. pulmonaria*. High light acclimatized thalli of *L. pulmonaria* synthesize melanins in the upper cortex, reducing cortical transmission of photosynthetic active light (Gauslaa & Solhaug 2001), which may give protection against photoinhibition in high light sites (Mafole *et al.* 2019b). The strategy can sustain similar growth rates in lichens transplanted to sites that vary widely in forest canopy openness (Gauslaa & Goward 2020). However, the photosynthetic efficiency in low light is reduced (Mafole *et al.* 2017). Melanic *Bryoria* hair lichens are more resistant to photoinhibition in the dry state than the yellow usnic acid-containing species *Usnea* and *Alectoria*. This difference probably explains why *Bryoria* species are mainly found in the upper canopy, whereas usnic acid-containing species are more frequent in the lower canopy (Färber *et al.* 2014).

Melanins affect the energy budgets of lichens. For instance, the dark, melanic thalli of some hair lichens absorb more light and may even melt snow in winter, feeding the lichens with water (Coxon & Coyle 2003). The thallus temperature of *L. pulmonaria* may be 3 °C higher in melanized than in pale thalli (McEvoy *et al.* 2007a). Melanic fungi are more frequent in cold areas (Gostinčar *et al.* 2012) where the melanin-induced heating might represent a competitive advantage. However, melanization of the fungal partner of a lichen might be a trade-off between protection against high light photoinhibition and avoidance of high temperature damage. This might explain the lower frequency of dark melanic lichens in hot areas where heat damage can be a problem.

The cyanobacterial lichens in the genera *Collema*, *Gonohymenia* and *Peltula* growing in high light sites contain the yellow-brown pigment scytonemin as a screening compound (Büdel *et al.* 1997). In contrast to secondary lichen compounds produced by the mycobiont discussed above, scytonemin is synthesized only by their cyanobacterial photobionts (Büdel *et al.* 1997). Scytonemin is located extracellularly in the outer sheath of some cyanobacteria, and it is also found in free-living cyanobacteria (Garcia-Pichel & Castenholz 1991). Its absorbance spectrum shows that it has high screening efficiency for UV radiation. In addition, it has high screening potential, especially for blue wavelengths of the electromagnetic spectrum (Garcia-Pichel & Castenholz 1991) the most photoinhibitory range of visible radiation.

A frequent strategy in lichens when dry is to increase reflectance (see e.g. Gauslaa 1984) and decrease cortical transmission (Ertl 1951; Gauslaa & Solhaug 2001; McEvoy *et al.* 2007b). It seems likely that the absence of water between cortical hyphae will increase reflection from hyphal surfaces. This is analogous to the situation in higher plants where infiltrating leaves with water reduced the optical path of light, thereby increasing transmission (Vogelmann 1993). Therefore, hydration will reduce the reflection of the upper cortex, and at the same time light transmission will increase.

Finally, some lichens display 'structural avoidance' of high light by curling their thalli. The edges of *L. pulmonaria* thalli, for example, will curl inwards during drying, making them less susceptible to high light photoinhibition in the dry state (Barták *et al.* 2006). An extreme example of structural avoidance is the vagrant lichen *Xanthoparmelia hueana* in the Namib Desert. In the dry state it curls, exposing only the highly melanic lower


side, whereas when moistened it uncurls and exposes the green upper side (Büdel & Scheidegger 2008).

Avoidance of ROS Formation by Regulating Light-Use Efficiency

Despite possessing a variety of light screening pigments, there are times when the amount of light that reaches the photobiont exceeds that which can be used in photosynthesis. This can be problematic because, as discussed above, excess light energy results in elevated levels of ROS produced by chlorophyll ($^1\text{O}_2$) and electron transport chains (O_2^- and H_2O_2), thus leading to photo-oxidative damage (Roach & Krieger-Liszka 2019). Photobionts use several processes to regulate the efficiency at which light energy is used, which are collectively referred to as non-photochemical quenching (NPQ). Estimating NPQ in lichens with chlorophyte photobionts (chlorobionts, eukaryotic algae) is relatively simple with standard chlorophyll fluorescence devices (see Kalaji *et al.* (2014) for details). For cyanobacterial photobionts (cyanobionts), specialized devices using red excitation light are required. Nevertheless, it is important to keep in mind that several NPQ components can occur simultaneously (Roach & Na 2017), each affecting chlorophyll fluorescence and requiring careful interpretation of the measurements.

The light energy needed to excite PSII is harvested in completely different ways in cyano- and chlorobionts. Cyanobacteria possess phycobilin proteins (also called phycobilisomes) made up of light-absorbing tetrapyrrole-containing phycocyanin and allophycocyanin pigments (Fig. 1A). These have been functionally replaced in chlorobionts by light-harvesting complex (LHC) antenna proteins (Fig. 1B), which contain carotenoid and chlorophyll pigments. Safely converting excess light energy to heat is referred to as thermal dissipation, or qE (Müller *et al.* 2001). Activation of qE is rapid, and it can increase within seconds to minutes (Niyogi & Truong 2013; Erickson *et al.* 2015), although on transition to darkness it may take longer to reduce or 'relax' (Kromdijk *et al.* 2016). Regardless of the photobiont, lichen desiccation is associated with induction of regulated thermal dissipation of excess light, a trait found in all desiccation-tolerant organisms (Calatayud *et al.* 1997; Heber *et al.* 2000; Komura *et al.* 2010; Wieners *et al.* 2018).

Considering the major structural differences between the phycobilisomes that occur in cyanobionts and the LHC of chlorobionts, it is not surprising that major differences in the regulation of light harvesting exist between these two photobionts (Fig. 1). Cyanobacteria contain an Orange Carotenoid Protein (OCP). OCP is coded by a highly conserved gene that is present in most of the known cyanobacterial genomes, including *Nostoc*, a common lichen photobiont (Boulay *et al.* 2008; Kerfeld *et al.* 2017). Excess energy collected by phycobilisomes can be rapidly dissipated by OCP (Wilson *et al.* 2008), thereby preventing excess ROS formation by photosystem reaction centres (Fig. 1A). OCP binds various carotenoids and, like zeaxanthin, can function as an antioxidant (Sedoud *et al.* 2014). PSI reaction centres can rapidly dissipate excess energy via charge recombination, even from the PSII antenna in a process described as 'spillover' or 'state transitions' (Fig. 1A). However, due to the absence of LHC in cyanobacteria, cyanobacteria-type state transitions are mechanistically unrelated to state transitions of algae (Calzadilla *et al.* 2019), which involve the migration of LHC between photosystems (see below). Cyanobacteria also possess chlorophyll-binding (CAB) proteins such as IsiA, which is produced under stress conditions



Feature	Cyanobionts	Chlorobionts
Light harvesting instrument	phycobilisomes	light harvesting complexes
Antenna pigments	(allo)phycocyanin	carotenoids, chlorophyll
Reaction centre pigments	chlorophyll a, phaeophytin	chlorophyll a, phaeophytin
PSII photoinhibition repair	rapid	rapid
qE NPQ	OCP	LHCSR
Zeaxanthin NPQ	-	some species
State transition NPQ	-	yes
Spillover (PSI quenching)	yes	yes (desiccation)
PSI electron donor	cytochrome c ₆	plastocyanin
Flavodiirons	yes	yes
HLIP – ELIP proteins	yes	yes
IsiA PSI antenna protein	yes	-

Fig. 1. Regulation of the light use efficiency in lichen photobionts, as depicted on cross-sections of thylakoid membranes that host the various photosynthetic protein complexes. In cyanobacteria (cyanobionts) (A) stromal phycobilisomes (blue and red) assist as light harvesting antenna for PSII, with excess energy thermally dissipated (qE) by the orange carotenoid protein (OCP-qE). In contrast, in chlorophyte photobionts (eukaryotic algae, chlorobionts) (B) thylakoid membrane-embedded light-harvesting complexes (e.g. LHCSR) assist in harvesting light, while other LHC-type proteins dissipate excess energy (e.g. LHCSR) upon protonation, and the xanthophyll cycle contributes to thermal dissipation in some photobionts (zeaxanthin-qE). Photoinhibition (qI) is a universal attribute of PSII, lowering charge separations in PSII, also affecting light use efficiency. Electrons released by PSII enter the photosynthetic electron transport chain (PETC, black-dashed line), first transported by plastoquinone (PQ) to the cytochrome b₆f complex (Cyt. b₆f). Subsequently, electrons are transported by a cytochrome (Cyt. c₆) in cyanobacteria, and by plastocyanin (PC) in eukaryotes, to PSI, and eventually reduce NADP⁺ to NADPH. The PSI reaction centre, P700, is also an excellent quencher via charge recombination, which may facilitate removal of excess energy from PSII via ‘spill-over’, particularly when PSII and PSI come in close contact during desiccation. In cyanobacteria, IsiA proteins can assist PSI with harvesting light, while in eukaryotes LHCSR can migrate between PSII and PSI, during state transitions. At the donor side of PSI, flavodiiron proteins (FLV) can take electrons and reduce O₂ to H₂O, averting excess reducing power. (C) Similarities and differences between the two photobionts listed. ELIP – HLIP = early high light-inducible proteins; NPQ = processes to regulate the efficiency at which light energy is used; PSI and PSII = photosystems I and II.

and transfers light energy to PSI, counteracting PSI photoinhibition (Havaux *et al.* 2005). However, the CAB/LHC proteins in cyanobacteria that are most closely related to those from eukaryotic algae are the early high light-inducible proteins (ELI/HLIP) which bind chlorophylls and assist in biogenesis of other chlorophyll-binding proteins (e.g. PSII) and assist in other light-stress-associated processes (Komenda & Sobotka 2016).

In contrast to the phycobilisomes of cyanobacteria, eukaryotic algae have LHC antenna proteins and, as a result, dissipate excess energy using strategies similar to those found in plants. The enzyme-catalyzed xanthophyll cycle responds to changing light intensities, whereby the carotenoid violaxanthin is enzymatically converted to zeaxanthin in a pH-regulated process that occurs during increases in light intensity. Zeaxanthin is involved in thermal dissipation in some but not all chlorobionts (Demmig-Adams *et al.* 1990; Míguez *et al.* 2017b). In such cases, an alternative role of zeaxanthin, further to antioxidant and structural roles, is the facilitation of recovery of PSII activity during rehydration (Štepičová *et al.* 2008; Verhoeven *et al.* 2018).

LHC proteins have diversified into several isoforms, each having a unique function in light harvesting and NPQ (Büchel 2015). Thus, rather than a typical light-harvesting role, the LHC-stress related (LHCSR or LHCX) proteins accumulate under stress to thermally dissipate excess energy (Peers *et al.* 2009). In chlorobionts, LHCSR proteins are thought to dissipate excess energy in LHCI-PSII complexes, but LHCSR3 may also protect PSI from photoinhibition (Bergner *et al.* 2015; Roach *et al.* 2020). This is probably related to state transitions, an NPQ mechanism that is important in *Chlamydomonas reinhardtii*, for instance, for acclimation to high light, in which LHC migrates between PSII and PSI (Allorent *et al.* 2013). Presumably, LHCSR3 can protect PSI by quenching LHCI when it is diverting energy to PSI after transition to state II (Girolomoni *et al.* 2019). The downstream effect of state transitions is a change in the relative activities of PSII and PSI, and therefore redox poise of the photosynthetic electron transport chain which is important for efficient photosynthesis (Rochaix 2011). Although work with lichens is just beginning, measurements of slow and rapid chlorophyll fluorescence kinetics in chlorophycean photobionts strongly suggest that state transitions are highly active (Mishra *et al.* 2015; Marečková & Barták 2016).

There have been limited studies on the ecophysiology of NPQ. As discussed above, an inevitable consequence of being poikilohydric and growing in generally harsh environments, is that lichen photobionts frequently absorb more light energy than they can use in carbon fixation. For example, *Lecanora muralis* and *Fulgensia fulgens* were shown to be metabolically active for only a third of the year and could carry out photosynthesis for less than half that time (Lange 2002; Lange & Green 2008). The NPQ mechanisms discussed above require moist thalli with active metabolism. When lichen thalli are exposed to high direct solar radiation for a significant time they will desiccate and remain dry and mainly metabolically inactive for most of the time exposed to high light. However, chlorophyll in the dry thalli will still absorb excess light which needs mechanisms not dependent on metabolic activity to dissipate the light in a safe way (Verhoeven *et al.* 2018). These mechanisms are poorly known but at least two different quenching mechanisms in desiccated lichens have been proposed and discussed in a series of papers by Ulrich Heber (see e.g. Heber *et al.* 2010; Heber 2012). First, there may be direct quenching in PSII, possible from charge recombination. The second, and possibly more important,

quenching mechanism in dry lichen thalli is spillover from PSII to PSI (see Fig. 1) when the two photosystems come in closer contact during desiccation (Slavov *et al.* 2013). The chlorophyll-containing reaction centre of PSI is an excellent quencher of excess light energy.

Studies with model algae and cyanobacteria have revealed a plethora of mechanisms required for regulation of light energy. Unfortunately, no studies appear to have been carried out on OCP-based dissipation in cyanobionts, or on specific roles for LHCSR in chlorobionts. Nonetheless, for chlorobionts two general conclusions can be drawn. First, unlike in plants, xanthophyll pigments are not always involved in thermal dissipation. Second, in general, high levels of thermal dissipation have been found in lichens that need photoprotection. For example, Calatayud *et al.* (1997) showed that NPQ tends to be induced in drying thalli of *Parmelina quercina*. Vráblíková *et al.* (2006) studied seasonal variation of NPQ in *Xanthoria parietina* sampled in one location in Norway for one year. NPQ rapidly increased from early spring until summer solstice, suggesting a higher need for photoprotection in the season with the highest solar irradiance. As discussed in our conclusions below, further studies are needed, for example to test seasonal variations in NPQ in more species and to separate out when diurnally and seasonally each component is required.

Finally, as with any tolerance mechanism, it is important to realize that there is a 'cost' to photosynthetic organisms of increasing NPQ. This cost is a reduction in photosynthetic capacity (Demmig-Adams *et al.* 2012), especially following a reduction in light intensity (Kromdijk *et al.* 2016), because NPQ takes a finite time to 'relax'. The photobiont must carry out a balancing act, on the one hand efficiently utilizing every possible photon to fix carbon when light is limited, while on the other hand dissipating energy when there is an excess. Careful regulation of the transitions between these two alternative states of the photosynthetic system is essential for optimizing both productivity and safety in continuously changing environments.

Scavenging ROS by Enzymatic and Non-Enzymatic Antioxidants

Enzymatic antioxidants

When light absorption overwhelms the capacity of quenching mechanisms, ROS formation in the photosynthetic apparatus will result. In higher plants, a multi-level H₂O₂ removal system exists in chloroplasts, including many thiol peroxidases (e.g. peroxiredoxins), and the ascorbate-glutathione pathway (also known as the Halliwell-Asada cycle). This latter pathway effectively removes ROS from chloroplasts and other cellular locations using ascorbate, glutathione (GSH) and NADPH along with the enzymes linking them (Pandey *et al.* 2015; Hasanuzzaman *et al.* 2019). In higher plant chloroplasts, superoxide dismutase (SOD) dismutates O₂⁻ to H₂O₂ which is then broken down to H₂O and O₂ in a reaction catalyzed by chloroplastic forms of ascorbate peroxidase (APX), with most reactions occurring in the stroma. In the process, ascorbate is converted to monodehydroascorbate (MDHA), some of which disproportionates to dehydroascorbate (DHA). Ascorbate is regenerated from DHA by dehydroascorbate reductase (DHAR) in a reaction involving the oxidation of GSH to oxidized glutathione (GSSG). GSH is recovered from GSSG by glutathione reductase (GR), with NADPH providing the reducing power. Monodehydroascorbate reductase (MDHAR) reduces any MDHA that has not disproportionated directly back to ascorbate. In higher

plants the effectiveness of this cycle may be limited by the sensitivity of APX to H_2O_2 (Kitajima 2008), particularly at low ascorbate concentrations. However, APX may be less sensitive in free-living algae such as *Chlamydomonas reinhardtii* (Takeda *et al.* 1997), which also have much lower ascorbate contents than plants (Gest *et al.* 2013). Interestingly, for higher plants, Maruta *et al.* (2016) have proposed an alternative view that the more H_2O_2 resistant chloroplastic peroxiredoxins protect sensitive sites. The sensitivity of APX to H_2O_2 allows the formation of high levels of H_2O_2 under stressful conditions, with H_2O_2 functioning as a signalling molecule.

Whether a classical chloroplast ascorbate-glutathione pathway occurs in free-living cyanobacteria and chlorophycean algae remains unclear. Apparently, some cyanobacteria possess APX-like activity and similar thiol-based redox recycling mechanisms (Tel-Or *et al.* 1985; Miyake *et al.* 1991). Haghjou *et al.* (2009) studied the role of the ascorbate-glutathione pathway in the free-living green alga *Dunaliella salina*. While all of the components of the pathway appeared to be present, high light stress only increased the activity of SOD and the amount of ascorbate and GSH, but had little effect on the activity of the other enzymes involved in the pathway. By contrast, exposing *C. reinhardtii* to high light for 1 h increased APX (Roach *et al.* 2015) and GR activity (Lin *et al.* 2018), and GSH content (Lin *et al.* 2018; Roach *et al.* 2018). The thorough database survey of Maruta *et al.* (2016) indicated that unicellular green algae contain only chloroplastic monofunctional APXs that lack any transmembrane domains, suggesting that they might occur in the stroma. However, the concentrations of ascorbate in green algae is typically around 100–400 μ M, much lower than those found in higher plants (Gest *et al.* 2013). For the other enzymes involved in the ascorbate-glutathione pathway, two GR isoforms have been found in *C. reinhardtii*, thought to be localized in the cytoplasm and chloroplast (Serrano & Llobell 1993), and certainly microalgae contain SOD isoforms that are chloroplastic (Wolfe-Simon *et al.* 2005). However, the location of other enzymes in the ascorbate-glutathione pathway in green algae (e.g. MDHAR and DHAR) remains unclear. However, even if in algae most detoxification occurs in the cytoplasm, it seems likely that, as in higher plants, cytosolic ROS detoxification imparts ‘cross compartment protection’ of organelles during periods of stress (Davletova *et al.* 2005). Finally, catalase is important in algae for tolerating high H_2O_2 levels and may contribute to H_2O_2 -mediated light stress signalling, although its cellular location is unclear (Michelet *et al.* 2013). Taken together, the available data suggest that free-living algae can readily break down H_2O_2 produced by the photosystems, but whether they possess the classical chloroplast ascorbate-glutathione pathway found in higher plants is uncertain.

In lichens, only fragmentary information is available on the role of enzymatic antioxidants in scavenging high light-induced ROS. One reason for this is that investigations are complicated by the high ratio of fungal to algal biomass in the thallus. Lichen mycobionts certainly contain GR (Kraner 2002; Kraner *et al.* 2005) and other key enzymes such as SOD (Weissman *et al.* 2005). While Vrábliková *et al.* (2005) showed that high light reduces total thallus GSH levels in *Umbilicaria antarctica* and *Lasallia pustulata*, the location of the GSH (mycobiont or photobiont) was unclear. In a more detailed study, Kraner *et al.* (2005) quantified GSH in an isolated photobiont of *Cladonia vulcani*, although the changes of GSH in the isolated symbionts in response to desiccation were clearly different to those in intact thalli. In theory, ascorbate metabolism should be

easier to study, because fungi do not contain ascorbate and the associated enzymes (Smirnoff 2018). Unfortunately, we currently lack a clear overview of the ROS scavenging enzymes in lichen photobionts. Transcripts and proteins possess unique sequences depending on whether they are from the fungus or the photobiont, and therefore meta-transcriptomics/proteomics will be very useful tools in future studies, allowing photobiont- and mycobiont-specific processes to be distinguished.

Non-enzymatic antioxidants

Apart from their roles in the ascorbate-glutathione cycle, ascorbate and GSH probably act as antioxidants on their own, or as part of other protective pathways. Indirect evidence for the general importance of ascorbate comes from the observation that a depleted ascorbate level in the mutant vacuolar transporter chaperone 2 (VTC2) deficient free-living alga *Chlamydomonas reinhardtii* probably led to elevated NPQ values (Vidal-Meireles *et al.* 2020). Ascorbate is also a substrate for violaxanthin de-epoxidase during the formation of zeaxanthin (Yamamoto & Higashi 1978), although not in *C. reinhardtii* (Vidal-Meireles *et al.* 2020). GSH is a powerful ROS scavenger in free-living algae; for example, Roach *et al.* (2018) showed that GSH conjugates very quickly with electrophiles produced during photo-oxidative stress. While ascorbate and GSH appear highly likely to be important in scavenging light-induced ROS in photobionts, the role of other low molecular weight antioxidants remains speculative. These include a great variety of secondary metabolites with strong in vitro antioxidant activity (Kosanić *et al.* 2011; Thandhani *et al.* 2011). However, as discussed above, classical lichen secondary metabolites mostly occur as crystals on the cell walls of the mycobiont (Molnar & Farkas 2010) and would seem unlikely to directly scavenge ROS produced inside the photobiont. Similarly, while fungal melanins are also powerful antioxidants, they are located in the cell walls of the mycobiont (Mafole *et al.* 2019a). More plausibly, some photobionts such as *Trentepohlia* contain high concentrations of carotenoids, such as β -carotene. While carotenes can act as light screening pigments (Kharkongor & Ramanujam 2015), it seems likely that in addition they behave as lipid-soluble antioxidants. Certainly, pigments from free-living algae have recently been shown to provide health benefits due to their antioxidant potential (Sathasivam & Jang-Seu 2018). Furthermore, Kraner *et al.* (2005) showed that desiccation reduces β -carotene in the photobiont of *Cladonia vulcani*, presumably as a consequence of oxidative stress. However, the ability of β -carotene to protect algal cells by scavenging light-induced ROS appears not to have been tested directly, either in free-living algae or photobionts. Future studies need to test whether tolerance to high light is correlated with the levels of molecules having the potential to scavenge ROS. For example, chloroplasts of alpine plants contain up to ten times the amount of ascorbate and glutathione found in lowland individuals of the same species (Streb *et al.* 1997), and it would be interesting to test whether the photobionts of high-altitude lichens also contain elevated levels of these low molecular weight antioxidants.

Repair

The PSII repair cycle

In photosynthesizing organisms, an important target for light stress is the PSII complex in which electron transport can be

impaired due to damage of the catalytic Mn cluster of the water oxidizing complex (Vass *et al.* 2014). The D1 and D2 proteins, which form the backbone of the reaction centre complex, appear particularly sensitive and are also damaged by UV-B exposure (Li *et al.* 2018). Some damage appears to occur even under moderate light intensities; therefore, photosynthesizing organisms must continuously repair the damage to these proteins. The balance between damaging processes and restoration of the structural and functional integrity of PSII complexes determines the extent of the PSII damage. The 'PSII-repair cycle', occurring in both chloroplasts and cyanobacteria, involves proteolytic removal of the damaged D1 and D2 proteins, production of new subunits, incorporation of them into the PSII complex, re-ligation of redox cofactors and finally activation of the reaction centre (Nath *et al.* 2013; Vass *et al.* 2014). In free-living chlorophycean algae adapting to high light, chloroplasts can gradually acquire greater capacity for such repair (Kim *et al.* 1993). Similarly, in the algal symbionts of corals, application of an inhibitor of PSII repair, lincomycin, in combination with light stress has a greater effect on coral adapted to high rather than low light (Jeans *et al.* 2013). Perhaps surprisingly, the PSII repair cycle appears not to have been studied in lichen photobionts. Although such a cycle is only likely to operate in hydrated lichens (Buffoni Hall *et al.* 2003), upregulation of this cycle might help them to acclimate to high light.

Recent studies have revealed that for efficient PSII repair, photolyase-mediated repair of DNA is required (Vass *et al.* 2013). DNA is highly sensitive to high light stress, which can cause the formation of polymerase-blocking lesions in the molecule. Unrepaired DNA damage interrupts the PSII repair cycle at the step of gene transcription, and thus inhibits *de novo* D1 and D2 protein synthesis. Most of these DNA lesions are cyclobutane pyrimidine dimers which can be effectively reversed to native functional nucleotides by the photolyase enzymes. Photolyases and cryptochromes form an almost ubiquitous family of blue light photoreceptors involved in the repair and maintenance of DNA integrity or regulatory control (Franz *et al.* 2018). The roles of these enzymes in UV-induced signalling and DNA repair were shown in the free-living cyanobacterium *Synechocystis* (Vass *et al.* 2014) and the green alga *Chlamydomonas reinhardtii* (Franz *et al.* 2018). Interestingly, mutant *Synechocystis* that lacks the cryptochrome Syn-CRY, also displayed reduced levels of proteins involved in CO₂ fixation (Vass *et al.* 2014). Possibly this mutant with a reduced capacity to fix CO₂ has a decreased rate of PSII repair due to an enhanced accumulation of ROS and, as a result, the inhibition of synthesis of PSII proteins and especially the D1 protein.

As described above, LHC proteins are important in photoprotection in plants and algae by dissipating excess light energy. Interestingly, some of these proteins also appear to be involved in the regulation of chlorophyll synthesis and in the assembly and repair of PSII and PSI, possibly by mediating the insertion of newly synthesized pigments into the photosynthetic reaction centres (Rochaix & Bassi 2019).

The extraordinary ability of lichens to survive harsh environments that include light stress have been proved in astrobiological experiments. Within the framework of the 'Lithopanspermia' space experiment, the lichen *Aspicilia fruticulosa* from the Guadalajara steppic highlands was exposed to open space for 10 days. While the space vacuum and cosmic radiation did not impair the metabolic activity of the lichen, solar electromagnetic radiation, especially in the wavelength range between 100 and

200 nm, reduced chlorophyll a yield fluorescence. Interestingly, however, there was a complete recovery after 72 h of reactivation (Raggio *et al.* 2011). All samples showed positive rates of net photosynthesis and dark respiration in a gas exchange experiment. The authors concluded that *A. fruticulosa* can repair any space-induced damage to its photosynthetic apparatus. In other experiments, the lichen *Xanthoria elegans* was exposed to space conditions and simulated Mars-analogue conditions for 18 months. According to the LIVE/DEAD staining results, the lichen photobiont showed an average viability rate of 71%, whereas the even more resistant lichen mycobiont showed a rate of 84% (Brandt *et al.* 2015). Successful recovery of photosynthetic activity if properly re-activated was demonstrated.

Given their typical habitats, lichens often experience multiple stresses from high light, UV exposure and desiccation (see above sections). Therefore, it is likely that cross-tolerance repair-based mechanisms operate to allow lichens to tolerate these stresses, similar to those found in the desert cyanobacterium *Chroococcidiopsis* after exposure to Mars-like UV flux and long-term desiccation. Expression analysis of the genes responsible for repair of UV-induced DNA damage, including a photolyase encoding gene (*phrA*), showed that repair of UV-induced DNA damage contributes to the repair of desiccation-induced damage (Mosca *et al.* 2019).

Chlorophagy

As outlined above, during photoinhibition chloroplasts become an active site of ROS formation. Recent studies have shown that autophagy, a process that functions in eukaryotes for the intracellular degradation of cytoplasmic components, participates in the removal of damaged chloroplasts (Nakamura & Izumi 2018). TEM images suggest that entire chloroplasts can be engulfed by the autophagosomes, which eventually fuse with the central vacuole where the chloroplasts are digested. Although not yet observed in lichen photobionts, chlorophagy-like breakdown of components of the chloroplast can occur in green microalgae (Gorelova *et al.* 2019) and involves the target of rapamycin (TOR) kinase signalling, as in other eukaryotes and higher plants (Pérez-Pérez *et al.* 2017). Future studies need to test whether similar processes occur in lichen photobionts.

Conclusions

In this review, we have outlined mechanisms that enable lichens to tolerate high light stress, including mechanisms that are important in free-living relatives of lichen photobionts but that have not yet been studied in lichens. However, some potential mechanisms are unlikely to be important in lichens. First, one potential avoidance mechanism is the Mehler reaction, in which PSI reduces molecular oxygen, leading to the formation of ATP and O₂⁻ without NADPH. The latest view is that this process saturates at relatively low irradiances and therefore, while clearly operating, probably only makes a minor contribution to avoidance (Foyer 2018). Second, chlororespiration, another potential avoidance mechanism, is a process in plant chloroplasts that involves a respiratory electron transport chain within the thylakoid membrane. Recent work suggests that chlororespiration is probably not important in protecting algae from continuous high light (Nawrocki *et al.* 2019). Interestingly, however, Nawrocki *et al.* (2019) present data suggesting that chlororespiration may prevent photoinhibition under conditions of rapidly fluctuating light. As

these conditions are exactly those that may be experienced by lichens growing under a dense canopy and deriving most of their light energy from 'sun flecks' (Coxson & Stevenson 2007), chlororespiration is a process worthy of more detailed study.


Research on high light stress in lichens is now entering an exciting phase. Even a cursory reading of the literature on light stress in plants and algae reveals that there is great interest in elucidating precise mechanisms of energy dissipation (Kaiser *et al.* 2019), and signalling pathways during high light stress (Gollan & Avo 2020). Much of this research is driven by the need to improve the performance of crop plants. Studies on lichen photobionts are just beginning but might potentially inform work on other photosynthetic organisms. For example, recent modelling studies on crop plants have suggested that slow reductions in NPQ as plants transition from high to low light may limit yield (for review see Kaiser *et al.* (2019)). Various attempts have been made to genetically engineer faster reductions in NPQ, for example by upregulating zeaxanthin epoxidase or controlling thylakoid proton gradients. However, it may be instructive to compare NPQ in lichens that grow in environments that experience extremely rapid fluctuations of light, for example 'sun-fleck' species, with those that grow in habitats with more stable light conditions. Potentially, sun-fleck species might display more rapid relaxation of NPQ. Understanding the mechanisms whereby this occurs might help studies aimed at generating more efficient crop plants.

At a broader scale, ecophysiological observations reported in the older literature can now be investigated at a more biochemical level. For example, Kershaw & MacFarlane (1980) reported that populations of *Peltigera aphthosa* collected from the dense shade of spruce are extremely sensitive to quite modest levels of light, while populations collected from open habitats are much more tolerant. It should now be possible to explain whether screening, energy dissipation, ROS scavenging or repair are responsible for the greater tolerance of a population. Future work will benefit from modern molecular biological techniques (e.g. meta-transcriptomics) that can clearly identify changes in activity of the genes of the photobiont separately from those of the mycobiont. Furthermore, it will be possible to test the relative importance of these strategies in different types of lichens growing in diverse habitats. For example, it could be predicted that the adaptations found in *Lobaria pulmonaria*, which grows in habitats where the maximum light level is *c.* 100 $\mu\text{mol m}^{-2} \text{s}^{-1}$ and is often less (Gauslaa & Solhaug 2000), will be different from those in lichens that form desert soil crust communities. More information is needed about the ability of lichens to 'harden' to changing levels of light stress (i.e. to display phenotypic plasticity). It could be predicted that high plasticity might occur in lichens that grow in habitats with regular changes in light, such as temperate corticolous lichens subjected to regular seasonal changes in solar radiation and canopy cover. By comparison, lichens growing in habitats with much less temporal variation in light, such as those from exposed sites in sub-tropical or tropical areas, could be predicted to display less ability to harden.

Finally, it is worth qualifying our suggestions, made throughout this review, that future studies on light stress in lichens will benefit from the results of studies using other photosynthetic organisms, particularly free-living algae and cyanobacteria. It is important to remember that photobionts display higher stress tolerance when part of a functioning lichen thallus than when grown as isolated cultures (Kraner *et al.* 2005), although the reasons for this increased tolerance remain unclear. Future work will need to

address exactly how symbiosis influences tolerance to high light in photobionts. Intriguingly, in a completely different symbiosis, photosynthetic sea slugs have been shown to induce protective changes to the light reactions of the chloroplasts they steal from algae (Havurinne & Tyystjärvi 2020). Conversely, lichen photobionts may be subjected to stresses absent in free-living algae. For example, they may have an increased sensitivity to photoinhibition as a lichen thallus dries slowly (Calatayud *et al.* 1997), and paradoxically also when a thallus is oversaturated as a result of limited diffusion of CO₂. Lichens are symbiotic organisms and their photobionts probably possess unique adaptations to high light stress.

Acknowledgements. RB thanks the Russian Government Program of Competitive Growth of Kazan Federal University for partial financial support. FM thanks the Russian Science Foundation (grant no. 18-14-00198) and the Russian government assignment of FRC Kazan Scientific Center of RAS for financial support. TR thanks the NRF and the OeAD (project ZA 14/2019) for funding for a bilateral agreement between Austria and South Africa. All the authors are grateful to Yngvar Gauslaa for numerous useful discussions over many years, and valuable comments on the present manuscript.

Author ORCIDs.  Richard Peter Beckett, 0000-0002-0530-4244; Farida Minibayeva, 0000-0003-0827-181X; Knut Asbjørn Solhaug, 0000-0003-3368-5205; Thomas Roach, 0000-0002-0259-0468.

References

- Ahmadjian V (1995) Lichens are more important than you think. *Bioscience* **45**, 124.
- Allorent G, Tokutsu R, Roach T, Peers G, Cardol P, Girard-Bascou J, Seigneurin-Berny D, Petroustos D, Kuntz M, Breyton C, *et al.* (2013) A dual strategy to cope with high light in *Chlamydomonas reinhardtii*. *Plant Cell* **25**, 545–557.
- Barták M, Solhaug KA, Vrábliková H and Gauslaa Y (2006) Curling during desiccation protects the foliose lichen *Lobaria pulmonaria* against photoinhibition. *Oecologia* **149**, 553–560.
- Beckett RP, Kraner I and Minibayeva F (2008) Stress physiology and the symbiosis. In Nash TH, III (ed.), *Lichen Biology*, 2nd edition. Cambridge: Cambridge University Press, pp. 134–151.
- Bergner SV, Scholz M, Trompelt K, Barth J, Gabelein P, Steinbeck J, Xue HD, Clowes S, Fucile G, Goldschmidt-Clermont M, *et al.* (2015) STATE TRANSITION7-dependent phosphorylation is modulated by changing environmental conditions, and its absence triggers remodeling of photosynthetic protein complexes. *Plant Physiology* **168**, 615–634.
- Boulay C, Abasova L, Six C, Vass I and Kirilovsky D (2008) Occurrence and function of the orange carotenoid protein in photoprotective mechanisms in various cyanobacteria. *Biochimica et Biophysica Acta* **1777**, 1344–1354.
- Brandt A, de Vera J-P, Onofri S and Ott S (2015) Viability of the lichen *Xanthoria elegans* and its symbionts after 18 months of space exposure and simulated Mars conditions on the ISS. *International Journal of Astrobiology* **14**, 411–425.
- Büchel C (2015) Evolution and function of light harvesting proteins. *Journal of Plant Physiology* **172**, 62–75.
- Büdel B and Scheidegger C (2008) Thallus morphology and anatomy. In Nash TH, III (ed.), *Lichen Biology*, 2nd edition. Cambridge: Cambridge University Press, pp. 40–68.
- Büdel B, Karsten U and Garcia-Pichel F (1997) Ultraviolet absorbing scytonemin and mycosporine-like amino acid derivatives in exposed rock-inhabiting cyanobacterial lichens. *Oecologia* **112**, 165–172.
- Buffoni Hall RS, Paulsson M, Duncan K, Tobin AK, Widell S and Bornman JF (2003) Water- and temperature-dependence of DNA damage and repair in the fruticose lichen *Cladonia arbuscula* ssp. *mitis* exposed to UV-B radiation. *Physiologia Plantarum* **118**, 371–379.
- Calatayud A, Deltoro VI, Barreno E and del Valle-Tascon S (1997) Changes in *in vivo* chlorophyll fluorescence quenching in lichen thalli as a function of water content and suggestion of zeaxanthin-associated photoprotection. *Physiologia Plantarum* **101**, 93–102.

- Calzadilla PI, Zhan J, Sétif P, Lemaire C, Solymosi D, Battchikova N, Wang Q and Kirilovsky D (2019) The cytochrome *b₆f* complex is not involved in cyanobacterial state transitions. *Plant Cell* **31**, 911–931.
- Carniel FC, Zanelli D, Bertuzzi S and Tretiach M (2015) Desiccation tolerance and lichenization: a case study with the aeroterrestrial microalga *Trebouxia* sp. (Chlorophyta). *Planta* **242**, 493–505.
- Challabathula D, Zhang QW and Bartels D (2018) Protection of photosynthesis in desiccation-tolerant resurrection plants. *Journal of Plant Physiology* **227**, 84–92.
- Cho SM, Lee H, Hong SG and Lee J (2020) Study of ecophysiological responses of the Antarctic fruticose lichen *Cladonia borealis* using the PAM fluorescence system under natural and laboratory conditions. *Plants* **9**, 85.
- Cowan IR, Lange OL and Green TGA (1992) Carbon-dioxide exchange in lichens: determination of transport and carboxylation characteristics. *Planta* **187**, 282–294.
- Coxson DS and Coyle M (2003) Niche partitioning and photosynthetic response of alecioroid lichens from subalpine spruce-fir forest in north-central British Columbia, Canada: the role of canopy microclimate gradients. *Lichenologist* **35**, 157–175.
- Coxson DS and Stevenson SK (2007) Influence of high-contrast and low-contrast forest edges on growth rates of *Lobaria pulmonaria* in the inland rainforest, British Columbia. *Forest Ecology and Management* **253**, 103–111.
- Davletova S, Rizhsky L, Liang H, Shengqiang Z, Oliver DJ, Coutu J, Shulaev V, Schlauch K, Mittler R (2005) Cytosolic ascorbate peroxidase 1 is a central component of the reactive oxygen gene network of *Arabidopsis*. *Plant Cell* **17**, 268–281.
- Demmig-Adams B, Adams WW, III, Czygan F-C, Schreiber U and Lange OL (1990) Differences in the capacity for radiationless energy dissipation in the photochemical apparatus of green and blue-green algal lichens associated with differences in carotenoid composition. *Planta* **180**, 582–589.
- Demmig-Adams B, Cohu CM, Muller O and Adams WW, III (2012) Modulation of photosynthetic energy conversion efficiency in nature: from seconds to seasons. *Photosynthesis Research* **113**, 75–88.
- Derks A, Schaven K and Bruce D (2015) Diverse mechanisms for photoprotection in photosynthesis. Dynamic regulation of photosystem II excitation in response to rapid environmental change. *Biochimica et Biophysica Acta* **1847**, 468–485.
- Dietz S, Büdel B, Lange OL and Bilger W (2000) Transmittance of light through the cortex of lichens from contrasting habitats. *Bibliotheca Lichenologica* **75**, 171–182.
- Erickson E, Wakao S and Niyogi KK (2015) Light stress and photoprotection in *Chlamydomonas reinhardtii*. *Plant Journal* **82**, 449–465.
- Ertl L (1951) Über die Lichtverhältnisse in Laubflechten. *Planta* **39**, 245–270.
- Färber L, Solhaug KA, Esseen PA, Bilger W and Gauslaa Y (2014) Sunscreening fungal pigments influence the vertical gradient of pendulous lichens in boreal forest canopies. *Ecology* **95**, 1464–1471.
- Fernandez-Marin B, Kranner I, San Sebastian M, Artetxe U, Laza JM, Vilas JL, Pritchard HW, Nadajaran J, Miguez F, Becerril JM, *et al.* (2013) Evidence for the absence of enzymatic reactions in the glassy state. A case study of xanthophyll cycle pigments in the desiccation-tolerant moss *Syntrichia ruralis*. *Journal of Experimental Botany* **64**, 3033–3043.
- Foyer CH (2018) Reactive oxygen species, oxidative signaling and the regulation of photosynthesis. *Environmental and Experimental Botany* **154**, 134–142.
- Franz S, Ignatz E, Wenzel S, Zielosko H, Putu EPGN, Maestre-Reyna M, Tsai MD, Yamamoto J, Mittag M and Essen LO (2018) Structure of the bifunctional cryptochrome aCRY from *Chlamydomonas reinhardtii*. *Nucleic Acids Research* **46**, 8010–8022.
- Garcia-Pichel F and Castenholz RW (1991) Characterization and biological implications of scytonemin, a cyanobacterial sheath pigment. *Journal of Phycology* **27**, 395–409.
- Gauslaa Y (1984) Heat resistance and energy budget in different Scandinavian plants. *Holarctic Ecology* **7**, 1–78.
- Gauslaa Y and Goward T (2020) Melanic pigments and canopy-specific elemental concentration shape growth rates of the lichen *Lobaria pulmonaria* in unmanaged mixed forest. *Fungal Ecology* **47**, 100984.
- Gauslaa Y and Solhaug KA (1996) Differences in the susceptibility to light stress between epiphytic lichens of ancient and young boreal forest stands. *Functional Ecology* **10**, 344–354.
- Gauslaa Y and Solhaug KA (2000) High-light-intensity damage to the foliose lichen *Lobaria pulmonaria* within a natural forest: the applicability of chlorophyll fluorescence methods. *Lichenologist* **32**, 271–289.
- Gauslaa Y and Solhaug KA (2001) Fungal melanins as a sun screen for symbiotic green algae in the lichen *Lobaria pulmonaria*. *Oecologia* **126**, 462–471.
- Gauslaa Y, Coxson DS and Solhaug KA (2012) The paradox of higher light tolerance during desiccation in rare old forest cyanolichens than in more widespread co-occurring chloro- and cephalolichens. *New Phytologist* **195**, 812–822.
- Gauslaa Y, Alam MA, Lucas P-L, Chowdhury DP and Solhaug KA (2017) Fungal tissue *per se* is stronger as a UV-B screen than secondary fungal extrolites in *Lobaria pulmonaria*. *Fungal Ecology* **26**, 109–113.
- Gest N, Gautier H and Stevens R (2013) Ascorbate as seen through plant evolution: the rise of a successful molecule? *Journal of Experimental Botany* **64**, 33–53.
- Girolomoni L, Cazzaniga S, Pinnola A, Perozeni F, Ballottari M and Bassi R (2019) LHCSR3 is a nonphotochemical quencher of both photosystems in *Chlamydomonas reinhardtii*. *Proceedings of the National Academy of Sciences of the United States of America* **116**, 4212–4217.
- Gollan PJ and Aro E-M (2020) Photosynthetic signalling during high light stress and recovery: targets and dynamics. *Philosophical Transactions of the Royal Society B* **375**, 20190406.
- Gorelova O, Baulina O, Ismagulova T, Kokabi K, Lobakova E, Selyakh I, Semenova L, Chivkunova O, Karpova O, Scherbakov P, *et al.* (2019) Stress-induced changes in the ultrastructure of the photosynthetic apparatus of green microalgae. *Protoplasma* **256**, 261–277.
- Gostinčar C, Muggia L and Grube M (2012) Polyextremotolerant black fungi: oligotrophism, adaptive potential, and a link to lichen symbioses. *Frontiers in Microbiology* **3**, 390.
- Green TGA, Büdel B, Meyer A, Zellner H and Lange OL (1997) Temperate rainforest lichens in New Zealand: light response of photosynthesis. *New Zealand Journal of Botany* **35**, 493–504.
- Haghjoui MM, Shariati M and Smirnov N (2009) The effect of acute high light and low temperature stresses on the ascorbate-glutathione cycle and superoxide dismutase activity in two *Dunaliella salina* strains. *Physiologia Plantarum* **135**, 272–280.
- Hasanuzzaman M, Borhannuddin Bhuyan MHM, Anee TI, Parvin K, Nahar K, Al Mahmud J and Fujita M (2019) Regulation of ascorbate-glutathione pathway in mitigating oxidative damage in plants under abiotic stress. *Antioxidants* **8**, 384.
- Havaux M, Guedeney G, Hagemann M, Yermenko N, Matthijs HCP and Jeanjean R (2005) The chlorophyll-binding protein IsiA is inducible by high light and protects the cyanobacterium *Synechocystis* PCC6803 from photooxidative stress. *FEBS Letters* **579**, 2289–2293.
- Havurinne V and Tyystjärvi E (2020) Photosynthetic sea slugs induce protective changes to the light reactions of the chloroplasts they steal from algae. *Elife* **9**, e57389.
- Heber U (2012) Conservation and dissipation of light energy in desiccation-tolerant photoautotrophs, two sides of the same coin. *Photosynthesis Research* **113**, 5–13.
- Heber U, Bilger W, Bligny R and Lange OL (2000) Phototolerance of lichens, mosses and higher plants in an alpine environment: analysis of photoreactions. *Planta* **211**, 770–780.
- Heber U, Bilger W and Shuvalov VA (2006) Thermal energy dissipation in reaction centres and in the antenna of photosystem II protects desiccated poikilohydric mosses against photo-oxidation. *Journal of Experimental Botany* **57**, 2993–3006.
- Heber U, Bilger W, Turk R and Lange OL (2010) Photoprotection of reaction centres in photosynthetic organisms: mechanisms of thermal energy dissipation in desiccated thalli of the lichen *Lobaria pulmonaria*. *New Phytologist* **185**, 459–470.
- Huneck S and Yoshimura I (1996) *Identification of Lichen Substances*. Berlin: Springer-Verlag.
- Huner NPA, Öquist G and Sarhan F (1998) Energy balance and acclimation to light and cold. *Trends in Plant Science* **3**, 224–230.
- Jairus K, Löhmus A and Löhmus P (2009) Lichen acclimatization on retention trees: a conservation physiology lesson. *Journal of Applied Ecology* **46**, 930–936.

- Jeans J, Campbell DA and Hoogenboom MO (2013) Increased reliance upon photosystem II repair following acclimation to high-light by coral-dinoflagellate symbioses. *Photosynthesis Research* **118**, 219–229.
- Kaiser E, Galvis VC and Armbruster U (2019) Efficient photosynthesis in dynamic light environments: a chloroplast's perspective. *Biochemical Journal* **476**, 2725–2741.
- Kalaji HM, Schansker G, Ladle RJ, Goltsev V, Bosa K, Allakhverdiev SI, Brestic M, Bussott F, Calatayud A, Dabrowski P, *et al.* (2014) Frequently asked questions about *in vivo* chlorophyll fluorescence: practical issues. *Photosynthesis Research* **122**, 121–158.
- Kerfeld CA, Melnicki MR, Sutter M and Dominguez-Martin MA (2017) Structure, function and evolution of the cyanobacterial orange carotenoid protein and its homologs. *New Phytologist* **215**, 937–951.
- Kershaw KA (1985) *Physiological Ecology of Lichens*. Cambridge: Cambridge University Press.
- Kershaw KA and MacFarlane JD (1980) Physiological-environmental interactions in lichens. X. Light as an ecological factor. *New Phytologist* **84**, 687–701.
- Kharkongor D and Ramanujam P (2015) Spatial and temporal variation of carotenoids in four species of *Trentepohlia* (*Trentepohliales*, Chlorophyta). *Journal of Botany* **2015**, 201641.
- Kim JH, Nemson JA and Melis A (1993) Photosystem II reaction center damage and repair in *Dunaliella salina* (green alga). Analysis under physiological and irradiance-stress conditions. *Plant Physiology* **103**, 181–189.
- Kitajima S (2008) Hydrogen peroxide-mediated inactivation of two chloroplastic peroxidases, ascorbate peroxidase and 2-cys peroxidoxin. *Journal of Photochemistry and Photobiology B* **84**, 1404–1409.
- Komenda J and Sobotka R (2016) Cyanobacterial high-light-inducible proteins — protectors of chlorophyll-protein synthesis and assembly. *Biochimica et Biophysica Acta (BBA) – Bioenergetics* **1857**, 288–295.
- Komura M, Yamagishia A, Shibata Y, Iwasaki I and Itoha S (2010) Mechanism of strong quenching of photosystem II chlorophyll fluorescence under drought stress in a lichen, *Physciella melanchla*, studied by subpicosecond fluorescence spectroscopy. *Bioenergetics* **1797**, 331–338.
- Kosanić M, Ranković B and Vukojević J (2011) Antioxidant properties of some lichen species. *Journal of Food Science and Technology* **48**, 584–590.
- Kranner I (2002) Glutathione status correlates with different degrees of desiccation tolerance in three lichens. *New Phytologist* **154**, 451–460.
- Kranner I, Zorn M, Turk B, Wornik S, Beckett RP and Batic F (2003) Biochemical traits of lichens differing in relative desiccation tolerance. *New Phytologist* **160**, 167–176.
- Kranner I, Cram WJ, Zorn M, Wornik S, Yoshimura I, Stabentheiner E and Pfeifhofer HW (2005) Antioxidants and photoprotection in a lichen as compared with its isolated symbiotic partners. *Proceedings of the National Academy of Sciences of the United States of America* **102**, 3141–3146.
- Kromdijk J, Glowacka K, Leonelli L, Gabilly ST, Iwai M, Niyogi KK and Long SP (2016) Improving photosynthesis and crop productivity by accelerating recovery from photoprotection. *Science* **354**, 857–861.
- Lange OL (2002) Photosynthetic productivity of the epilithic lichen *Lecanora muralis*: long-term field monitoring of CO₂ exchange and its physiological interpretation. I. Dependence of photosynthesis on water content, light, temperature, and CO₂ concentration from laboratory measurements. *Flora* **197**, 233–249.
- Lange OL and Green TGA (2008) Diel and seasonal courses of ambient carbon dioxide concentration and their effect on productivity of the epilithic lichen *Lecanora muralis* in a temperate, suburban habitat. *Lichenologist* **40**, 449–462.
- Leisner JMR, Green TGA and Lange OL (1997) Photobiont activity of a temperate crustose lichen: long-term chlorophyll fluorescence and CO₂ exchange measurements in the field. *Symbiosis* **23**, 165–182.
- Levitt J (2012) *Responses of Plants to Environmental Stresses. Volume I. Chilling, Freezing and High Temperature*, 2nd edition. London: Academic Press.
- Li L, Aro EM and Millar AH (2018) Mechanisms of photodamage and protein turnover in photoinhibition. *Trends in Plant Science* **23**, 667–676.
- Lin T, Rao M, Lu H, Chiou C, Lin S, Chao H, Zheng L, Cheng H and Lee T (2018) A role for glutathione reductase and glutathione in the tolerance of *Chlamydomonas reinhardtii* to photo-oxidative stress. *Physiologia Plantarum* **162**, 35–48.
- Liu J, Lu Y, Hua W and Last RL (2019) A new light on photosystem II maintenance in oxygenic photosynthesis. *Frontiers in Plant Science* **10**, 975.
- Mafole TC, Chiang C, Solhaug KA and Beckett RP (2017) Melanisation in the old forest lichen *Lobaria pulmonaria* (L) Hoffm. reduces the efficiency of photosynthesis. *Fungal Ecology* **29**, 103–110.
- Mafole TC, Solhaug KA, Minibayeva FV and Beckett RP (2019a) Occurrence and possible roles of melanic pigments in lichenized ascomycetes. *Fungal Biology Reviews* **33**, 159–165.
- Mafole TC, Solhaug KA, Minibayeva FV and Beckett RP (2019b) Tolerance to photoinhibition within lichen species is higher in melanised thalli. *Photosynthetica* **57**, 96–102.
- Marečková M and Barták M (2016) Effects of short-term low temperature stress on chlorophyll fluorescence transients in Antarctic lichen species. *Czech Polar Reports* **6**, 54–65.
- Maruta T, Sawa Y, Shigeoka S and Ishikawa T (2016) Diversity and evolution of ascorbate peroxidase functions in chloroplasts: more than just a classical antioxidant enzyme? *Plant and Cell Physiology* **57**, 1377–1386.
- McEvoy M, Gauslaa Y and Solhaug KA (2007a) Changes in pools of depositions and melanins, and their function, during growth and acclimation under contrasting natural light in the lichen *Lobaria pulmonaria*. *New Phytologist* **175**, 271–282.
- McEvoy M, Solhaug KA and Gauslaa Y (2007b) Solar radiation screening in usnic acid-containing cortices of the lichen *Nephroma arcticum*. *Symbiosis* **43**, 143–150.
- Michelet L, Roach T, Fischer, BB, Bedhomme M, Lemaire SD and Krieger-Liszskay A (2013) Down-regulation of catalase activity allows transient accumulation of a hydrogen peroxide signal in *Chlamydomonas reinhardtii*. *Plant Cell and Environment* **36**, 1204–1213.
- Míguez F, Fernández-Marín B, Becerril J-M and García-Plazaola JI (2017a) Diversity of winter photoinhibitory responses: a case study in co-occurring lichens, mosses, herbs and woody plants from subalpine environments. *Physiologia Plantarum* **160**, 282–296.
- Míguez F, Schiefelbein U, Karsten U, García-Plazaola JI and Gustavs L (2017b) Unraveling the photoprotective response of lichenized and free-living green algae (Trebouxiphyceae, Chlorophyta) to photoinhibition stress. *Frontiers in Plant Science* **8**, 1144.
- Mishra A, Hájek J, Tuháčková T and Barták M (2015) Features of chlorophyll fluorescence transients can be used to investigate low temperature induced effects on photosystem II of algal lichens from polar regions. *Czech Polar Reports* **5**, 99–111.
- Miyake C, Michihata F and Asada K (1991) Scavenging of hydrogen peroxide in prokaryotic and eukaryotic algae: acquisition of ascorbate peroxidase during the evolution of cyanobacteria. *Plant and Cell Physiology* **32**, 33–43.
- Molnar K and Farkas E (2010) Current results on biological activities of lichen secondary metabolites: a review. *Zeitschrift für Naturforschung Section C* **65**, 157–173.
- Mosca C, Rothschild LJ, Napoli A, Ferré F, Pietrosanto M, Fagiarone C, Baqué M, Rabbow E, Rettberg P and Billi D (2019) Over-expression of UV-damage DNA repair genes and ribonucleic acid persistence contribute to the resilience of dried biofilms of the desert cyanobacterium *Chroococcidiopsis* exposed to Mars-like UV flux and long-term desiccation. *Frontiers in Microbiology* **10**, 2312.
- Müller P, Li X-P and Niyogi KK (2001) Non-photochemical quenching. A response to excess light energy. *Plant Physiology* **125**, 1558–1566.
- Nakamura S and Izumi M (2018) Regulation of chlorophagy during photoinhibition and senescence: lessons from mitophagy. *Plant and Cell Physiology* **59**, 1135–1143.
- Nath K, Jajoo A, Poudyal RS, Timilsina R, Park YS, Aro EM, Nam HG and Lee CH (2013) Towards a critical understanding of the photosystem II repair mechanism and its regulation during stress conditions. *FEBS Letters* **587**, 3372–3381.
- Nawrocki WJ, Buchert F, Joliet P, Rappaport F, Bailleul B and Wollman FA (2019) Chlororespiration controls growth under intermittent light. *Plant Physiology* **179**, 630–639.
- Niyogi KK and Truong TB (2013) Evolution of flexible non-photochemical quenching mechanisms that regulate light harvesting in oxygenic photosynthesis. *Current Opinion in Plant Biology* **16**, 307–314.

- Noctor G, Mhamdi A and Foyer C (2014) The roles of reactive oxygen metabolism in drought: not so cut and dried. *Plant Physiology* **164**, 1636–1648.
- Ohnishi N, Allakhverdiev SI, Takahashi S, Higashi S, Watanabe M, Nishiyama Y and Murata N (2005) Two-step mechanism of photodamage to photosystem II: step 1 occurs at the oxygen-evolving complex and step 2 occurs at the photochemical reaction center. *Biochemistry* **44**, 8494–8499.
- Öquist G and Huner NPA (2003) Photosynthesis of overwintering evergreen plants. *Annual Review of Plant Biology* **54**, 329–355.
- Osmond B, Badger M, Maxwell K, Bjorkman O and Leegood R (1997) Too many photos: photorespiration, photoinhibition and photooxidation. *Trends in Plant Science* **2**, 119–121.
- Pandey P, Singh J, Achary VMM and Reddy MK (2015) Redox homeostasis via gene families of ascorbate-glutathione pathway. *Frontiers in Environmental Science* **3**, 25.
- Peers G, Truong TB, Ostendorf E, Busch A, Elrad D, Grossman AR, Hippler M and Niyogi KK (2009) An ancient light-harvesting protein is critical for the regulation of algal photosynthesis. *Nature* **462**, 518–521.
- Pérez-Pérez ME, Couso I and Crespo JL (2017) The TOR signaling network in the model unicellular green alga *Chlamydomonas reinhardtii*. *Biomolecules* **7**, 54.
- Phinney NH, Gauslaa Y and Solhaug KA (2019) Why chartreuse? The pigment vulpinic acid screens blue light in the lichen *Letharia vulpina*. *Planta* **249**, 709–718.
- Piccotto M and Tretiach M (2010) Photosynthesis in chlorolichens: the influence of the habitat light regime. *Journal of Plant Research* **123**, 763–775.
- Pospišil P (2016) Production of reactive oxygen species by photosystem II as a response to light and temperature stress. *Frontiers in Plant Science* **7**, 1950.
- Raggio J, Pintado A, Ascaso C, de la Torre R, de los Ríos A, Wierzbos J, Horneck G and Sancho LG (2011) Whole lichen thalli survive exposure to space conditions: results of Lithopanspermia experiment with *Aspicilia fruticulosa*. *Astrobiology* **11**, 281–292.
- Roach T and Krieger-Liszka A (2019) Photosynthetic regulatory mechanisms for efficiency and prevention of photo-oxidative stress. *Annual Plant Reviews Online* **2**, 273–306.
- Roach T and Na CS (2017) LHCSR3 affects de-coupling and re-coupling of LHCII to PSII during state transitions in *Chlamydomonas reinhardtii*. *Scientific Reports* **7**, 43145.
- Roach T, Na CS and Krieger-Liszka A (2015) High light-induced hydrogen peroxide production in *Chlamydomonas reinhardtii* is increased by high CO₂ availability. *Plant Journal* **81**, 759–766.
- Roach T, Baur T, Stöggel W and Krieger-Liszka A (2017) *Chlamydomonas reinhardtii* responding to high light: a role for 2-propenal (acrolein). *Physiologia Plantarum* **161**, 75–87.
- Roach T, Stöggel W, Baur T and Kranner I (2018) Distress and eustress of reactive electrophiles and relevance to light stress acclimation via stimulation of thiol/disulphide-based redox defences. *Free Radical Biology and Medicine* **122**, 65–73.
- Roach T, Na CS, Stöggel W and Krieger-Liszka A (2020) The non-photochemical quenching protein LHCSR3 prevents oxygen-dependent photoinhibition in *Chlamydomonas reinhardtii*. *Journal of Experimental Botany* **71**, 2650–2660.
- Rochaix J-D (2011) Regulation of photosynthetic electron transport. *Biochimica et Biophysica Acta* **1807**, 375–383.
- Rochaix J-D and Bassi R (2019) LHC-like proteins involved in stress responses and biogenesis/repair of the photosynthetic apparatus. *Biochemical Journal* **476**, 581–593.
- Sahu N, Singh SN, Singh P, Mishra S, Karakoti N, Bajpai R, Behera SK, Nayaka S and Upreti DK (2019) Microclimatic variations and their effects on photosynthetic efficiencies and lichen species distribution along elevational gradients in Garhwal Himalayas. *Biodiversity and Conservation* **28**, 1953–1976.
- Sathasivam R and Jang-Seu K (2018) A review of the biological activities of microalgal carotenoids and their potential use in healthcare and cosmetic industries. *Marine Drugs* **16**, 26.
- Sedoud A, López-Igual R, Ur Rehman A, Wilson A, Perreau F, Boulay C, Vass I, Krieger-Liszka A and Kirilovsky D (2014) The cyanobacterial photoactive orange carotenoid protein is an excellent singlet oxygen quencher. *Plant Cell* **26**, 1781–1791.
- Serrano A and Lloell A (1993) Occurrence of 2 isoforms of glutathione-reductase in the green alga *Chlamydomonas reinhardtii*. *Planta* **190**, 199–205.
- Slavov C, Reus M and Holzwarth AR (2013) Two different mechanisms cooperate in the desiccation-induced excited state quenching in *Parmelia* lichen. *Journal of Physical Chemistry B* **117**, 11326–11336.
- Smirnov N (2018) Ascorbic acid metabolism and functions: a comparison of plants and mammals. *Free Radical Biology and Medicine* **122**, 116–129.
- Solhaug KA and Gauslaa Y (1996) Parietin, a photoprotective secondary product of the lichen *Xanthoria parietina*. *Oecologia* **108**, 412–418.
- Solhaug KA and Gauslaa Y (2012) Secondary lichen compounds as protection against excess solar radiation and herbivores. *Progress in Botany* **73**, 283–304.
- Solhaug KA, Gauslaa Y, Nybakken L and Bilger W (2003) UV-induction of sun-screening pigments in lichens. *New Phytologist* **158**, 91–100.
- Solhaug KA, Larsson P and Gauslaa Y (2010) Light screening in lichen cortices can be quantified by chlorophyll fluorescence techniques for both reflecting and absorbing pigments. *Planta* **231**, 1003–1011.
- Stålfelt MG (1938) Der Gasaustausch der Flechten. *Planta* **29**, 11–31.
- Štepičová J, Gauslaa Y, Cempírková-Vráblíková H and Solhaug KA (2008) Irradiance prior to and during desiccation improves the tolerance to excess irradiance in the desiccated state of the old forest lichen *Lobaria pulmonaria*. *Photosynthetica* **46**, 286–290.
- Streb P, Feierabend J and Bligny R (1997) Resistance to photoinhibition of photosystem II and catalase and antioxidative protection in high mountain plants. *Plant, Cell and Environment* **20**, 1030–1040.
- Takeda T, Ishikawa T and Shigeoka S (1997) Metabolism of hydrogen peroxide by the scavenging system in *Chlamydomonas reinhardtii*. *Physiologia Plantarum* **99**, 49–55.
- Tel-Or E, Huflejt M and Packer L (1985) The role of glutathione and ascorbate in hydroperoxide removal in cyanobacteria. *Biochemical and Biophysical Research Communications* **132**, 533–539.
- Thadhani VM, Choudhary MI, Ali S, Omar I, Siddique H and Karunaratne V (2011) Antioxidant activity of some lichen metabolites. *Natural Product Research* **25**, 1827–1837.
- Triantaphylidēs C, Krischke M, Hoerberichts FA, Ksas B, Gresser G, Havaux M, Van Breusegem F and Mueller MJ (2008) Singlet oxygen is the major reactive oxygen species involved in photooxidative damage to plants. *Plant Physiology* **148**, 960–968.
- Vass I-Z, Kós PB, Sass L, Nagy CI and Vass I (2013) The ability of cyanobacterial cells to restore UV-B radiation induced damage to Photosystem II is influenced by photolyase dependent DNA repair. *Photochemistry and Photobiology* **89**, 384–390.
- Vass I-Z, Kós PB, Knoppová J, Komenda J and Vass I (2014) The cry-DASH cryptochrome encoded by the *sl11629* gene in the cyanobacterium *Synechocystis* PCC 6803 is required for Photosystem II repair. *Journal of Photochemistry and Photobiology B: Biology* **130**, 318–326.
- Verhoeven A, Garcia-Plazaola JI and Fernandez-Marin B (2018) Shared mechanisms of photoprotection in photosynthetic organisms tolerant to desiccation or to low temperature. *Environmental and Experimental Botany* **154**, 66–79.
- Vidal-Meireles A, Tóth D, Kovács L, Neupert J and Tóth SZ (2020) Ascorbate deficiency does not limit nonphotochemical quenching in *Chlamydomonas reinhardtii*. *Plant Physiology* **182**, 597–611.
- Vogelmann TC (1993) Plant-tissue optics. *Annual Review of Plant Physiology and Plant Molecular Biology* **44**, 231–251.
- Vráblíková H, McEvoy M, Solhaug KA, Barták M and Gauslaa Y (2006) Annual variation in photoacclimation and photoprotection of the photobiont in the foliose lichen *Xanthoria parietina*. *Journal of Photochemistry and Photobiology B: Biology* **83**, 151–162.
- Vráblíková H, Barták M and Wonisch A (2005) Changes in glutathione and xanthophyll cycle pigments in the high light-stressed lichens *Umbilicaria antarctica* and *Lasallia pustulata*. *Journal of Photochemistry and Photobiology B: Biology* **79**, 35–41.
- Weissman L, Garty J and Hochman A (2005) Characterization of enzymatic antioxidants in the lichen *Ramalina lacera* and their response to rehydration. *Applied and Environmental Microbiology* **71**, 6508–6514.

- Wieners PC, Mudimu O and Bilger W** (2018) Survey of the occurrence of desiccation-induced quenching of basal fluorescence in 28 species of green microalgae. *Planta* **248**, 601–612.
- Williams L, Colesie C, Ullmann A, Westberg M, Wedin M and Büdel B** (2017) Lichen acclimation to changing environments: photobiont switching vs. climate-specific uniqueness in *Psora decipiens*. *Ecology and Evolution* **7**, 2560–2574.
- Wilson A, Punginelli C, Gall A, Bonetti C, Alexandre M, Routaboul JM, Kerfeld CA, van Grondelle R, Robert B, Kennis JTM, et al.** (2008) A photoactive carotenoid protein acting as light intensity sensor. *Proceedings of the National Academy of Sciences of the United States of America* **105**, 12075–12080.
- Wolfe-Simon F, Grzebyk D, Schofield O and Falkowski PG** (2005) The role and evolution of superoxide dismutases in algae. *Journal of Phycology* **41**, 453–465.
- Yamamoto H and Higashi R** (1978) Violaxanthin de-epoxidase: lipid composition and substrate specificity. *Archives of Biochemistry and Biophysics* **190**, 514–522.
- Zavafer A, Chow WS and Cheah MH** (2015) The action spectrum of Photosystem II photoinactivation in visible light. *Journal of Photochemistry and Photobiology B: Biology* **152**, 247–260.