

## Correspondence

### ATTEMPTED SUICIDE

DEAR SIR,

The offer by Dr. Kreitman and his colleagues of 'parasuicide' as an alternative to 'attempted suicide' (*Journal*, June 1969, pp. 746-747) raises more problems than it solves. I think firstly that they are wrong in suggesting without qualification that 'the great majority of patients so designated are not in fact attempting suicide'. The acceptance by these patients of the risk of death is too definite to be dismissed in this way, even if the proportion at serious risk is possibly a minority. When Stengel and Cook showed that attempted suicide was not simply failed suicide and nothing else, they emphasized the complexity of motivations both in the suicide and the attempted suicide. Indeed, as Kreitman *et al.* rightly point out in rejecting the unfortunate term 'self-poisoning', there is a very real association between 'attempted suicide' and 'completed suicide'. The move to 'parasuicide' seems to jettison too much from this view.

Secondly, it would be misleading to regard attempted suicide as a simulation of suicide. If 'parasuicide' is meant to suggest an act of mimicry, it presumably implies simulation intended by the patient, and the total absence of serious self-damaging motivation. For how many cases can this be acceptably proved? Both the patient and the psychiatrist would have to agree the conclusion, and at least some psychiatrists find this situation relatively rare in their routine work with patients who have attempted suicide.

Thirdly, if we are to have categories both of attempted suicide and parasuicide I suspect this will lead to considerable uncertainty in clinical practice and meaningless results in epidemiology.

Lastly and least important, 'para' is of Greek etymology, and 'suicide' Latin. Some would object to mixing roots from different languages. I would not myself think that an important objection if the term helped in other respects, but it does not seem to do so. It may indeed secure about as much agreement as 'para-psychology'.

H. MERSKEY.

*The National Hospital for Nervous Diseases,  
Queen Square, London, W.C.1.*

#### REFERENCES

- STENDEL, E. and COOK, N. G. (1958). 'Attempted Suicide: Social Significance and Effects.' Maudsley Monograph. London.

### COMMUNITY PSYCHIATRY

DEAR SIR,

I am stimulated by the proud note struck in Dr. Maurice Silverman's letter (*Journal*, July 1969, p. 863) to reflect on the controversy which has persisted over the past decade regarding the number of beds required to provide an adequate psychiatric service for any given population. The trend, like that of the female hemline, is clearly towards the creation of the mini-unit. In the absence of major therapeutic advances or of improved manpower during this time, such reductions often have to be achieved through the ingenuity of the psychiatrist, by substituting chairs for beds in his day hospital, redefining the specialty within narrower limits, or restricting the demand by educating the community to tolerate higher thresholds of misery within their midst.

With appropriate acknowledgement, may I offer for your readers' consideration the formulation of a new law which states that bed-space contracts so as to decrease the patients available to fill it. Perhaps while waiting for a biochemical breakthrough we have discovered the final solution to the problem of psychiatric illness.

I am led to the conclusion that there are other factors influencing the organization of medical care besides the needs of the patient. If some of these are the needs of the doctor it would be interesting to investigate what these are, and why in the psychiatrist's case there should be such a powerful drive to reduce the number of beds under his care.

SIDNEY LEVINE

*Oldham and District General Hospital,  
Rochdale Road,  
Oldham, Lancs.*

### TRAINING OF PSYCHIATRISTS

DEAR SIR,

Dr. Fraser (*Journal*, August 1969, p. 979) would have a good point in favour of some subnormals being the concern of psychiatrists if, in fact, they were only referred for 'deviant behaviour or abnormal psychological development' and were discharged as soon as these had been corrected. In practice,

I doubt whether this is true in Fife or in many other areas; when it obtains there seems little point in the main residential or community provisions for the mentally retarded remaining under continuous psychiatric (or medical) supervision. In the meantime, it appears that we are conducting a rather expensive holding operation.

T. L. PILKINGTON.

*Edinburgh University Department of Psychiatry,  
Morningside Park,  
Edinburgh EH10 5HF.*

#### NURSES FOR CHILDREN'S UNITS

DEAR SIR,

The setting up of in-patient units for children and adolescents is creating an increasing demand for staff with special experience in the management and treatment of disturbed children. At present there is no specific training or career prospects for such staff. The qualification for psychiatric nurses is obtained by three years' training in a mental hospital with adult patients, and for paediatric nurses by three years' training in a general hospital, working with children with severe physical illnesses, the majority of whom will be infants. Neither of these courses of training prepare staff for work with disturbed children or adolescents, and they may even lead to methods and attitudes which will be disadvantageous. People with an aptitude for working with disturbed children do not necessarily have an aptitude for, or interest, in working with adult patients or physically ill children, and there is no reason why they should. The prospect of three years' training in a course which has very little bearing on the work in which they are interested may put off a large proportion of very good candidates.

At present the grade of Assistant Nurse is all that we can offer people with no nursing qualifications, regardless of what their qualifications are in other fields. Our advertisements for assistants in my Children's In-patient Unit are always answered by

a large number of people, and the applicants often include people with teaching qualifications and University degrees. Such people are sometimes prepared to work for a year as an Assistant Nurse, because of the valuable experience, but one cannot expect anybody who is good enough to do the work to remain as an Assistant Nurse for long. The result is that these excellent people with a genuine aptitude for the work have to leave, either to train in nursing or to go into some other field of work with children, such as teaching.

In my view a suitable pay structure and career should be available for people wishing to work in residential care of emotionally and behaviourally disturbed children. The qualifications following a suitable period of training should be recognized in all fields of residential work, hospital, approved schools and children's departments, and should lead to good pay and career prospects. I suggest that our Association might give urgent attention to the problems of establishing courses, qualifications and career structure for this highly specialized work.

CHRISTOPHER J. WARDLE.

*The Dryden Clinic,  
Exe Vale Hospital,  
Dryden Road, Wonford, Exeter.*

#### A REQUEST

DEAR SIR,

The Research and Clinical Section would like to present at their Open Meetings research which is either in progress or completed but not yet published. Anyone wishing to present a paper at the Quarterly Meetings of the R.M.P.A. is invited to write to the Clinical Secretary of this Section, Dr. B. M. Barraclough, at this address, providing a short summary of the method and results.

PETER SAINSBURY.

*Chairman of the Executive Committee of  
the Research and Clinical Section.  
MRC Clinical Psychiatry Research Unit,  
Graylingwell Hospital, Chichester, Sussex.*