

Home ranges of Guiana dolphins (*Sotalia guianensis*) (Cetacea: Delphinidae) in the Cananéia estuary, Brazil

JÚLIA EMI DE FARIA OSHIMA¹, MARCOS CÉSAR DE OLIVEIRA SANTOS¹, MARIEL BAZZALO²,
PAULO ANDRÉ DE CARVALHO FLORES³ AND FABIANO DO NASCIMENTO PUPIM⁴

¹Projeto Atlantis, Laboratório de Biologia da Conservação de Cetáceos, Departamento de Zoologia, Instituto de Biociências da Universidade Estadual Paulista ‘Júlio de Mesquita Filho’ (UNESP), Campus Rio Claro, Avenida 24-A, 1515, Bela Vista, Rio Claro, SP, 13506-900, Brazil, ²Carrera de Doctorado, Universidad de Buenos Aires, Argentina, ³Centro Nacional de Pesquisa e Conservação de Mamíferos Aquáticos—CMA, ICMBio, Rod. Mauricio Sirotsky Sobrinho, s/n, km 02, Jurerê, Florianópolis, SC, 88053-700, Brazil, ⁴Programa de Pós-Graduação em Geociências e Meio Ambiente, IGCE, Universidade Estadual Paulista ‘Júlio de Mesquita Filho’ (UNESP), Rio Claro, Avenida 24-A, 1515, Bela Vista, Rio Claro, SP, 13506-900, Brazil

Home ranges of seven Guiana dolphins (Sotalia guianensis) (Van Bénédén, 1864) were studied in the Cananéia estuary (~25°03'S 47°55'W), south-eastern Brazil. Boat-based observations were conducted from May 2000 to July 2003 in ~132 km² of protected inner waters. The photo-identification technique was used to follow naturally marked individuals through time and space. From a total of 138 catalogued individuals, five males and two females presented 20+ sightings and were used for home range estimation. Sightings were plotted and analysed in a Geographic Information System (GIS). With the ‘Home Range Tools’ extension the fixed kernel density estimator with band width (smoothing parameter) chosen via least squares cross-validation was performed for each individual. The fixed kernel method was used to estimate the non-parametric utility distribution of each dolphin, keeping band width (h) constant for a data set. The first polygons created by these parameters had an amoeboid shape and in some cases more than one centre of activity. The 95% home range estimated outlines varied from 1.6 to 22.9 km² (7.9 ± 8.3 km²). This large interval shows strong evidences on individual variation in S. guianensis’ home ranges. Several individuals showed small home ranges when compared to other cetacean species. An overlap of home ranges of different sizes and shapes were observed for Guiana dolphins with large range movements. Centres of activity were concentrated in the main entrance of the Cananéia estuary. This was a first attempt to understand the way S. guianensis uses the Cananéia estuary and such data are essential for conservation and management purposes.

Keywords: *Sotalia guianensis*, home range, habitat use, kernel density estimator

Submitted 26 January 2010; accepted 14 June 2010; first published online 1 September 2010

INTRODUCTION

Home range was described by Burt (1943) as ‘the area traversed by the individual in its normal activities of food gathering, mating, and caring of young’. Recently, Powell (2000) defined home range as a ‘fairly confined area where animals enact their day-to-day activities’. The size of an animal’s home range is usually correlated with habitat heterogeneity, productivity and the biological requirements of a species (McNab, 1963). Most animals do not use their entire home range with equal intensity, but tend to concentrate their time in particular areas (Dixon & Chapman, 1980; Samuel *et al.*, 1985). These particular sectors of an animal’s home range are called core areas and usually are associated with greater density of resource. As a consequence, they will be used more than expected by random use (e.g. Samuel *et al.*, 1985; Powell, 2000).

Methods to estimate home ranges have been used for decades. From location data, most home range estimators produce a utility distribution (UD) describing the intensity of use of different areas by an animal (Powell, 2000). Thus, the UD maps intensity of use and can be transformed to a probabilistic model of home range which describes the relative amount of time an animal spends in any place (Seaman & Powell, 1996; Powell, 2000). Both parametric and non-parametric methods have been used to estimate the UD (Worton, 1989). The kernel density estimator (see Silverman, 1986), introduced to ecologists as a home range estimator by Worton (1989), is one of the best non-parametric statistical methods to estimate probability densities because of its robustness and large applicability (Seaman & Powell, 1996; Seaman *et al.*, 1999; Powell, 2000).

A considerable amount of studies regarding cetacean use of area are based on the application of the photo-identification technique (Würsig & Würsig, 1977; Hammond *et al.*, 1990), which allows the identification of individuals through time and space. Naturally marked individuals provide important information on cetacean ranging patterns (e.g. Connor & Smolker, 1985; Wells *et al.*, 1987; Bigg *et al.*, 1990; Würsig

Corresponding author:
J.E. Oshima
Email: juliaoshima@yahoo.com.br

& Harris, 1990; Herzing, 1997; Marten & Psarakos, 1999; Santos *et al.*, 2001). Flores & Bazzalo (2004) were the first to estimate home ranges of Guiana dolphins, *Sotalia guianensis* (Van Bénédén, 1864), in a long-term study conducted in Baía Norte (27°S), Santa Catarina State, Brazil. Although several other studies described Guiana dolphins' range patterns and habitat use (e.g. Rossi-Santos *et al.*, 2006; Azevedo *et al.*, 2007; Wedekin *et al.*, 2007), most used group location to detect habitat 'preferences' or ranges of the investigated populations. Individuals within a particular population can differ greatly in ranging patterns (Defran *et al.*, 1999) and may shift between local site-fidelity and longer movements away from the site where they were first identified (Würsig & Würsig, 1979; Wells *et al.*, 1990; Würsig & Harris, 1990; Bearzi *et al.*, 1997). A low number of studies explored individual home ranges of *S. guianensis* using the minimum convex polygon (e.g. Flores & Bazzalo, 2004; Hardt, 2005; Rossi-Santos *et al.*, 2007; Batista, 2008) and the kernel density estimator (e.g. Flores & Bazzalo, 2004; Hardt, 2005). It would be recommendable to invest more efforts to conduct such studies along this poorly known cetacean species distribution.

Sotalia guianensis dwell in coastal and estuarine waters of the western South Atlantic, from southern Brazil to Honduras (Flores & da Silva, 2009). In the Cananéia estuary (25°S), located in south-eastern Brazil, *S. guianensis* has been the focus of several studies since the end of the 1990s (e.g. Santos *et al.*, 2000, 2001; Santos & Rosso, 2007, 2008). Due to its ecological relevance, the Cananéia estuary was designated as a protected reserve in the 1980s (Schaeffer-Novelli *et al.*, 1990). However, in recent years part of this ecosystem has been threatened by non-regulated tourism activities and destruction of mangroves for housing. Understanding the way *S. guianensis* uses its area is an important tool to plan management and conservation actions. Therefore, the aim of this study was to investigate spatial patterns of use and estimate core areas and home range sizes of Guiana dolphins in the Cananéia estuary.

MATERIALS AND METHODS

Study area

The Cananéia estuary (25°03'S 47°55'W) is located on the southern coast of São Paulo State, Brazil, and the surveyed area included the waters encircling the Cananéia Island (Figure 1). The 132 km² surveyed area was divided into three smaller sub-sets known as A₀, A₁–A₄, and A₅. There are no physical barriers isolating the quoted sub-areas, which were stratified in order to cover the entire estuarine area where dolphins can be found in a lesser amount of days. Based on their sizes and daylight time available for observations, sub-areas A₁ to A₄ were surveyed on the same day, and A₀ and A₅ on two other separate days. A₁–A₄ were surveyed from 2000 on, A₀ from 2001 on and A₅ from 2002 on.

Data collection

Field work was conducted from May 2000 to July 2003. Efforts were unevenly distributed in time and space. Small motor-powered boats (15 and 30 hp) were used to survey subareas in zigzag movements to find as many groups of dolphins as

possible. Surveys were conducted in Beaufort sea states from 0 to 2. When an individual dolphin or a group of dolphins was found, the boat approached the animal(s) in a parallel orientation for the photo-identification efforts. Group/individual position was collected with a GPS using datum WGS 84. On most occasions, all individuals in a group including calves were photographed using a 35-mm reflex camera with a 300 mm zoom lens and ISO-400 colour printed films. Pictures were taken at distances ranging from 2 to 15 m. Photo analyses protocols followed the methodology described by Santos & Rosso (2008). Individuals with distinct notches on the dorsal fin border which allowed re-sightings were catalogued. Catalogued individuals were identified as females based on long-term observations (≥ 5 identifications in different months) during which they were in close companionship with calves (echelon position *sensu* Mann & Smuts, 1999) yet isolated from other individuals. In contrast, animals were identified as 'possible males' after long-term observations (≥ 5 identifications in different months over a 3-year period) during which they were not observed with calves (Santos & Rosso, 2008). The sex of the KN#86 individual was determined using molecular genetic procedures based on skin samples obtained with an adapted 80-pound crossbow. Details on the DNA analyses are described in Santos & Rosso (2008).

Data analysis

Using ArcGIS[®] 9.2 edition tools, a cartographic base was created in digital format by manual vectorial edition from remote sensing products ETM+ /Landsat-7, orbit 220/77, dated from 26 September 1999, projected coordinate system WGS 84–22S zone. A shape file was created for each photo-identified Guiana dolphin with 20+ geographical locations collected in distinct dates. If a dolphin was photographed more than once in a day, only the first sighting was used for analyses to avoid auto-correlation of data. The home range for each individual was calculated using the extension Home Range Tools (HRT) version 1.1 in the ArcGIS[®] 9.2. The HRT software is an ArcGIS[®] version of the Home Range Extension (HRE) for ArcVIEW 3.1. The HRT extends ArcGIS[®] to analyse home ranges of animals and is available from <http://blue.lakeheadu.ca/hre/> (Rodgers *et al.*, 2007). The HRT extension also permits the determination of the output UD raster form, so raster cell size was defined as 100 m and scaling factor as 1,000,000. The fixed kernel density estimator using the least squares cross-validation (LSCV) to choose the band width was performed to calculate the utilization UD. The fixed kernel keeps band width (h) constant for a data set (see Powell, 2000). Isopleths of 95%, 50% and 25% (UD contours) of the home ranges were generated to compare sizes of the areas and their possible cores. The home range areas usually had parts estimated on land due to the proximity of several location points to shallow waters and mangrove borders. Areas which overlapped with land were extracted using the tool ERASE from the ArcGIS[®].

RESULTS

A total of 374 groups of Guiana dolphins, ranging from 2 to 60 dolphins per group, were photographed in 87 survey days (Table 1). A total of 29,327 photographs were analysed,




Fig. 1. Map showing the Cananéia estuary in the southern limit of São Paulo State, Brazil. Subareas A1 – A4, A0 and A5 were surveyed in distinct days from 2000 to 2003. Black dots are the limits of subareas.

from which 6,312 (21.5%) were considered useful for identification purposes.

From a total of 138 catalogued individuals, only 7 (5.1%) were photographed on 20+ distinct occasions and thus were selected for home range analyses. Their identification numbers and sighting history are detailed in Table 2.

The 95% home range estimated outlines varied from 1.6 to 22.9 km² (Table 3). The largest ranging area was 22.9 km² and the smallest one was 1.6 km², regarding a male and a female, respectively. Estimated areas for males varied from 2.3 to 22.9 km², and for females varied from 1.6 to 3.9 km². In both cases these areas were estimated using the 95% UD. Data on 50% and 25% UD are shown in Table 3. All home

range polygons are shown in Figure 2 A–G. The polygons created had usually an amoeboid shape and in some cases more than one centre of activity.

DISCUSSION

Although the presented results were based on a relatively small number of monitored dolphins ($N = 7$), it is the first assay estimating individual Guiana dolphins home ranges in the Cananéia estuary based on a short-term photo-identification effort from 2000 to 2003. Those results were gathered in a large area of ~132 km², which can be fully monitored in three distinct days using one boat, where the local abundance of dolphins reaches ~300 to 400 individuals and using developed films and older reflex cameras which rendered almost 30,000 photographs to be analysed. It is also important to consider the low rate of useful photographs (21.5%) when dealing with a shy small cetacean species such as *S. guianensis*. In a catalogue composed of 138 individuals, only seven adults showed 20+ sightings in distinct dates after 87 fielding days. Based on this scenario, in this first

Table 1. Efforts invested in photo-identification studies of Guiana dolphins (*Sotalia guianensis*) in the Cananéia estuary, Brazil. The number of survey days, sighted groups and photographs taken are described by subarea between 2000 and 2003.

Year/subareas	Ao	A1 – A4	A5	Overall	
2000	Days	–	12	–	12
	Groups	–	95	–	95
	Photographs	–	7,067	–	7,067
2001	Days	4	21	–	25
	Groups	16	99	–	115
	Photographs	950	9,347	–	10,297
2002	Days	10	16	4	30
	Groups	25	40	21	86
	Photographs	1,671	3,167	944	5,782
2003	Days	10	4	6	20
	Groups	30	22	26	78
	Photographs	3,102	1,720	1,359	6,181
2000–2003	Days	24	53	10	87
	Groups	71	256	47	374
	Photographs	5,723	21,301	2,303	29,327

Table 2. Gender and number of locations recorded for seven Guiana dolphins (*Sotalia guianensis*) photo-identified between 2000 and 2003 in the Cananéia estuary, Brazil.

Dolphin ID	Sex	2000	2001	2002	2003	Overall
KN#15	M	1	8	10	5	24
KN#30	F	4	10	9	4	27
KN#43	M	6	10	3	1	20
KN#83	M	6	20	7	3	36
KN#86	M	3	11	9	3	26
KN#147	F	7	11	3	2	23
KN#186	M	6	5	7	2	20

Table 3. Home ranges of seven Guiana dolphins (*Sotalia guianensis*) individually identified from 2000 to 2003 in the Cananéia estuary, Brazil. Columns indicate respectively individual identification, sex and the home range area (km²) represented by the 95%, the 50% and 25% utility distributions (UDs).

Individuals	Sex	95% UD	50% UD	25% UD
KN#15	M	16.35	4.91	1.54
KN#30	F	3.89	0.81	0.35
KN#43	M	2.32	0.45	0.17
KN#83	M	2.96	0.63	0.24
KN#86	M	5.27	1.18	0.31
KN#147	F	1.60	0.31	0.11
KN#186	M	22.86	8.60	4.17
Mean		7.89	2.41	0.98
SD		8.30	3.16	1.49

assay using *S. guianensis* sightings in the Cananéia estuary, emphasis was given in investing efforts on careless steps to evaluate home range areas (see Data Analysis section), as well as comparing the observed home range sizes with several other studies using the same technique and methods. As only two females were included in the analyses, comparisons involving gender were avoided based on small sample size.

The kernel density estimator was chosen for these first observations in Cananéia because it is considered the best estimator available for estimating home ranges (see Seaman & Powell, 1996; Seaman *et al.*, 1999; Powell, 2000). It produces an unbiased density estimate directly from data and it is not influenced by grid size or placement (Silverman, 1986). The minimum convex polygon estimator is still being used and remains as a commonly applied technique mostly for comparisons to previous studies and due to its ease of calculation (see Seaman & Powell, 1996; Seaman *et al.*, 1999; Flores & Bazzalo, 2004). However, comparisons to previous work are unreliable given the extreme sensitivity to sample size by the minimum convex polygon, which also ignores the information provided by interior data points and produces only crude outlines of animals' home ranges (Powell, 2000).

It is possible to observe differences of the values of the estimated areas in four different regions of the Guiana dolphins' distribution in Brazil. Individual home range areas observed for Guiana dolphins in the Cananéia estuary varied from 1.6 to 22.9 km². The results presented by Flores & Bazzalo (2004) in Baía Norte showed home ranges varying from 5.4 to 21.5 km² with the minimum convex polygon, and from 12.6 to 19.6 km² using the kernel density estimator. In Baía da Babitonga (26°S), Santa Catarina State, home ranges of Guiana dolphins varied from 1.6 to 25.7 km² using the minimum convex polygon, and from 4.3 to 91.5 km² with the kernel estimator (Hardt, 2005). Using the minimum convex polygon, Batista (2008) found a variation from 1 to 8.7 km² on home ranges observed for Guiana dolphins found in the Rio Paraguaçu estuary (12°S), Bahia State, Brazil. The use of different estimators and different minimum number of sightings may have partially influenced those differences. Other factors such as local habitat characteristics, food availability in time and space, population abundance and individual variations based on age and sex may also have influenced the observed differences. Standardization of analyses will surely be of some help in further comparisons. Differences among home range sizes have already been discussed for bottlenose

dolphins. Gubbins (2002) found smaller home ranges when studying *Tursiops truncatus* in South Carolina, compared with bottlenose dolphins observed in Texas (Würsig & Lynn, 1996) and Florida (Wells, 1991). Gubbins (2002) assumed that the smaller home ranges observed could indicate abundant food uniformly distributed in space and time.

The minimum number of sightings to estimate individual *S. guianensis* home ranges varied from 10 (Hardt, 2005; Batista, 2008) to 33 (Flores & Bazzalo, 2004) and maximum from 18 (Batista, 2008) to 53 (Flores & Bazzalo, 2004). In the present study, the numbers of sightings ranged from 20 to 36. Mares *et al.* (1980) and Schoener (1981) suggested that a minimum number of 20–25 sightings per individual would be important to ensure independence of home range area from sample size. Home range estimations are significantly affected by sample size and by the estimator used. Thus, a greater sample size can result in a more precisely estimated home range (see Seaman & Powell, 1996; Seaman *et al.*, 1999; Powell, 2000; Owen *et al.*, 2002). Several of the relatively small home ranges found in this study may be underestimated. Further investigations are possible to be conducted with a broader sample size in the Cananéia estuary as the monitoring programme has continued.

When comparing the quoted studies on small cetacean home ranges, it was possible to detect that *T. truncatus* showed larger home ranges when compared to the areas regarding *S. guianensis*. Are those differences simply the product of the influence of species' size? Detailed investigations should be conducted to compare the size of home ranges of different cetacean species in order to detect the main factors driving such differences. Previous studies on ecological aspects regarding *S. guianensis* showed individual site fidelity in several areas of the species distribution (e.g. Santos *et al.*, 2001; Azevedo *et al.*, 2007). It is likely that there is a tendency for individuals of this species to use relatively smaller areas when compared to other cetacean species such as bottlenose dolphins.

The UD's showed location points concentrated in the main entrance of the Cananéia estuary, which could reflect the emphasis on efforts given to the subareas A1–A4 in the first year of the survey. Efforts were evenly distributed since 2002 in order to minimize the effects of a non-uniformly distributed investigation. After expanding the monitored areas, no consistent differences on sighting locations were observed for the seven monitored individuals presented herein.

Home ranges may change in time and space and such dynamics require continuous observations for a better understanding of the ecological factors underlying their establishment. Animals are capable of having cognitive maps of where they live (Peters, 1978) and such maps must change as the animal learns about its environment, hence, the maps change with time. As new resources are discovered and old ones disappear, the cognitive map may change, as well as the home range (Powell, 2000). Thus, home ranges may represent a transitory situation of a determined length of time and this is why a continuous monitoring of this population of Guiana dolphins inhabiting the Cananéia estuary is important.

Using satellite tags should be an important step to explore the dynamics regarding *S. guianensis*' home ranges, rendering precise short-term information on location points with a smaller amount of efforts when compared to photo-identification in a large habitat and considering a population




Fig. 2. (A–G) Home ranges of seven Guiana dolphins (*Sotalia guianensis*) monitored from 2000 to 2003 in the Cananéia estuary, Brazil.

of hundreds of dolphins. The use of coastal waters is another important issue that deserves further attention because movements outside the estuary may represent important components for several monitored dolphins. Thus, further

investigations in coastal waters could add important pieces to the puzzle on Guiana dolphins' home ranges. Regarding conservation issues, these results suggest that awareness for tourists and boat owners should be important as the main

entrance of the Cananéia estuary usually shelters large aggregations of Guiana dolphins and hosts the main area of usage of several monitored dolphins. Coastal dolphins can be good indicators for examination of pollutant or disease vectors in nearshore habitats (Moore, 2008). Therefore, considering the small range of *S. guianensis* distribution and its fidelity to some estuaries and shallow waters, this species could be seen as sentinel of marine ecosystem changes (e.g. Moore, 2008) which provides an important tool to guide conservation and management activities in the Cananéia estuary and in other parts of its distribution.

ACKNOWLEDGEMENTS

Logistic support was provided by the Instituto Oceanográfico da Universidade de São Paulo. We also thank the referees who provided important comments for the improvement of the final text. The Whale and Dolphin Conservation Society, the Society for Marine Mammalogy, the Earthwatch Institute and the Cetacean Society International financially supported Guiana dolphins' photo-identification studies in the Cananéia estuary. Marcos Santos received grants from Fundação de Amparo à Pesquisa do Estado de São Paulo (FAPESP) to conduct his doctoral studies (process number 01/05128-8) and receives grants from 'Programa Jovem Pesquisador—FAPESP' (process numbers 05/59439-5 and 05/54149-9). Júlia Emi de Faria Oshima received grants from Programa de Recursos Humanos da ANP para o setor de petróleo e gás—PRH 05-ANP/MCT/FINEP (process number 48610002667/99-16) and grants and fellowship from Fundação de Amparo à Pesquisa do Estado de São Paulo (FAPESP) (process number 2008/56543-4).

REFERENCES

- Azevedo A.F., Oliveira A.M., Viana S.C. and Sluys M.V. (2007) Habitat use by marine tucuxis (*Sotalia guianensis*) (Cetacea: Delphinidae) in Guanabara Bay, south-eastern Brazil. *Journal of the Marine Biological Association of the United Kingdom* 87, 201–205.
- Batista R.L.G. (2008) *Uso de área e associação entre os botos-cinza Sotalia guianensis (Van Benédén, 1864) do estuário do Rio Paraguaçu—BA*. MSc dissertation. Universidade Santa Cruz, Ilhéus, BA, Brazil.
- Bearzi G., Notarbartolo-Di-Sciara G. and Politi E. (1997) Social ecology of bottlenose dolphins in Kvarneric (northern Adriatic Sea). *Marine Mammal Science* 13, 650–668.
- Bigg M.A., Olesiuk P.F., Ellis G.M., Ford J.K. and Balcomb K.C. (1990) Social organization and genealogy of resident killer whales (*Orcinus orca*) in the coastal waters of British Columbia and Washington State. In Hammond P.S., Mizroch S.A. and Donovan G.P. (eds) *Individual recognition of cetaceans use of photo-identification and other techniques to estimate population parameters*. International Whaling Commission, Special Issue 12, Cambridge, pp. 383–405.
- Burt W.H. (1943) Territoriality and home range concepts as applied to mammals. *Journal of Mammalogy* 24, 346–352.
- Connor R.C. and Smolker R. (1985) Habituated dolphins (*Tursiops* sp.) in Western Australia. *Journal of Mammalogy* 66, 398–400.
- Defran R.H., Weller D.W., Kelly D.L. and Espinosa M.A. (1999) Range characteristics of Pacific coast bottlenose dolphins (*Tursiops truncatus*) in the Southern California Bight. *Marine Mammal Science* 15, 381–393.
- Dixon K.R. and Chapman J.A. (1980) Harmonic mean measure of animal activity areas. *Ecology* 61, 1040–1044.
- Flores P.A.C. and Bazzalo M. (2004) Home ranges and movement patterns of the marine tucuxi dolphin *Sotalia fluviatilis*, in Baía Norte, Southern Brazil. *Latin American Journal of Aquatic Mammals* 3, 37–52.
- Flores P.A.C. and da Silva V.M.F. (2009) Tucuxi and Guiana dolphin. *Sotalia fluviatilis* and *S. guianensis*. In Perrin W.F., Würsig B. and Thewissen J.G.M. (eds) *Encyclopedia of marine mammals*. San Diego, CA: Academic Press, pp. 1188–1192.
- Gubbins C. (2002) Use of home ranges by resident bottlenose dolphins (*Tursiops truncatus*) in a South Carolina estuary. *Journal of Mammal Research* 83, 178–187.
- Hammond P.S., Mizroch S.A. and Donovan G.P. (eds) (1990) Individual recognition of cetaceans: use of photo-identification and other techniques to estimate population parameters. *Reports of the International Whaling Commission Special issue 12*. Cambridge, UK.
- Hardt F.A.S. (2005) *Padrões de residência do golfinho Sotalia guianensis (CETACEA, DELPHINIDAE) na Baía da Babitonga, litoral norte de Santa Catarina, Brasil*. MSc dissertation. Universidade Federal do Paraná, Curitiba, PR, Brazil.
- Herzing D.L. (1997) The life of free ranging Atlantic spotted dolphins (*Stenella frontalis*): age classes, color phases and female reproduction. *Marine Mammal Science* 13, 576–595.
- Mann J. and Smuts B.B. (1999) Behavioral development of wild bottlenose dolphin newborns. *Behaviour* 136, 529–566.
- Mares M.A., Willig M.R. and Bitar N.A. (1980) Home range size in eastern chipmunks, *Tamias stratus*, as a function of number of captures: statistical biases of inadequate sampling. *Journal of Mammalogy* 61, 661–669.
- Marten K. and Psarakos S. (1999) Long-term site fidelity and possible long-term associations of wild spinner dolphins (*Stenella longirostris*) seen off Oahu, Hawaii. *Marine Mammal Science* 15, 1329–1336.
- McNab B.K. (1963) Bioenergetics and the determination of home range size. *American Naturalist* 97, 133–140.
- Moore S.E. (2008) Marine mammals as ecosystem sentinels. *Journal of Mammalogy* 89, 534–540.
- Owen E.C.G., Wells R.S. and Hofman S. (2002) Raging and association patterns of paired and unpaired adult male Atlantic bottlenose dolphins, *Tursiops truncatus*, in Sarasota, Florida, provide no evidence for alternative male strategies. *Canadian Journal of Zoology* 80, 2072–2089.
- Peters R. (1978) Communication, cognitive mapping, and strategy in wolves and hominids. In Hall R.L. and Sharp H.S. (eds) *Wolf and man: evolution in parallel*. New York: Academic Press, pp. 95–108.
- Powell R.A. (2000) Animal home ranges and territories and home range estimators. In Boitani L. and Fuller T.K. (eds) *Research and techniques in animal ecology: controversies and consequences*. New York: Columbia University Press, pp. 65–110.
- Rodgers A.R., Carr A.P., Beyer H.L., Smith L. and Kie J.G. (2007) *HRT: Home Range Tools for ArcGIS*. Ontario Ministry of Natural Resources, Centre for Northern Forest Ecosystem Research, Thunder Bay, Ontario, Canada. <http://blue.lakeheadu.ca/hre> (accessed on 10 October 2008).
- Rossi-Santos M.R., Wedekin L.L. and Sousa-Lima R.S. (2006) Distribution and habitat use of small cetaceans off Abrolhos Bank, eastern Brazil. *Latin American Journal of Aquatic Mammals* 5, 23–28.
- Rossi-Santos M.R., Wedekin L.L. and Monteiro-Filho E.L.A. (2007) Residence and site fidelity of *Sotalia guianensis* in the Caravelas

- River Estuary, eastern Brazil. *Journal of the Marine Biological Association of the United Kingdom* 87, 207–212.
- Samuel M.D., Pierce D.J. and Garton E.O.** (1985) Identifying areas of concentrated use within the home range. *Journal of Animal Ecology* 54, 711–719.
- Santos M.C. de O. and Rosso S.** (2007) Ecological aspects of marine tucuxi dolphins (*Sotalia guianensis*) based on group size and composition in the Cananéia estuary, southeastern Brazil. *Latin American Journal of Aquatic Mammals* 6, 71–82.
- Santos M.C. de O. and Rosso S.** (2008) Social organization of marine tucuxi dolphins, *Sotalia guianensis*, in the Cananéia estuary of southeastern Brazil. *Journal of Mammalogy* 88, 347–355.
- Santos M.C. de O., Rosso S., Siciliano S., Zerbini A., Zampiroli E., Vicente A.F. and Alvarenga F.** (2000) Behavioral observations of the marine tucuxi dolphin (*Sotalia fluviatilis*) in São Paulo estuarine waters, Southeastern Brazil. *Aquatic Mammals* 26, 260–267.
- Santos M.C. de O., Acuña L.B. and Rosso S.** (2001) Insights on site fidelity and calving intervals of the marine tucuxi dolphin (*Sotalia fluviatilis*) in south-eastern Brazil. *Journal of the Marine Biological Association of the United Kingdom* 81, 1049–1052.
- Schaeffer-Novelli Y., Mesquita H.S.L. and Cintrón-Molero G.** (1990) The Cananéia lagoon estuarine system, São Paulo, Brazil. *Estuaries* 13, 193–203.
- Schoener T.W.** (1981) An empirically based estimate of home range. *Theoretical Population Biology* 20, 281–325.
- Seaman D.E. and Powell R.A.** (1996) An evaluation of the accuracy of kernel density estimators for home range analysis. *Ecology* 77, 2075–2085.
- Seaman D.E., Millsaugh J.J., Kernohan B.J., Brundige G.C., Raedeke K.J. and Gitzen R.A.** (1999) Effects of sample size on kernel home range estimates. *Journal of Wildlife Management* 63, 739–747.
- Silverman B.W.** (1986) *Density estimation for statistics and data analysis*. London: Chapman and Hall.
- Wedekin L.L., Daura-Jorge F.G., Piacentini V.Q. and Simões-Lopes P.C.** (2007) Seasonal variations in spatial usage by the estuarine dolphin, *Sotalia guianensis* (van Bénédén, 1864) (Cetacea; Delphinidae) at its southern limit of distribution. *Brazilian Journal of Biology* 67, 1–8.
- Wells R.S.** (1991) The role of long-term study in understanding the social structure of a bottlenose dolphin community. In Pryor K. and Norris K. (eds) *Dolphin societies—discoveries and puzzles*. Berkeley, CA: University of California Press, pp. 199–226.
- Wells R.S., Scott M.D. and Irvine A.B.** (1987) The social structure of free-ranging bottlenose dolphins. In Genoways H.H. (ed.) *Current mammalogy*. New York: Plenum Press, pp. 247–305.
- Wells R.S., Hansen L.J., Baldrige A., Dohl T.P., Kelly D.L. and DeFran R.H.** (1990) Northward extension of the range of bottlenose dolphins along the California coast. In Leatherwood S. and Reeves R.R. (eds) *The bottlenose dolphin*. San Diego, CA: Academic Press, pp. 421–431.
- Worton B.J.** (1989) Kernel methods for estimating the utilization distribution in home-range studies. *Ecology* 70, 164–168.
- Würsig B. and Würsig M.** (1977) The photographic determination of group size, composition, and stability of coastal porpoises (*Tursiops truncatus*). *Science* 198, 755–756.
- Würsig B. and Würsig M.** (1979) Behavior and ecology of bottlenose porpoises, *Tursiops truncatus*, in the South Atlantic. *Fishery Bulletin* 77, 399–442.
- Würsig B. and Harris G.** (1990) Site and association fidelity in bottlenose dolphins off Argentina. In Leatherwood S. and Reeves R.R. (eds) *The bottlenose dolphin*. San Diego, CA: Academic Press, pp. 361–365.
- and
- Würsig B. and Lynn S.K.** (1996) *Movements, site fidelity, and respiration patterns of bottlenose dolphins on the central Texas coast*. National Oceanic and Atmospheric Administration, Technical Memorandum NMFS-EFSC-383:1–111.

Correspondence should be addressed to:

J.E.F. Oshima
 Projeto Atlantis, Laboratório de Biologia da
 Conservação de Cetáceos, Departamento de Zoologia
 Instituto de Biociências da Universidade
 Estadual Paulista ‘Júlio de Mesquita Filho’ (UNESP)
 Campus Rio Claro, Avenida 24-A, 1515, Bela Vista
 Rio Claro, SP, 13506-900, Brazil
 email: juliaoshima@yahoo.com.br