

Introduction: Patients with concussion frequently present to the emergency department (ED). Studies of athletes and children indicate that concussion symptoms are often more severe and prolonged in females compared with males. To-date, study of sex-based concussion differences in general adult populations have been limited. This study examined sex-based differences in concussion outcomes. **Methods:** Adult (>17 years) patients presenting to one of three urban EDs in Edmonton, Alberta with Glasgow coma scale score 13 within 72 hours of a concussive event were recruited by on-site research assistants. Follow-up calls at 30 and 90 days post ED discharge captured extent of PCS using the Rivermead Post-Concussion questionnaire (RPQ), effect on daily living activities measured by the Rivermead Head Injury Questionnaire (RHIQ), and overall health-related quality of life using the 12-item Short Form Health Survey (SF-12). Dichotomous and categorical variables were compared using Fishers exact test; continuous variables were compared using t-tests or Mann-Whitney tests, as appropriate. **Results:** Overall, 130/250 enrolled patients were female. The median age was 35 years; men trended towards being younger (median = 32 years; IQR: 23, 45) than women (median = 40 years; IQR: 22, 52). Compared to women, more men were single (56% vs. 38% (p=0.007) and employed (82% vs. 71% (p=0.055). Men and women experienced different injury mechanisms (p=0.007) with more women reporting injury due to a fall (44% vs. 26%), while more men were injured at work (16% vs. 7%) or due to an assault (11% vs. 3%). Men had a higher return to ED rate (13% vs. 5%; p=0.015). Women had higher RPQ scores at baseline (p<0.001) and 30-day follow-up (p=0.001); this difference was not significant by 90 days (p=0.099). While women reported on the RHIQ at 30 days that their injury affected their usual activities significantly more than men (Median = 5, IQR: 0, 11 vs. median = 0.5, IQR: 0.5, 7; p=0.004), both groups had similar scores on the SF-12 physical composite and mental composite scales at all three measurement points. **Conclusion:** In a general ED concussion population, demographic differences exist between men and women. Based on self-reported and objective outcomes, womens usual activities may be more affected by concussion and PCS than men. Further analysis of these differences is required in order to identify different treatment options and ensure adequate care and treatment of injury.

Keywords: concussion, sex-based differences, injury

LO47

Incidence of intracranial bleeding in anticoagulated emergency patients with minor head injury: a meta-analysis

K. de Wit, MBChB, BSc, MD, MSc, H. Minas, BSc, W. Arthur, BSc, M. Turcotte, BSc, Msc, M. Eventov, BSc, S. Mason, MBBS MD, D. Nishijima, MD, MAS, M. Li, MD, G. Versm e, MD, McMaster University, Hamilton, ON

Introduction: The proportion of Canadians receiving anticoagulation medication is increasing. Falls in the elderly are the most common cause of minor head injury and an increasing proportion of these patients are prescribed anticoagulation. Emergency department (ED) guidelines advise performing a CT head scan for all anticoagulated head injured patients, but the risk of intracranial hemorrhage (ICH) after a minor head injury (patients who have a Glasgow coma score (GSC) of 15) is unclear. We conducted a systematic review and meta-analysis to determine the point incidence of ICH in anticoagulated ED patients presenting with a minor head injury. **Methods:** We systematically searched Pubmed, EMBASE, Cochrane database, DARE, google scholar and conference abstracts (May 2017). Experts were contacted. Meta-Analyses and Systematic Reviews of Observational Studies (MOOSE) guidelines were followed with two authors reviewing titles,

four authors reviewing full text and four authors performing data extraction. We included all prospective studies recruiting consecutive anticoagulated ED patients presenting with a head injury. We obtained additional data from the authors of the included studies on the subset of GCS 15 patients. We performed a meta-analysis to estimate the point incidence of ICH among patients with a GCS score of 15 using a random effects model. **Results:** A total of five studies (and 4,080 GCS 15, anticoagulated patients) from the Netherlands, Italy, France, USA and UK were included in the analysis. One study contributed 2,871 patients. Direct oral anticoagulants were prescribed in only 60 (1.5%) patients. There was significant heterogeneity between studies with regards to mechanism of injury, CT scanning and follow up method (I² = 93%). The random effects pooled incidence of ICH was 8.9% (95% CI 5.0-13.8%). **Conclusion:** We found little data to reflect contemporary anticoagulant prescribing practice. Around 9% of warfarinized patients with a minor head injury develop ICH. Future studies should evaluate the safety of selective CT head scanning in this population.

Keywords: head injury, computed tomography scan, anticoagulation

LO48

Does FAST change management of blunt trauma patients?

R. Thavanathan, MD, I. G. Stiell, MD, MSc, O. Levac-Martinho, BScN, J. Worrall, MD, B. W. Ritcey, MD, University of Ottawa, Department of Emergency Medicine, Ottawa, ON

Introduction: Despite widespread use of FAST in trauma, there is a lack of data supporting its usefulness. We sought to identify the impact of FAST on clinical management of blunt trauma patients. **Methods:** This health records review was conducted at a single large academic Level 1 trauma center emergency department. Patients with a suspicion of acute blunt traumatic abdominal injury were identified from our health records database. Data were collected regarding FAST utilization, CT scan utilization and timing, need for definitive management, disposition, and length of stay (LOS). **Results:** 285 patients were included, 152 (53.3%) received a FAST examination, with 33 (22%) having a direct impact on clinical management. CT was performed in 112 (73.6%) of the FAST group, with mean time to imaging of 147.4 minutes, time to trauma team assessment of 21.5 minutes, and ED-LOS of 8.6 hours. In the non-FAST group, 33 (24.8%) received a CT, with time to imaging of 133 minutes, time to trauma team assessment of 133 minutes, and ED-LOS of 13.8 hours. 75.6% of the FAST group required admission and 9.2% required definitive management; admission was needed for 38.3% of the non-FAST group and 2.2% required definitive management. **Conclusion:** This is the first study to assess patient outcomes with respect to FAST in the era of early whole body CT in trauma. Although FAST does not directly impact care for the majority of blunt trauma patients, it demonstrates usefulness in some patients by directing CT utilization and expediting disposition from the ED.

Keywords: focused assessment with sonography for trauma (FAST), blunt abdominal trauma, point of care ultrasound

LO49

Achieving just outcomes: forensic evidence collection in sexual assault cases

K. Sampsel, MD, MSc, K. Muldoon, MPH, PhD, A. Drumm, BSc, T. Leach, M. Heimerl, BA, MSW, University of Ottawa, Department of Emergency Medicine, Ottawa, ON

Introduction: Achieving just outcomes in sexual assault cases is one of the most serious and complex problems facing the health care and justice systems. The objectives of this analysis were to determine the