

Exploring the spatially resolved initial mass function in SAMI star-forming galaxies

Diego Salvador,¹ Andrew M. Hopkins,¹ Matt Owers,^{1,2} Themiya Nanayakkara,³ and Scott M. Croom^{2,4}

¹School of Mathematical and Physical Sciences, 12 Wally's Walk, Macquarie University, NSW 2109, Australia.

²ARC Centre of Excellence for All Sky Astrophysics in 3 Dimensions (ASTRO 3D), Australia.

³Centre for Astrophysics and Supercomputing, Swinburne University of Technology, P.O. Box 218, Hawthorn, 3122, VIC, Australia.

⁴Sydney Institute for Astronomy, School of Physics, University of Sydney, NSW 2006, Australia.

Author for correspondence: D. Salvador, Email: diego.salvador@hdr.mq.edu.au.

Abstract

The initial mass function (IMF) is a construct that describes the distribution of stellar masses for a newly formed population of stars. It is a fundamental element underlying all of star and galaxy formation, and has been the subject of extensive investigation for more than 60 years. In the past few decades there has been a growing, and now substantial, body of evidence supporting the need for a variable IMF. In this light, it is crucial to investigate the IMF's characteristics across different spatial scales and to understand the factors driving its variability. We make use of spatially resolved spectroscopy to examine the high-mass IMF slope of star-forming galaxies within the SAMI survey. By applying the Kennicutt method and stellar population synthesis models, we estimated both the spaxel-resolved (α_{res}) and galaxy-integrated (α_{int}) high-mass IMF slopes of these galaxies. Our findings indicate that the resolved and integrated IMF slopes exhibit a near 1:1 relationship for $\alpha_{int} \gtrsim -2.7$. We observe a wide range of α_{res} distributions within galaxies. To explore the sources of this variability, we analyse the relationships between the resolved and integrated IMF slopes and both the star formation rate (SFR) and SFR surface density (Σ_{SFR}). Our results reveal a strong correlation where flatter/steeper slopes are associated with higher/lower SFR and Σ_{SFR} . This trend is qualitatively similar for resolved and global scales. Additionally, we identify a mass dependency in the relationship with SFR, though none was found in the relation between the resolved slope and Σ_{SFR} . These findings suggest an scenario where the formation of high-mass stars is favoured in regions with more concentrated star formation. This may be a consequence of the reduced fragmentation of molecular clouds, which nonetheless accrete more material.

1. INTRODUCTION

Galaxy evolution arises from the complex interactions between gravitational, hydrodynamical, and radiative processes (e.g. Somerville & Davé, 2015). The evolutionary mechanisms governed by these processes are, to a large extent, dependent on the distribution of stellar masses (Hopkins, 2018). This “initial mass function” (IMF) characterises this distribution within a given volume following a star formation event. As mass is the primary determinant of a star's evolutionary path, the IMF is crucial for comprehending star formation (Kennicutt, 1998; Hoversten & Glazebrook, 2008; Elmegreen, 2009; Lee et al., 2009; Meurer et al., 2009; Gunawardhana et al., 2011), chemical enrichment (Portinari et al., 1997; Tornatore et al., 2007; Komiya, 2011; Goswami et al., 2021), feedback processes (Dib et al., 2010; Smith, 2021), mass-to-light ratios (Portinari et al., 2004; Cappellari et al., 2012; Mehrgan et al., 2024), and dark matter content (Rocca-Volmerange & Guiderdoni, 1990; Treu et al., 2010).

The IMF is commonly built as a piecewise power law, with each segment defined by a number α that denotes the slope of the power law within that mass range. This parameter reflects the distribution of stars such that $dN/dM \propto m^\alpha$ (Salpeter, 1955; Kroupa et al., 1993; Kroupa, 2001). Treating the IMF as an observable is particularly challenging as most massive stars ($\gtrsim 30M_\odot$) have typically evolved off the main sequence

within 1 Myr, while lower-mass stars ($\lesssim 30M_\odot$) still remain in the main sequence (Kroupa et al., 2013). Consequently, it is not possible to capture the complete mass distribution at a single point in time. Instead, the IMF is commonly derived by observing and analysing several stellar populations across various stages of evolution, allowing for a statistical estimation of the overall distribution (Kroupa, 2001; Kroupa et al., 2013; Chabrier et al., 2014; Hopkins, 2018; Li et al., 2023).

Given the statistical nature of the IMF, several methods have been developed to estimate it. These include the Kennicutt method (Kennicutt, 1983; Kennicutt et al., 1994) for star forming galaxies, which uses the H α line equivalent width ($EW_{H\alpha}$) and an optical colour (such as $B - V$ or $g - r$) to derive the IMF. Other techniques involve dynamical approaches, such as gravitational lensing (Ferreras et al., 2010; Treu et al., 2010; Leier et al., 2016) and velocity dispersion in early-type galaxies (Ferreras et al., 2012; Zaritsky et al., 2012; Esdaile et al., 2021). Additionally, spectral features are also employed to infer the IMF, including the 8183 – 8195 Å Na I doublet (van Dokkum & Conroy, 2010; Smith et al., 2015), the 9916 Å Wing–Ford FeH molecular band (Hardy, 1990; van Dokkum & Conroy, 2010; Conroy & van Dokkum, 2012; La Barbera et al., 2016; Vaughan et al., 2018), the 8600 μm Ca II triplet (Saglia et al., 2002; Vazdekis et al., 2003) and the TiO molecular band (Martín-Navarro et al., 2015a; La Barbera et al., 2016,

2017) for non-star forming galaxies, and the 2300 Å CO index (Mieske & Kroupa, 2008) and the $^{13}\text{C}/^{18}\text{O}$ ratio (Zhang *et al.*, 2018; Brown & Wilson, 2019) for star forming galaxies.

It is commonly assumed, either for convenience or lack of strong evidence otherwise, that the IMF is universal (Scalo, 1986; Kroupa, 2001; Bastian *et al.*, 2010; Hopkins, 2013), but there is strong conjecture around that point (Kennicutt, 1998; Hoversten & Glazebrook, 2008; Meurer *et al.*, 2009; Hoversten & Glazebrook, 2010; Treu *et al.*, 2010; van Dokkum & Conroy, 2010; Gunawardhana *et al.*, 2011; Narayanan & Davé, 2012; Nanayakkara *et al.*, 2017; Smith, 2014). A Universal IMF is supported by the observation that star formation occurs in a variety of settings with differing densities and chemical compositions, yet most IMF estimations converge on a slope of $\alpha = -2.35$, known as the Salpeter slope. However, this notion has been debated since the early 1960s, with arguments supporting a varying IMF (Schmidt, 1963). More recent studies have highlighted the necessity for varying IMFs to account for differences in metallicity (Martín-Navarro *et al.*, 2015b; Zonoozi *et al.*, 2016), $[\text{Mg}/\text{Fe}]$ content (van Dokkum & Conroy, 2010; Conroy & van Dokkum, 2012), SFR (Gunawardhana *et al.*, 2011), radial distance (Martín-Navarro *et al.*, 2015a), velocity dispersion (Ferreras *et al.*, 2012; Zaritsky *et al.*, 2012; Conroy *et al.*, 2013; Pernet *et al.*, 2024), $F_{\text{H}\alpha}/F_{\text{UV}}$ flux ratio (Meurer *et al.*, 2009), surface brightness (Hoversten & Glazebrook, 2010), stellar mass-to-light ratios (Cappellari *et al.*, 2012), stellar orbits (Poci *et al.*, 2022), environment (Geha *et al.*, 2013; Chabrier *et al.*, 2014) and redshift (Nanayakkara *et al.*, 2017). Moreover, a variable IMF could be invoked to explain phenomena such as the G-dwarf problem (Worthey *et al.*, 1996) and the thickness and tilt of the fundamental plane of elliptical galaxies (Graves & Faber, 2010).

Several approaches for quantifying IMF shapes in star forming galaxies are now well established (Kennicutt, 1983; Kennicutt *et al.*, 1994; Buat *et al.*, 1987; Meurer *et al.*, 2009), and models such as the IGIMF (Kroupa & Weidner, 2003) have been developed to explore variations found using these approaches. Given the advent of large scale resolved spectroscopy datasets, such as SAMI (Croom *et al.*, 2012), CALIFA (Sánchez *et al.*, 2012), and MaNGA (Bundy *et al.*, 2015), it is now possible to apply such techniques to explore how the IMF may vary within galaxies. Notably, the work of Martín-Navarro *et al.* (2015a, 2019), La Barbera *et al.* (2016); La Barbera *et al.* (2019), Parikh *et al.* (2018) and Barbosa *et al.* (2021) stands out for their resolved IMF estimations using spectral features in early-type galaxies and Martín-Navarro *et al.* (2023) on the late-type galaxy NGC 3351. Most of these studies focus on low-mass stars in non-star-forming galaxies, thus reflecting an already established IMF.

In this work, we explore resolved IMF measurements within star forming galaxies using integral field spectroscopy (IFS) data from the SAMI survey DR3 (Croom *et al.*, 2021). We perform spaxel-scale measurements of the high-mass IMF slope using the Kennicutt method (Kennicutt, 1983; Kennicutt *et al.*, 1994). This is the first large-scale exploration of resolved IMF measurements in star forming galaxies. This approach provides a novel perspective on the IMF slope distribution within galax-

ies. It enables comparisons between local IMF slope estimations and those derived from the traditional integrated galaxy light. Because the SAMI sample is drawn in part from the Galaxy And Mass Assembly (GAMA) galaxy survey (Driver *et al.*, 2022) we can compare directly to analogous global IMF measurements for those galaxies (Gunawardhana *et al.*, 2011). Our goal is to establish a new method for extracting new information about the variation of the IMF within galaxies, and trying to understand and link this information with the already-existing global galaxy integrated light IMF measurements. Additionally, we compare the resolved IMF measurements with other resolved physical quantities, such as star formation rate (SFR), star formation rate density (Σ_{SFR}) and the galaxy stellar mass, to investigate potential underlying mechanisms driving the IMF shapes. These can again be compared against other published results (Lee *et al.*, 2009; Meurer *et al.*, 2009; Gunawardhana *et al.*, 2011; Weidner *et al.*, 2013). Our analyses will enhance our understanding of the interplay between the IMF and various galactic properties, shedding light on the mechanisms driving galaxy evolution.

The layout of this paper is as follows. In § 2 we describe the SAMI data products and the sample we use. In § 3 we describe the Kennicutt method and the population synthesis model required to make the IMF slope estimations. In § 4 we describe our results. In § 5 we address the limitations of this work, contextualises our results within the framework of existing research, and explores their implications. Finally, in § 6 we summarise our findings and key remarks.

Throughout we assume cosmological parameters of $H_0 = 70 \text{ km s}^{-1} \text{ Mpc}^{-1}$, $\Omega_M = 0.3$, $\Omega_\Lambda = 0.7$ and $\Omega_k = 0$.

2. DATA

We use the data products provided by the SAMI Galaxy Survey Data Release 3 (Croom *et al.*, 2021). The survey's data were collected using the Sydney-AAO Multi-object Integral field spectrograph (SAMI; Croom *et al.* 2012), located at the 3.9-metre Anglo-Australian Telescope (AAT) at Siding Spring Observatory. SAMI consists of 13 fibre bundles, called hex-abundles, inserted into a custom field plate of 1-degree diameter on the sky at the corrected prime focus of the 3.9-metre aperture of the AAT. Each of these IFUs consists of 61 elements of 1.6 arcsec each, resulting in a field of view (FoV) of 15 arcsec for each IFU. This setup allows the instrument to perform multiplexed observations, with 12 IFUs observing galaxies and one calibration star simultaneously.

The SAMI Galaxy Survey (Croom *et al.*, 2012; Bryant *et al.*, 2015; Croom *et al.*, 2021), was conducted between 2013 and 2018, collecting data on 3068 galaxies covering a broad range of stellar masses (10^7 – $10^{12} M_\odot$) and environments (field, galaxy groups, and clusters) at $z < 0.095$. Targets were primarily selected from 144 deg^2 equatorial regions from the GAMA survey (Driver *et al.*, 2009), and 8 cluster regions described in Owers *et al.* (2017).

The SAMI IFU allows for spatially resolved spectroscopy of each galaxy, providing spectra for several regions across the galaxy (Bryant *et al.*, 2015). The primary SAMI data

products are two data cubes, each formatted as a $2048 \times 50 \times 50$ array, with 2048 representing the wavelength axis and 50×50 representing spatial coordinates. The wavelength ranges for the blue and red cubes are $3630 - 5800 \text{ \AA}$ and $6250 - 7460 \text{ \AA}$, respectively. Each spaxel (spatial pixel) has a pixel scale of 0.5 arcseconds, with a spectral (wavelength) pixel scale of 1.04 \AA per pixel for the blue cubes and 0.57 \AA per pixel for the red cubes. The spectral resolutions are $R = 1808$ ($\sigma = 70.4 \text{ km s}^{-1}$) for the blue arm and $R = 4304$ ($\sigma = 29.6 \text{ km s}^{-1}$) for the red arm. The data reduction and cubing process is described elsewhere (Allen et al., 2014; Sharp et al., 2015; Green et al., 2018; Scott et al., 2018; Croom et al., 2021). The point spread function for each observation, with an average FWHM of 2.06 arcsec, is determined by fitting a Moffat profile to the flux distribution of the secondary standard star (Scott et al., 2018).

We also use other SAMI DR3 data products, such as two-dimensional maps of total emission line fluxes for $H\alpha$ and $H\beta$, gas velocity, gas velocity dispersion, star formation rate (SFR), star formation rate density (Σ_{SFR}), and SFR masks for star-forming spaxels. Line flux maps are created by summing all the flux based on a single component Gaussian fit associated with a given emission line in each spaxel. $H\alpha$ maps are extinction-corrected using the Balmer decrement and the (Cardelli et al., 1989) extinction law. Gas velocity and velocity dispersion maps are obtained by fitting 11 strong optical emission lines with a single Gaussian component using LZIFU (LaZy-IFU, Ho et al. 2016). SFR maps (in $M_{\odot} \text{ yr}^{-1}$) are derived from the extinction-corrected $H\alpha$ maps, assuming $\text{SFR} = L(H\alpha) \times \left(\frac{7.9}{1.53}\right) \times 10^{-42} [M_{\odot} \text{ yr}^{-1}]$ (Medling et al., 2018). Σ_{SFR} (in $M_{\odot} \text{ yr}^{-1} \text{ kpc}^{-2}$) maps are provided similarly. Each spaxel emission is classified as star-forming, in a SFR mask, according to the BPT/VO87 diagnostics from Kewley et al. (2006), based on Kewley et al. (2001) and Kauffmann et al. (2003) $[\text{OIII}]/H\beta$, $[\text{NII}]/H\alpha$, $[\text{SII}]/H\alpha$, and $[\text{OI}]/H\alpha$ flux ratio dividing lines.

We also use other physical quantities available in the SAMI DR3 tables. From `InputCatGAMADR3` and `InputCatClustersDR3` we extract stellar masses and effective radius, while $H\alpha_{R_e}$ and $H\beta_{R_e}$ fluxes, SFR_{R_e} , and spectroscopic redshift are retrieved from `EmissionLine1compDR3`. Note that the R_e sub-index indicates that these quantities are estimated from an elliptical aperture with a semi-major axis of 1 effective radius (R_e), where R_e was measured using a single-Sérsic fit (Kelvin et al., 2012; Owers et al., 2019). We also use the blue and red $1 R_e$ aperture spectra to calculate the galaxy integrated light colours and equivalent widths.

As we aim to calculate the g - and r -band fluxes from the SAMI data using g and r SDSS filters, it is important to note that part of the bluest region of the r band filter range ($5380 - 7230 \text{ \AA}$) falls outside the wavelength coverage of the SAMI red data cube ($6240 - 7460 \text{ \AA}$). To address this, we construct a truncated r band, r_t (Figure 1), which covers the rest-frame wavelength range from 6240 \AA to 7460 \AA . For consistency, we use this truncated band r_t throughout the rest of the paper.

In this work we select all galaxies in the SAMI survey DR3

with $7 \leq \log(M/M_{\odot}) \leq 12$ and a $10^{-3} \leq \text{SFR} \leq 10^2$ within $0 < z < 1.12$. The Kennicutt method is generally unsuitable for passive galaxies due to their minimal or absent $H\alpha$ and UV emissions^a from high-mass young stars (Kennicutt, 1983). Therefore, we restrict our analysis to galaxies identified as non-passive based on the spectroscopic classifications by Owers et al. (2019). Additionally, our analysis exclusively considers spaxels classified as star-forming, as indicated by the SFR mask in the SAMI data products. The resulting sample contains 1344 objects. Figure 2 provides a visual summary of the physical properties of our sample.

3. METHODS

We make use of the Kennicutt diagnostic (Kennicutt, 1983; Kennicutt et al., 1994; Hoversten & Glazebrook, 2008, 2010; Gunawardhana et al., 2011; Nanayakkara et al., 2017) to estimate the IMF slope by comparing the $\text{EW}_{H\alpha}$ line equivalent width with the $g - r$ colour. The $\text{EW}_{H\alpha}$ is the ratio of the $H\alpha$ flux, primarily contributed by gas ionisation by young massive stars ($> 10 M_{\odot}$), to the continuum at 6563 \AA , sensitive to the older stellar populations. This ratio is, thus, sensitive to the high-mass star population. In contrast, the $g - r$ colour is sensitive to the low-mass star population, as cooler, low-mass stars emit less blue light than red light, resulting in a larger $g - r$ value. This makes the $\text{EW}_{H\alpha}$ vs $g - r$ colour diagnostic an effective probe for the stellar IMF slope in star-forming galaxies.

We calculate $\text{EW}_{H\alpha}$ as the ratio of the $H\alpha$ flux from the emission-line fits (LZIFU) to the continuum, which is estimated following the method outlined by Cardiel et al. (1998). To correct the observed $\text{EW}_{H\alpha}$ for obscuration, we follow the prescription in Gunawardhana et al. (2011), which recovers the flux absorbed by dust by considering the gas and continuum colour excess derived from Balmer decrement ($H\alpha/H\beta$) and the obscuration curve from Cardelli et al. (1989). We use Balmer decrement maps provided by SAMI, and thus we only consider this spaxels in our sample. It is important to note that the emission line fluxes in SAMI have already been corrected for underlying stellar absorption, so no additional correction is required. Colours are corrected for dust extinction using the reddening curve from Calzetti (2001).

We employ the PÉGASE.3 population synthesis tool (Fioc & Rocca-Volmerange, 2019), based on the original PÉGASE software described in Fioc & Rocca-Volmerange (1997), to construct galaxy evolutionary tracks in the $\text{EW}_{H\alpha}$ vs $g - r_t$ space. PÉGASE is a spectral synthesis code that models the spectral evolution and chemical composition of galaxy components. Given an input IMF form and specified low- and high-mass slopes, it generates galaxy spectra, from which it computes the normalised attenuated monochromatic continuum luminosity, in-band fluxes, and equivalent widths at various galactic ages. PÉGASE models the $\text{EW}_{H\alpha}$ emission by assuming that young,

^aPost-starburst galaxies may exhibit some $H\alpha$ emission contribution from asymptotic giant branch (AGB) stars. However, given that this phase of stellar evolution is relatively short-lived, its impact on the overall $H\alpha$ emission can be considered negligible.




Figure 1. *a)* The normalised filter response functions for the *g*, *r*, and *r_t* bands. *b)* An example spectrum of a SAMI galaxy at $z = 0.025$. The blue and red regions denote the wavelength ranges of the blue and red SAMI data cubes, respectively. The vertical red lines indicate the positions of the rest-frame wavelength of the H β (left) and H α (right) emission lines. Bands have also been shifted to the rest-frame.




Figure 2. This figure presents key properties of galaxies in our SAMI sample. Panel *a)* illustrates the stellar mass distribution, while panel *b)* shows the *r*-band magnitude, panel *c)* highlights the effective radius of galaxies, and panel *d)* displays the logarithmic star formation rate (SFR) as a function of redshift (z). Panel *e)* maps the SFR-stellar mass relation, with points colour-coded by effective radius, demonstrating that the galaxies in our sample follow the typical star-forming main sequence. Our sample predominantly consists of high-stellar-mass galaxies due to the SAMI survey's mass selection criteria, its focus on low redshift, and its exclusion of compact or very small systems.

massive stars emit Lyman continuum photons, which ionise the surrounding gas. This results in nebular emission lines, including $EW_{H\alpha}$. Nebular emission is calculated using grids generated by the Cloudy code (Ferland et al., 2023). These models simulate dust-free HII regions as a function of ISM metallicity and the number of Lyman continuum photons emitted by the ionising source. The geometry used in these calculations is radiation-bounded and spherical, with constant hydrogen density throughout the ionised cloud. This allows us to create a group of evolutionary tracks in the $EW_{H\alpha}$ vs $g-r$ colour space, each of them associated with a different high-mass IMF slope. We have chosen PÉGASE as it is one of the few stellar population synthesis (SPS) tools that allows to treat the IMF as a free parameter and modify its formalism, low and high mass range.

To adequately cover the range of IMF slopes in the $EW_{H\alpha}$ vs $g-r$ colour space, we compute several evolutionary tracks using PÉGASE. We consider a fixed low-mass slope of 0.5 and a variable high-mass slope ranging from -4 to -1.5 in increments of 0.1. We consider the lower and upper stellar mass limits of our models to be $1M_{\odot}$ and $120M_{\odot}$, respectively. Other model assumptions include no infall, no extinction, no galactic winds, a non-evolving stellar metallicity of 0.02, and an exponentially declining star formation rate with an e-folding time of 1.1 Gyr. PÉGASE calculates spectra at fixed ages ranging from 0 to 20 Gyr, with increasing time steps. However, we only consider ages greater than 100 Myr, as it takes approximately that time for PÉGASE to stabilise the spectral energy distribution (SED) shape. All generated tracks are shown in panel *b*) of Figure 3. The location of the evolutionary tracks depends on several model input parameters, such as star formation histories, high-mass cutoffs and metallicities. It is not our objective to explore variations with these parameters, as they have been thoroughly investigated in prior works (Hoversten & Glazebrook, 2008, 2010; Gunawardhana et al., 2011; Nanayakkara et al., 2017, 2020). Note that as we use a truncated r colour, r_t , the evolutionary tracks are shifted towards larger (bluer) $g-r$ values, as the r magnitudes are fainter due to less flux in the r_t band compared to the full r . A comparison between three PÉGASE tracks using the original r band and the truncated r_t band is depicted in the panel *a*) of Figure 3.

The SAMI blue and red data cubes enable us to estimate the $EW_{H\alpha}$ and the $g-r_t$ colour for all star-forming spaxels in every galaxy within our sample. We developed a method to assign an IMF slope to each spaxel based on its position in the $\log(EW_{H\alpha})$ vs $g-r_t$ space. We use the evolutionary tracks from PÉGASE and their associated IMF slopes to determine the IMF slope for each data point through an inverse distance weighted interpolation (IDWI) algorithm (Shepard, 1968). IDWI is a spatial interpolation method that interpolates between a group of points by considering the distance from known points to the point to be interpolated. Let x_i be a point in the parameter space, associated with a slope value u_i . Then, the interpolated value u for an arbitrary point x is given by:

$$u(x) = \frac{\sum_{i=1}^N \omega_i(x) u_i}{\sum_{i=1}^N \omega_i(x)}, \quad (1)$$

where

$$\omega_i = \frac{1}{d(x_i, x)^m}, \quad (2)$$

and $d(x_i, x)$ is the distance between the points x and x_i . When $d(x_i, x) = 0$, then $u(x) = u_i$. The power parameter m typically ranges between 1 and 4. In this work we use $m = 3$ as higher values of m give greater importance to the closest values to the interpolated point.

In panel *a*) of Figure 4 we show how all the spaxels of our sample distribute in the $\log(EW_{H\alpha})$ vs $g-r_t$ space, colour-coded by their high-mass IMF slope. Our approach provides a novel perspective on the IMF distribution and variation in star-forming galaxies, highlighting that most data points cluster around $\alpha_{res} = -2$ and $\alpha_{res} = -2.35$ (panel *b*) of Figure 4), and that the spaxel distribution closely resembles that of galaxies (see Figure 4 in Gunawardhana et al. 2011). Notice that a group of spaxels fall outside the region enclosed by the outermost evolutionary tracks. For this set of spaxels our interpolation is inaccurate, as the IMF slope value should be extrapolated. However, only $\lesssim 8\%$ of the spaxels have this problem and thus our results should not be severely affected.

To estimate the uncertainties of the IMF slopes, we use the $g-r_t$ colour uncertainty shown in the bottom left corner of Figure 4. We randomly select 1,000 points in the $\log(EW_{H\alpha})$ versus $g-r_t$ plane, located within a rectangular box that contains the area with the highest density of points (regions with density greater than 0.288 in panel *b*) of Figure 4). For each point (x, y) , we apply the IDWI method to calculate the IMF slope at the coordinates $(x - err_{g-r_t}, y)$ and $(x + err_{g-r_t}, y)$, where err_{g-r_t} is the colour error. We then find the difference between these two slope estimates. Since the uncertainty in $EW_{H\alpha}$ is negligible and the models mainly vary along the $g-r_t$ axis, we do not consider variations in $EW_{H\alpha}$ when estimating uncertainty. The uncertainty in α_{res} is determined as the standard deviation of the differences calculated from the 1,000 points, yielding an uncertainty of $\Delta\alpha_{res} \sim 0.167$.

Since SAMI spaxels cover a spatial scale ranging from ~ 70 to ~ 700 parsecs, they encompass a larger area than a typical individual star-forming region. Consequently, concerns regarding the sampling of the initial mass function (IMF) are minimised.

We also apply the same procedure to the galaxy-integrated light measurements. Using the $1R_e$ aperture SAMI red and blue cubes, we estimate the $EW_{H\alpha}$ equivalent widths and $g-r_t$ colours, and use the same IDWI criterion to assign an IMF slope to each galaxy. We compare the distribution of spaxels versus the integrated measurements in the $\log(EW_{H\alpha})$ vs $g-r_t$ space in Figure 5. It is noteworthy that the spaxels within the same galaxy do not necessarily cluster around the integrated measurements. Additionally, both spaxels and integrated measurements exhibit a range of IMF slope values, both flatter and steeper than the Salpeter slope. These observations suggest that the IMF slope of a galaxy and the locally resolved IMF slope within that galaxy are not necessarily in a one-to-one relation. We explore the variability in α_{res} further in the next section.

We denote the IMF slope estimated for individual resolved spaxels as α_{res} , while the IMF slope derived from global or integrated galaxy light measurements from the $1R_e$ spectra is referred to as α_{int} . In future sections, $\bar{\alpha}$ will refer to the IMF slope corresponding to the coordinates of maximum spaxel density in the $\log(EW_{H\alpha})$ vs $g - r_t$ space.

4. RESULTS

Understanding the spatial distribution of the IMF slope within galaxies is essential for grasping star formation processes and galaxy evolution. The IMF slope provides insights into the stellar mass distribution, reflecting variations in star formation efficiency and the star formation history of a galaxy. By examining how the IMF slope varies across different regions of a galaxy, we can gain a deeper understanding of the underlying physical conditions and processes influencing star formation.

To address these questions, we have produced resolved IMF slope maps for each galaxy in our sample using the SAMI IFU and the IMF slope estimation method described in § 3. A selection of these maps is presented in Figure 6, along with the spatial distribution maps of other quantities. The IMF slope maps exhibit diverse spatial distributions and morphologies. For example, galaxy 271562 exhibits a radial gradient where the IMF slope is flatter (more positive) in the centre and steeper (more negative) toward the periphery. In contrast, galaxy 220515 shows the opposite trend, with a steeper IMF slope at the centre and a flatter slope toward the edges. Notably, their SFR maps differ: 271562 shows no radial gradient, whereas 220515's SFR map closely mirrors the gradient seen in its IMF slope. This underscores a relation between α_{res} and SFR which will be explored further below. Additionally, galaxy 517868 contains a localised region with steeper IMF slopes compared to the rest of the galaxy. In contrast, the maps for galaxy 618116 are relatively flat, with no clear patterns or distinct features. Lastly, galaxy 376478 lacks central star formation and exhibits a relatively flat IMF slope map with values around $\alpha_{res} \sim -2.6$. Notably, both galaxies 618116 and 376478 have very high SFRs but display steep α_{res} slopes. This suggests that when star formation is widely dispersed, the available gas is converted predominantly into numerous low-mass stars rather than a few high-mass stars, leading to more negative α_{res} values. The features observed in these maps of the α_{res} distribution are consistently reflected in the SFR, $g - r_t$, and $EW_{H\alpha}$ maps.

Overall, the IMF slope spatial distribution appears neither uniform (i.e., consistent across the galaxy, typical standard deviation of $\sigma = 0.31$) nor random (i.e., lacking any ordered pattern or smooth transition). The underlying causes of these distinct IMF slope morphologies must be analysed on a case-by-case basis.

In this section we explore the relationship between the resolved and integrated IMF slopes, as well as the link between the resolved and integrated IMF slopes and stellar mass, SFR, and Σ_{SFR} .

4.1 Links between α_{res} and α_{int}

A key innovation and advantage of our work is the capability to estimate both the global and the spatially resolved IMF-slope. This enables us to address important questions about the relationship between these two quantities. Specifically, we aim to determine whether they are related, whether they are consistent with each other, or whether one can provide context for interpreting the other.

To relate the integrated IMF-slope of each galaxy to its spatially resolved IMF-slopes, we use a representative value for the galaxy's typical resolved IMF-slope. We have chosen to use the IMF-slope associated with the point of maximum density ($\bar{\alpha}$) in the distribution of spaxels in the $\log(EW_{H\alpha})$ vs $g - r_t$ parameter space for each galaxy^b. Although we also considered alternative representative measures, such as the mean and median of all resolved slopes, our results using these alternatives did not show significant differences. However, $\bar{\alpha}$ is preferred because it reflects the central tendency of the resolved IMF-slope distribution. This quantity provides a robust indicator of the typical IMF-slope value across different regions of the galaxy, whereas the mean and median can be influenced by outliers or skewed distributions.

The comparison between the integrated IMF-slope (α_{int}) and the resolved IMF-slope ($\bar{\alpha}$) is illustrated in Figure 7. A notable observation is that many data points cluster close to the one-to-one line (indicated by the black dotted line). For most galaxies, $\bar{\alpha}$ tends to be flatter than the integrated slope. However, when the integrated slope is flatter than approximately $\alpha_{int} > -2.7$, the relationship between the two slopes becomes closer to the one-to-one line.

This observation is further illustrated in Figure 8. We find that 71% of the galaxies have a flatter $\bar{\alpha}$ than their integrated slope ($\alpha_{int} - \bar{\alpha} < 0$), while 29% of the galaxies show the opposite trend, with a flatter $\bar{\alpha}$. Additionally, approximately 93% of the galaxies have $|\alpha_{int} - \bar{\alpha}| < 0.5$, and $|\alpha_{int} - \bar{\alpha}| < 0.25$ for about 81%. A difference of this magnitude is comparable to our typical uncertainties in the estimate of α_{res} ($\Delta\alpha_{res} \sim 0.167$), and so is unlikely to be significant. This only indicates that the distribution of α_{res} closely aligns with the value of α_{int} , rather than implying that there is no variation in the slope within galaxies. This comparison validates our approach using resolved data, allowing us to effectively compare our findings with previous results where spatially resolved data is unavailable.

We also differentiate, in Figure 7, between galaxies with central star formation (represented by pink circles) and those without it (represented by black crosses). Galaxies with central star formation are defined as those in which the central $0.5R_e$ region contains spaxels with a S/N at $4100\text{\AA} > 3\text{pix}^{-1}$, and at least 10% of these spaxels classified as star-forming or intermediate (those where emission likely results from a combination of star-forming and non-star forming mechanisms). Spaxel classification is based on the detection of specific emission lines,

^bIn this approach, we calculate the maximum density coordinates of the spaxels of each galaxy in the $\log(EW_{H\alpha})$ vs $g - r_t$ parameter space, and we then assign a slope value to this point using the IDWI method.




Figure 3. *a)* A comparison of the PEGASE evolutionary tracks generated using the original SDSS g and r band filters (dashed lines) with those generated using the truncated r_t filter (solid lines). Each colour represents a track with a different IMF slope: red for $\alpha = -3.0$, green for $\alpha = -2.35$, and blue for $\alpha = -2.0$. The truncated $g-r_t$ colour is shifted to the right compared to the original $g-r$ colour due to the reduced flux in the r_t band. *b)* The $\log(\text{EW}_{\text{H}\alpha})$ versus $g-r_t$ parameter space, populated by all generated PEGASE evolutionary tracks with the truncated filters. Each track corresponds to a different high-mass IMF slope, ranging from $\alpha = -1.5$ (top track) to $\alpha = -4.0$ (bottom track) in increments of 0.01.




Figure 4. Panel (a) presents the $\log(\text{EW}_{\text{H}\alpha})$ versus $g-r_t$ colour space, populated by spaxels from all galaxies in the sample. Each spaxel is colour-coded according to its high-mass IMF slope (α_{res}), assigned through inverse distance weighted interpolation criterion. The black lines correspond to evolutionary tracks from the PEGASE models, with each track representing a different high-mass IMF slope (flatter IMFs to the top-right). The thick black tracks highlight slopes of -2 , -2.35 , and -3 from top to bottom. A red cross in the bottom left corner indicates representative uncertainties along each axis. Panel (b) shows the density contours of the spaxel distribution from panel (a), emphasising that the bulk of the data points lie within the region covered by the PEGASE models. Notably, a significant portion of the data falls between the tracks associated with slopes of -3 and -2.35 .

using either the $[\text{NII}] (6583\text{\AA})/\text{H}\alpha$ versus $[\text{OIII}] (5007\text{\AA})/\text{H}\beta$ BPT diagram, or just the $[\text{NII}] (6583\text{\AA})/\text{H}\alpha$ ratio. The spectral classification methodology for spaxels and galaxies is described in detail in Owers et al. (2019).

In our sample, 182 objects exhibit no central star formation,

while 1,162 objects do. Most galaxies without central star formation sit outside the bulk of the data points. For these galaxies, the integrated light is likely dominated by emission from older stellar populations rather than recent star formation. Consequently, the Kennicutt method may be less reliable in




Figure 5. The distribution of spaxels from six galaxies in our sample within the $\log(\text{EW}_{\text{H}\alpha})$ versus $g-r_t$ colour parameter space. Spaxels from the same galaxy are represented as squares of the same colour, while the integrated galaxy measurements of $\log(\text{EW}_{\text{H}\alpha})$ and $g-r_t$ (obtained using the $1-R_e$ aperture spectra) are shown as stars. The black lines correspond to three PEGASE tracks with $\alpha = -2$ (top), $\alpha = -2.35$ (middle) and $\alpha = -3$ (bottom).

these cases, potentially leading to the inferred steeper IMF slope.

We investigate the resolved IMF slope distributions in greater detail for six galaxies in Figure 9. These galaxies are randomly selected as a representative collection to illustrate the distributions of α_{res} within galaxies. In this figure, α_{int} is represented by a coloured vertical line, while the Salpeter slope is indicated by a black line for reference. Each galaxy exhibits a distinct range of slope distributions, with some showing broader or narrower slope ranges. Notably, the peak of the slope distribution does not necessarily align with either the Salpeter slope or the integrated slope.

To understand why the breadth of slope distributions varies among galaxies, we analyse the relationship between the IMF slope and other resolved properties of the galaxies in the next section. If these properties are related to the IMF slope, they could help explain the observed differences in the shapes of the slope distributions.

4.2 Relation between α_{res} and other resolved properties

Given the direct link between the IMF slope and the star formation process, it is reasonable to anticipate that it may be associated with parameters such as the SFR and the Σ_{SFR} . This section aims to quantify the relationship between these quantities more systematically, looking for an IMF slope driver.

In Figure 10 and Figure 11, we analyse the relationship

between α_{int} and the global SFR, as well as α_{res} and α_{int} with the resolved and global Σ_{SFR} , respectively. We do not compare α_{res} and the SFR per spaxel as this is fundamentally giving the same information than Σ_{SFR} . To facilitate this analysis, we divide our sample into bins based on SFR and Σ_{SFR} . Bins span $\log(\text{SFR}) = [-5.5, -1.5]$ and $\log(\Sigma_{\text{SFR}}) = [-4.5, -0.5]$ in steps of 0.5. For each bin, we determine a representative IMF slope ($\bar{\alpha}_{\text{res}}$ and $\bar{\alpha}_{\text{int}}$) by identifying the maximum density coordinates of all spaxels (for resolved IMF slopes) or galaxies (for integrated IMF slopes) within that bin. The IMF slope at each coordinate is then assigned using the IDWI criterion, as outlined in § 3.

It is evident that the IMF slope exhibits a strong correlation with both the SFR and Σ_{SFR} , with a trend toward flatter $\bar{\alpha}_{\text{int}}$ at higher values of SFR and Σ_{SFR} . In Figure 10, we additionally examined the IMF slope as a function of SFR across three stellar mass bins: $M < 10^9 M_{\odot}$, $10^9 M_{\odot} \leq M \leq 10^{10} M_{\odot}$, and $M > 10^{10} M_{\odot}$. Our analysis revealed that although all mass bins display a similar qualitative trend, lower-mass galaxies tend to have flatter slopes at a given SFR compared to mid-mass and high-mass galaxies.

When comparing our results to other published work, we find that our measurements fill the gap between the low-SFRs probed by Weidner *et al.* (2013) and the higher SFRs from Lee *et al.* (2009) and Gunawardhana *et al.* (2011). It is important to note that our integrated measurements are derived from a $1 R_e$ aperture, whereas previous studies use a variety of photometric measurements that would correspond to a range of different effective apertures. This may introduce subtle systematics in this comparison, but given the challenges in any similar IMF analysis, these are unlikely to be a significant contributor.

As illustrated by the black line in Figure 11, there is a clear relationship between $\bar{\alpha}_{\text{res}}$ (panel *a*) and $\bar{\alpha}_{\text{int}}$ (panel *b*) with Σ_{SFR} , where higher Σ_{SFR} values are associated with flatter IMF slopes. Similar to the slope-SFR relationship, the integrated IMF slope exhibits a mass dependency with respect to Σ_{SFR} : high-mass galaxies tend to have steeper slopes compared to mid- and low-mass galaxies for a given Σ_{SFR} . However, this mass dependency is not observed in the resolved IMF slopes with respect to Σ_{SFR} . This suggests that Σ_{SFR} may be an underlying physical driver of the IMF. This points to a scenario where the IMF is more closely linked to local parameters such as Σ_{SFR} rather than global properties like galaxy mass. Further exploration of the cause of this relationship is discussed below in § 5.

By fitting a linear model to the relationship between resolved IMF slopes and Σ_{SFR} across all mass ranges (depicted by the solid pink line), we derive the following general equation:

$$\alpha_{\text{res}} = 0.26 \log(\Sigma_{\text{SFR}}) - 2.03 \pm 0.036, \quad (3)$$

where Σ_{SFR} is in units of $M_{\odot} \text{ yr}^{-1} \text{ kpc}^{-2}$. To estimate the uncertainty in Equation 3, we employed a Jackknife resampling method. Specifically, we randomly removed 10% of the data points from each bin of Σ_{SFR} and recalculated the linear fit multiple times. We then use the standard deviation of slopes (σ_A) and intercepts (σ_B) from these repeated fits to

estimate the total uncertainty of the linear fit as $\text{ERR}_{\alpha_{\text{res}}} = \sqrt{[\log(\Sigma_{\text{SFR}}) \cdot \sigma_A]^2 + \sigma_B^2}$.

In Figure 12, we further examine the distribution of individual spaxels in the α_{res} -SFR space. As expected, panel *b*) of Figure 12 shows that spaxels are clustered around the binned relation lines of $\bar{\alpha}_{\text{res}}$ with $\log(\text{SFR})$, as depicted in Figure 11. Notably, a mass dependence emerges within the α_{res} -SFR relationship. The steep α_{res} values observed around $\log(\Sigma_{\text{SFR}}) \sim -5$ likely arise from limitations in our slope-estimation methods and the coverage of our SPS models.

This figure underscores that the observed slope-SFR relation in previous figures is not merely an artefact of the binning method but reflects a genuine physical connection between these quantities. Spaxels follow a moderate correlation with SFR, with a Spearman correlation coefficient of $\rho_{\text{SFR}} = 0.415$ ($p \sim 0$). Although these relation exhibit significant scatter, much of this variability is likely due to measurement uncertainties. For example, the colour error bar in Figure 4 spans several tracks, corresponding to $\Delta\alpha_{\text{res}} \sim 0.167$. Despite this, we focus more on the observed trends rather than the exact numerical values. The strong correlations depicted in Figures 10 and 11 suggest a general relationship between the IMF slope and both SFR and Σ_{SFR} , applicable to both resolved and global slopes.

4.3 Radial dependency of α_{res}

To investigate a potential radial dependency of α_{res} , we calculated the average value of α_{res} for spaxels within each radial bin. Each bin is defined by a radius that encompasses 20%, 40%, 60%, 80%, and 100% of the spaxels. The results are presented in Figure 13, which shows the radial profiles of α_{res} for galaxies in three distinct mass bins. For visualisation purposes, each panel displays profiles for a representative subset of 75 randomly selected galaxies.

Using the $\log(\Sigma_{\text{SFR}})$ -radius (R) relations from Medling et al. (2018) for galaxies on the main sequence (see their Figure 7), combined with our derived relationship between $\log(\Sigma_{\text{SFR}})$ and α_{res} (equation 3), we predict the relationship between radius and α_{res} . This predicted relation is the weak radial dependence shown as the red lines in Figure 13. While some galaxies do exhibit variations in α_{res} with radius, the majority of objects have relatively flat radial profiles with minimal or no variation. The predicted shape is only marginally consistent with our measured values of α_{res} for $R/R_e < 2$. Given the uncertainties in our measurements, though, this relation may still be consistent with what we find, but there is clearly no strong signal across the sample investigated here.

Additionally, Figure 13 includes the mean α_{res} values calculated across six radial bins, ranging from $R/R_e = 0$ to 3 in increments of 0.5. These mean values align somewhat more closely with the predicted relation (red lines), suggesting that there may be a weak underlying radial trend in α_{res} that is not immediately evident in individual profiles, although this is probably not quite as strong as predicted.

5. DISCUSSION

5.1 Limitations on the method

Given the innovative aspects of our approach, it is crucial to recognise the inherent limitations and potential sources of uncertainty within our methodology.

The accuracy of our results is highly dependent on the capabilities of the SPS tool employed. We opted to use the PÉGASE SPS tool primarily due to its flexibility in modifying the IMF functional form, adjusting lower and upper mass ranges, SFHs, and slopes. However, PÉGASE does not account for certain stellar properties, such as binary interactions and rotation (Leitherer et al., 1999; Stanway et al., 2016; Eldridge & Stanway, 2012; Eldridge et al., 2017), which could limit its accuracy in comparison to more complete SPS tools and may introduce some degree of uncertainty in our IMF estimation.

As discussed by Conroy et al. (2013), systematic uncertainties in SPS models can lead to variations in IMF estimates. Nevertheless, previous studies by Gunawardhana et al. (2011) and Nanayakkara et al. (2017, 2020) have demonstrated that results obtained with PÉGASE are not significantly different from those generated by other SPS tools, such as STARBURST99 (Leitherer et al., 1999). The underlying limitation remains that spectral synthesis models are constrained by the accuracy of the stellar models they incorporate (Hoversten & Glazebrook, 2010). Therefore, while the specific quantitative results may vary when using different population synthesis tools, we expect the qualitative trends observed in our work to remain robust.

Given that this work is among the first to apply this methodology for estimating the IMF on a spaxel-by-spaxel basis, it is still too early to fully understand whether the methods for estimating the IMF encounter additional limitations when applied at more resolved scales. Our IMF estimation method may require a higher signal-to-noise ratio (S/N) on smaller scales to retrieve reliable results. One advantage of conducting the analysis with integrated light is that the effects of different stellar populations are smoothed out, as noted by Hoversten & Glazebrook (2008). However, a more localised approach may require special attention to the distinct stellar populations present. As a galaxy evolves, the stellar population present in any given spaxel is likely to consist of a mix of populations that formed with different ages, metallicities, and potentially IMF shapes. This could significantly impact the accuracy of our estimation of α_{res} . Conversely, the similar results that we see, broadly speaking, between α_{res} and α_{int} are reassuring, suggesting that this effect is not introducing a significant bias or systematic. Also, in this analysis we estimate the high mass IMF slope, which is dominated by short lived high mass stars. The lifetime of such stars is negligible compared to the rotation speed of galaxies that would drive population mixing. This strongly mitigates against any biases arising through the possibility of population mixing, which may be a more substantial concern when analysing the low mass, long-lived, stellar populations. Further studies are needed to explore these effects in greater detail.

Another limitation of our IMF estimates lies in the method we use to assign a slope to each spaxel or galaxy based on




Figure 6. This figure showcases maps of five example galaxies, each represented by a column. From top to bottom, the rows display the following quantities: α_{res} , $\log(\text{SFR})$, $g - r_t$ colour, and $\log(\text{EW}_{\text{H}\alpha})$. These galaxies were selected to highlight diverse spatial features. Galaxies 220515 and 271562 exhibit clear radial gradients, while 517868 shows a non-central region with steeper IMF slopes. In contrast, galaxy 618116 has a relatively uniform (flat) distribution. Finally, galaxy 376478, which lacks central star formation, displays a range of steep IMF slopes. These examples illustrate the structural diversity and star formation characteristics present in the sample.




Figure 7. This figure illustrates the relationship between the high-mass IMF slope ($\bar{\alpha}$) and the integrated IMF slope (α_{int}), focusing on galaxies where at least 50% of the spaxels are classified as star-forming. Pink circles represent galaxies with central star formation, while black crosses indicate galaxies without central star formation. The black dotted line denotes the one-to-one relationship, serving as a reference for comparison. Additionally, the green and blue lines show the best linear fits for galaxies with and without central star formation, respectively. The red cross in the corner indicates the typical uncertainties in the IMF. This plot highlights the differences between local and global IMF slopes.




Figure 8. The distribution of differences between the integrated IMF slope (α_{int}) and the $\bar{\alpha}$ across our sample.

its location in the $\log(EW_{H\alpha})$ vs $g - r_t$ space. Although we have a range of spectral evolution models that encompass the majority of the objects of interest, a certain percentage falls outside the region covered by these models. For those spaxels, our interpolation model becomes unreliable, as extrapolation would be required. However, extrapolated points may deviate significantly from what theoretical models would predict. It is possible to generate additional models to cover the portions of the $\log(EW_{H\alpha})$ vs $g - r_t$ space that are not currently spanned by including different parameters, such as varying SFHs and metallicities. Nevertheless, as mentioned above, this issue affects only a small fraction of our data (less than 8% of all spaxels) and does not significantly impact our overall conclusions. In a similar vein, Nanayakkara et al. (2020) demonstrated that incorporating stochastic starburst events into the SFHs of galaxies can lead to higher $EW_{H\alpha}$ values. This would result in a modified distribution of the models in the $\log(EW_{H\alpha})$ versus $g - r_t$ space. This suggests that some portion of the variation in the inferred IMF slopes seen in our results may actually stem from differences in SFHs, metallicities or other properties. Incorporating the full suite of SFHs and metallicities as additional parameters would significantly increase the complexity of any investigation, and is beyond the scope of this current analysis.

Finally, we note that we compared the resolved slope of star-forming spaxels with the slope obtained from the integrated light of galaxies, which includes both star-forming and non-star-forming spaxels, and thus different stellar populations. In future work, we aim to explore methods to estimate the slope in these non-star-forming spaxels to construct complete




Figure 9. This figure displays the distribution of the high-mass IMF resolved slope (α_{res}) for the six coloured galaxies presented in Figure 5. Each distribution is represented by a kernel density estimation (KDE) shown as a black line. The vertical black line indicates the Salpeter slope ($\alpha = -2.35$), while the vertical coloured lines represent the integrated slopes for each galaxy. In the top left corner of each panel, the mean and standard deviation of the slope distribution are provided. Additionally, the α_{res} map for each galaxy is displayed in the top right corner, offering a visual context for the slope distributions.




Figure 10. This figure illustrates the relationship between the integrated high-mass IMF slope ($\bar{\alpha}_{\text{int}}$) and the logarithm of the star formation rate ($\log(\text{SFR})$) across all galaxies in the sample. The colour of each line denotes different stellar mass bins, highlighting the variations in slope with respect to star formation activity. Uncertainties are determined using a Jackknife resampling method. Additionally, results from previous studies by Lee *et al.* (2009), Gunawardhana *et al.* (2011), and Weidner *et al.* (2013) are represented in purple for comparison. The Salpeter slope is shown as an orange line for reference.




Figure 11. This figure depicts the relationship between the resolved high-mass IMF slope (panel a) and the integrated high-mass IMF slope (panel b) against $\log(\Sigma_{\text{SFR}})$. We make use of eight uniform bins of Σ_{SFR} , associating an IMF slope to the coordinate of maximum density within the distribution of spaxels/galaxies in each bin. The colour of each line corresponds to different stellar mass bins. Uncertainties in the relationships are derived through Jackknife resampling resampling within each bin. The pink line in both panels represents the best linear fit for the relationship between the resolved $\bar{\alpha}$ and $\log(\Sigma_{\text{SFR}})$ considering all stellar masses. The orange line indicates the Salpeter slope for reference.




Figure 12. Panel (a) displays the resolved high-mass IMF slope for each spaxel as a function of $\log(\text{SFR})$. The black lines represent the $\bar{\alpha}_{\text{res}}-\log(\Sigma_{\text{SFR}})$ relations derived from Figure 11. The red cross in the bottom right corner indicates the uncertainties along both axes. Panel (b) illustrates the contours of the spaxel distributions from panel (a). It is evident that the majority of the data points align closely with the black lines. The steep $\bar{\alpha}_{\text{res}}$ values observed around $\log(\Sigma_{\text{SFR}}) \sim -5$ likely arise from limitations in our slope-estimation methods and the coverage of our SPS models. For reference, the orange line indicates the Salpeter slope.




Figure 13. Radial profiles of the resolved high-mass IMF slope (α_{res}) for 75 randomly selected galaxies across different mass bins. In each panel, the black lines represent the average α_{res} values calculated for spaxels within defined radial bins. These radial bins are determined by radii that encompass annuli spanning 20%, 40%, 60%, 80%, and 100% of the total spaxels in each galaxy. The green lines represent the mean α_{res} values of the radial profiles across 6 radial bins, ranging from $R/R_e = 0$ to 3 in increments of 0.5. The red lines indicate the predicted relationship between radius (R) and α_{res} , derived from the $\Sigma_{\text{SFR}}-R$ relation of Medling et al. (2018) and our $\Sigma_{\text{SFR}}-\alpha_{\text{res}}$ relation (as outlined in equation 3). The purple cross in the corner represents the uncertainties in the IMF. For reference, the orange line denotes the Salpeter slope.

IMF slope maps and make more robust comparisons.

5.2 Understanding the link between α_{res} and α_{int}

The IMF has often been assumed to be universal, typically adopting the Salpeter slope ($\alpha = -2.35$), likely because it provides a simple approach and reasonably accurate assumption. However, there is an overwhelming body of evidence suggesting the necessity of considering variable IMFs.

These findings suggest that the IMF is influenced by various properties, driving its variability (Martín-Navarro et al., 2019; Barbosa et al., 2021; Martín-Navarro et al., 2021, 2023). Given that star formation conditions within a galaxy can vary significantly (e.g., different metallicities, magnetic field intensities, turbulence), a critical question arises: at what scale does the IMF begin to vary? Specifically, do the local variations in the IMF (e.g., at the scale of molecular clouds or star clusters) average out when considering the global scale (e.g., an entire galaxy), or does the global IMF robustly depend on local variations, leading to IMF differences between galaxies? The more fundamental question is whether there is a (continuous) connection between the local and global IMF.

The qualitative consistency observed between $\bar{\alpha}_{res}$ and $\bar{\alpha}_{int}$ in their relationships with SFR and Σ_{SFR} (Figures 10 and 11) suggests a fundamental connection between the local and global IMF properties within galaxies. While the explicit nature of this relationship requires further exploration, our findings indicate that local variations in the IMF contribute to the formation of the global IMF signature.

As illustrated in Figure 8, the difference between $\bar{\alpha}$ and α_{int} in our sample follows a distribution characterised by a mean of $\mu = -0.124$ and a standard deviation of $\sigma = 0.27$. The typical standard deviation of the resolved slope within a galaxy is approximately 0.34. It is important to note that galaxies lacking central star formation generally exhibit lower values of α_{int} ($\alpha_{int} < -2.7$) while maintaining consistent $\bar{\alpha}$ values (Figure 7). This phenomenon is likely attributable to the influence of central pixels on the α_{int} value. These non-star-forming central pixels are likely contributing light from older stellar populations to the overall galaxy light. Given that the Kennicutt method relies on the strength of the H α line and the $g-r_t$ colour, this older stellar population may introduce bias into our IMF slope estimation. Supporting this explanation is the observation that, in galaxies with central star formation, $\bar{\alpha}$ and α_{int} tend to align more closely with the one-to-one relation.

It is important to acknowledge the assumptions made to facilitate the comparison between the resolved and integrated IMF slopes. First, we represent all the resolved slopes by a single value, $\bar{\alpha}$, for each galaxy, which does not represent a direct comparison between the full distribution of α_{res} with α_{int} . Future studies could explore more appropriate methods to address this limitation. Second, our analysis focuses solely on spaxels associated with star formation, excluding those without such activity. This introduces a potential bias when comparing the slopes inferred from this sample to those estimated from the integrated galaxy light, which includes contributions from

older stellar populations and may bias our IMF estimation less reliable. To mitigate this bias, we ensured that all galaxies in our sample have over 50% of their spaxels classified as star forming.

Despite all these limitations, we still find a reasonable agreement between α_{res} and α_{int} . Our results suggest that local IMF variations significantly contribute to the global IMF signature, highlighting the importance of considering variable IMFs in understanding star formation across different scales.

5.3 The connection with SFR and Σ_{SFR}

One of our key findings is the relationship between the IMF slope (both resolved and integrated) and the SFR and Σ_{SFR} , where galaxies with higher SFR and Σ_{SFR} exhibit flatter IMF slopes. Since the IMF is directly linked to the star formation process, the observed relation between the IMF slope and these star formation indicators is likely associated with various factors critical to star formation. Star formation begins with the fragmentation and collapse of cold molecular clouds (Krumholz et al., 2019). Perturbed velocity fields and turbulence induce shocks, creating regions of compressed gas. As the density increases, gravitational collapse eventually overcomes turbulent support (McKee & Ostriker, 2007). Consequently, the physical conditions of the ISM—such as turbulence, feedback, outflows, gas cooling/heating, magnetic fields, and metallicity—are expected to significantly impact star formation efficiency (Bate & Bonnell, 2005; Gouliermis, 2018; Grasha et al., 2017).

The Jeans mass of an unperturbed molecular cloud represents the threshold at which the system becomes gravitationally bound. In regions with a high star formation rate (SFR $> 3-5M_{\odot}yr^{-1}$), radiation from hot stars can warm the surrounding dust, which then thermally couples with the gas. This process leads to an increase in the Jeans mass in high-density and high-temperature regions, as more mass is required to overcome the thermal and kinetic energy of the dust and gas, making star formation more difficult. Consequently, the SFR and Jeans mass are closely interconnected. As early as 1992, Larson (1992, 1998, 2005) proposed a connection between the IMF and the Jeans mass. An increase in the Jeans mass results in fewer clumps forming within a star-forming cloud, but these clumps will concentrate more mass. This leads to a flatter high-mass IMF slope (Bonnell et al., 2006). This scenario helps explain why galaxies 618116 and 376478 in Figure 6 exhibit steep α_{res} values despite having high SFRs. When the Jeans mass is low, star formation becomes more efficient, resulting in the formation of numerous low-mass stars while inhibiting the formation of high-mass stars. This process consequently steepens the slope of the IMF.

Bate & Bonnell (2005) conducted hydrodynamical simulations to model the formation of stellar systems from the collapse of molecular clouds. These outcomes were compared to earlier calculations (Bate et al., 2003), where the Jeans mass was a factor of three higher than the original value of $1 M_{\odot}$. The comparison revealed that denser clouds tend to produce a higher proportion of brown dwarfs and exhibit higher velocity dispersions. The study found that the interaction between ac-

cretion and ejection processes could replicate low-mass IMFs, while variations in the magnitude of dispersion in the accretion rates of individual objects would primarily affect the slope of the high-mass IMF. Specifically, the high-mass IMF slope becomes flatter in denser star-forming regions. This finding aligns closely with our results, as well as those of Gunawardhana *et al.* (2011). Additionally, Bonnell *et al.* (2006) determined that the “knee” of the IMF is approximately determined by the Jeans mass, and that a barotropic equation of state can produce a realistic IMF for high Jeans masses.

Hydrodynamical simulations by Narayanan & Davé (2012) suggest that variations in the IMF, particularly through the scaling of Jeans mass with the IMF’s characteristic mass, could potentially reconcile discrepancies between observed and model-predicted star formation rates. However, they acknowledge that these differences could also be addressed without assuming a varying IMF. Magnetic fields play a critical role in star formation, as shown by Sharda *et al.* (2020), who found that strong magnetic fields can suppress gas cloud fragmentation, reducing the formation of low-mass stars. Despite this, other studies indicate that the impact of magnetic fields on star formation may not be as significant as radiative feedback (Bate, 2011; Myers *et al.*, 2013) and thermodynamic processes Lee & Hennebelle (2018).

We find that the high-mass slope of the IMF is primarily influenced by the SFR surface density, which may be explained by the formation of high mass stars altering the Jeans mass and fragmentation within molecular clouds. This is consistent with the Kennicutt-Schmidt law (Schmidt, 1959; Kennicutt, 1989), which posits a strong correlation between SFR and gas density. While it is beyond the scope of this work to determine which parameter or process predominantly suppresses star formation, evidence indicates that magnetic fields and radiative feedback have a non-negligible impact on star formation.

The observed stellar mass dependency in the slope-SFRD relation for integrated measurements, as shown in Figure 11, supports the idea that the IMF slope is determined by local processes. If the resolved IMF slopes depended on the galaxy’s total stellar mass, we would need to explain how different regions across a galaxy adjust their IMF values to ensure alignment with such a global mass dependency trend. The presence of a mass dependency in the α_{int} vs Σ_{SFR} relation suggests that high-mass galaxies tend to have steeper IMF slopes, likely due to their older stellar populations, arising from local drivers. Gunawardhana *et al.* (2011) find a relationship between their α_{int} and specific star formation rate, which may also indicate an underlying local mass dependency.

There is a well-established relationship between Σ_{SFR} and galactocentric radius, as demonstrated by Medling *et al.* (2018). In this work, we have identified a relationship between the IMF slope and Σ_{SFR} , leading us to anticipate a corresponding relationship between the IMF slope and galactocentric radius. Indeed, studies by Martín-Navarro *et al.* (2015a, 2019), Davis & McDermid (2017), Vaughan *et al.* (2018), and van Dokkum *et al.* (2017) have shown that massive early-type galaxies exhibit a radial dependency of the IMF, with α_{res} becoming steeper towards the outskirts of the galaxy. Our results,

though, suggest a lack of significant radial dependencies in the spatial distribution of the IMF slope. We believe this is due to the high scatter in the α_{res} vs Σ_{SFR} relation (solid black line in panel a) in Figure 12), which likely weakens any radial dependencies. Further work is needed to more precisely quantify the spatial variations of the IMF slope. Recent work by Martín-Navarro *et al.* (2023), which presents a low-mass IMF slope map for a star-forming galaxy, also indicates minimal radial dependency of the IMF slope. Radial IMF gradients have only been observed in quiescent ETGs so far, and only probe the low-mass end of the IMF, a regime distinct from the high-mass slope we focus on here. Further work is needed to verify the existence of radial gradient of the IMF high-mass slope of star forming galaxies. Given the precision of our α_{res} estimations, such gradients would only be detectable if they are sufficiently pronounced to overcome the limitations of our current accuracy.

6. SUMMARY

We have conducted the most extensive systematic analysis to date on the distribution of the high-mass IMF slope with spatially resolved spectroscopic galaxy data. We have used data from star-forming galaxies taken from the SAMI survey. The high-mass IMF slope is estimated using the Kennicutt method. To achieve this, we employ the PÉGASE population synthesis tool to generate several galaxy evolutionary tracks, each corresponding to a different IMF slope. We then use a spatial interpolation technique (IDWI) to assign a slope to spaxels and galaxies based on their location in the $\log(EW_{H\alpha})$ vs $g - r_i$ colour space.

Our study examines the relationship between the spaxel-scale resolved IMF slope and the global integrated galaxy light IMF slope, as well as the variations in the resolved IMF slope across different galaxies. We aim to explain these variations by exploring the connection between the IMF slope and galaxy properties such as SFR and Σ_{SFR} . Our findings indicate that:

- The IMF of local regions within a galaxy are largely consistent with the global IMF of the galaxy, particularly for galaxies with $\alpha_{int} \gtrsim -2.7$. However, the steepest IMF slopes derived from global measurements may be overestimated due to the contribution of spaxels with flatter IMF slopes in the inner regions of galaxies, particularly in those without central star formation.
- We attribute the higher difference between $\bar{\alpha}$ and α_{int} for steeper global IMF slopes to the reliability of the Kennicutt method, which depends on strong H α emission. Galaxies with higher star formation rates tend to have both a flatter IMF slope and stronger H α lines, thereby enhancing the accuracy of IMF slope estimation. In contrast, galaxies with lower star formation rates may lack strong H α lines, leading to greater uncertainty in our IMF estimates.
- We find a variety of values of α_{res} within galaxies, (Figure 9). While $\bar{\alpha}$ is typically not significantly different from the integrated IMF, the resolved IMF slope distribution can be wide, narrow, or even bimodal. Some galaxies have

a radial gradient with α_{res} , although the population as a whole does not show a strong radial trend.

- We find a strong correlation between the resolved and integrated IMF slopes with SFR and Σ_{SFR} (Figure 11). Specifically, a flatter/steeper IMF slope is associated with higher/lower SFR and Σ_{SFR} , both for local and global scales. We introduce Equation 3 to link α_{res} to Σ_{SFR} .
- The $\bar{\alpha}_{int}$ -SFR relation (Figure 10) exhibit a mass dependency, where lower stellar mass galaxies tend to have shallower slopes at a given SFR. This mass dependency is also evident in the relation between $\bar{\alpha}_{int}$ and Σ_{SFR} (Figure 11), but it goes away when considering $\bar{\alpha}_{res}$. We suggest that regions with higher star formation density, in particular those characterised by more concentrated star formation, are likely to produce a larger number of high-mass stars, leading to a flattening of the high-mass IMF slope. This scenario would require an increase in the Jeans mass, which could be driven by factors such as turbulence, magnetic fields or other ISM processes.

The aim of this work is to present an analysis of the high-mass IMF slope on a spaxel-by-spaxel scale, with the goal of understanding the relationship between the IMF at this scale and the global IMF of galaxies. Additionally, we examine how the variability of the resolved slope correlates with other galaxy properties such as SFR and Σ_{SFR} . Comparisons between recent JWST observations and semi-analytical models provide evidence supporting a top-heavy IMF at $z > 10$ (Yung et al., 2024). These findings support our results, as the early universe features higher star formation density, and our study indicates a flatter high-mass IMF for in such environments. Future work will focus on enhancing the rigour of our approach by considering the SFH as a variable and exploring methods to estimate both the low-mass and high-mass resolved IMF slopes in regions with and without star formation. This could potentially allow us to further investigate differences in the distribution of the resolved slopes in passive and star forming galaxies across various environments. With this information, we can gain deeper insights into the origin of the IMF and its relationship not only with galaxy properties but also with their environment.

Acknowledgement

The SAMI Galaxy Survey is based on observations made at the Anglo-Australian Telescope. The Sydney-AAO Multi-object Integral field spectrograph (SAMI) was developed jointly by the University of Sydney and the Australian Astronomical Observatory. The SAMI input catalogue is based on data taken from the Sloan Digital Sky Survey, the GAMA Survey and the VST ATLAS Survey. The SAMI Galaxy Survey is supported by the Australian Research Council Centre of Excellence for All Sky Astrophysics in 3 Dimensions (ASTRO 3D), through project number CE170100013, the Australian Research Council Centre of Excellence for All-sky Astrophysics (CAASTRO), through project number CE110001020, and other participating institutions. The SAMI Galaxy Survey website is <http://sami-survey.org/>.

References

- Allen, J. T., Croom, S. M., Konstantopoulos, I. S., et al. 2014, *Monthly Notices of the Royal Astronomical Society*, 446, 1567–1583
- Barbosa, C. E., Spiniello, C., Arnaboldi, M., et al. 2021, *A&A*, 649, A93
- Bastian, N., Covey, K. R., & Meyer, M. R. 2010, *ARA&A*, 48, 339
- Bate, M. R. 2011, *Monthly Notices of the Royal Astronomical Society*, 419, 3115–3146
- Bate, M. R., & Bonnell, I. A. 2005, *MNRAS*, 356, 1201
- Bate, M. R., Bonnell, I. A., & Bromm, V. 2003, *MNRAS*, 339, 577
- Bonnell, I. A., Clarke, C. J., & Bate, M. R. 2006, *MNRAS*, 368, 1296
- Brown, T., & Wilson, C. D. 2019, *The Astrophysical Journal*, 879, 17
- Bryant, J. J., Owers, M. S., Robotham, A. S. G., et al. 2015, *MNRAS*, 447, 2857
- Buat, V., Donas, J., & Deharveng, J. M. 1987, *A&A*, 185, 33
- Bundy, K., Bershad, M. A., Law, D. R., et al. 2015, *ApJ*, 798, 7
- Calzetti, D. 2001, *PASP*, 113, 1449
- Cappellari, M., McDermid, R. M., Alatalo, K., et al. 2012, *Nature*, 484, 485
- Cardelli, J. A., Clayton, G. C., & Mathis, J. S. 1989, *ApJ*, 345, 245
- Cardiel, N., Gorgas, J., Cenarro, J., & Gonzalez, J. J. 1998, *A&AS*, 127, 597
- Chabrier, G., Hennebelle, P., & Charlot, S. 2014, *The Astrophysical Journal*, 796, 75
- Conroy, C., Dutton, A. A., Graves, G. J., Mendel, J. T., & van Dokkum, P. G. 2013, *ApJ*, 776, L26
- Conroy, C., & van Dokkum, P. G. 2012, *ApJ*, 760, 71
- Croom, S. M., Lawrence, J. S., Bland-Hawthorn, J., et al. 2012, *MNRAS*, 421, 872
- Croom, S. M., Owers, M. S., Scott, N., et al. 2021, *MNRAS*, 505, 991
- Davis, T. A., & McDermid, R. M. 2017, *MNRAS*, 464, 453
- Dib, S., Shadmehri, M., Padoan, P., et al. 2010, *MNRAS*, 405, 401
- Driver, S. P., Norberg, P., Baldry, I. K., et al. 2009, *Astronomy and Geophysics*, 50, 5.12
- Driver, S. P., Bellstedt, S., Robotham, A. S. G., et al. 2022, *MNRAS*, 513, 439
- Eldridge, J. J., & Stanway, E. R. 2012, *MNRAS*, 419, 479
- Eldridge, J. J., Stanway, E. R., Xiao, L., et al. 2017, *PASA*, 34, e058
- Elmegreen, B. G. 2009, in *IAU Symposium*, Vol. 254, *The Galaxy Disk in Cosmological Context*, ed. J. Andersen, Nordströara, B. m, & J. Bland-Hawthorn, 289–300
- Esdaille, J., Glazebrook, K., Labbé, I., et al. 2021, *The Astrophysical Journal Letters*, 908, L35
- Ferland, G. J., Chatzikos, M., Guzmán, F., et al. 2023, *The 2017 Release of Cloudy*, arXiv:1705.10877
- Ferreras, I., Barbera, F. L., Rosa, I. G. d. l., et al. 2012, *Monthly Notices of the Royal Astronomical Society: Letters*, 429, L15–L19
- Ferreras, I., Saha, P., Leier, D., Courbin, F., & Falco, E. E. 2010, *MNRAS*, 409, L30
- Fioc, M., & Rocca-Volmerange, B. 1997, *A&A*, 326, 950
- . 2019, arXiv e-prints, arXiv:1902.02198
- Geha, M., Brown, T. M., Tumlinson, J., et al. 2013, *The Astrophysical Journal*, 771, 29
- Goswami, S., Slemer, A., Marigo, P., et al. 2021, *A&A*, 650, A203
- Gouliermis, D. A. 2018, *PASP*, 130, 072001
- Grasha, K., Calzetti, D., Adamo, A., et al. 2017, *The Astrophysical Journal*, 840, 113
- Graves, G. J., & Faber, S. M. 2010, *ApJ*, 717, 803
- Green, A. W., Croom, S. M., Scott, N., et al. 2018, *MNRAS*, 475, 716
- Gunawardhana, M. L. P., Hopkins, A. M., Sharp, R. G., et al. 2011, *MNRAS*, 415, 1647
- Hardy, E. 1990, in *European Southern Observatory Conference and Workshop Proceedings*, Vol. 35, *European Southern Observatory Conference and Workshop Proceedings*, ed. B. J. Jarvis & D. M. Terndrup, 209
- Ho, I.-T., Medling, A. M., Groves, B., et al. 2016, *Astrophysics and Space Science*, 361, doi:10.1007/s10509-016-2865-2
- Hopkins, A. M. 2018, *PASA*, 35, e039

- Hopkins, P. F. 2013, *Monthly Notices of the Royal Astronomical Society*, 433, 170–177
- Hoversten, E. A., & Glazebrook, K. 2008, *ApJ*, 675, 163
- . 2010, arXiv e-prints, arXiv:1011.3818
- Kauffmann, G., Heckman, T. M., Tremonti, C., et al. 2003, *MNRAS*, 346, 1055
- Kelvin, L. S., Driver, S. P., Robotham, A. S. G., et al. 2012, *MNRAS*, 421, 1007
- Kennicutt, R. C., J. 1983, *ApJ*, 272, 54
- Kennicutt, Robert C., J. 1989, *ApJ*, 344, 685
- . 1998, *ARA&A*, 36, 189
- Kennicutt, Robert C., J., Tamblyn, P., & Congdon, C. E. 1994, *ApJ*, 435, 22
- Kewley, L. J., Dopita, M. A., Sutherland, R. S., Heisler, C. A., & Trevena, J. 2001, *ApJ*, 556, 121
- Kewley, L. J., Groves, B., Kauffmann, G., & Heckman, T. 2006, *MNRAS*, 372, 961
- Komiya, Y. 2011, *ApJ*, 736, 73
- Kroupa, P. 2001, *MNRAS*, 322, 231
- Kroupa, P., Tout, C. A., & Gilmore, G. 1993, *MNRAS*, 262, 545
- Kroupa, P., & Weidner, C. 2003, *ApJ*, 598, 1076
- Kroupa, P., Weidner, C., Pflamm-Altenburg, J., et al. 2013, in *Planets, Stars and Stellar Systems. Volume 5: Galactic Structure and Stellar Populations*, ed. T. D. Oswalt & G. Gilmore, Vol. 5, 115
- Krumholz, M. R., McKee, C. F., & Bland-Hawthorn, J. 2019, *ARA&A*, 57, 227
- La Barbera, F., Vazdekis, A., Ferreras, I., et al. 2017, *MNRAS*, 464, 3597
- . 2016, *MNRAS*, 457, 1468
- Larson, R. B. 1992, *MNRAS*, 256, 641
- . 1998, *MNRAS*, 301, 569
- . 2005, *MNRAS*, 359, 211
- La Barbera, F., Vazdekis, A., Ferreras, I., et al. 2019, *Monthly Notices of the Royal Astronomical Society*, 489, 4090–4110
- Lee, J. C., Gil de Paz, A., Tremonti, C., et al. 2009, *ApJ*, 706, 599
- Lee, Y.-N., & Hennebelle, P. 2018, *A&A*, 611, A89
- Leier, D., Ferreras, I., Saha, P., et al. 2016, *Monthly Notices of the Royal Astronomical Society*, 459, 3677–3692
- Leitherer, C., Schaerer, D., Goldader, J. D., et al. 1999, *The Astrophysical Journal Supplement Series*, 123, 3–40
- Li, J., Liu, C., Zhang, Z.-Y., et al. 2023, *Nature*, 613, 460–462
- Martín-Navarro, I., La Barbera, F., Vazdekis, A., Falcón-Barroso, J., & Ferreras, I. 2015a, *MNRAS*, 447, 1033
- Martín-Navarro, I., Vazdekis, A., La Barbera, F., et al. 2015b, *ApJ*, 806, L31
- Martín-Navarro, I., Lyubenova, M., van de Ven, G., et al. 2019, *A&A*, 626, A124
- Martín-Navarro, I., Pinna, F., Coccato, L., et al. 2021, *A&A*, 654, A59
- Martín-Navarro, I., de Lorenzo-Cáceres, A., Gadotti, D. A., et al. 2023, arXiv e-prints, arXiv:2312.13355
- McKee, C. F., & Ostriker, E. C. 2007, *ARA&A*, 45, 565
- Medling, A. M., Cortese, L., Croom, S. M., et al. 2018, *MNRAS*, 475, 5194
- Mehrgan, K., Thomas, J., Saglia, R., et al. 2024, *ApJ*, 961, 127
- Meurer, G. R., Wong, O. I., Kim, J. H., et al. 2009, *ApJ*, 695, 765
- Mieske, S., & Kroupa, P. 2008, *ApJ*, 677, 276
- Myers, A. T., McKee, C. F., Cunningham, A. J., Klein, R. I., & Krumholz, M. R. 2013, *ApJ*, 766, 97
- Nanayakkara, T., Glazebrook, K., Kacprzak, G. G., et al. 2017, *MNRAS*, 468, 3071
- Nanayakkara, T., Brinchmann, J., Glazebrook, K., et al. 2020, *ApJ*, 889, 180
- Narayanan, D., & Davé, R. 2012, *MNRAS*, 423, 3601
- Owers, M. S., Allen, J. T., Baldry, I., et al. 2017, *MNRAS*, 468, 1824
- Owers, M. S., Hudson, M. J., Oman, K. A., et al. 2019, *ApJ*, 873, 52
- Parikh, T., Thomas, D., Maraston, C., et al. 2018, *MNRAS*, 477, 3954
- Pernet, E., Böcker, A., & Martín-Navarro, I. 2024, arXiv e-prints, arXiv:2406.04394
- Poci, A., McDermaid, R. M., Lyubenova, M., et al. 2022, *MNRAS*, 514, 3660
- Portinari, L., Chiosi, C., & Bressan, A. 1997, *Galactic chemical enrichment with new metallicity dependent yields*, arXiv:astro-ph/9711337
- Portinari, L., Sommer-Larsen, J., & Tantalo, R. 2004, *MNRAS*, 347, 691
- Rocca-Volmerange, B., & Guiderdoni, B. 1990, *MNRAS*, 247, 166
- Saglia, R. P., Maraston, C., Thomas, D., Bender, R., & Colless, M. 2002, *ApJ*, 579, L13
- Salpeter, E. E. 1955, *ApJ*, 121, 161
- Sánchez, S. F., Kennicutt, R. C., Gil de Paz, A., et al. 2012, *A&A*, 538, A8
- Scalo, J. M. 1986, *Fund. Cosmic Phys.*, 11, 1
- Schmidt, M. 1959, *ApJ*, 129, 243
- . 1963, *ApJ*, 137, 758
- Scott, N., van de Sande, J., Croom, S. M., et al. 2018, *MNRAS*, 481, 2299
- Sharda, P., Federrath, C., & Krumholz, M. R. 2020, *MNRAS*, 497, 336
- Sharp, R., Allen, J. T., Fogarty, L. M. R., et al. 2015, *MNRAS*, 446, 1551
- Shepard, D. 1968, in *Proceedings of the 1968 23rd ACM National Conference, ACM '68* (New York, NY, USA: Association for Computing Machinery), 517–524
- Smith, M. C. 2021, *MNRAS*, 502, 5417
- Smith, R. J. 2014, *MNRAS*, 443, L69
- Smith, R. J., Alton, P., Lucey, J. R., Conroy, C., & Carter, D. 2015, *Monthly Notices of the Royal Astronomical Society: Letters*, 454, L71–L75
- Somerville, R. S., & Davé, R. 2015, *Annual Review of Astronomy and Astrophysics*, 53, 51–113
- Stanway, E. R., Eldridge, J. J., & Becker, G. D. 2016, *MNRAS*, 456, 485
- Tornatore, L., Borgani, S., Dolag, K., & Matteucci, F. 2007, *MNRAS*, 382, 1050
- Treu, T., Auger, M. W., Koopmans, L. V. E., et al. 2010, *ApJ*, 709, 1195
- van Dokkum, P., Conroy, C., Villaume, A., Brodie, J., & Romanowsky, A. J. 2017, *ApJ*, 841, 68
- van Dokkum, P. G., & Conroy, C. 2010, *Nature*, 468, 940
- Vaughan, S. P., Davies, R. L., Zieleniewski, S., & Houghton, R. C. W. 2018, *MNRAS*, 475, 1073
- Vazdekis, A., Cenarro, A. J., Gorgas, J., Cardiel, N., & Peletier, R. F. 2003, *MNRAS*, 340, 1317
- Weidner, C., Kroupa, P., Pflamm-Altenburg, J., & Vazdekis, A. 2013, *MNRAS*, 436, 3309
- Worthey, G., Dorman, B., & Jones, L. A. 1996, *AJ*, 112, 948
- Yung, L. Y. A., Somerville, R. S., Finkelstein, S. L., Wilkins, S. M., & Gardner, J. P. 2024, *MNRAS*, 527, 5929
- Zaritsky, D., Colucci, J. E., Pessev, P. M., Bernstein, R. A., & Chandar, R. 2012, *ApJ*, 761, 93
- Zhang, Z.-Y., Romano, D., Ivison, R. J., Papadopoulos, P. P., & Matteucci, F. 2018, *Nature*, 558, 260
- Zonoozi, A. H., Haghi, H., & Kroupa, P. 2016, *ApJ*, 826, 89