

Original Article

Cite this article: Teixeira RJ, Machado JC, Faria S, Remondes-Costa S, Brandão T, Branco M, Moreira S, Pereira MG (2020). Brief emotional screening in oncology: Specificity and sensitivity of the emotion thermometers in the Portuguese cancer population. *Palliative and Supportive Care* **18**, 39–46. <https://doi.org/10.1017/S1478951519000208>

Received: 8 September 2018

Revised: 13 March 2019

Accepted: 19 March 2019

Keywords:

Emotion thermometers; Distress; Cancer; Portugal; Validation

Author for correspondence:

M. Graça Pereira, University of Minho, School of Psychology, Department of Applied Psychology, Campus de Gualtar, 4710-057 Braga, Portugal. Email: gracep@psi.uminho.pt

Brief emotional screening in oncology: Specificity and sensitivity of the emotion thermometers in the Portuguese cancer population

Ricardo João Teixeira, PH.D.¹, José C. Machado, PH.D.², Sara Faria, M.S.¹, Sónia Remondes-Costa, PH.D.³, Tânia Brandão, PH.D.^{4,5}, Margarida Branco, B.S.⁶, Sara Moreira, B.S.⁶ and M. Graça Pereira, PH.D.¹

¹University of Minho, School of Psychology, Braga, Portugal; ²University of Minho, Social Sciences Institute, Braga, Portugal; ³University of Trás-os-Montes e Alto Douro, Department of Education and Psychology, Vila Real, Portugal; ⁴CIP, Departamento de Psicologia, Universidade Autónoma de Lisboa Luís de Camões, Lisboa, Portugal; ⁵CPUP, Center for Psychology at University of Porto, Porto, Portugal and ⁶Liaison Psychiatry and Health Psychology Unit - Psychiatry and Mental Health Service, Oporto Hospital Centre, Porto, Portugal

Abstract

Objective. This study aimed to determine the cutoff and the specificity and sensitivity of the Emotion Thermometers (ET) in a Portuguese sample of cancer patients.

Method. A total of 147 patients (mean age = 49.2; SD = 12.6) completed the ET, the Brief Symptom Inventory (BSI), and the Subjective Experiences of Illness Suffering Inventory. Data were collected in a cancer support institution and in a major hospital in the North of Portugal.

Result. The optimal cutoff for the Anxiety Thermometer was 5v6 (until 5 and 6 or more), which identified 74% of the BSI-anxiety cases and 70% of noncases. The Depression Thermometer cutoff was 4v5 (until 4 and 5 or more), which identified 85% of BSI-depression cases and 82% of noncases. Cutoff for the Anger Thermometer was 4v5 (until 4 and 5 or more), which identified 83% of BSI-hostility cases and 73% of noncases; for the Distress Thermometer, the optimal cutoff was 4v5 (until 4 and 5 or more), which identified 84% of the suffering cases and 73% of noncases. Finally, for the Help Thermometer, it was 3v4 (until 3 and 4 or more), which helped to identify 93% of the suffering cases and 64% of noncases.

Significance of results. Results supported the Portuguese version of the ET as an important screening tool for identifying the emotional distress in cancer patients.

Introduction

The way cancer patients respond emotionally to the diagnosis and treatment of cancer is a long-standing concern in the psycho-oncology field. During the cancer trajectory, patients tend to exhibit signs and symptoms of emotional distress (Holland & Mastrovito, 1980). The presence of stress and anxiety and depressive symptoms among cancer patients is well documented in the literature. Studies indicate that depressive symptoms are significantly more common in depressed and nondepressed patients, regardless of the stage of the disease (Mitchell et al., 2012). In fact, approximately 16% of the cancer patients experience major depression, and about 22% exhibit minor depression and dysthymia combined (Mitchell et al., 2011). High stress levels are also reported in different phases of the cancer trajectory—at the beginning of chemotherapy treatments (Faul et al., 2010) as well as at the completion of the treatment (Garafalo et al., 2009).

Psychological morbidity may interfere with the effectiveness of cancer treatments, compromising significantly patients' survival. The emotional stress triggered by the disease seems to affect the immunosuppressive function of the immune system, favoring the progression of malignancy (McDonald et al., 2013). Also, the emotional stress as well as the depressive symptoms may also increase the risk of recurrence (Antoni et al., 2009; Spiegel & Giese-Davis, 2003). Also, empirical evidence has suggested that when patients have stress management skills or are helped to develop these skills, the anxiety and depression symptoms as well as the stress levels tend to decrease, and their mental health tends to improve (Faul et al., 2010). The evaluation, monitoring, and treatment of emotional distress have been recognized as an important practice with clinical and economic benefits for cancer patients and their families. Some studies pointed to the importance of this practice to improve the effectiveness of the oncology treatment (e.g., Bultz & Holland, 2006).

The assessment of emotional distress in clinical settings was initially focused on the identification of symptomatic cases of anxiety, stress, and depression in cancer patients. However, this assessment procedure immediately began to be questioned. In fact, the instruments used (despite their recognized validity and reliability) were too long (Mitchell, 2010a). Moreover, the inclusion of items about the experience of somatic symptoms created problems in differentiating signs and symptoms of emotional disorders and the common side effects of treatments for cancer (Jacobsen & Heather, 2008). Also, this type of assessment conferred a psychopathological and psychiatric connotation to the emotional distress experienced by the cancer patient resulting from the oncological disease (Mitchell, 2010b). Therefore, a scientific term more suited to represent the emotional experience of people affected by cancer diagnosis was developed and is currently defined as “emotional distress” (National Comprehensive Cancer Network, 2007).

Based on this new concept, emotional distress is now recognized as the sixth vital sign and, for that reason, should be periodically monitored among cancer patients, along with the other vital signs such as blood pressure, temperature, or pain (Bultz & Holland, 2006). Thus, the National Comprehensive Cancer Network developed specific guidelines for the screening of emotional distress among cancer patients in clinical settings. The first brief instrument recommended for screening the emotional distress in clinical settings was the Distress Thermometer (DT). Different studies have proven the reliability and the accuracy of this instrument to identify emotional distress in cancer patients as compared to other longer instruments (i.e., Psychological Distress Inventory, Hospital Anxiety and Depression Scales, Brief Symptoms Inventory [BSI]). Furthermore, the DT has revealed the same precision of these longer measures, but its advantages are its brevity and ease of administration and scoring. For these reasons, it has been considered as a more cost-effective way for screening emotional distress among cancer patients and is therefore preferred by the researchers as an appropriate instrument to be used in clinical settings (Gessler *et al.*, 2008; Gil *et al.*, 2005; Mitchell, 2007; VanHouse *et al.*, 2015).

However, more recently, other instruments, such as the Emotion Thermometers (ET; Mitchell *et al.*, 2010a) has been recognized to have greater scientific sustainability with regard to the precision in the screening of emotional distress in cancer patients. The ET is a combination of five visual analog scales in the form of four predictor domains (distress, anxiety, depression, anger) and one outcome domain (need for help). In comparison to the DT, the ET gives a more precise evaluation of the level of emotional distress (Mitchell *et al.*, 2010a). When applied alone, the Anxiety Thermometer can identify higher levels of emotional suffering (93.9% of accuracy) compared with the assessment obtained by the DT (54.4% of accuracy), thus revealing an increased sensitivity for identifying emotional complications in cancer patients (Mitchell *et al.*, 2010a). Anxiety and Depression Thermometers also have a statistically significant relationship with the anxiety and the depression subscales of the Hospital Anxiety and Depression Scales (Beck *et al.*, 2016). Against the Beck Depression Inventory, the ET has also shown good accuracy (>80%) for diagnosing depression in cancer patients (Schubart *et al.*, 2015).

In Portugal, a preliminary study to validate the ET has been performed in 104 cancer patients (the majority were middle-age women) recruited from the outpatient clinic of the Portuguese League Against Cancer. Descriptive analyses showed that

participants scored higher in the Anxiety and the Depression thermometers, followed by the distress and the anger thermometers. Correlational analyses revealed that all of the five ET were associated with the anxiety, depression, and hostility subscales of the BSI (personal communication).

The ET represents an advance in screening for anxiety and depression in the psycho-oncology field (Mitchell *et al.*, 2010a, 2010b). The reasons to recommend the ET as an appropriate instrument for use in clinical settings is its ability to screen for emotional distress in cancer patients and allow a quick and appropriate referral for an adequate therapeutic response (Gusani *et al.*, 2009; Mitchell, 2007, 2010a, 2010b; Schubart *et al.*, 2010, 2015). Despite the presence of guidelines and recommendations for the use of appropriate screening tools, the emotional distress of patients often goes unnoticed by oncologists in their clinical practice (Jacobsen, 2007). Although the international recommendations for the need to screen the emotional distress of cancer patients, this practice is still not consistently implemented (Fallowfield *et al.*, 2001; Pirl *et al.*, 2007). In fact, since 2004, the Canadian Strategy for Cancer Control (Rebalance Focus Action Group, 2005) has worked on the development of a treatment protocol, which includes the integration of a periodic monitoring of the patients' emotional distress in the physician's daily clinical routine (Bultz & Holland, 2006; Carlson & Bultz, 2003). This highlights the need of validating ET across countries to include the emotional screening as a regular practice in the care offered to cancer patients.

In Portugal, the National Comprehensive Cancer Network recommendations for screening the emotional suffering of cancer patients is not yet widely implemented given the lack of information about the validity (in terms of specificity and sensitivity) of the ET among Portuguese cancer patients. Thus, the purpose of this study was to determine the cutoff and the specificity and sensitivity of the emotion thermometers in the Portuguese oncology population to enable an early detection of emotional disorders among cancer patients, and to facilitate and accelerate the referral for psychosocial services, thereby contributing to the improvement of cancer care in Portugal. Therefore, the present study presented the diagnostic cutoff scores of each thermometer scale (distress, anxiety, depression, anger, and help). The establishment of cutoff scores is an important step to contribute to consolidate the emotional distress screening as a regular practice in Portugal and to help health care professionals to make faster and important decisions regarding the patients who may need psychosocial support.

Methods

Participants

Participants were 147 cancer patients attending a cancer support institution or a major hospital in Portugal. Ninety-eight (66.7%) patients were female and 48 (32.7%) were male. The mean age of the sample was 49.2 years ($SD = 12.6$; median = 50), and the mean disease duration was 6.8 months ($SD = 7.3$; median = 5), suggesting that this was predominantly an early cancer sample. A total of 101 (68.7%) patients were active and 23 (15.6%) were retired. Ninety-one (61.9%) patients were married, 23 (15.6%) were single, 19 (12.9%) were divorced/separated, and 10 (6.8%) were widowers. The mean years of education was 9.4 years ($SD = 4.0$; median = 9).

The most common diagnosis was breast cancer ($n = 54$, 36.7%), followed by bowel cancer ($n = 15$, 10.2%). The mean number of chemotherapy sessions was 4.1 ($SD = 2.2$; median = 4). Forty-two (28.6%) had metastasis and 115 (78.2%) were being treated with QT-EV. Eighty-five (76.6%) had already undergone surgery, 42 (37.8%) had been administered intravenous chemotherapy, and 32 (28.8%) had undergone radiotherapy. The aim of the current treatment was to provide a cure for 36 (24.5%), maintain adjuvant therapy for 32 (21.8%), focus on neoadjuvant therapy for 26 (17.7%), and provide palliative care for 11 (7.5%) patients.

Instruments

ET

The ET (Mitchell et al., 2010a) is used in clinical practice to assess the severity of emotional disorders in cancer patients. It consists of five thermometers assessed on an 11-point Likert scale (range 0–10) and includes four emotional domains, DT, Anxiety (AnxT), Depression (DepT), and Anger (AngT), and one nonemotional domain, Need for Help (HT). All the domains include a visual analog scale. A high score indicates higher levels of distress, anxiety, depression, anger, and more need for help. The Cronbach's alpha of the original ET (Mitchell et al., 2010a) was .91, whereas in the present study was .93.

BSI

The BSI (Derogatis & Spencer, 1982) is a self-report questionnaire that allows assessing clinical symptoms standards. The questionnaire comprises 53 items scored on a 5-point Likert scale (range 0–4). For the purpose of this study, only the depression, anxiety, and hostility subscales were used. A high score on these dimensions indicates higher levels of depression, anxiety, and hostility. The Cronbach's alpha of the original scale ranged from .71 to .85 (Derogatis & Spencer, 1982). This measure was translated and adapted to the Portuguese population by Canavarro (1999) that found an adequate internal consistency (alphas between .62 and .80). In the present study, the alphas were .92, .93, and .90 for the subscales of depression, anxiety, and hostility, respectively.

The Subjective Experiences of Illness Suffering Inventory (SEISI)

The SEISI was developed by McIntyre and Gameiro (1999) in a sample of Portuguese cancer patients for assessing the suffering associated with their oncological disease. The questionnaire includes 44 items scored on a 5-point Likert scale (range 1–5). It assesses five domains: physical suffering, psychological suffering, existential suffering, socio-relational suffering, and positive experience of suffering. Only the total score was used in the present study, with higher scores indicating higher suffering. Alphas in the original version ranged from .69 to .85 (.93 for the total score) (McIntyre & Gameiro, 1999). In this study, the alpha was .97.

Procedure

Data were collected at the Portuguese League against Cancer or at the Central Hospital of Oporto (both institutions are located at the North of Portugal). Psychologists working in these institutions invited cancer patients to take part in the study. These cancer patients were receiving active treatment, but none was hospitalized. Participation was voluntary, and all patients signed an informed consent. Participants were asked to indicate the number that best described their levels of distress, anxiety,

depression, anger, and need for help, in the thermometer, over the past 7 days. The study was approved by the review board of the Portuguese League against Cancer and the Ethical Committee of the Central Hospital. It follows all principles outlined in the Declaration of Helsinki.

Results

Thermometer variables

M and SD were obtained for the five scales. The mean score on the DT was 5.05 ($SD = 2.51$), 5.02 ($SD = 2.70$) on AnxT, 4.19 ($SD = 2.80$) on the DepT. On the AngT, the M obtained was 4.07 ($SD = 3.09$), and on the HT the M was 5.04 ($SD = 2.96$). Significant positive large correlations ($p < 0.001$) were found among all the thermometers (ranging from $r = .66$ to $r = .85$) (Table 1 and Figure 1).

Diagnostic validity

Validity using the BSI-anxiety for AnxT

Using a cutoff of 1.753 in BSI-anxiety as the criterion (the cutoff found in the Portuguese population; Canavarro, 1999), 50 patients (34.0%) were identified as experiencing clinically significant anxiety. The received operating characteristic (ROC) analysis for AnxT found an area under the curve (AUC) of .81 ($SE = .04$; $p < 0.001$; $CI_{95\%} [.74, .88]$), indicating an excellent discrimination (Figure 2). The optimal cutoff point equally favoring sensitivity (Ss; 74%) and specificity (Sp; 70%) was 5v6 (until 5 and 6 or more). Table 2 shows the rule in accuracy (positive predictive value [PPV]) and summary measures and rule-out accuracy (negative predictive value [NPV]).

Validity using the BSI-depression for DepT

Using a cutoff of 1.828 in BSI-depression as the criterion (the cutoff found in the Portuguese population; Canavarro, 1999), 58 patients (39.5%) were identified as experiencing clinically significant depression. The ROC analysis for DepT obtained the AUC of .89 ($SE = .03$; $p < .001$; $CI_{95\%} [.83, .94]$), indicating an excellent discrimination (Figure 2). The optimal cutoff point equally favoring Ss (85%) and Sp (82%) was 4v5 (until 4 and 5 or more). Table 2 shows the rule in accuracy (PPV) and summary measures and rule out accuracy (NPV).

Validity using the BSI-hostility for AngT

Using a cutoff of 1.411 in the BSI-hostility as the criterion (the cutoff found in the Portuguese population; Canavarro, 1999), 42 patients (28.6%) were identified as experiencing clinically significant hostility. The ROC analysis for AngT obtained the AUC of .84 ($SE = .04$; $p < .001$; $CI_{95\%} [.77, .92]$), indicating an excellent discrimination (Figure 2). The optimal cutoff point equally favoring Ss (83%) and Sp (56%) was 4v5 (until 4 and 5 or more). Table 2 shows the rule in accuracy (PPV) and summary measures and rule out accuracy (NPV).

Validity using the emotional suffering score for DT


Using a cutoff of 2.75 in suffering score as the criterion (because the SEISI has not a cutoff for clinical diagnosis, we have used the weighted mean obtained for the Portuguese population; Gameiro, 1999), 80 patients (54.4%) were identified as experiencing clinically significant suffering. The ROC analysis for DT obtained the AUC of .87 ($SE = .03$; $p < 0.001$; $CI_{95\%} [.81, .93]$), indicating

Table 1. Descriptive statistics and correlations of the emotion thermometers

Thermometers	Mean (SD)	DT	AnxT	DepT	AngT	HT
DT	5.05 (2.51)	1				
AnxT	5.02 (2.70)	.780*	1			
DepT	4.19 (2.80)	.852*	.674*	1		
AngT	4.07 (3.09)	.756*	.695*	.742*	1	
HT	5.04 (2.96)	.750*	.656*	.747*	.709*	1

Note: AngT, Anger thermometer; AnxT, Anxiety thermometer; DepT, Depression thermometer; DT, Distress thermometer; HT = Need for Help thermometer.

* $p < 0.05$.

**Fig. 1.** Distribution of scores on the emotion thermometers.

an excellent discrimination (Figure 2). The optimal cutoff point equally favoring Ss (84%) and Sp (79%) was 4v5 (until 4 and 5 or more). Table 2 shows the rule in accuracy (PPV) and summary measures and rule out accuracy (NPV).

Validity defined the suffering score for HT

Using a cutoff of 2.75 in suffering score as the criterion (weighted mean obtained for the Portuguese population; Gameiro, 1999), 80 patients (54.4%) were identified as experiencing clinically significant suffering. The ROC analysis for HT obtained the AUC of .83 ($SE = .03$; $p < .001$; $CI_{95\%} [.77; .90]$), indicating an excellent discrimination (Figure 2). The optimal cutoff point equally favoring Ss (93%) and Sp (64%) was 3v4 (until 3 and 4 or more). Table 2 shows the rule in accuracy (PPV) and summary measures and rule out accuracy (NPV).

Discussion

The presence of anxiety, depression, and anger among cancer patients, and their negative consequences for the process of adaptation to the cancer experience, highlights the importance of identifying appropriate tools for distress screening. The literature recommends the use of brief instruments capable of offering relevant information to help healthcare professionals in developing a differential diagnosis between normal symptomatology and

pathological symptomatology (Mitchell, 2010b). Studies have shown that visual analog scales have superior metrical characteristics in comparison to discrete scales (e.g., Reips & Funke, 2008). The DT has been widely used in Portugal; however, a reliable measure to evaluate other emotional changes in cancer patients is needed. This is the first study examining the ET tool (Mitchell *et al.*, 2010a) in a Portuguese sample of cancer patients. The main aim was to establish the diagnostic validity of the ET with the BSI (anxiety, depression, and hostility subscales) and a global measure of suffering (SEISI). Previous studies have not examined an Anxiety Thermometer, Depression Thermometer, and Anger Thermometer against a standardized instrument to evaluate symptoms of psychopathology (BSI), or compared the results from the ET's Distress and Help thermometers against a measure of suffering.

The results of the present study showed that the criterion validity of the ET was established using ROC analysis with BSI subscales, which is considered a gold standard. The AUC of AnxT against BSI anxiety subscale was .81, whereas the AUC of the DepT thermometer against BSI depression subscale was .89. The AUC of AngT against BSI hostility subscale was .84. The AUC of the DT against the total score on SEISI was .87. Finally, the AUC of HT thermometer against the total score on SEISI was .83. Also, all the ETs were positively intercorrelated, as found in previous studies (Hinze *et al.*, 2019; Mitchell *et al.*,




Fig. 2. ROC curves. AngT, Anger Thermometer; AnxT, Anxiety Thermometer; AUC, area under the curve; CI, confidence interval; DepT, Depression Thermometer; DT, Distress Thermometer; HT, Need for Help; ROC, receiver operating characteristic.

2010a), suggesting that the set of thermometers measure a more general “emotional distress” construct. However, although intercorrelations were high, they were not high enough for considering the ETs as a unidimensional measure, but a measure assessing different dimensions of emotional distress. Also, it is possible that patients may have difficulties in qualifying what is distress, anxiety, or depression, as suggested by previous studies (e.g., Beck et al., 2016) leading to high intercorrelations among ETs.

Overall, the AUC values indicated that ETs have an excellent discrimination. These findings are in line with the results obtained by Mitchell et al. (2010a), which examined the discriminative value of the ETs among a sample of British cancer patients. Concerning the cutoffs for the Portuguese, using the BSI-Anxiety as the criterion, the optimal thermometers were AngT and HT; using the BSI-Depression as the criterion, the optimal thermometers were DepT and HT; and using the BSI-Hostility as the criterion, the optimal thermometers were AngT and DT. However, when the total suffering score was used as the criterion, the optimal thermometers were AngT and DT. When the total suffering score was used as the criterion, the optimal thermometers were AngT and AnxT. As found in the Mitchell’s et al. study (2010a), the AngT showed to be better/optimal for detecting overall distress. Yet, the author recommended a combination of items to be more accurate (Mitchell

et al., 2010b). In the present study, only the DepT showed to be optimal for detecting depression, because it presented a greater sensibility and specificity. The ETs have been found to have a good performance in detecting major depression (against Diagnostic and Statistical Manual of Mental Disorders, 4th edition, depression) within other contexts (e.g., cardiovascular disease, epilepsy) (Mitchell et al., 2012; Rampling et al., 2012).

Many studies emphasize the need of evaluating emotion deregulation in oncology settings, mainly distress, anxiety, and depression (Cordes et al., 2014; Pandey et al., 2006). However, although the literature has recognized the importance of anger assessment in oncology settings (Penedo et al., 2006; Philip et al., 2007), anger has been rarely examined. To our knowledge, this is the first study evaluating anger/hostility in a Portuguese sample of cancer patients. The use of the AngT should be of great utility because is an imperative deregulator of coping for individuals facing a cancer diagnosis and treatment. Anger is often one of the first emotional reactions a person has to a cancer diagnosis. Many cancer patients wonder “Why me?”; this can lead to feelings of angry and frustration. According to Lown (2007), anger may represent a disruption in the doctor–patient relationship, and it is important for physicians to adjust their behavior and communication accordingly. In this study, however, the AngT has the lowest PPV. From all the ETs, the AngT presented the lowest mean. It is possible that among these participants, anger is not so prevalent,

Table 2. Validity of the thermometers

	Method	Sensitivity	Specificity	PPV	NPV	PSI	Youden index
AnxT Using Anxiety Score With Cutoff 5v6	DT (5v6)	.740	.732	.587	.845	.433	.472
	AnxT (5v6)	.740	.701	.561	.840	.400	.441
	DepT (5v6)	.620	.804	.620	.804	.424	.424
	AngT (5v6)	.680	.835	.680	.835	.515	.515
	HT (5v6)	.820	.619	.526	.870	.395	.439
DepT Using Depression Score With Cutoff 4v5	DT (4v5)	.897	.629	.612	.903	.515	.526
	AnxT (4v5)	.845	.584	.570	.852	.422	.429
	DepT (4v5)	.845	.820	.754	.890	.644	.665
	AngT (4v5)	.810	.820	.746	.869	.615	.631
	HT (4v5)	.966	.607	.615	.964	.580	.572
AngT Using Hostility Score With Cut-off 4v5	DT (4v5)	.929	.562	.459	.952	.410	.490
	AnxT (4v5)	.881	.533	.430	.918	.348	.414
	DepT (4v5)	.786	.695	.508	.890	.398	.481
	AngT (4v5)	.833	.733	.556	.917	.472	.567
	HT (4v5)	.905	.495	.418	.929	.346	.400
DT Using Suffering Score With Cut-off 4v5	DT (4v5)	.838	.731	.788	.790	.579	.569
	AnxT (4v5)	.825	.701	.767	.770	.538	.526
	DepT (4v5)	.663	.821	.815	.671	.486	.483
	AngT (4v5)	.700	.896	.889	.714	.603	.596
	HT (4v5)	.850	.657	.747	.786	.533	.507
HT Using Suffering Score With Cut-off 3v4	DT (3v4)	.925	.612	.740	.872	.612	.537
	AnxT (3v4)	.925	.657	.763	.880	.643	.582
	DepT (3v4)	.750	.746	.779	.714	.494	.496
	AngT (3v4)	.813	.861	.867	.792	.658	.663
	HT (3v4)	.925	.642	.755	.878	.633	.657

AngT, Anger thermometer; AnxT, Anxiety thermometer; DepT, Depression thermometer; DT, Distress thermometer; HT, Need to Help thermometer; NPV, negative predictive value; PPV, positive predictive value; PSI, predictive summary index.

which can lead to the existence of a high proportion of false positives. Probably, and to reduce false positives, the value of the AngT should be more restricted. Further research is needed with regard to the AngT.

The need for help of cancer patients should also be a concern for healthcare professionals. Indeed, more important than identifying distress and its prevalence, is to provide care for cancer patients in need (Mitchell, 2010a). In fact, HT was the one with the highest PPV values: when using with suffering scores, reinforcing its utility for identifying cancer patients who need most psychosocial support. This is even more important because studies have shown that only one-third of cancer patients with significant distress (using the DT), report wanting professional help for emotional problems (Graves et al., 2007). According to Baker-Glenn and Mitchell (2008), the most usual requests are face-to-face psychological support and complementary therapies. In Portugal, cancer patients who are undergoing treatment in public and private hospitals do not always get the psychological help they need. This is often due to a lack of effective communication between health professionals and patients. So, the HT can be a useful tool to identify and refer to adequate psychosocial support cancer patients who need help.


Having a screening tool capable of detecting the emotional distress experienced by cancer patients is of high importance given its role in protecting patients from future adjustment disorders (Iscoe et al., 1991) and on reducing costs associated with the lack of treatment of these disorders (Boberg et al., 2003). The ET has several advantages related to its brevity and ease administration and scoring. In fact, it can be administrated by health professionals with no mental health training (e.g., nurses). This is an advantage because cancer services do not always have the support of a mental health professional; and patients have much more contact with non-mental health professionals (even in cancer services that have the support of mental health professionals). When patients screen positive, healthcare professionals can refer them for further assessment by mental healthcare professionals (Beck et al., 2016). Because the administration of ETs only requires that patients circle the number that best describe their levels of distress, anxiety, depression, anger, and need for help, and because clinical cutoffs are established for scoring ETs, we do not anticipate any limitation if ETs are administered by non-mental healthcare professionals.

There are some limitations in this study. The sample size is modest and patients were predominantly in the early stages of

the disease. Future studies should be conducted with larger samples and with cancer patients in different stages of the disease (especially late-stage cancers). However, for early-stage cancer, ETs seem to be a valid screen tool. Moreover, the ETs were not compared with the DT. Further research is needed with Portuguese cancer patients to study if ETs expand on the strengths of the DT, as found in a previous study (Mitchell et al., 2010b). Finally, the SEISI does not have a clinical cutoff to establish a diagnosis, which can limit the findings. Future studies should validate ETs against other diagnostic instruments.

Some implications for clinical practice can be derived from this study. For health professionals working in oncological settings, the AnxT, DepT, AngT and HT provide an easy way to assess and identify patient's emotional needs.

Distress has been referred to as the sixth vital sign in cancer care, requiring that providers assess and treat it with the same importance as any physical illness (Bultz & Carlson, 2006; Mitchell et al., 2012; Mitchell & Coyne, 2009). Screening identifies unmet needs, but screening alone is an ineffective strategy requiring the availability of psychosocial services (Schubart et al. 2015). In the United States, the American College of Surgeons (2012) now requires that cancer centers implement screening programs for psychosocial distress using validated instruments with established clinical cutoffs. In Portugal, further studies are needed to evaluate how health professionals use the information collected with the ET to improve patient care and how this information affects long-term patient outcomes.

Author ORCID.  Ricardo João Teixeira, 0000-0003-3820-1660; José C. Machado, 0000-0003-2552-118X; Sara Faria, 0000-0001-9792-5816; Sónia Remondes-Costa, 0000-0003-4282-3964; Tânia Brandão, 0000-0001-7865-2445; M. Graça Pereira, 0000-0001-7987-2562

References

- American College of Surgeons** (2012) *Commission on cancer. Cancer program standards 2012: Ensuring patient-centered care. V1.2.1*. Chicago: American College of Surgeons.
- Antoni MH, Lechner S, Diaz A, et al.** (2009) Cognitive behavioral stress management effects on psychosocial and physiological adaptation in women undergoing treatment for breast cancer. *Brain, Behavior, and Immunity* 23(5), 580–591.
- Baker-Glenn EA and Mitchell AJ** (2008) Screening for perceived need for help using the latest screening tools - what do we know about who wants help? *Psycho-Oncology* 17, S56–S57.
- Beck KR, Tan SM, Lum SS, et al.** (2016) Validation on the emotion thermometers and hospital anxiety and depression scales in Singapore: Screening cancer patients for distress, anxiety and depression. *Asia-Pacific Journal of Clinical Oncology* 12(2), e241–e249.
- Boberg EW, Gustafson DH, Hawkins RP, et al.** (2003) Assessing the unmet information, support and care delivery needs of men with prostate cancer. *Patient Education and Counseling* 49(3), 233–242.
- Bultz BD and Carlson LE** (2006) Emotional distress: The sixth vital sign - future directions in cancer care. *Psycho-Oncology* 15(2), 93–95.
- Bultz BD and Holland JC** (2006) Emotional distress in patients with cancer: The sixth vital sign. *Psychosocial Oncology* 3(5), 311–314.
- Canavarró MC** (1999) Inventário de Sintomas Psicopatológicos - BSI [Brief Symptom Inventory - BSI]. In MR Simões, M Gonçalves, and LS Almeida (Eds.), *Testes e provas psicológicas em Portugal [Psychological tests in Portugal]*, Vol. II, Braga, Portugal: Apport.
- Carlson LE and Bultz BD** (2003) Cancer distress screening. Needs, models, and methods. *Journal of Psychosomatic Research* 55(5), 403–409.
- Cordes MC, Scherwath A, Ahmad T, et al.** (2014) Distress, anxiety and depression in patients with brain metastases before and after radiotherapy. *BMC Cancer* 14(731).
- Derogatis LR and Spencer PM** (1982) *Administration and procedures: BSI manual I*. Baltimore: Clinical Psychometric Research.
- Fallowfield L, Ratcliffe D, Jenkins V, et al.** (2001) Psychiatric morbidity and its recognition by doctors in patients with cancer. *British Journal of Cancer* 84(8), 1011–1015.
- Faul JA, Heather SJ, Williams C, et al.** (2010) Relationships of stress management skill to psychological distress and quality of life in adults with cancer. *Psycho-Oncology* 19(1), 102–109.
- Gameiro MH** (1999) *O Sofrimento na doença [Suffering in illness]*. Coimbra, Portugal: Quarteto Editora.
- Garafalo JP, Choppala S, Hamann H, et al.** (2009) Uncertainty during the transition from cancer patient to survivor. *Cancer Nursing* 32(4), 8–14.
- Gessler S, Law J, Daniells E, et al.** (2008) Screening for distress in cancer patients: Is the thermometer distress a valid measure in UK and does it measure change over time? A prospective validation study. *Psycho-Oncology* 17(6), 538–547.
- Gil F, Grassi L, Travado L, et al.** (2005) Use of distress depression thermometers to measure psychosocial morbidity among southern European cancer patients. *Supportive Care in Cancer* 13(8), 600–606.
- Graves KD, Arnold SM, Love CL, et al.** (2007) Distress screening in a multidisciplinary lung cancer clinic: Prevalence and predictors of clinically significant distress. *Lung Cancer* 55(2), 215–224.
- Gusani NJ, Schubart JR, Wise J, et al.** (2009) Cancer survivorship: A new challenge for surgical and medical oncologists. *Journal of General Internal Medicine* 24(2), 456–458.
- Hinz A, Mitchell AJ, Dégi CL, et al.** (2019) Normative values for the distress thermometer (DT) and the emotion thermometers (ET), derived from a German general population sample. *Quality of Life Research* 28, 277–282.
- Holland JC and Mastrovito R** (1980) Psychological adaptation to breast cancer. *Cancer* 15(46), 1045–1052.
- Iscoe N, Williams JI, Szalai JP, et al.** (1991) Prediction of psychosocial distress in patients with cancer. In D. Osoba (Ed.), *Effect of cancer on quality of life* (pp. 41–59). London: CRC Press.
- Jacobsen PB** (2007) Screening for psychological distress in cancer patients: Challenges and opportunities. *Journal of Clinical Oncology* 25(29), 4526–4527.
- Jacobsen PB and Heather SJ** (2008) Psychological interventions for anxiety and depression in adult cancer patients achievements and challenges. *Cancer Journal for Clinicians* 58(4), 214–230.
- Lown B** (2007) Difficult conversations: Anger in the clinician-patient/family relationship. *Southern Medical Journal* 100(1), 33–39.
- McDonald GP, O'Connell M, and Lutgendorf SK** (2013) Psychoneuroimmunology and cancer: a decade of discovery, paradigm shifts, and methodological innovations. *Brain, Behavior and Immunity* 30(Suppl 15), S1–S9.
- McIntyre T and Gameiro M** (1999) Inventário de Experiências Subjectivas de Sofrimento na Doença [Illness Subjective Suffering Inventory]. In A Soares, S Araújo, & A Caires (Eds.), *Avaliação Psicológica: Formas e Contextos [Psychological Assessment: Forms and Contexts]* (pp. 723–736). Braga: APPORT.
- Mitchell AJ** (2007) Pooled results from 38 analyses of the accuracy of distress thermometer and other ultra-short methods of detecting cancer-related mood disorders. *Journal of Clinical Oncology* 25(29), 4670–4681.
- Mitchell AJ** (2010a) Screening procedures for psychosocial distress. In JC Holland, WS Breitbart, PB Jacobsen, et al. (Eds.), *Psycho-Oncology* (2nd ed., pp. 389–398). New York: Oxford University Press.
- Mitchell AJ** (2010b) Short screening tools for cancer-related distress: A review and diagnostic validity meta-analysis. *Journal of the National Comprehensive Cancer Network* 8(4), 487–494.
- Mitchell AJ and Coyne JC** (2009) *Screening for depression in clinical practice: An evidence-based guide*. New York: Oxford University Press.
- Mitchell AJ, Baker-Glenn EA, Granger L, et al.** (2010a) Can the Distress Thermometer be improved by additional mood domains? *Part I. Initial validation of the Emotion Thermometers tool*. *Psycho-Oncology* 19(2), 125–133.
- Mitchell AJ, Baker-Glenn EA, Park B, et al.** (2010b) Can the Distress Thermometer be improved by additional mood domains? *Part II. What is the optimal combination of Emotion Thermometers?* *Psycho-Oncology* 19(2), 134–140.

- Mitchell AJ, Chan M, Bhatti H, et al.** (2011) Prevalence of depression, anxiety, and adjustment disorder in oncological, haematological, and palliative-care settings: A meta-analysis of 94 interview-based studies. *Lancet Oncology* **12**(2), 160–174.
- Mitchell AJ, Lord K, and Symonds P** (2012) Which symptoms are indicative of DSM-IV depression in cancer settings? An analysis of the diagnostic significance of somatic and non-somatic symptoms. *Journal of Affective Disorders* **138**(1–2), 137–148.
- Mitchell AJ, Morgan JP, Petersen D, et al.** (2012) Validation of simple visual-analogue thermometer screen for mood complications of cardiovascular disease: The Emotion Thermometers. *Journal of Affective Disorders* **136**, 1257–1263.
- National Comprehensive Cancer Network** (2007) NCCN clinical practice guidelines in oncology: distress management (V.1).
- Pandey M, Sarita GP, Devi N, et al.** (2006) Distress, anxiety, and depression in cancer patients undergoing chemotherapy. *World Journal of Surgical Oncology* **4**(68).
- Penedo FJ, Dahn JR, Kinsinger D, et al.** (2006) Anger suppression mediates the relationship between optimism and natural killer cell cytotoxicity in men treated for localized prostate cancer. *Journal of Psychosomatic Research* **60**(4), 423–427.
- Philip J, Gold M, Schwarz M, et al.** (2007) Anger in palliative care: A clinical approach. *Internal Medicine Journal* **37**(1), 49–55.
- Pirl WF, Muriel A, Hwang V, et al.** (2007) Screening for psychosocial distress: A national survey of oncologists. *Supportive Oncology* **5**(10), 499–504.
- Rampling J, Mitchell AJ, Von Oertzen T, et al.** (2012) Screening for depression in epilepsy clinics. A comparison of conventional and visual-analogue methods. *Epilepsia* **53**, 1713–1721.
- Rebalance Focus Action Group** (2005) A position paper: Screening key indicators in cancer patients—Pain as a 5th vital sign and emotional distress as a 6th vital sign. *Canadian Strategy for Cancer Control Bulletin* **7**(4).
- Reips UD and Funke F** (2008) Interval-level measurement with visual analogue scales in Internet-based research: VAS Generator. *Behavior Research Methods* **40**(3), 699–704.
- Schubart JR, Mitchell A, Dietrich L, et al.** (2015) Accuracy of the Emotion Thermometers (ET) screening tool in patients undergoing surgery for upper gastrointestinal malignancies. *Journal of Psychosocial Oncology* **33**(1), 1–14.
- Schubart JR, Wise J, Deshaies I, et al.** (2010) Quality of life assessment in postoperative patients with upper GI malignancies. *Journal of Surgical Research* **163**(1), 40–46.
- Spiegel D and Giese-Davis J** (2003) Depression and cancer: Mechanisms and disease progression. *Biological Psychiatry* **54**(3), 269–282.
- VanHouse L, Black LL, Doty K, et al.** (2015) An analysis of the distress thermometer problem list and distress in patients with cancer. *Supportive Care in Cancer* **23**(5), 1225–1232.