

This is a “preproof” accepted article for *Invasive Plant Science and Management*. This version may be subject to change in the production process, *and does not include access to supplementary material*. DOI: 10.1017/inp.2024.41

Running footer: Control of *Elaeagnus pungens*

A comparison of different chemical control application methods for managing *Elaeagnus pungens* in South Carolina

Molly N. Darr¹, Louis B. Nottingham¹, Janet W. Steele², Timothy L. Evans³, and David R. Coyle⁴

¹Assistant Professor (MND ORCID 0000-0003-1510-6556, LBN ORCID 0000-0003-2558-6418), Washington State University, Northwest Research and Extension Center, Mount Vernon, WA, USA; ²Area Forestry and Wildlife Agent, Clemson University Cooperative Extension, Orangeburg, SC, USA; ³Director of Land Conservation, Audubon South Carolina, Charleston, SC, USA, ⁴Associate Professor (ORCID 0000-0003-3074-0440), Clemson University, Clemson, SC, USA.

Author for correspondence: Molly Darr, Assistant Professor, Washington State University, Northwest Research and Extension Center, 16650 WA-536, Mount Vernon, WA 98273. (E-mail: molly.darr@wsu.edu)

Abstract: Several *Elaeagnus* species (autumn olive [*E. umbellata* (Thunb.)], Russian olive [*E. angustifolia* (L.)], and thorny olive [*E. pungens* (Thunb.)]) are invasive in North America. *Elaeagnus pungens* is prevalent throughout much of the southeastern United States, commonly overtaking wooded and natural areas, bottomlands, and roadsides. While many management methods including several herbicide treatments have been evaluated, efficacy of these methods can vary based on the size and density of the target plants. Further, personal communication with land managers revealed a lack of information that incorporated application effort, duration and associated cost into treatment efficacy and usefulness. We evaluated three herbicide application methods using the free acid formulation of triclopyr in an *E. pungens* infested forest in South Carolina, United States to determine the effectiveness of each application method. We estimated pre-treatment *E. pungens* biomass and destructively harvested all live material post treatment to obtain actual biomass values. Foliar herbicide application was ineffective, but both cut stump and basal bark application nearly eliminated *E. pungens* in the treatment plots. The basal bark application took slightly more time to complete than cut stump treatments but was described as less physically demanding by applicators. Based on treatment efficacy and time required, the basal bark application method seems most prudent for controlling *E. pungens* in these areas. These results will help land managers more effectively use their resources for invasive woody plant control.

Keywords: forest management; herbicide; invasive species; thorny olive; woody shrub

Management Implications: *Elaeagnus pungens* is an aggressive invasive shrub in North America that can take over natural areas and impact the ecology of the local environment. Material and labor costs associated with controlling this shrub can be high, and land managers need effective methods to reduce or eliminate *E. pungens* growth. Management of *E. pungens* can be uniquely challenging for several reasons. The thorns of the felled plant material or sprawling branches can puncture tires of equipment used in management activities, and cause injury to people working in these areas. The thick growth of particularly aggressive stands of *E. pungens* can result in a damp, shady forest floor and a lack of adequate fuel, precluding prescribed burning treatments. We evaluated three herbicide application methods using triclopyr as the free acid formulation (Trycera[®]): foliar, basal bark, and cut stump, in central South Carolina to determine the fastest, easiest management approach with maximum control efficacy. We prioritized treatment methods that could be applied with limited personnel (2-3 applicators) and without motorized vehicles. We evaluated these factors to help land managers more effectively use their resources, limit the introduction of herbicides into the environment, and determine a reliably effective method to kill *E. pungens* plants. Basal bark and cut stump treatments were much more effective in reducing *E. pungens* biomass in our plots than the foliar treatment. Between basal bark and cut stump treatment methods, we recommend basal bark due to its relative ease of application compared to the cut stump application method and equal efficacy in terms of controlling *E. pungens*. This approach should be used as part of an integrated management strategy, and further studies should evaluate other management tactics to reduce seedling germination success and control the vigorous resprouting associated with *E. pungens* growth.

Introduction

An expansion in global trade and travel has increased the number of invasive species impacting natural and managed systems worldwide (Liebhold 2012; Seebens et al. 2017; Essl et al. 2020) and led to a drastic rise in global costs associated with invasive species management (Crystal-Ornelas et al. 2021). In the United States alone, invasive species costs ranged between US\$ 18.2 and 78.9 billion between 1970 and 2020 (Diagne et al. 2021; Zenni et al. 2021). Additionally, natural and managed systems can be negatively impacted as native species are often outcompeted by unchecked invasive species populations, which can lead to a reduction in biodiversity and the alteration of entire trophic cascades (Schmitz et al. 2000; Beschta and Ripple 2009; Kimbro et al. 2009). Economic and ecological impacts associated with invasive species introduction will only continue to escalate over time without significant regulatory intervention.

The southeastern United States, with its long growing season and warm, humid climate is an ecologically diverse region with a rich forestry history, economy, and production potential (Napton et al. 2010; Carter et al. 2021). However, forests in this region are also impacted by many invasive plant species (Oswalt et al. 2015). Invasive plants are adept at establishing and flourishing in areas where management activities (e.g., logging, clearing, burning, fire suppression, and reforestation) and the indirect effects of climate change disrupt forest ecosystems (Holmes et al. 2009). Several invasive shrub species in the region are known to outcompete native species and dominate the forest understory (Maynard-Bean et al. 2020); these species include Chinese privet *Ligustrum sinense* (Lour.), bush honeysuckles (*Lonicera mackii* [Rubr.] Maxim; *L. morrowii* [Gray]; *L. tatarica* [L.]; *L. x bella* [Zabel]) and silverberry or olives (*Elaeagnus* spp.).

Elaeagnus pungens (Thunb.), colloquially known as thorny olive, silverthorn, thorny elaeagnus, spiny oleaster, or silverberry, is a broadleaved evergreen shrub native to Japan and China (Figure 1). It was introduced as an ornamental species in 1830 and later promoted for wildlife (Davison 1942), and now occurs throughout the southeastern United States and in parts of the northeastern United States (Miller 2006). *Elaeagnus pungens* is a multi-stemmed, freely branched, dense shrub that can reach 7.5 m tall and 4.5 m wide, and can use other species to

support branch growth and climb opportunistically (Serviss and Tumilson 2021) (Figure 1a,f). Once established, this shrub produces copious, fast-growing branch “whips”, which allow the plant to quickly increase in size and overtake neighboring vegetation (Figure 1a). Additionally, root suckering and prolific stem sprouts can lead to dense understory growth in a forested setting (Miller 2006). The stems produce thorns and the leaves are tough and waxy with a dark green surface and an underside that is silvery with brown scales (Figure 1b,c,e). In the southeastern United States, *E. pungens* benefits from an especially lengthy growing season that allows it to disperse and outcompete native plants (Riffe 2018). Specifically, *E. pungens* flowers and fruits for approximately ten months out of the year (Dirr 1990, Miller 2006). The fruit, fleshy red drupes, are readily consumed by birds and other animals which disperse them across large areas (Davison 1942), allowing *E. pungens* to quickly spread across the landscape (Figure 1c,d). *Elaeagnus pungens* grows densely in a variety of environmental conditions (open sun, forested settings, frequently flooded areas, disturbed sites, etc.), and is often a problem in fence rows, roadside margins, waste areas, and open woodlands.

Despite demonstrated deleterious ecological impacts, *Elaeagnus* spp. are still commonly sold in nurseries and online, and are cultivated for hedges, screening, natural barriers, bank stabilization along highways, and landscape uses (Beaury et al. 2021; Fertakos et al. 2023). *Elaeagnus* spp. have been shown to reduce the abundance of native species, facilitate the establishment of other non-native plants, and cause long-lasting impacts on local soil characteristics and flora (Collette and Pither 2015; Katz et al. 2020). Their presence impacts soil microbial communities (Malinich et al. 2017) and alters stream biogeochemical cycles (Mineau et al. 2011). *Elaeagnus* spp. can be aggressive invaders in forests (Yates et al. 2004) and negatively impact native tree seedling and sapling abundance (Lázaro-Lobo et al. 2021). Even though birds feed on *Elaeagnus* fruit (Davison 1942), one study showed a decrease in cavity nesting birds in areas invaded by *Elaeagnus*, presumably due to a lack of overstory trees (Fischer et al. 2012). Further, planting *Elaeagnus* in highway medians led to an increase in bird mortality by automobile strikes (Watts and Paxton 2000).

In his book titled *Manual of Woody Landscape Plants*, Dirr (1990) described *E. pungens* in its natural form as “a genuine horror” and went on to say, “fast does not adequately describe the speed at which it grows”. Because of *E. pungens*’ prolific and hardy nature, employing a

long-term, ecosystem-wide strategy prioritizing prevention, active monitoring and prompt eradication would likely be more successful than site-specific, local control measures. However, ecosystem-wide strategies are not always feasible or possible. Several local control methods have been evaluated with varying efficacy. Prescribed fire can be an effective management tool to reduce seed viability and kill young *Elaeagnus* plants (Muscha et al. 2023) but established plants will typically resprout following a fire (Michielsen et al. 2017) or cutting (Corns and Schraa 1965).

Herbicides are a common, established, and often effective management tactic for invasive plants (Pile Knapp et al. 2023), and formulations with the active ingredient triclopyr are known for their efficacy on woody plant species (e.g., Bovey 1965; Bovey and Whisenant 1991; DiAllesandro 2012; Enloe et al. 2016; Enloe et al. 2023) including *Elaeagnus* (Edgin and Ebinger 2001). Though most research on *Elaeagnus* management has occurred in the western United States and focused on *Elaeagnus* control in the context of rangeland management, a combination of cutting and stump treatment with triclopyr was shown to be effective for *Elaeagnus* management in former coal mines in the Appalachian region of the United States (Franke et al. 2018). However, relatively few studies have examined best management practices for forested lands in this region. We conducted informal interviews with land managers in the Southeast to determine the most common practices used to control woody understory growth in this region, whether motorized vehicles were used, the number of personnel most often available, along with the operator's assessment of the tools at their disposal. We determined that many land managers lacked information that incorporated time, effort, and cost of application with treatment efficacy when controlling woody understory growth. Our objective was to determine an effective chemical management method for a dense *E. pungens* infestation in a southern hardwood dominated forested area. We prioritized methods that would be easily accepted and utilized by land managers. Using the active ingredient triclopyr in the free acid formulation, we tested foliar spray, basal bark, and cut stump application methods with the goal of quantifying local *Elaeagnus* control and ease of application.

Materials and Methods

Site Description

Our study was conducted on a 214 ha (529 ac) forested tract in Calhoun County, South Carolina, United States (33° 38' 11.9364" N, 80° 42' 21.4056" W). Over 26 ha (65 ac) of the understory in this area is dominated by *E. pungens* (voucher specimens are deposited in the Clemson University Herbarium [accessible as Molly Darr #1 (CLEMS0083037, CLEMS0083038, CLEMS0083039), Molly Darr #2 (CLEMS0083040, CLEMS0083041, CLEMS0083042, CLEMS0083043), and Molly Darr #3 (CLEMS0083044, CLEMS0083045)]. The region's humid subcontinental climate has long, warm summers and mild winters with a mean maximum temperature from 2018 to 2023 of 37.3 °C (99.2 °F), a mean minimum temperature of -7.8 °C (46.04 °F), and an average annual precipitation of 110 cm (43 in) (USNWS 2023). Soils in these areas are classified as "southern Coastal Plain" and consist of Faceville fine sandy loam and Ailey-Vaucluse complex soil series (USDA 2019) (Table 1). The study site consisted of a hardwood dominated riparian bottomland stand and an adjacent upland old-field sweetgum (*Liquidambar styraciflua* [L.]) stand. The overstory of the bottomland area was composed primarily of sweetgum, with winged elm (*Ulmus alata* [Michx.]), loblolly pine (*Pinus taeda* [L.]), white oak (*Quercus alba* [L.]), and ash (*Fraxinus* spp.) interspersed. The overstory of the upland area was a mixture of white oak, southern red oak (*Quercus falcata* [Michx.]), loblolly pine, American beech (*Fagus grandifolia* [Ehrh.]), hickory (*Carya* spp.), and sweetgum (Table 1). Treatment plots were assigned in a randomized complete block design and grouped into three 0.4 ha (1 ac) blocks, each at a separate geographic location on the study site (Zar 2010) (Table 1). Each block contained 16 1m² quadrats, and each quadrat within a block was randomly assigned one of three treatments (foliar spray, basal bark, and cut stump) or an untreated control (UTC). All four treatments had four replicates per block; therefore, each treatment had 12 replicates across the entire experiment.

Pretreatment measurements

Pretreatment measurements were conducted to obtain a baseline measurement of *Elaeagnus* plant abundance and biomass to confirm that no preexisting differences occurred among treatment assignments. On 11 March 2020 we measured the total number of *E. pungens* plants, along with the basal circumference (cm) of every *E. pungens* plant within each 1m² quadrat (16 quadrats per block, 48 total quadrats across three blocks). Basal stem circumference was measured at 15 cm (6 in) above ground level. Ideally, pretreatment measurements would

have used direct measurements of biomass, as was performed for post treatment measurements (see below). However, a true measurement of biomass requires destructive sampling, which could not be performed prior to treatment applications for this experiment. Instead, since basal stem diameter has a strong relationship with total aboveground biomass for many woody shrubs and trees (Telfer 1969; Reeves and Lenhart 1988) we employed a non-destructive sampling method to estimate pre-treatment quadrat biomass by using basal stem circumference as an indicator of biomass. To verify that basal stem circumference was an accurate indicator of biomass for a species with such variable branching, we destructively sampled *E. pungens* plants outside of our study locations to test this relationship. We cut down representative samples of all the circumference size classes we had collected in the baseline measurements (3 to 50 cm). Basal stem circumference of 45 plants was measured at 15 cm (6 in) above ground level, with 15 plants collected outside of each study location (block). After recording the basal stem circumference of each individual plant, plants were cut, placed in paper bags, and transported to the laboratory where they were dried at 65 °C to a constant weight. We calculated the relationship between stem circumference and biomass using a power function regression analysis and observed a strong relationship between basal stem circumference and biomass ($y = 1.173x^{3.1079}$; $R^2 = 0.813$), suggesting that basal stem circumference is an adequate indicator of biomass for pretreatment quadrat comparisons.

Herbicide Treatments

We applied triclopyr herbicide using the free acid product formulation (Trycera[®], 343.92 g ai L⁻¹, Helena Agri-Enterprises LLC., Collierville, TN, United States) as foliar spray, basal bark, and cut stump application methods between January and September 2021 (Table 2). We used *E. pungen*'s annual cycle of growth initiation, timing of flowering (October to December) and fruiting (March to July) to determine the time of year that is most effective for each treatment in this environment (Dirr 1990, Ferrell et al. 2019, Miller 2006, Miller et al. 2013).

The foliar spray treatment was applied when *E. pungens* was actively growing new leaves, but the plant was not yet flowering. We used a 15 L backpack pump sprayer to apply the solution (.2% Trycera[®], Table 2) directly to the leaves and stems and attempted to spray to runoff. The basal bark application was made during fall, and the herbicide solution (25%

Trycera[®], Table 2) was applied to the lower 30 to 40 cm (12 to 16 in) of every *E. pungens* stem using a 15 L backpack pump sprayer. For cut stump applications, stems were cut with hand clippers or a chain saw about 15 cm (6 in) above ground level and treated directly per label directions. We used a 15 L backpack pump sprayer to apply the solution (100% Trycera[®], Table 2) directly to the cut stump, covering the entire wood surface. No action was taken in the untreated control plots.

Post-treatment measurements

On 16 May 2022 the total biomass of surviving *E. pungens* plants was destructively sampled in each treatment plot. These post treatment biomass measurements were taken the season following treatment application, 12, 8, and 16 months after the foliar spray, basal bark, and cut stump treatments, respectively. Because treatments were administered at different times of year (as is necessary for each treatment to be effective), the time window between application and sampling was not even among treatments (Ferrell et al. 2019, Miller et al. 2013). We ensured plants in each treatment had a full overwintering cycle to respond to herbicide application, and the time allotted was adequate to show the desired effect (Shaner 2014). We prioritized sampling all treatments on the same day to produce measurements that are not biased by seasonality.

On the day of sampling, all living *E. pungens* plants within each quadrat were clipped 15 cm (6 in) above ground level, including live foliage and woody material. Dead *E. pungens* material was left in the field. Living plant material was identified by retention of leaves and woody tissue with living cambium. Dead material was colorless, brittle, and leafless. The freshly clipped *E. pungens* stems were bagged and transported to the laboratory the following day, where the drying process began immediately. Once all bags were dried, the plant material was weighed to determine post-treatment biomass for each quadrat using the same process described above. All measurements for all treatments occurred on the same day.

Data Analysis

Statistical analyses were performed in SAS 9.4 using the ‘proc glimmix’ procedure (SAS Institute, Inc., 2023). All tests were performed using generalized linear models with treatment (‘UTC’, ‘foliar spray’, ‘basal bark’, and ‘cut stump’) as the fixed effect independent variables.

For the pretreatment assessment, two response variables were tested, mean stem circumference per m² quadrat and mean number of stems per m² quadrat. For the post-treatment assessment, the response variable tested was mean biomass per m² quadrat. The model included block (i.e., location) as a random effect and treatment as the fixed effect. Various distributions (Gaussian, Poisson, negative binomial, or lognormal) were examined for each response variable and selected based on optimal qualities: random spread in residual/ predicted plots, linear pattern in residual/ quantile plots, and low AICc values. A lognormal distribution was used for both pretreatment tests and the post treatment test. Treatment effects within each model were considered significant at $P < 0.05$. Significant models were then analyzed by Tukey's HSD to determine if differences occurred among individual treatments, and significance was accepted at $P < 0.05$.

Results and Discussion

Prior to application of treatments, neither the mean basal circumference (cm) nor the mean number of *E. pungens* stems per m² ($F_{3,42} = 1.29$; $P = 0.292$) was different among treatment plots ($F_{3,42} = 0.42$; $P = 0.738$) (Figure 2a,b), demonstrating that no preexisting bias existed among plots in terms of plant size or abundance prior to treatment applications. After treatment applications, biomass (kg m² of *E. pungens* was significantly different among treatments ($F_{3,42} = 26.63$; $P < 0.001$). Both cut stump and basal bark treatments resulted in significantly lower *E. pungens* biomass (0.012 ± 0.004 kg m² and 0.006 ± 0.005 kg m², respectively) compared with the foliar spray and untreated control (2.27 ± 1.39 kg m² and 2.97 ± 1.41 kg m², respectively) (Figure 2c). The post-treatment biomass in the cut stump and basal bark treatments were not different from each other, nor were those in the foliar spray and UTC treatments (Figure 2c).

Herbicides are used around the world for vegetation management in forest ecosystems and responsible use requires a constant refinement of application techniques to ensure the most efficient and effective management methods are being used (Little et al. 2006; Pile Knapp et al. 2023). One of our goals was to evaluate the realistic usability of these application techniques from a land manager's perspective (Kettenring and Adams 2011). Natural resource land managers typically consider several different facets of invasive plant management techniques

when determining which is most appropriate or useful for their specific situation (Lindenmayer et al. 2015; Kerr et al. 2016). In our study, we considered treatment effort, duration, and associated cost in addition to *E. pungens* mortality and related reduction in biomass to identify the optimal treatment method, and to that end our study provided immediate and useful results for managers. These data could be included in a decision tree to help guide management activities in similar areas (e.g., Lindenmayer et al. 2015).

Each treatment method we evaluated had pros and cons. While foliar herbicide application is often one of the fastest and least physically demanding application methods, it was the least effective treatment method in our study (Table 2, Figure 2c). Foliar applications of triclopyr as the free acid formulation had the most immediate and dramatic visual effect, but the resulting superficial visual crown reduction was misleading. Upper foliage was killed within 5 months of treatment (personal observation), and resprouting was evident around the plant base within 8 months of treatment (personal observation). In many cases, the bottom half of the plant remained healthy, and growth continued normally. Both cut stump and basal bark application methods significantly impacted *E. pungens* mortality (Figure 2c). Cut stump application took less time but was more physically demanding than the basal bark application method, and required three people to be on site while both foliar spray and basal bark applications were completed with two people (TL Evans, personal communication) (Table 2). The cut stump method required the applicators to cut through the base of the plant and drag the plant material out of the way to reach additional plants in other quadrats and move through the treatment area. This method left a great deal of large, sprawling, dead *E. pungens* branches on the forest floor with intertwined sprouts, making physical navigation difficult. Further, the untreated tops of recently felled *E. pungens* may still hold viable seeds for a period of time post-treatment. Some studies have shown the use of triclopyr through cut stump applications yielded high percentages of resprouts in plants prone to root suckering (DiTomaso and Kyser 2007; Fogliatto et al. 2020). Anecdotally, more resprouting appeared to be present in quadrats treated with the cut stump method than those treated with basal bark applications, but more research is needed to determine sprouting potential for *E. pungens* while using this method. While the basal bark application method required a slightly longer application time than cut stump or foliar spray, basal spray application was equally effective, less physically demanding on the applicator than cut stump applications and

could easily be performed by one person if necessary (TL Evans, personal communication) (Table 2).

Herbicide application should be conducted in a manner which minimizes negative impacts to non-target flora and fauna. To do this requires a combination of empirical data (e.g., Gibson et al. 2019) and knowledge of how active ingredients work. Selective treatments with triclopyr instead of a broad-spectrum herbicide (e.g., glyphosate) were used to avoid non-target impacts to native grasses. Other commonly used herbicides in forested settings (e.g., picloram) are soil active, and other non-target species may be injured through root absorption. Additionally, Trycera[®] carries an aquatic label allowing basal bark and cut stump applications in aquatic sites, making triclopyr a logical choice for this invasive species removal effort.

Recently, Yannelli et al. (2022) listed 15 emerging challenges and opportunities for vegetation science, one of which was halting forest degradation by targeted restoration in prioritized ecosystems. Accomplishing such a goal requires a thorough knowledge of methods to reduce or eliminate unwanted vegetation to facilitate the restoration of desired species. The details surrounding management costs (including finances, time, and labor) all factor into a land manager's decision-making process when determining when or if to engage in management activities for invasive plants (Zhang et al. 2023). Though some may posit that straightforward studies like these which evaluate a single active ingredient on a single target species are too basic to have broader global applicability, we argue that these studies are crucial in the restoration of degraded forest ecosystems worldwide. Our study provides several essential details for land managers dealing with *Elaeagnus* spp. in temperate systems.

The use of *E. pungens* in ornamental, hedgerow, wildlife, and roadside plantings are the primary causes for its current widespread distribution in North America. Restricting its sale and use for landscape and roadside plantings would contribute positively to reducing its spread. As with any invasive flora, maintaining healthy natural landscapes through the cultivation of native plant communities and weed prevention and control is more effective than strictly attempting to control invasion. However, cost-effective management options are necessary for land reclamation and ecosystem restoration in invaded areas (Kimball et al. 2015; Meli et al. 2017). This study addressed a common management question and provided strong evidence for two

effective application methods for *Elaeagnus* management in a forested setting. As many invasive shrubs in this region typically fill the same ecological niche and can be managed with similar techniques, our findings likely apply to other woody invasive species in North American natural areas (e.g., *Lonicera*, *Ligustrum*, *Rhamnus spp.*) for which triclopyr is already known to be an effective herbicide (Bisikwa et al. 2020, Delanoy and Archibold 2007, DiAllesandro 2012, Enloe et al. 2017, Enloe et al. 2018, Harrington and Miller 2017, Hogan et al. 2024, Mervosh and Gumbart 2015). We advocate for additional research to further develop management techniques for *Elaeagnus* and other invasive woody flora, as this knowledge will directly and positively impact natural area restoration efforts worldwide.

Acknowledgements

We thank Audubon South Carolina and the Columbia Chapter of the Audubon Society for project support. This work could not have been completed without the help of an amazing volunteer crew (Dave Schuetrum, Scott Weitecha, and Hank Stallworth, Audubon South Carolina Staff, Mark Musselman, Rick Armstrong, and Jessica Sharp-Miner).

Funding

Helena Agri-Enterprises, LLC graciously provided the product used in this study. This is Technical Contribution No. 7345 of the Clemson University Experiment Station. This material is based upon work supported by the NIFA/USDA, under project number SC-1050622,1700622. Any opinions, findings, conclusions, or recommendations expressed in this publication are those of the author(s) and do not necessarily reflect the view of the USDA.

Competing Interests

Competing interests: The author(s) declare none.

References

- Beaury EM, Patrick M, Bradley BA (2021) Invaders for sale: the ongoing spread of invasive species by the plant trade industry. *Front Ecol Environ* 19:550-556, [10.1002/fee.2392](https://doi.org/10.1002/fee.2392)
- Beschta RL, Ripple WJ (2009) Large predators and trophic cascades in terrestrial ecosystems of the western United States. *Conserv Biol* 142:2401-2414, [10.1016/j.biocon.2009.06.015](https://doi.org/10.1016/j.biocon.2009.06.015)
- Bisikwa J, Natukunda MI, Becker RL (2020) Effect of method and time of management on European Buckthorn (*Rhamnus cathartica* L.) growth and development in Minnesota. *J Sci Agric* 14:113-123, [10.25081/jsa.2020.v4.6457](https://doi.org/10.25081/jsa.2020.v4.6457)
- Bovey RW (1965) Control of Russian olive by aerial applications of herbicides. *J Range Manage* 18:194-195
- Bovey RW, Whisenant SG (1991) Control of honey mesquite with clopyralid, triclopyr, or clopyralid: triclopyr mixtures. *J Range Manage* 44:52-55
- Carter DR, Allen HL, Fox TR, Albaugh TJ, Rubilar RA, Campoe OC, Cook RL (2021) A 50-year retrospective of the forest productivity cooperative in the Southeastern United States: Regionwide trials. *J For* 119:73-85, [10.1093/jofore/fvaa046](https://doi.org/10.1093/jofore/fvaa046)
- Collette LKD, Pither J (2015) Russian-olive (*Elaeagnus angustifolia*) biology and ecology and its potential to invade northern North American riparian ecosystems. *Inv Plant Sci Manage* 8:1-14, [10.1614/IPSM-D-14-00037.1](https://doi.org/10.1614/IPSM-D-14-00037.1)
- Corns WG, Schraa RJ (1965) Mechanical and chemical control of silverberry (*Elaeagnus communtata* Bernh.) on native grassland. *J Range Manage* 18:15-19, [10.2307/3895741](https://doi.org/10.2307/3895741)
- Crystal-Ornelas R, Hudgins EJ, Cuthbert RN, Haubrock PJ, Fantle-Lepczyk J, Angulo E, Kramer AM, Ballesteros-Mejia L, Leroy B, Leung B, López-López E, Diagne C, Courchamp F (2021) Economic costs of biological invasions within North America. *NeoBiota* 67:485-510, [10.3897/neobiota.67.58038](https://doi.org/10.3897/neobiota.67.58038)
- Davison VE (1942) Use of certain *Elaeagnus* species. *J Ornithol* 59:581–583, [10.2307/4079479](https://doi.org/10.2307/4079479)
- Delanoy L, Archibold OW (2007) Efficacy of control measures for European buckthorn (*Rhamnus cathartica* L.) in Saskatchewan. *Environ Manage* 40: 709-718, [10.1007/s00267-006-0409-1](https://doi.org/10.1007/s00267-006-0409-1)
- Diagne C, Turbelin AJ, Moodley D, Novoa A, Leroy B, Angulo E, Adamjy T, Dia AKMC, Taheri A, Tambo J, Dobigny G, Courchamp F (2021) The economic costs of biological invasions in Africa: a growing but neglected threat? *NeoBiota* 67:11-51, [10.3897/neobiota.67.59132](https://doi.org/10.3897/neobiota.67.59132)

DiAllesandro A (2012) An assessment of foliar application of triclopyr of varying concentrations for managing glossy buckthorn (*Rhamnus frangula*) seedlings and resprouts (Michigan). *Ecol Rest* 30:18-19

DiTomaso JM, Kyser GB (2007) Control of *Ailanthus altissima* using stem herbicide application techniques. *AUF* 33:55-63, [10.48044/jauf.2007.007](https://doi.org/10.48044/jauf.2007.007)

Dirr MA (1990) Manual of woody landscape plants: their identification, ornamental characteristics, culture, propagation and uses. 4th ed. Champaign IL: Stipes Publishing Co. Pp 390-391

Edgin B, Ebinger JE (2001) Control of autumn olive (*Elaeagnus umbellata* Thunb.) at Beall Woods Nature Preserve, Illinois, USA. *Nat Areas J* 21:386–388

Enloe SF, O’Sullivan SE, Loewenstein NJ, Brantley EF, Lauer DK (2016) Triclopyr application timing and concentration influence low-volume basal bark efficacy on Chinese privet (*Ligustrum sinense*). *Inv Plant Sci Manage* 9:235-241, [10.1614/IPSM-D-16-00028.1](https://doi.org/10.1614/IPSM-D-16-00028.1)

Enloe SF, O’Sullivan SE, Loewenstein NJ, Brantley E, Lauer DK (2018) The influence of treatment timing and shrub size on Chinese privet (*Ligustrum sinense*) control with cut stump herbicide treatments in the southeastern United States. *IPSM* 11: 49-55, [10.1017/inp.2018.3](https://doi.org/10.1017/inp.2018.3)

Enloe SF, Leary JK, Lastinger C, Lauer DK (2023) Reduced hack and squirt treatment with aminocyclopyrachlor and aminopyralid for invasive shrub control. *Inv Plant Sci Manage* 16:64-72, [10.1017/inp.2023.10](https://doi.org/10.1017/inp.2023.10)

Essl F, Latombe G, Lenzner B, Pagad S, Seebens H, Smith K, Wilson JRU, Genovesi P (2020) The convention on biological diversity (CBD)’s post-2020 target on invasive alien species – what should it include and how should it be monitored? *NeoBiota* 62:99-121, [10.3897/neobiota.62.53972](https://doi.org/10.3897/neobiota.62.53972)

Ferrell J, Langeland K, Sellers B (2019) Herbicide application techniques for woody plant control. University of Florida Extension Publications SS-AGR-260. <https://edis.ifas.ufl.edu/publication/ag245>. Accessed Dec 12. 2024

Fertakos ME, Beaury EM, Ford NR, Kinlock NL, Adams DW, Bradley BA (2023) Historical plant sales (HPS) database: Documenting the spatiotemporal history of plant sales in the conterminous U.S. *Ecol* 104:e4106, [10.1002/ecy.4106](https://doi.org/10.1002/ecy.4106)

Fischer RA, Valente JJ, Guilfoyle MP, Kaller MD, Jackson SS, Ratti JT (2012) Bird community response to vegetation cover and composition in riparian habitats dominated by russian olive (*Elaeagnus angustifolia*). *Northwest Sci* 86:39-52, [10.3955/046.086.0104](https://doi.org/10.3955/046.086.0104)

Fogliatto S, Milan M, Vidotto F (2020) Control of *Ailanthus altissima* using cut stump and basal bark herbicide applications in an eighteenth-century fortress. *Weed Res* 60:425-434, [10.1111/wre.12447](https://doi.org/10.1111/wre.12447)

Franke M, Zipper C, Barney J (2018) Native hardwood tree seedling establishment following invasive autumn-olive (*Elaeagnus umbellata*) removal on a reclaimed coal mine. *Inv Plant Sci Manage* 11:155-161, [10.1017/inp.2018.19](https://doi.org/10.1017/inp.2018.19)

Gibson DJ, Shupert LA, Liu X (2019) Do no harm: efficacy of a single herbicide application to control an invasive shrub while minimizing collateral damage to native species. *Plants* 8:426, [10.3390/plants8100426](https://doi.org/10.3390/plants8100426)

Harrington TB and Miller JH (2005) Effects of application rate, timing, and formulation of glyphosate and triclopyr on control of Chinese privet (*Ligustrum sinense*). *Weed Technol* 19:47-54, [10.1614/WT-03-220R2](https://doi.org/10.1614/WT-03-220R2)

Hogan KFE, Baker K, Back EM, Barber NA (2024) Basal bark herbicide treatment of *Lonicera maackii* (Amur honeysuckle) is effective regardless of application timing, with limited nontarget effects on native plant diversity. *Ecol solu evid* 5:e12332, [10.1002/2688-8319.12332](https://doi.org/10.1002/2688-8319.12332)

Holmes TP, Aukema JE, Von Holle B, Liebhold A, Sills E (2009) Economic impacts of invasive species in forests. *Ann NY Acad Sci* 1162:18-38, [10.1111/j.1749-6632.2009.04446.x](https://doi.org/10.1111/j.1749-6632.2009.04446.x)

Katz GL, Tuttle GM, Denslow MW, Norton AP (2020) Legacy effects of russian olive (*Elaeagnus angustifolia* L.) in a riparian ecosystem three years post-removal. *Wetland* 40:1897-1907, [10.1007/s13157-020-01385-3](https://doi.org/10.1007/s13157-020-01385-3)

Kettenring KM, Adams CR (2011) Lessons learned from invasive plant control experiments: a systematic review and meta-analysis. *J Appl Ecol* 48:970-979, [10.1111/j.1365-2664.2011.01979.x](https://doi.org/10.1111/j.1365-2664.2011.01979.x)

Kerr NZ, Baxter PWJ, Salguero-Gómez R, Wardle GM, Buckley YM (2016) Prioritizing management actions for invasive populations using cost, efficacy, demography and expert opinion for 14 plant species world-wide. *J Appl Ecol* 53:305-316, [10.1111/1365-2664.12592](https://doi.org/10.1111/1365-2664.12592)

Kimball S, Lulow M, Sorenson Q, Balazs K, Fang Y-C, Davis SJ, O'Connell M, Huxman TE (2015) Cost-effective ecological restoration. *Rest Ecol* 23:800-810, [10.1111/rec.12261](https://doi.org/10.1111/rec.12261)

Kimbro DL, Grosholz ED, Baukus AJ, Nesbitt JN, Travis NM, Attoe S, Coleman-Hulbert C (2009) Invasive species cause large-scale loss of native California oyster habitat by disrupting trophic cascades. *Oecologia* 160:563-575, [10.1007/s00442-009-1322-0](https://doi.org/10.1007/s00442-009-1322-0)

Lázaro-Lobo A, Lucardi RD, Ramirez-Reyes C, Ervin GN (2021) Region-wide assessment of fine-scale associations between invasive plants and forest regeneration. *For Ecol Manage* 483:118930, [10.1016/j.foreco.2021.118930](https://doi.org/10.1016/j.foreco.2021.118930)

Liebhold AM (2012) Forest pest management in a changing world. *Int J Pest Manage* 58:289-295, [10.1080/09670874.2012.678405](https://doi.org/10.1080/09670874.2012.678405)

Lindenmayer DB, Wood J, MacGregor C, Buckley YM, Dexter N, Fortescue M, Hobbs RJ, Catford JA (2015) A long-term experimental case study of the ecological effectiveness and cost effectiveness of invasive plant management in achieving conservation goals: Bitou bush control in Booderee National Park in eastern Australia. *PLoS One* 10:e0128482, [0.1371/journal.pone.0128482](https://doi.org/10.1371/journal.pone.0128482)

Little KM, Willoughby I, Wagner RG, Adams P, Frochot H, Gava J, Gous S, Lautenschlager RA, örlander G, Sankaran KV, Wei RP (2006) Towards reduced herbicide use in forest vegetation management. *South African For J* 207:63-79, [10.2989/10295920609505254](https://doi.org/10.2989/10295920609505254)

Malinich E, Lynn-Bell N, Kourtev PS (2017) The effect of the invasive *Elaeagnus umbellata* on soil microbial communities depends on proximity of soils to plants. *Ecosphere* (Wash DC) 8:e01827, [10.1002/ecs2.1827](https://doi.org/10.1002/ecs2.1827)

Maynard-Bean E, Kaye M, Wagner T, Burkhart EP (2020) Citizen scientists record novel leaf phenology of invasive shrubs in eastern U.S. forests. *Biol Invasions* 22:3325–3337, [10.1007/s10530-020-02326-1](https://doi.org/10.1007/s10530-020-02326-1)

Meli P, Herrera FF, Melo F, Pinto S, Aguirre N, Musálem K, Minaverri C, Ramírez W, Brancalion PHS (2017) Four approaches to guide ecological restoration in Latin America. *Rest Ecol* 25:156-163, [10.1111/rec.12473](https://doi.org/10.1111/rec.12473)

Mervosh TL, Gumbart D (2015) Cutting and herbicide treatments for control of Oriental bittersweet, Pale swallow-wort and Morrow's honeysuckle. *Natural Areas J* 35:256-265, [10.3375/043.035.0206](https://doi.org/10.3375/043.035.0206)

Michielsen, M, Szemák L, Fenesi A, Nijs I, Ruprecht E (2017) Resprouting of woody species encroaching temperate European grasslands after cutting and burning. *Appl Veg Sci* 20:388-396, [10.1111/avsc.12300](https://doi.org/10.1111/avsc.12300)

Miller JH (2006) Nonnative invasive plants of southern forests: a field guide for identification and control Asheville NC: US Department of Agriculture, Forest Service, Southern Research Station. 15 p

Miller JH, Manning ST, Enloe SF (2013) A management guide for invasive plants in southern forests. Asheville NC: US Department of Agriculture Forest Service, Southern Research Station. 120 p

Mineau MM, Baxter CV, Marcarelli AM (2011) A non-native riparian tree (*Elaeagnus angustifolia*) changes nutrient dynamics in streams. *Ecosystems* 14:353–365, [10.1007/s10021-011-9415-0](https://doi.org/10.1007/s10021-011-9415-0)

Muscha JM, Vermeire LT, Angerer JP (2023) Fire reduces Russian olive seed germination and seedling survival with increasing fuel load. *Rest Ecol* 31:e13904, [10.1111/rec.13904](https://doi.org/10.1111/rec.13904)

Napton DE, Auch RF, Headley R, Taylor JL (2010) Land changes and their driving forces in the Southeastern United States. *Reg Environ Change* 10:37-53, [10.1007/s10113-009-0084-x](https://doi.org/10.1007/s10113-009-0084-x)

National Weather Service (2024) National Oceanic and Atmospheric Administration. <https://www.weather.gov/wrh/Climate?wfo=cae>. Accessed October 19, 2023

Oswalt CM, Fei S, Guo Q, Iannone III BV, Oswalt SN, Pijanowski BC, Potter KM (2015) A subcontinental view of forest plant invasions. *NeoBiota* 24:49-54, [10.3897/neobiota.24.8378](https://doi.org/10.3897/neobiota.24.8378)

Pile Knapp LS, Coyle DR, Dey DC, Fraser JS, Hutchinson T, Jenkins MA, Kern CC, Knapp BO, Maddox D, Pinchot C, Wang GG (2023) Invasive plant management in eastern North American Forests: A systematic review. *For Ecol Manage* 550:121517, [10.1016/j.foreco.2023.121517](https://doi.org/10.1016/j.foreco.2023.121517)

Reeves HC, Lenhart JD (1988) Fuel weight prediction equations for understory woody plants in Eastern Texas. *Texas J Sci* 40:49-53

Riffe EC (2018) A study of autumn olive (*Elaeagnus umbellata*) phenology and associated physiological traits that may facilitate its invasion of the understory of a southern Appalachian forest. M.S. thesis. Boone, NC: Appalachian State University. 75 p

SAS Institute Inc. (2023) *SAS/STAT® 15.3 User's Guide*. Cary, NC

Seebens H, Essl F, Blasius B (2017) The intermediate distance hypothesis of biological invasions. *Ecol Lett* 20:158-165, [10.1111/ele.12715](https://doi.org/10.1111/ele.12715)

Serviss BE, Tumilson R (2021) Guide to the naturalized, escaped, and adventive woody flora of Arkansas. *Phytoneuron* 29: 10-51

Schmitz OJ, Hamback PA, Beckerman AP (2000) Trophic cascades in terrestrial systems: A review of the effects of carnivore removals on plants. *Am Nat* 155:141-153, [10.1086/303311](https://doi.org/10.1086/303311)

Shaner DL, ed (2014) *Herbicide handbook*. 10th ed. Lawrence, KS: Weed Science Society of America. Pp 459-461

Telfer ES (1969) Weight-diameter relationships for 22 woody plant species. *Can J Bot* 47:1851-1855, [10.1139/b69-271](https://doi.org/10.1139/b69-271)

United States Department of Agriculture (2019) Natural Resources Conservation Service, Web Soil Survey. <https://websoilsurvey.nrcs.usda.gov/app/>. Accessed November 8, 2023

Watts BD, Paxton BJ (2000) The influence of thorny elaeagnus on automobile-induced bird mortality. Center for Conservation Biology Technical Report Series. Williamsburg, VA: College of William and Mary. 18 p

Yannelli FA, Bazzichetto M, Conradi T, Pattison Z, Andrade B, Anibaba QA, Bonari G, Chelli S, Cuk M, Damasceno G, Fantinato E, Geange S, Guuroh R, Musa Holle MJ, Kůzmič F, Lembrechts J, Mosyftiani A, Šikuljak T, Teixeira J, Tordoni E, Pérez-Valladares CX, Sperandii MG (2022) Fifteen emerging challenges and opportunities for vegetation science: A horizon scan by Early Career Researchers. *J Veg Sci* 3:e13119, [10.1111/jvs.13119](https://doi.org/10.1111/jvs.13119)

Yates ED, Levia DF, Williams CL (2004) Recruitment of three non-native invasive plants into a fragmented forest in southern Illinois. *For Ecol Manage* 190:119-130, [10.1016/j.foreco.2003.11.008](https://doi.org/10.1016/j.foreco.2003.11.008)

Zar JH (2010) *Biostatistical Analysis*. 5th Ed. Essex: Pearson Education Limited. Pp 103-106

Zenni RD, Essl F, García-Berthou E, McDermott SM (2021) The economic costs of biological invasions around the world. In: Zenni RD, McDermott S, García-Berthou E, Essl F (Eds) *The economic costs of biological invasions around the world*. *NeoBiota* 67:1-9, [10.3897/neobiota.67.69971](https://doi.org/10.3897/neobiota.67.69971)

Zhang B, Zhai L, Ervin GN, Coyle DR. 2023. Effective and timely use of models to inform on-the-ground management of invasive plants. *Biol Invasions* 25:2089-2102, [10.1007/s10530-023-03043-1](https://doi.org/10.1007/s10530-023-03043-1)

Table 1: Site characteristics for each experimental block in the *Elaeagnus pungens* management study conducted in Calhoun Co., South Carolina, United States.

Study location (Block)	1	2	3
Coordinates	33° 38' 16.9404" N, 80° 42' 26.0028" W	33° 38' 21.3108" N, 80° 42' 47.8296" W	33° 38' 26.502" N, 80° 42' 23.6232" W
Soil series	Faceville fine sandy loam, 2 to 6 percent slopes	Ailey-Vaucluse complex, 6 to 15 percent slopes	Faceville fine sandy loam, 2 to 6 percent slopes
Species composition	<p><u>Overstory:</u> <i>Liquidambar styraciflua</i></p> <p><u>Midstory:</u> <i>Elaeagnus pungens</i></p> <p><u>Understory:</u> None</p>	<p><u>Overstory:</u> <i>Ulmus alata</i>, <i>Liquidambar styraciflua</i>, <i>Pinus taeda</i>, <i>Quercus alba</i>, <i>Fraxinus</i> spp.</p> <p><u>Midstory:</u> <i>Elaeagnus pungens</i>, <i>Ligustrum sinense</i>, <i>Acer negundo</i>, <i>Morus rubra</i>, <i>Celtis laevigata</i></p> <p><u>Understory:</u> <i>Elaeagnus pungens</i>, <i>Ligustrum sinense</i>, <i>Vitis rotundifolia</i> Michx., various native grasses</p>	<p><u>Overstory:</u> <i>Quercus alba</i>, <i>Quercus falcata</i>, <i>Pinus taeda</i>, <i>Fagus grandifolia</i>, <i>Liquidambar styraciflua</i>, <i>Carya</i> spp.</p> <p><u>Midstory:</u> <i>Cornus florida</i> L., <i>Elaeagnus pungens</i></p> <p><u>Understory:</u> <i>Vitis rotundifolia</i>, <i>Elaeagnus pungens</i>, <i>Smilax rotundifolia</i> L., various herbaceous species</p>
Basal area (BA)	BA: 33.5 to 39.6 m ² ha ⁻¹	BA: 18.3 to 24.4 m ² ha ⁻¹	BA: 18.3 to 27.4 m ² ha ⁻¹
Average root collar diameter (rcd) of dominant overstory species	dbh: 15.2 to 20.3 cm	dbh: 35.6 to 51 cm	dbh: 25.4 to 40.6 cm

Table 2: Field application details for experimental treatments: foliar spray, cut stump, and basal bark.

Treatment	Foliar spray	Cut stump	Basal bark
Date applied	14 May 2021	7 January 2021	10 September 2021
Temperature relative humidity (RH) ⁻¹	24°C 35% RH ⁻¹	14°C 79% RH ⁻¹	29°C 76% RH ⁻¹
Dilution type	Formulated product ¹ diluted in water	Formulated product ¹ undiluted	Formulated product ¹ diluted in petroleum-oil carrier
Application dilution (L Trycera ¹ + L dilutant)	0.3L + 14.7L H ₂ O	(undiluted)	0.95L + 2.85L oil
Concentration (g triclopyr L ⁻¹ applied solution)	6.9g L ⁻¹	344 g L ⁻¹	86.0g L ⁻¹
Application duration	4 hours	5 hours	6 hours

¹Trycera[®], 344 g ai triclopyr L⁻¹.



Figure 1. Characteristics of *Elaeagnus pungens* in Calhoun Co., South Carolina, United States: (A) dense, sprawling growth; (B) leaf surface is dark green and waxy; (C) leaf undersides are silver and reflective; (D) fruit are red drupes; (E) thorns 2.5 to 5 cm in length grow on branches; (F) growth is multi-stemmed and freely branched. Photos A, D, E, and F by M.N. Darr; photos B and C by D.R. Coyle.




Figure 2. (A) Stem size pre-treatment comparison from an *Elaeagnus pungens* management study in Calhoun Co., South Carolina, United States: mean (\pm SE) stem circumference of *Elaeagnus pungens* per m² plot for each experimental treatment. (B) Stand density pretreatment comparison: mean (\pm SE) number of *Elaeagnus pungens* stems per m² plot for each experimental treatment. (C) Above round plant biomass post-treatment comparison: mean (\pm SE) *Elaeagnus pungens* dry weight per m² plot for each experimental treatment. *Elaeagnus pungens* biomass was significantly greater in the untreated control and foliar spray treatment compared to the basal bark and cut stump application methods. Means sharing the same letter are not significantly different from each other.