

Animal Research Paper

Cite this article: Beltrán MC, Manzur A, Rodríguez M, Díaz JR, Peris C (2018). Effect of mid-line or low-line milking systems on lipolysis and milk composition in dairy goats. *The Journal of Agricultural Science* **156**, 848–854. <https://doi.org/10.1017/S0021859618000771>

Received: 15 January 2018
Revised: 30 July 2018
Accepted: 6 September 2018
First published online: 2 October 2018

Key words:

Goat milk; lipolysis; milking system; mid-line milking; low-line milking

Author for correspondence:

M.C. Beltrán, E-mail: mbeltran@dca.upv.es

Effect of mid-line or low-line milking systems on lipolysis and milk composition in dairy goats

M. C. Beltrán¹, A. Manzur², M. Rodríguez¹, J. R. Díaz³ and C. Peris¹

¹Institut de Ciència i Tecnologia Animal, Universitat Politècnica de València, Camí de Vera, s/n. 46022 València, Spain; ²Departamento de Rumiantes, Facultad de Medicina Veterinaria y Zootecnia, Universidad Autónoma de Chiapas, Ctra. Tuxtla-Ejido Emiliano Zapata, km 8, Mexico and ³Departamento de Tecnología Agroalimentaria, Escuela Politécnica Superior de Orihuela, Universidad Miguel Hernández, Ctra. Beniel, Km 3.2, 03312 Orihuela, Spain

Abstract

Two experiments were carried out to investigate how milking in mid-line (ML) affects the lipolysis level and milk composition in goat livestock, in comparison with low-line (LL) milking. The first experiment took place, in triplicate, on an experimental farm. For each replicate, a crossover design (62 goats, two treatments, ML and LL, in two periods each lasting 4 days) was used. Milk samples were taken daily at 0 and 24 h after milking. In the first experimental replicate, some enzymatic coagulation cheeses were made, which were assessed by a panel of tasters at 50 and 100 days of maturation. In the second experiment, the lipolysis level and composition of tank milk from 55 commercial dairy goat farms (25 ML and 30 LL) were analysed, in milk samples taken in three different weeks. The results of the first experiment showed that ML milking increased free fatty acid (FFA) concentration in raw goat's milk significantly (0.71 v. 0.40 mmol/l, respectively). However, in the milk samples taken from commercial farms the FFA concentration remained unaffected by the milking pipeline height (0.59 v. 0.58 mmol/l for ML and LL, respectively). No significant differences were found in the milk composition, nor in the sensory characteristics in the cured cheeses, which suggests that factors other than the milking height are able to influence the level of lipolysis under commercial conditions. Therefore, ML milking should not be discouraged, provided that the correct functioning and management of the milking operation and milk storage on the farm is guaranteed.

Introduction

Milk fat lipolysis consists of enzymatic hydrolysis of the triglycerides of fat globules, which leads to an increase in the concentration of free fatty acids (FFAs) and, consequently, of the acidity of the fat. In raw milk, lipolysis is mainly caused by the action of lipoprotein lipase (LPL), a natural milk enzyme synthesized in secretory epithelial cells (Chen *et al.*, 2003), although it may be partly haematic in origin (Chilliard *et al.*, 2003). Other enzymes with significant lipolytic activity are lipases of microbial origin, mainly from psychotropic germs (Ouattara *et al.*, 2004), and somatic cell lipases (Gargouri *et al.*, 2008).

Different physiological, genetic and nutritional factors have been identified that may influence the LPL activity in raw milk (Deeth, 2006; Chilliard *et al.*, 2014). The good correlation between LPL activity and lipolysis could be due to a higher degree of association of LPL with the fat phase in goat's milk (Chilliard *et al.*, 1984), thus increasing the enzyme–substrate interactions, contrary to what occurs in cow's milk (Chilliard *et al.*, 2003). On the other hand, it has also been shown that the factors that impair fat globule membrane integrity, such as excessive shaking or abrupt changes in storage temperature, can increase the lipolysis level in milk by exposing the triglycerides to lipase action (Meffe, 1994; Chen *et al.*, 2003).

It should be noted that the hydrolytic release of FFA from triglycerides can have negative consequences for the dairy industry. First, it may affect the technological properties of milk, causing fat loss and delays in the growth of starter cultures used in the production of fermented products such as cheese or yoghurt (IDF 1991; Collomb and Spahni, 1995). It may also give rise to the appearance of off-flavours, described as rancid, butyric, astringent or even bitter, in milk and its by-products (Le Mens *et al.*, 1997; Deeth 2006). For this reason, FFA concentration is often used as an indicator of the organoleptic quality of milk, which is occasionally included in inter-professional regulations for payment by quality (Pirisi *et al.*, 2007; Skeie 2014). Additionally, another possible negative effect of lipolysis is that it could affect the analytical results of milk composition obtained with infrared equipment (IDF 2000), as the release of FFA from triglycerides due to lipase action changes the readings (absorbances) from the equipment in certain wavelengths that affect the determination of fat (fat wavelength A, 5.7 µm) and protein (protein wavelength: 6.5 µm). Similarly, Robertson *et al.* (1981) stated that in cow's milk

an increase in FFA of 1 meq/l resulted in analyses with infrared equipment showing a decrease in fat (-0.033%) and an increase in crude protein content ($+0.019\%$). These analytical changes are also reflected in the IDF standard for cow's milk analysis with mid-infrared based equipment (IDF 2000).

Although lipolysis progresses during milk storage, most of it occurs in the first 24 h of refrigeration (Wiking *et al.*, 2003; Ouattara *et al.*, 2004), before it reaches the dairy industry. Therefore, the mechanical stress that affects milk in the milking systems becomes a crucial aspect to preserve milk supply and quality.

Several studies in cattle have demonstrated the relationship between certain milking conditions and the increase in lipolysis in milk (Pillay *et al.*, 1980; Escobar and Bradley, 1990; Abeni *et al.*, 2005). One of these factors is milking with a milk line located above the animals' standing level, i.e. in mid-line (ML) or high-line (HL), with elevations lower or higher than 1.25 m, respectively (ISO, 2007a). In this type of setup, the milk is mixed with air so that it rises through the long milk tube, forming bubbles and, therefore, being submitted to more turbulent agitation. In cattle, it has been shown that HL milking, in comparison with low-line milking (LL), increases milk lipolysis (Gudding and Lorentzen, 1982; Mikulová, 2011); the higher the milkline height or the air intake caused by the milking cluster, the greater the increase (Judge *et al.*, 1977; Meffe, 1994; Rasmussen *et al.*, 2006). However, information about this is scarce in small ruminants, even though the use of ML milking has become increasingly popular in recent years. This is because, with an equal number of milking stalls, the installation of an ML usually cuts initial investment by around 25–35% (Díaz *et al.*, 2004) compared with LL. To the best of our knowledge, there are only two studies: one comparing ML *v.* LL in sheep (Díaz *et al.*, 2004) and another comparing HL *v.* LL in goats (Morand-Fehr *et al.*, 1983), neither of which found statistically significant differences in FFA concentrations in milk. Further studies along these lines would be warranted to determine whether the use of ML or HL milking can negatively affect the quality of goat's milk; a crucial aspect when considering that goat's milk is mainly used to manufacture cheese, whose sensory characteristics might be affected by an increase in the FFA concentration as a result of excessive lipolysis in the milk. In this sense, Morgan *et al.* (2001) noted a high risk of obtaining lactic coagulation cheeses with unacceptable sensory characteristics when the FFA concentration is ≥ 1 g oleic acid/100 g milkfat (3.5 meq/100 g milkfat) in goat's milk. However, there is no information on the effect of lipolysis on the sensory quality of cheeses made by enzymatic coagulation, a processing technique widely used in traditional goat cheese-making of Mediterranean countries. There are no studies evaluating the effect of lipolysis in goat's milk on the results of analyses performed with infrared equipment by milk quality laboratories. To this end, the aim of the current study was to assess the influence of ML milking system on lipolysis and components of goat's milk that are determined routinely with infrared equipment, taking milk from LL milking as a reference. The effect of ML milking on the sensory features of enzymatic coagulation goat cheeses was also evaluated.

Material and methods

Experimental procedure

To meet the aforementioned objectives, two experiments were carried out: the first at the dairy goat experimental farm of the Universitat Politècnica de València (UPV, Valencia, Spain) and

the second on commercial dairy goat farms, whose bulk milk was routinely analysed at the Interprofessional Dairy Laboratory of the Valencian Community Region (LILCOVAL, Valencia, Spain).

First experiment

This experiment was carried out in triplicate on the Universitat Politècnica de València (UPV) experimental farm. Each replicate experiment was designed as follows: 62 Murciano-Granadina breed goats, halfway through the lactation period (4 ± 1.0 month of lactation), were used. The animals were divided into two groups of 28 goats each, according to production level and lactation number, with each group randomly assigned to ML or LL milking for an initial 4-day period. Then, the treatments (ML and LL) were switched between the two groups for a second experimental 4-day period.

The goats were milked once on a daily basis (08:30 h) following a routine which included machine stripping, manual teat-cup removal and iodine post-dipping solution.

The milking parlour (2×12) had two milking pipelines installed with six clusters in ML and 12 clusters in LL. The ML milkline, dead in type, had an internal diameter of 52 mm and length of 520 cm and was located at 112 cm above the goat standing level. The LL milkline, looped type, was 52 mm in diameter, 1500 cm in length and located at 40 cm below the standing level.

The Almatic™ cluster G50 from DeLaval (Tumba, Sweden) was used in the current study. However, in the case of ML milking, a claw from DeLaval cluster SG-TF80 (claw volume 100 ml) was incorporated.

A different nominal vacuum was set (40 kPa in ML and 37 kPa in LL) so that the average teat-end vacuum, in the absence of milk flow during milking, was similar in both types of milking systems. The effective reserve (ML: 750 litres/min; LL: 950 litres/min) complied with international recommendations (higher than 512 and 804 litres/min in ML and LL, respectively; extra air for automatic teat-cup valves: 32 litres/min; ISO, 2007b). The pulsation rate (90 cycles/min) and ratio (60:40) were the same in ML and LL. The air intakes in the milking cluster (7.5 litres/min) were produced in the inlets at the base of the teat-cup liners.

The milk from the two groups of animals was stored separately in tanks that were empty at the start of each experimental period and which accumulated the milk obtained during each 4-day set. Bulk milk samples (250 ml) were taken on a daily basis from each batch of animals immediately after milking (0 h) and 24 h later, just before starting the next milking, to determine the following variables: FFA content, main milk components (fat, protein, lactose and dry matter), pH, freezing point, somatic cell count (SCC) and total bacterial count (TBC). For the FFA analysis, a 30 ml aliquot was separated after sampling, adding hydrogen peroxide (0.02%) following IDF recommendations (1991). In addition, the bulk milk accumulated during each 4-day period was used to prepare enzymatic coagulation cheeses, whose sensory characteristics were assessed by a test panel after 50 and 100 days' maturation. The cheese-making could only be performed in the first experimental replicate.

Second experiment

Of the 200 commercial dairy goat farms whose milk is usually analysed by the Interprofessional Dairy Laboratory of the Valencian Community Region (LILCOVAL, Valencia, Spain), 55 were chosen at random, 25 with ML milking and 30 with LL. All these commercial exploitations produce milk from Murciano-Granadina goats and carry out a daily milking routine

Table 1. Average values of parameters in bulk milk samples of Murciano-Granadina breed goats according to the type of milking used (ML: mid-line; LL: low-line) and refrigerated storage time (0 and 24 h post-milking) obtained under experimental farm conditions

Parameter	Milking type			Refrigerated storage time (h)		
	ML	LL	S.E.M.	0	24	S.E.M.
FFA (mmol/l)	0.71	0.40	0.015***	0.50	0.61	0.015***
Fat (g/kg)	45.4	45.5	0.18 ^{N.S.}	45.6	45.3	0.18 ^{N.S.}
Protein (g/kg)	33.3	33.3	0.08 ^{N.S.}	33.3	33.3	0.08 ^{N.S.}
Lactose (g/kg)	45.0	44.9	0.06 ^{N.S.}	44.9	44.9	0.06 ^{N.S.}
Dry mater (g/kg)	132.6	132.7	0.27 ^{N.S.}	132.8	132.5	0.27 ^{N.S.}
pH	6.74	6.74	0.008 ^{N.S.}	6.76	6.73	0.008*
Freezing point (°C)	-0.554	-0.554	0.0016 ^{N.S.}	-0.553	-0.555	0.0016 ^{N.S.}
SCC log	6.11	6.11	0.009 ^{N.S.}	6.11	6.11	0.009 ^{N.S.}
TBC log	5.31	5.47	0.086 ^{N.S.}	5.29	5.49	0.090 ^{N.S.}

FFA, free fatty acids (mmol/l); SCC log, somatic cell count (cell/ml) logarithm; TBC log, total bacterial count (cfu/ml) logarithm. S.E.M., standard error of the mean.

* $P < 0.05$ and *** $P < 0.001$. Degrees of freedom for milking type and refrigerated storage time are respectively, 1 and 1.

including machine stripping, manual teat-cup removal and some of them use an iodine post-dipping solution.

Milking parlours had one or two milking platforms (0.28 and 0.62, respectively), most of them having between 12 and 24 stalls per platform. In ML, there was usually a milking cluster for every 2–4 stalls, whereas the most frequent setup in LL was one milking cluster for every 1–2 stalls.

Bulk milk samples from commercial farms were taken in 50 ml flasks containing azidiol as a preservative (133 µl/40 ml milk), as stipulated by the Spanish legislation (Ministerio de la Presidencia, 2011). Milk sampling was performed weekly during three consecutive weeks, between April and May (one sample/week and farm) to determine the same variables as in the first experiment.

Cheese procedure

Four batches of cheese were prepared in a commercial artisan cheese factory, with milk collected by LL and ML milking from each of the two periods considered in the first replicate of the experiment.

Pasteurized goat's milk (74 °C, 15 s) inoculated with starter cultures and spiked with calcium chloride was coagulated by rennet at 32 ± 1.0 °C. After coagulation (40 min), the curd was cut and gently shaken for 20–30 min while the temperature was steadily increased until reaching a maximum of 38 °C. After moulding, the cheeses (900–1000 g) were pressed for 2 h under increasing pressure until a pH value of 5.3–5.4 was reached. Next, cheeses were salted immersed in brine (22 Bè) for 4 h and then placed in an airing chamber (4 °C, 75% RH) for 48 h. Finally, the cheeses were kept in a ripening chamber (10–12 °C, 80–85% RH) for a 100-day period.

Analytical procedure

FFA quantification was performed, in duplicate, at the Interprofessional Dairy Laboratory of Cantabria Region (LILC, Santander, Spain) using the copper soap method (IDF, 1991). The SCC was analysed with Fossomatic 5000 equipment (Foss, Hillerød, Denmark) and the chemical composition of the milk (fat, protein, lactose and dry matter) was analysed using Milko Scan FT 6000 infrared equipment (Foss). The freezing point

and the pH of goat's milk were determined by reference methods using a thermistor cryoscope (Cryostar, Fungger-Gerber, Germany) and a conventional pH meter (Crison Instruments, Barcelona, Spain), respectively. The TBC of the milk samples was determined from the standard plate count at 30 °C (ISO, 2013).

Sensory analysis of cheeses made with milk from ML and LL milking was performed by a panel of 62 tasters (balanced 0.50 by gender and aged from 20 to 55 years) through a triangular test (ISO, 2004). At each tasting session, the judges analysed two successive triads with cheeses from both experimental periods. These tests were performed repeatedly at 50 and 100 days of maturation, obtaining a total of 248 evaluations.

Statistical analysis

In the first experiment, the milk quality variables (FFA content, main milk components (fat, protein, lactose and dry matter), pH, freezing point, SCC and TBC) were analysed using PROC GLM procedures in SAS 9.2, with a model that included the following fixed effects: milkline (ML and LL), post-milking time (0 and 24 h), replication of the trial (1–3), day of the period (1–4), their respective interactions and the effect of the group of animals within each replicate.

The variables of the second experiment were analysed using PROC MIXED procedures in SAS 9.2, as per Littell *et al.* (1998), using a model that considered the fixed effects of the milkline (ML and LL), the week of sampling (1–3) and their interaction and the random effect of the farm (1–55). In both analyses, when an interaction was non-significant, the corresponding interaction term was pooled with the error.

The data obtained from the sensorial analysis of the cheeses (frequency of hits in the triangular test) were analysed statistically based on the binomial distribution of the parameter $P = 1/3$ with n responses (ISO, 2004).

Results

Goat's milk quality parameters

In the first experiment it was observed that ML milking caused a significant increase ($P < 0.001$) in the FFA concentration in goat's




Fig. 1. FFA concentrations (average value \pm s.e.m., expressed as mmol/l) in the bulk milk of Murciano-Granadina breed goats according to the type of milking (ML: mid-line; LL: low-line) in each replicate of the study conducted under experimental conditions.




Fig. 2. Evolution of FFA concentration (mmol/l) in bulk milk of Murciano-Granadina breed goats during a 4-day storage period (mean values of milk from ML and LL milking).

milk compared with LL milking (Table 1). This was the case in milk samples taken immediately after milking (ML: 0.64 ± 0.020 mmol/l; LL: 0.35 ± 0.020 mmol/l) as well as in those taken after 24 h of refrigerated storage (ML: 0.77 ± 0.020 mmol/l; LL: 0.45 ± 0.020 mmol/l). A higher level of lipolysis in ML than in LL was also found in each of the three experimental replicates, although in the last replicate a smaller difference was observed (milkline \times replicate interaction significant, $P < 0.05$; Fig. 1).

The time elapsed since milking also significantly affected the FFA concentration in goat's milk, in such a way that the milk samples taken after 24 h in refrigerated tank storage presented higher values ($P < 0.001$) than those taken immediately after milking (Table 1). Concerning the evolution of lipolysis with storage days, Fig. 2 shows how the release of FFA in bulk milk tended to increase during the 4-day study period, although the differences were only significant ($P < 0.05$) when the values of the first day were compared with those of the following days. All the interactions considered in the model (except the day \times milkline interaction, described above) were non-significant.

Table 2. Quality parameters (average value \pm s.e.m.) for goat's milk in commercial farm bulk samples according to the type of milking used (ML: mid-line; LL: low-line)

Variable	Milking type		Sig.
	ML (n = 25)	LL (n = 30)	
FFA (mmol/l)	0.59 ± 0.049	0.58 ± 0.045	N.S.
Fat (g/kg)	62 ± 1.4	60 ± 1.3	N.S.
Protein (g/kg)	41.5 ± 0.96	41.0 ± 0.88	N.S.
Lactose (g/kg)	46.3 ± 0.28	46.0 ± 0.26	N.S.
Dry matter (g/kg)	158 ± 2.2	156 ± 2.0	N.S.
pH	6.74 ± 0.013	6.73 ± 0.012	N.S.
Freezing point ($^{\circ}$ C)	-0.556 ± 0.0026	-0.559 ± 0.0024	N.S.
SCC log	6.19 ± 0.046	6.18 ± 0.042	N.S.
TBC log	4.9 ± 0.12	4.7 ± 0.11	N.S.

FFA, free fatty acid (mmol/l); SCC log, somatic cell count (cell/ml) logarithm; TBC log, total bacterial count (cfu/ml) logarithm; N.S., not significant. Degrees of freedom for milking type are 53.

Table 3. Quality parameters of goat's milk from ML and LL milking used for the production of cured cheese in each of the two 4-day periods of the first experiment

Variable	Period 1		Period 2	
	ML	LL	ML	LL
FFA (mmol/l)	0.76	0.43	0.83	0.47
Fat (g/kg)	45.4	45.3	45.6	47.0
Crude protein (g/kg)	34.6	35.7	34.5	34.9
Lactose (g/kg)	45.4	44.6	45.3	44.3
Dry matter (g/kg)	134.3	132.6	134.3	135.4
SCC ($\times 1000$ cells/ml)	1242	1167	1320	1450
TBC (cfu/ml)	175	160	153	169

FFA, free fatty acids; SCC, somatic cell count; TBC, total bacterial count.

The rest of the milk variables analysed (gross composition, pH-value, freezing point, SCC and TBC) was unaffected by the milking pipeline height (Table 1). The time elapsed since milking only significantly affected ($P < 0.05$) the pH-value, which was higher (0.03) in milk samples taken immediately after milking than in those taken at 24 h post-milking. The storage day factor and all the interactions included in the statistical model did not significantly affect the aforementioned variables.

On the other hand, in the second experiment, carried out under commercial conditions, the level of lipolysis in the milk did not differ significantly between ML and LL milking (Table 2). Moreover, goat's milk from most commercial farms presented an FFA concentration between 0.2 and 0.8 mmol/l, regardless of the type of milking installation used (Fig. 3). Nor did the two groups of farms differ significantly in milk gross composition (fat, protein, lactose and total solids), pH-value, freezing point, SCC or TBC (Table 2). The milkline \times week interaction was also non-significant in all cases.




Fig. 3. Frequency distribution of FFA concentration (mmol/l) in goat milk samples from commercial farms according to the type of milking used (ML: mid-line; LL: low-line).

Table 4. Results of the sensorial analysis triangular test on cheeses made with milk obtained in the first experimental replicate from ML or LL milking, performed at 50 and 100 days of maturation

Triangular test	Ripening time (days)	
	50	100
Judges	62	62
Triads ^a	124	124
Hits	46	48
Sig. Lev.	N.S.	N.S.

^aTwo triads per judge (one in the first experimental period, and another in the second); N.S., non-significant differences.

Sensory analysis of goat's milk cheeses

The characteristics of the goat's milk used in each of the cured cheese manufactures are presented in Table 3 and the results of sensory analysis of the cheese samples in Table 4.

The outcomes show that the judges were not able to perceive significant differences between the two types of cheeses for either of the two maturing times considered (50 and 100 days). On the other hand, when tasters were successful in differentiating the two types of samples, 0.63 of the judgements considered that the cheeses from ML milking presented more intense flavour features (stronger, more acid or spicier) than the cheeses from LL milking, with the latter generally being the main reason for their choice.

Discussion

The average FFA concentration found in the current work for LL and ML milking in the first experiment (0.40 and 0.71 mmol/l, respectively; 0.88 and 1.5 meq/100 g milkfat), and in the second (0.58 and 0.59 mmol/l; 0.96 meq/100 g milkfat), falls within the range reported by other authors (Žan *et al.*, 2006; Strzalkowska *et al.*, 2010; Chilliard *et al.*, 2014) for goat's milk.

The three replicates performed under experimental conditions coincide in demonstrating that ML milking significantly increases the level of lipolysis in goat's raw milk, compared with LL milking, presenting an increase in the FFA concentration of between 62 and 92%. According to Meffe (1994), the height of the milkline

above the animals' standing level can accentuate deterioration of the membrane of the fat globules through two mechanisms. The first is that the rise of the milk through the long milk tube, mixed with air and in a totally turbulent regime, causes an increase in the air–milk interface (the more the higher the air/milk ratio), giving rise to a greater deformation and risk of rupture of the membrane of the fat globules in the said interface. The second mechanism, less important than the previous one, derives from the friction of the milk against the walls of the pipes, subjecting the fat globule to shear forces that can break its membrane. Therefore, as in ML milking installations the length of the long milk tube is usually almost double that in LL (in the current work 215 and 100 cm, respectively), it can be assumed that the cited risk will increase. Moreover, it must be noted that in the three assays performed, an increase in the FFA concentration in the milk after 24 h in the refrigerated storage tank (between 10 and 42% depending on the type of line and assay) was observed, which agrees with the findings of other authors in cows' milk (Wiking *et al.*, 2002; Ouattara *et al.*, 2004).

However, the fact that commercial farms did not show significant differences in the FFA concentration in goat's milk depending on the type of milking installation (ML *v.* LL) suggests that the other factors able to influence lipolysis (physiological, genetic and other features of the milking machine, among others; Deeth 2006; Chilliard *et al.*, 2014) are more important overall than the effect of the milk line height alone. In fact, some ML farms repeatedly had low FFA values (0.2–0.4 mmol/l), whereas other farms using LL always presented FFA concentrations higher than 0.8 mmol/l. Thus, it does not seem that the increase in lipolysis that can occur exclusively due to the effect of ML milking is a decisive argument to discourage this type of milking installation.

In any case, the FFA concentration in milk from most commercial farms considered in this study was lower than the threshold values applied in quality payment systems for goat's milk used in some regions of France (1.77 meq/100 g milkfat; Piriš *et al.*, 2007), and Norway (1.33 meq/l; Skeie 2014), regardless of the type of milking system employed. Thus, 0.97 of goat's milk samples from farms using ML milking presented a FFA concentration lower than the French threshold, more restrictive, whereas for farms with LL milking, this proportion was 0.94.

On the other hand, ML milking had no relevant effect on the analytical results of the different milk components analysed by

infrared equipment, in either experiment one or experiment two, as the differences found were quantitatively of low importance ($\leq 0.02\%$) and did not reach significance in any case. Similarly, in cows' milk, Kaylegian *et al.* (2007) also found changes in fat and protein values no higher than 0.01% when increasing FFA up to 0.2 meq/kg milk.

Regarding cheeses, no information is currently available on the maximum thresholds of lipolysis (FFA) in the goat's milk of indigenous breeds to avoid the deterioration of the sensory characteristics of the flavour of the cheeses, in particular those made by enzymatic coagulation. The organoleptic analysis results suggest that FFA values in milk of 0.79 mmol/l compared with 0.45 mmol/l (1.73 meq/100 g milkfat *v.* 0.97 meq/100 g milkfat) do not alter significantly the sensory characteristics of the enzymatic coagulation cured cheeses. Nevertheless, these FFA levels are sufficient for some consumers to be able to detect some more intense flavour features (i.e. stronger, more acid or spicier) in the cheeses made with milk from ML milking. In any case, these values are far from those reported by other authors as causing off-flavours in goat's cheeses (3.5 meq/100 g milkfat, Morgan *et al.*, 2001). However, it seems prudent not to directly extrapolate these results to the environment of the current work, given the differences in milk composition and the cheese manufacturing process. This topic, therefore, remains open for future studies.

Conclusion

In experimental farm conditions, it was found that ML milking significantly increased the FFA concentration in raw goat's milk compared with LL milking system. However, the results obtained on commercial farms failed to confirm these differences, which points to the existence of other factors (related to the animals, feeding or other conditions of the machine and/or milking routine used) that may have a greater influence on the level of lipolysis of the milk than the mere fact of milking in ML or LL. No differences were found in other milk quality parameters, or were sensory defects in the enzymatic coagulation cheeses perceptible by consumers. There is, therefore, no reason to discourage farmers from this type of milking setup, provided that the correct functioning and management of the milking operation and milk storage on the farm is guaranteed.

Financial support. This research received no specific grant from any funding agency, commercial or not-for-profit sectors.

Conflicts of interest. None.

Ethical standards. Not applicable.

References

- Abeni F, Degano L, Calza F, Giangiacoimo R and Pirlo G (2005) Milk quality and automatic milking: fat globule size, natural creaming and lipolysis. *Journal of Dairy Science* **88**, 3519–3529.
- Chen L, Daniel RM and Coolbear T (2003) Detection and impact of protease and lipase activities in milk and milk powders. *International Dairy Journal* **13**, 255–275.
- Chilliard Y, Selselet-Attou G, Bas P and Morand-Fehr P (1984) Characteristics of lipolytic system in goat milk. *Journal of Dairy Science* **67**, 2216–2223.
- Chilliard Y, Ferlay A, Rouel J and Lamberet G (2003) A review of nutritional and physiological factors affecting goat milk lipid synthesis and lipolysis. *Journal of Dairy Science* **86**, 1751–1770.
- Chilliard Y, Toral PG, Shingfield KJ, Rouel J, Leroux C and Bernard L (2014) Effects of diet and physiological factors on milk fat synthesis, milk fat composition and lipolysis in the goat: a short review. *Small Ruminant Research* **122**, 31–37.
- Collomb M and Spahni M (1995) Revue des méthodes de dosage des acides gras libres dans le lait et les produits laitiers. *LWT – Food Science and Technology* **28**, 355–379.
- Deeth HC (2006) Lipoprotein lipase and lipolysis in milk. *International Dairy Journal* **16**, 555–562.
- Díaz JR, Peris C, Rodríguez M, Molina MP and Fernández N (2004) Effect of milking pipeline height on machine milking efficiency and milk quality in sheep. *Journal of Dairy Science* **87**, 1675–1683.
- Escobar GJ and Bradley RL (1990) Effect of mechanical treatment on the free fatty acid content of raw milk. *Journal of Dairy Science* **73**, 2054–2060.
- Gargouri A, Hamed H and ElFeki A (2008) Total and differential bulk cow milk somatic cell counts and their relation with lipolysis. *Livestock Science* **113**, 274–279.
- Gudding R and Lorentzen P (1982) The influence of low-line and high-line milking plants on udder health and lipolysis. *Nordisk Veterinaermedicin* **34**, 153–157.
- IDF (1991) Routine methods for determination of free fatty acids in milk. In *Determination of Free Fatty Acids in Milk and Milk Products*. Bulletin of the IDF 265. Brussels, Belgium: International Dairy Federation, pp. 26–35.
- IDF (2000) *Whole Milk. Determination of Milkfat, Protein and Lactose Content. Guidance on the Operation of Mid-infrared Instruments (IDF Standard 141C)*. Brussels, Belgium: International Dairy Federation Standard.
- ISO (2004) *Sensory Analysis. Methodology. Triangle Test (ISO 4120:2004)*. Geneva, Switzerland: International Organization for Standardization.
- ISO (2007a) *Milking Machine Installations – Vocabulary (ISO 3918:2007)*. Geneva, Switzerland: International Organization for Standardization.
- ISO (2007b) *Milking Machine Installations – Construction and Performance (ISO 5707:2007)*. Geneva, Switzerland: International Organization for Standardization.
- ISO (2013) *Microbiology of the Food Chain – Horizontal Method for the Enumeration of Microorganisms-Part 1: Colony Count at 30 Degrees C by the Pour Plate Technique (ISO 4831-1:2013)*. Geneva, Switzerland: International Organization for Standardization.
- Judge FJ, Fleming MG, O'Shea J and Raftery TF (1977) Effect of milking pipeline height and excessive air admission at the claw on free fatty acid development in raw milk. *Irish Journal of Agricultural Research* **16**, 115–122.
- Kaylegian KE, Lynch JM, Fleming JR and Barbano DM (2007) Lipolysis and proteolysis of modified and producer milks used for calibration of mid-infrared milk analyzers. *Journal of Dairy Science* **90**, 602–615.
- Le Mens P, Heuchel V, Jubert G, Bodin JP, Sauvageot F and Humbert G (1997) *Caractérisation et Origine des Défauts de Flaveur dans les Fromages de Chèvre. Compte Rendu de Fin d'étude no. 9983110 - Prog. Aliment.* Paris, France: Institut de l'Élevage.
- Littell RC, Henry PR and Ammerman CB (1998). Statistical analysis of repeated measures data, using SAS procedures. *Journal of Animal Science* **76**, 1216–1231.
- Meffe N (1994) La lipolyse dans le lait du vache: Bien en comprendre les mécanismes et les causes pour mieux la prévenir. *Recueil de Médecin Vétérinaire, Spécial-Qualité du Lait* **170**, 399–409.
- Mikulová M (2011) Content of free fatty acids, lipolytic bacteria and somatic cells in relation to milking technology. *Journal of Agrobiology* **28**, 49–54.
- Ministerio de la Presidencia (2011) Real Decreto 752/2011, de 27 de mayo, por el que se establece la normativa básica de control que deben cumplir los agentes del sector de leche cruda de oveja y cabra, BOE-A-2011-9995. *Boletín Oficial del Estado* **137**, 58609–58631.
- Morand-Fehr P, Selselet Attou G, Bas P and Chilliard Y (1983) Factors in milking conducive to lipolysis in goats' milk. In *Proceedings of the 3rd International Symposium on Machine Milking of Small Ruminants (Eds Server-Cuesta)*. Valladolid, Spain, pp. 276–284.
- Morgan F, Bodin JP and Gaborit P (2001) Lien entre le niveau de lipolyse du lait de chèvre et la qualité sensorielle des fromages au lait cru ou pasteurisés. *Le Lait* **81**, 743–756.

- Ouattara GC, Jeon IJ, Hart-Thakur RA and Schmidt KA** (2004) Fatty acids released from milk fat by lipoprotein lipase and lipolytic psychrotrophs. *Journal of Food Science* **69**, C659–C664.
- Pillay VT, Myhr AN, Gray JI and Biggs DA** (1980) Lipolysis in milk. II. Effect of milking systems. *Journal of Dairy Science* **63**, 1219–1223.
- Pirisi A, Lauret A and Dubeuf JP** (2007) Basic and incentive payments for goat and sheep milk in relation to quality. *Small Ruminant Research* **68**, 167–178.
- Rasmussen MD, Wiking L, Bjerring M and Larsen HC** (2006) Influence of air intake on the concentration of free fatty acids and vacuum fluctuations during automatic milking. *Journal of Dairy Science* **89**, 4596–4605.
- Robertson NH, Dixon A, Nowers JH and Brink DPS** (1981) The influence of lipolysis, pH and homogenization on infrared readings for fat, protein and lactose. *South African Journal of Dairy Technology* **13**, 3–7.
- Skeie SB** (2014) Quality aspects of goat milk for cheese production in Norway: a review. *Small Ruminant Research* **122**, 10–17.
- Strzalkowska N, Józwik A, Bagnicka E, Krzyżewski J, Horbańczuk K, Pyzel B, Sloniewska D and Horbańczuk JO** (2010) The concentration of free fatty acids in goat milk as related to the stage of lactation, age and somatic cell count. *Animal Science Papers and Reports* **28**, 389–395.
- Wiking L, Frøst MB, Larsen LB and Nielsen JH** (2002) Effects of storage conditions on lipolysis, proteolysis and sensory attributes in high quality raw milk. *Milchwissenschaft* **57**, 190–194.
- Wiking L, Björk L and Nielsen JH** (2003) Influence of feed composition on stability of fat globules during pumping of raw milk. *International Dairy Journal* **13**, 797–804.
- Žan M, Stibilj V and Rogelj I** (2006) Milk fatty acid composition of goats grazing on alpine pasture. *Small Ruminant Research* **64**, 45–52.