

ARTICLE

Central limit theorem for components in meandric systems through high moments

Svante Janson¹  and Paul Thévenin²

¹Department of Mathematics, Uppsala University, Uppsala, Sweden and ²Institut für Mathematik, University of Vienna, Vienna, Austria

Corresponding author: Paul Thévenin; Email: paul.thevenin@univie.acat

(Received 24 March 2023; revised 13 February 2024; accepted 25 March 2024; first published online 29 April 2024)

Abstract

We investigate here the behaviour of a large typical meandric system, proving a central limit theorem for the number of components of a given shape. Our main tool is a theorem of Gao and Wormald that allows us to deduce a central limit theorem from the asymptotics of large moments of our quantities of interest.

Keywords: Probability; meandric systems; central limit theorem

2020 MSC Codes: Primary: 60C05

1. Model and main result

1.1 Definitions and some notation

Let $n \geq 1$ be an integer. A *meandric system* of size n is a collection of non-crossing loops in the plane that intersect the horizontal axis exactly at the points $[2n] := \{1, \dots, 2n\}$. We call these points the *vertices* of the meandric system; two meandric systems that differ only by a continuous deformation of the plane that fixes the horizontal axis are regarded as the same. Meandric systems were introduced, to our knowledge, by Di Francesco, Golinelli, and Guitter [1] and have recently become again a topic of interest [2, 4, 6]. A meandric system can be regarded as a set of n non-crossing arcs with endpoints $[2n]$ in the upper half-plane and another such set in the lower half-plane; a meandric system thus determines two non-crossing matchings (pair-partitions) of $[2n]$, one for each half-plane, and it is easily seen that this yields a bijection between meandric systems of size n and pairs of two non-crossing matchings of $[2n]$. In particular, since the number of non-crossing matchings of $[2n]$ is the Catalan number,

$$\text{Cat}_n := \frac{(2n)!}{n!(n+1)!}, \quad (1.1)$$

see, for example, [7, item 61], the number of meandric systems of size n is Cat_n^2 .

Each connected component of a meandric system is a single loop, intersecting the horizontal axis in a subset of $[2n]$, say $\{i_1 < \dots < i_{2k}\}$, which we call the *support* of the loop. Note that necessarily, there is an even number of vertices in the support and an even number of integers in each gap (i_j, i_{j+1}) , that is, $i_{j+1} - i_j$ is odd for $1 \leq j < 2k$. We say that two such loops have the same shape

Supported by the Knut and Alice Wallenberg Foundation.

if they differ only by a translation. Thus, we may normalise each shape to have leftmost vertex 1 and make the following formal definition:

Definition 1.1. A shape is a (connected) non-crossing loop S whose support is a set of integers $\{i_1 = 1 < i_2 < \dots < i_{2k} = 2\ell\}$, for some $k, \ell \geq 1$, such that $i_{j+1} - i_j$ is odd for all $1 \leq j \leq 2k - 1$.

Let M be a meandric system and C be a connected component of M . We say that C has shape S if C and S differ only by a translation.

Our main theorem is the following. We prove two special cases as Theorems 3.1 and 4.4 and prove the remaining, more difficult, case in Section 4.2. In the paper, $\xrightarrow{(d)}$ and $\xrightarrow{\mathbb{P}}$ denote, respectively, convergence in distribution and convergence in probability.

Theorem 1.2. Fix a shape S . Let M_n be a uniformly random meandric system of size n (i.e. on $\llbracket 1, 2n \rrbracket$) and denote by $X_{S,n}$ the number of connected components of M_n with shape S . Then, $X_{S,n}$ satisfies a central limit theorem: there exist $\mu_S, \sigma_S > 0$ such that

$$\frac{X_{S,n} - n\mu_S}{\sigma_S\sqrt{n}} \xrightarrow[n \rightarrow \infty]{(d)} \mathcal{N}(0, 1), \tag{1.2}$$

where $\mathcal{N}(0, 1)$ denotes the standard normal distribution.

Observe that the convergence $\frac{X_{S,n}}{n} \xrightarrow[n \rightarrow \infty]{\mathbb{P}} \mu_S$ for some constant μ_S was already obtained in ref. [4], with an explicit expression for μ_S .

2. Preliminaries

2.1 More notation

For integers $m \leq n$, $\llbracket m, n \rrbracket$ denotes the integer interval $[m, n] \cap \mathbb{Z}$. The size of $\llbracket m, n \rrbracket$ is its number of points, that is, $n - m + 1$. Note that $\llbracket n \rrbracket = \llbracket 1, n \rrbracket$.

For a component C of a meandric system, we denote by L_C (R_C), the leftmost (rightmost) point in the support of C . Furthermore, we say that the base of C is the interval $\llbracket L_C, R_C \rrbracket$ and let $\ell(C)$ denote the half-length of C , defined as half the size of its base, that is, $\ell(C) := \frac{1}{2}(R_C - L_C + 1)$. (Note that $\ell(C)$ always is an integer.) We use the same definitions for a shape S ; then $L_S = 1$, and thus $R_S = 2\ell(S)$.

For integers $N \geq k \geq 0$, we let

$$(N)_k := N(N - 1) \cdots (N - k + 1) = \frac{N!}{(N - k)!} = k! \binom{N}{k}, \tag{2.1}$$

the k -th descending factorial of N .

We use standard o and O notation. Furthermore, for two (positive) sequences a_n and b_n , $a_n \sim b_n$ means $a_n/b_n \rightarrow 1$ as $n \rightarrow \infty$, that is, $a_n = b_n(1 + o(1))$, and $a_n = \Theta(b_n)$ means that there exist constants $c > 0$ and C such that $c \leq a_n/b_n \leq C$ for sufficiently large n . Note that, for example, $a_{n,r} \sim b_{n,r}$ for $r = O(\sqrt{n})$ means that this holds for every sequence $r = r(n) = O(\sqrt{n})$, which is equivalent to $a_{n,r} \sim b_{n,r}$ uniformly for $r \leq C\sqrt{n}$, for any $C < \infty$; uniformity in r is thus automatic in such cases. We write ‘uniformly for $r = O(\sqrt{n})$ ’ for ‘uniformly for $r \leq C\sqrt{n}$, for any $C < \infty$ ’. Unspecified limits are as $n \rightarrow \infty$.

2.2 The key tool: Gao and Wormald’s theorem

Our proof relies on a theorem due to Gao and Wormald [5], stating that we can deduce a central limit theorem for a sequence of variables from the asymptotic behaviour of their high (factorial) moments. Let us recall this result.

Theorem 2.1 (Gao & Wormald [5]). Let $\mu_n s_n > -1$ and set $\sigma_n := \sqrt{\mu_n + \mu_n^2 s_n}$, where $0 < \mu_n \rightarrow \infty$. Suppose that $\sigma_n = o(\mu_n)$, $\mu_n = o(\sigma_n^3)$, and that a sequence $\{X_n\}$ of nonnegative random variables satisfies as $n \rightarrow \infty$:

$$\mathbb{E} [(X_n)_r] \sim \mu_n^r \exp\left(\frac{r^2 s_n}{2}\right). \tag{2.2}$$

uniformly for all integers r in the range $c\mu_n/\sigma_n \leq r \leq C\mu_n/\sigma_n$, for some constants $C > c > 0$. Then $(X_n - \mu_n)/\sigma_n$ converges in distribution to a standard normal variable as $n \rightarrow \infty$.

In other words, if high factorial moments of a variable asymptotically match those of a normal distribution, then convergence to the normal distribution holds.

2.3 Some lemmas

We state some simple lemmas that will be used later. The first is a well-known estimate that we often will use in the sequel.

Lemma 2.2.

1. If $0 \leq k \leq n/2$, then

$$(n)_k = n^k \exp\left(-\frac{k^2}{2n} + O\left(\frac{k^3}{n^2} + \frac{k}{n}\right)\right). \tag{2.3}$$

2. In particular, if $k = O(\sqrt{n})$, then

$$(n)_k = n^k \exp\left(-\frac{k^2}{2n} + o(1)\right) \sim n^k \exp\left(-\frac{k^2}{2n}\right). \tag{2.4}$$

3. More generally, if $0 \leq k \leq m$ with $m = O(\sqrt{n})$, then

$$(n - m + k)_k \sim n^k \exp\left(-\frac{m^2 - (m - k)^2}{2n}\right) = n^k \exp\left(-\frac{k(2m - k)}{2n}\right). \tag{2.5}$$

Proof. (i), (ii): This follows easily from a Taylor expansion of $\log(1 - i/n)$ for $0 \leq i < k$; we omit the details.

(iii): This follows from (ii) and $(n - m + k)_k = (n)_m / (n)_{m-k}$. □

As one consequence, we obtain the following asymptotics.

Lemma 2.3. Let $n \rightarrow \infty$ and $0 \leq r = O(\sqrt{n})$. Then

$$\frac{\text{Cat}_{n-r}}{\text{Cat}_n} \sim 2^{-2r}. \tag{2.6}$$

Proof. The definition (1.1) and Lemma 2.2 yield

$$\frac{\text{Cat}_{n-r}}{\text{Cat}_n} = \frac{(n)_r (n+1)_r}{(2n)_{2r}} \sim \frac{(n)_r^2}{(2n)_{2r}} = \frac{n^{2r}}{(2n)^{2r}} \exp\left(-2\frac{r^2}{2n} + \frac{(2r)^2}{4n} + o(1)\right) \sim 2^{-2r}. \tag{2.7}$$

□

We end this section with another elementary and well-known result.


Lemma 2.4. Let $m, n, k \geq 1$. The number of unordered k -tuples of disjoint intervals of size m in $[n]$ is given by

$$\binom{n - k(m - 1)}{k}. \tag{2.8}$$

Proof. By deleting all points except the leftmost in each chosen interval, we obtain a bijection between the set of such k -tuples of intervals and the set of k -tuples of distinct points in $[n - k(m - 1)]$. \square

3. The first example: components of half-length 1

As a warm-up, we consider first the simple case where S is the loop of half-length 1. For any $i \in [2n]$, we let Y_i be the indicator that the following holds:



Then,

$$X_{S,n} = \sum_{i=1}^{2n-1} Y_i \tag{3.1}$$

and thus, for every $r \geq 1$, summing over $1 \leq i_1 < \dots < i_r < 2n$,

$$\mathbb{E} [(X_{S,n})_r] = \mathbb{E} \left[r! \sum_{i_1 < \dots < i_r} Y_{i_1} \cdots Y_{i_r} \right] = r! \sum_{i_1 < \dots < i_r} \mathbb{E} [Y_{i_1} \cdots Y_{i_r}]. \tag{3.2}$$

The expectation in the last sum is non-zero if and only if the r subintervals $[[i_j, i_j + 1]]$ of $[[1, 2n]]$ are disjoint, so by Lemma 2.4, there are $\binom{2n-r}{r}$ non-zero terms. Each of the non-zero terms is $1/\text{Cat}_n^2$ times the number of meandric systems of size n that contain r given loops of half-length 1; by deleting these loops (and the vertices in them), we obtain a bijection between such meandric systems and the meandric systems of size $n - r$, and hence the number of them is Cat_{n-r}^2 . Consequently, (3.2) yields

$$\mathbb{E} [(X_{S,n})_r] = (2n - r)_r \frac{\text{Cat}_{n-r}^2}{\text{Cat}_n^2} = \frac{(2n)_{2r}}{(2n)_r} \cdot \frac{\text{Cat}_{n-r}^2}{\text{Cat}_n^2}. \tag{3.3}$$

In particular, using Lemmas 2.2 and 2.3, if $r = O(\sqrt{n})$, then

$$\mathbb{E} [(X_{S,n})_r] \sim (2n)^{2r-r} \exp\left(-\frac{4r^2}{4n} + \frac{r^2}{4n}\right) 2^{-4r} = \left(\frac{n}{8}\right)^r \exp\left(-\frac{3r^2}{4n}\right). \tag{3.4}$$

In other words, (2.2) holds (uniformly) for $0 \leq r \leq C\sqrt{n}$, for any fixed $C < \infty$, with

$$\mu_n := \frac{n}{8}, \tag{3.5}$$

$$s_n := -\frac{3}{2n}. \tag{3.6}$$

We have $\mu_n s_n = -3/16 > -1$, and thus

$$\sigma_n := \sqrt{\mu_n(1 + \mu_n s_n)} = \sqrt{\frac{13}{128}n}. \tag{3.7}$$

We thus have $\sigma_n = o(\mu_n)$ and $\mu_n = o(\sigma_n^3)$, and consequently Theorem 2.1 applies and yields:

Theorem 3.1. *If S is a simple loop of half-length 1, then*

$$\frac{X_{S,n} - n/8}{\sqrt{13n/128}} \xrightarrow[n \rightarrow \infty]{(d)} \mathcal{N}(0, 1). \tag{3.8}$$

This is Theorem 1.2 for this particular choice of S , with $\mu_S = 1/8$ and $\sigma_S^2 = 13/128$.




Figure 1. A component C with four bounded faces F_1, F_2, F_3, F_4 . In this example, we have $K(S) = \text{Cat}_1^2 \text{Cat}_2 \text{Cat}_3 = 10$, $c_+(S) = 1$, and $c_-(S) = 0$, where S is the shape of C .

4. Extension to any fixed shape

Let us now show how we can extend this result to any fixed shape S . We now let Y_i be the indicator that there is a component C of shape S such that $L_C = i$; note that (3.1) and (3.2) still hold.

Recall that $\ell(S)$ is the half-length of S , so S has base $\llbracket 1, 2\ell(S) \rrbracket$. We also define here three other constants $K(S), c_+(S), c_-(S)$ depending on S . To avoid heavy notation, we will drop the argument S in what follows and only denote them by K, c_+, c_- .

Definition 4.1. (See an example in Figure 1) Observe that a component C of shape S , taken along with the horizontal axis, splits the plane into two unbounded faces, each belonging to one of the half-planes, and a certain number of bounded faces. Let F_+ denote the unbounded face in the upper half-plane, F_- the one in the lower half-plane, and $\mathcal{F}(C)$ the set of bounded faces. For a face F , let $v(F)$ be the number of vertices in $\llbracket L_C, R_C \rrbracket$ that lie on the boundary of F but not on C , and observe that necessarily $v(F)$ is even. We then set

$$K(S) := \prod_{F \in \mathcal{F}(C)} \text{Cat}_{v(F)/2}, \tag{4.1}$$

$$c_+(S) := v(F_+)/2, \tag{4.2}$$

$$c_-(S) := v(F_-)/2. \tag{4.3}$$

Note that these constants do not depend on the set of vertices on which C is defined, but only on its shape S .

4.1 Strong shapes

We say that two components *overlap* if their bases overlap. Hence, if the components have the same shape S , and the leftmost points in their supports are i and j , they overlap if $|j - i| < 2\ell(S)$.

For simplicity, we study first the case when this cannot happen. We say that a shape S is *strong* if two different components of a meandric system that both have shape S cannot overlap. Thus, if S is strong, then $Y_i Y_j = 0$ when $|j - i| < 2\ell(S)$. The simple loop in Section 3 and the loop in Fig. 1 are examples of strong shapes. A shape that is not strong is called *weak*; an example is given in Fig. 2.

Proposition 4.2. Let S be a strong shape of half-length $\ell(S)$. Then, for all $r \geq 1$, we have

$$\mathbb{E} [(X_{S,n})_r] = (2n - 2r\ell(S) + r)_r K^r \frac{\text{Cat}_{n-r\ell(S)+rc_+} \text{Cat}_{n-r\ell(S)+rc_-}}{\text{Cat}_n^2}. \tag{4.4}$$




Figure 2. Two components of same shape overlapping. Here, $\mathbb{E}[Y_1 Y_7] > 0$, while $2\ell(S) = 10$.

Proof. We argue as in Section 3. As noted above, (3.2) still holds, and since S is strong, we have $Y_i Y_j = 0$ when $|j - i| < 2\ell(S)$. Hence, the number of non-zero terms in (3.2) is $\binom{2n-r(2\ell(S)-1)}{r}$ by Lemma 2.4. Again, all non-zero terms have the same value, which is $1/\text{Cat}_n^2$ times the number of ways that r given disjoint loops of shape S can be completed to a meandric system of size n . We can fill in the bounded faces of each component in K ways, and there are $2n - 2r\ell(S) + 2rc_{\pm}$ vertices left in the upper and lower components, respectively, so they may be filled in $\text{Cat}_{n-r\ell(S)+rc_{\pm}}$ ways. This yields (4.4). \square

By Lemmas 2.2 and 2.3, it follows from (4.4) that, (uniformly) for $r = O(\sqrt{n})$, we have

$$\mathbb{E}[(X_{S,n})_r] \underset{n \rightarrow \infty}{\sim} \left(\frac{2nK}{4^{2\ell(S)-c_+-c_-}} \right)^r \exp\left(-\frac{r^2}{4n} [(2\ell(S))^2 - (2\ell(S) - 1)^2] \right). \tag{4.5}$$

This is (2.2) with

$$\mu_n := \frac{2nK}{4^{2\ell(S)-c_+-c_-}}, \tag{4.6}$$

$$s_n := -\frac{(2\ell(S))^2 - (2\ell(S) - 1)^2}{2n} = -\frac{4\ell(S) - 1}{2n}. \tag{4.7}$$

In order to apply Theorem 2.1, we need to check that $\mu_n s_n > -1$, which boils down to the following.

Lemma 4.3. *We have*

$$K(4\ell(S) - 1) < 4^{2\ell(S)-c_+-c_-}. \tag{4.8}$$

Proof. Observe that we can bound K using the fact that $\text{Cat}_n \leq \frac{4^n}{n+1}$ for all n . It is easy to see that for given c_{\pm} , out of all possible choices of components with these values of c_{\pm} , K is largest if there is only one bounded face in each half-plane, and thus,

$$K \leq \text{Cat}_{\ell(S)-c_+-1} \text{Cat}_{\ell(S)-c_- - 1} \leq \frac{4^{2\ell(S)-c_+-c_- - 2}}{(\ell(S) - c_+)(\ell(S) - c_-)} \leq \frac{4^{2\ell(S)-c_+-c_- - 2}}{\ell(S)}, \tag{4.9}$$

since $c_+ + c_- \leq \ell(S) - 1$ (to see this, observe that a vertex cannot belong to both unbounded faces of S and that at least two vertices belong to C). This yields (4.8) directly. \square

It is clear that $\mu_n \rightarrow \infty$. Furthermore, we have just proved that $1 + \mu_n s_n$ is a positive constant. Thus $\sigma_n = \Theta(\sqrt{\mu_n})$, and hence $\sigma_n = o(\mu_n)$ and $\mu_n = o(\sigma_n^3)$. We can therefore apply Theorem 2.1 to obtain the central limit theorem in this case too:

Theorem 4.4. *Let S be a strong shape. Then*

$$\frac{X_{S,n} - n\mu_S}{\sigma_S\sqrt{n}} \xrightarrow[n \rightarrow \infty]{(d)} \mathcal{N}(0, 1), \tag{4.10}$$

where

$$\mu_S = \frac{2K}{4^{2\ell(S)-c_+-c_-}} \quad \text{and} \quad \sigma_S = \sqrt{\frac{2K}{4^{2\ell(S)-c_+-c_-}} \left(1 - \frac{K(4\ell(S) - 1)}{4^{2\ell(S)-c_+-c_-}}\right)}. \tag{4.11}$$

This proves Theorem 1.2 in the case when S is a strong shape, with explicit formulas for μ_S and σ_S .

4.2 Weak shapes

Finally, we study the case of a weak shape S . Thus, now there may be overlaps between two components of shape S , that is, two indices $i < j$ such that $|j - i| < 2\ell(S)$ and $Y_i Y_j = 1$, where Y_i is defined as before. See Fig. 2 for an example.

Let A^r be the set of all r -tuples $E := \{i_1, \dots, i_r\}$ with $1 \leq i_1 < \dots < i_r \leq 2n$. For any such r -tuple E , define an equivalence relation \sim_E on $\{1, \dots, r\}$ as the smallest one (for the inclusion of the equivalence classes) satisfying: for all $1 \leq k_1, k_2 \leq r$ such that $|i_{k_1} - i_{k_2}| < 2\ell(S)$, $k_1 \sim_E k_2$. We call the equivalence classes of \sim_E blocks. Furthermore, for $1 \leq j \leq r$, we let A_j^r be the set of r -tuples $E \in A^r$ that have exactly j blocks. Thus $A^r = \bigsqcup_{j=1}^r A_j^r$. Note that A_r^r is the set of r -tuples E such that all blocks are singletons. An r -tuple E corresponds to a collection $(C_k)_1^r$ of loops of shape S , shifted such that C_k has $L_{C_k} = i_k$. In particular, $E \in A_r^r$ if and only if these loops are non-overlapping.

Define, for all $1 \leq u \leq r$:

$$F_u := \binom{2n - 2u\ell(S) + u}{u} K^u \frac{\text{Cat}_{n-u\ell(S)+uc_+} \text{Cat}_{n-u\ell(S)+uc_-}}{\text{Cat}_n^2}. \tag{4.12}$$

By the argument in the proof of Proposition 4.2, $u! F_u$ is the contribution to $\mathbb{E} [(X_{S,n})_u]$ from u -tuples of non-overlapping components.

We have the following estimates:

Lemma 4.5. *Let S be a weak shape.*

(i) *For all $r \geq 1$,*

$$\mathbb{E} [(X_{S,n})_r] \geq r! F_r. \tag{4.13}$$

(ii) *For all $1 \leq u \leq r$,*

$$\sum_{E \in A_u^r} \mathbb{E} \left[\prod_{i \in E} Y_i \right] \leq \binom{r-1}{u-1} (2\ell(S))^{r-u} F_u. \tag{4.14}$$

(iii) *For each fixed $M \geq 0$, uniformly for $r = O(\sqrt{n})$ with $r \geq 2M$,*

$$\sum_{E \in A_{r-M}^r} \mathbb{E} \left[\prod_{i \in E} Y_i \right] = \Theta(r^M F_{r-M}) \tag{4.15}$$

and, if also $r \rightarrow \infty$,

$$\sum_{E \in A_{r-M}^r(1,2)} \mathbb{E} \left[\prod_{i \in E} Y_i \right] = (1 - o(1)) \sum_{E \in A_{r-M}^r} \mathbb{E} \left[\prod_{i \in E} Y_i \right], \tag{4.16}$$

where $A_{r-M}^r(1, 2)$ is the subset of A_{r-M}^r made only of blocks of sizes 1 or 2.

Proof of Lemma 4.5. (i): We rewrite (3.2) as

$$\mathbb{E} [(X_{S,n})_r] = r! \sum_{E \in A^r} \mathbb{E} \left[\prod_{i \in E} Y_i \right] = r! \sum_{u=1}^r \sum_{E \in A_u^r} \mathbb{E} \left[\prod_{i \in E} Y_i \right]. \tag{4.17}$$

The term with $u = r$ yields the contribution from r -tuples of non-overlapping components, which as noted after (4.12) is $r!F_r$.

(ii): For each r -tuple $E \in A_u^r$, keep in the product only the leftmost point of each block, observing that, for any sets $A \subseteq B \subseteq \llbracket 1, 2n \rrbracket$, we have $\mathbb{E} [\prod_{i \in B} Y_i] \leq \mathbb{E} [\prod_{i \in A} Y_i]$. Note that this set of leftmost points belongs to A_u^u . If the size of the i -th leftmost block is j_i , then for each set of leftmost points, the number of possible positions of the other $j_i - 1$ points in the block is at most $(2\ell(S))^{j_i-1}$, since each point after the first is within $2\ell(S)$ of the preceding one. Hence,

$$\sum_{E \in A_u^r} \mathbb{E} \left[\prod_{i \in E} Y_i \right] \leq \sum_{\substack{j_1 + \dots + j_u = r \\ j_1, \dots, j_u \geq 1}} \prod_{i=1}^u (2\ell(S))^{j_i-1} \cdot \sum_{E' \in A_u^u} \mathbb{E} \left[\prod_{i \in E'} Y_i \right] = \sum_{\substack{j_1 + \dots + j_u = r \\ j_1, \dots, j_u \geq 1}} \prod_{i=1}^u (2\ell(S))^{j_i-1} \cdot F_u. \tag{4.18}$$

Finally, this yields (4.14), since the number of allowed sequences (j_1, \dots, j_u) is $\binom{r-1}{u-1}$ and $\prod_{i=1}^u (2\ell(S))^{j_i-1} = (2\ell(S))^{r-u}$ for all of them.

(iii): We partition the set A_{r-M}^r as follows. Consider an $(r - M)$ -tuple $T := (T_1, \dots, T_{r-M})$ of integers ≥ 1 , of sum r , and consider also a function J which, to each $1 \leq i \leq r - M$, associates a T_i -tuple J_i of integers $1 =: j_{i,1} < j_{i,2} < \dots < j_{i,T_i}$ such that, for all $1 \leq k \leq T_i - 1$, $j_{i,k+1} - j_{i,k} < 2\ell(S)$, and, furthermore, the T_i loops of shape S that start at the vertices $j_{i,k}$ ($k = 1, \dots, T_i$) are disjoint so that they may occur together as components in a meandric system. (We call such pairs (T, J) *admissible*.) Denote by $A_{T,J}$ the subset of A_{r-M}^r made of r -tuples E such that the i -th leftmost block of E has size T_i , and if this block is $\{a_i^1, \dots, a_i^{T_i}\}$, then we have $a_i^{k+1} - a_i^k = j_{i,k+1} - j_{i,k}$ for all $1 \leq k \leq T_i - 1$. In other words, $A_{T,J}$ accounts for all r -tuples of components with $r - M$ blocks, where the sizes of the blocks are given, as well as the intervals between the starting points of each component of shape S in each block. Hence, A_{r-M}^r is the union $\bigcup A_{T,J}$ over all admissible pairs (T, J) .

Since we only consider $(r - M)$ -tuples T such that

$$r = \sum_{i=1}^{r-M} T_i = r - M + \sum_{i=1}^{r-M} (T_i - 1), \tag{4.19}$$

there at most M indices i with $T_i > 1$ and thus at least $r - 2M$ indices with $T_i = 1$. Note also that if $T_i = 1$, then trivially, $J_i = (1)$. Given an admissible pair (T, J) , we define the reduced pair $(\widehat{T}, \widehat{J})$ by deleting all T_i and J_i such that $T_i = 1$ from T and J ; thus $\widehat{T} := (T_i : 1 \leq i \leq r - M \text{ and } T_i > 1)$ and similarly for \widehat{J} . Consequently, \widehat{T} and \widehat{J} are both sequences of (the same) length $\leq M$. Since (4.19) implies that their entries are bounded (for a fixed M), there is only a finite set \mathcal{T} of reduced pairs $(\widehat{T}, \widehat{J})$, where \mathcal{T} depends on M and S but not on r .

Conversely, given an admissible reduced pair $(\widehat{T}, \widehat{J})$, with $\widehat{T} = (\widehat{T}_1, \dots, \widehat{T}_k)$, we can obtain $(\widehat{T}, \widehat{J})$ from $\binom{r-M}{k}$ different (admissible) pairs (T, J) . Note that here, by (4.19), since each $\widehat{T}_i \geq 2$,

$$k \leq \sum_{i=1}^k (\widehat{T}_i - 1) = \sum_{i=1}^{r-M} (T_i - 1) = M, \tag{4.20}$$

with equality if and only if $\widehat{T}_i = 2$ for all $i \leq k$.

We now want to understand the behaviour of $\sum_{E \in A_{T,J}} \mathbb{E} \left[\prod_{i \in E} Y_i \right]$ for an admissible pair (T, J) . In a way similar to Proposition 4.2 (using an extension of Lemma 2.4 to intervals of different lengths), we obtain

$$\sum_{E \in A_{T,J}} \mathbb{E} \left[\prod_{i \in E} Y_i \right] = \binom{2n - 2\tilde{\ell} + (r - M)}{r - M} \tilde{K} \frac{\text{Cat}_{n-d_+} \text{Cat}_{n-d_-}}{\text{Cat}_n^2}, \tag{4.21}$$

where, for any $E \in A_{T,J}$, $\tilde{\ell}$ is the sum of the half-lengths of the blocks, \tilde{K} accounts for the bounded faces defined by the horizontal axis and the loops defined by E , and d_+, d_- for the unbounded faces. (Note that these constants are the same for all $E \in A_{T,J}$, so they depend only on T and J .) Moreover, since at least $r - 2M$ of these blocks are singletons, and the remaining blocks are determined by \widehat{T} and \widehat{J} , we can write

$$\tilde{K} = K^{r-2M} K', \tag{4.22}$$

for some $K' > 0$ depending only on $(\widehat{T}, \widehat{J})$. Similarly,

$$\tilde{\ell} = (r - M)\ell(S) + \ell', \tag{4.23}$$

$$d_+ = (r - M)(\ell(S) - c_+) + e_+, \tag{4.24}$$

$$d_- = (r - M)(\ell(S) - c_-) + e_- \tag{4.25}$$

for some ℓ', e_+, e_- depending only on $(\widehat{T}, \widehat{J})$. In particular, for a fixed M , it follows that K', ℓ', e_+, e_- can only take a fixed number of values independently of n and r .

We compare (4.21) and F_{r-M} given by (4.12). First, by Lemma 2.2(iii),

$$\begin{aligned} \frac{\binom{2n-2\tilde{\ell}+(r-M)}{r-M}}{\binom{2n-2(r-M)\ell(S)+(r-M)}{r-M}} &= \frac{(2n - 2\tilde{\ell} + (r - M))_{r-M}}{(2n - 2(r - M)\ell(S) + (r - M))_{r-M}} \\ &\sim \exp\left(-\frac{r - M}{4n}((4\tilde{\ell} - r + M) - (4(r - M)\ell(S) - r + M))\right) = \exp(o(1)), \end{aligned} \tag{4.26}$$

since $\tilde{\ell} = r\ell(S) + O(1)$ by (4.23) and $r = o(n)$. Similarly, as a consequence of Lemma 2.3 and (4.24)–(4.25),

$$\frac{\text{Cat}_{n-d_{\pm}}}{\text{Cat}_{n-(r-M)(\ell(S)-c_{\pm})}} \sim 4^{-d_{\pm}+(r-M)(\ell(S)-c_{\pm})} = 4^{-e_{\pm}}. \tag{4.27}$$

Consequently, using also (4.22), we obtain from (4.21) and (4.12),

$$\frac{\sum_{E \in A_{T,J}} \mathbb{E} \left[\prod_{i \in E} Y_i \right]}{F_{r-M}} = C_{T,J}(1 + o(1)), \tag{4.28}$$

where $C_{T,J} > 0$ only depends on $(\widehat{T}, \widehat{J})$ and therefore only takes a finite number of values. In particular,

$$\sum_{E \in A_{T,J}} \mathbb{E} \left[\prod_{i \in E} Y_i \right] = \Theta(F_{r-M}), \tag{4.29}$$

and this holds uniformly for $r = O(\sqrt{n})$ and all admissible (T, J) .

By (4.20) and the discussion before it, there are $\binom{r-M}{k} = \Theta(r^k)$ admissible pairs (T, J) for each $(\widehat{T}, \widehat{J})$, where $k \leq M$, with equality when all $\widehat{T}_i = 2$. Note that since we assume that the shape S is weak, there exists at least one such admissible $(\widehat{T}, \widehat{J})$ with $\widehat{T} = (2, \dots, 2)$. Hence, summing (4.29) over all (T, J) yields (4.15).

Moreover, $A_{r-M}^r \setminus A_{r-M}^r(1, 2)$ is the union $\bigcup' A_{T,J}$ where we only sum over admissible pairs (T, J) with some $T_i \geq 3$; these correspond to reduced pairs $(\widehat{T}, \widehat{J})$ with some $\widehat{T}_i \geq 3$, and we see

from (4.20) that each such reduced pair has length $\leq M - 1$ and thus corresponds to $O(r^{M-1})$ admissible pairs. Consequently, summing (4.29) over all (T, J) of this type yields

$$\sum_{E \in A_{r-M}^r \setminus A_{r-M}^r(1,2)} \mathbb{E} \left[\prod_{i \in E} Y_i \right] = O(r^{M-1} F_{r-M}) = o(r^M F_{r-M}), \tag{4.30}$$

which yields (4.16) by (4.15). □

The next proposition shows that, in order to get the asymptotic behaviour of $\mathbb{E} [(X_{S,n})_r]$, we only need to take into account the configurations whose number of blocks that are not singletons is a given constant.

Proposition 4.6. *Fix a weak shape S . Then, there exists $\eta > 0$ such that, for any $\varepsilon > 0$, there exists $M > 0$ such that we have, uniformly for $r \leq \eta\sqrt{n}$,*

$$\sum_{u \leq r-M} \sum_{E \in A_u^r} \mathbb{E} \left[\prod_{i \in E} Y_i \right] \leq \varepsilon F_r \leq \varepsilon \frac{1}{r!} \mathbb{E} [(X_{S,n})_r]. \tag{4.31}$$

Remark 4.7. *For convenience, we assume here that r/\sqrt{n} is small. In fact, Proposition 4.6 can easily be extended to $r \leq C\sqrt{n}$ for any C (with M depending on C and ε), but we do not need for this.*

To prove this, we start with a lemma:

Lemma 4.8. *There exists $Q > 0$ depending only on the shape S such that, for n large enough, for all $u \leq \sqrt{n}$:*

$$\frac{F_{u+1}}{F_u} \geq Q \frac{n}{u}. \tag{4.32}$$

Proof. We just compute the ratio term by term, recalling (4.12). We have $\frac{K^{u+1}}{K^u} = K$. The ratio of the ratios of Catalan numbers converges uniformly to a positive constant. Finally, the ratio of binomial coefficients is, using Lemma 2.2(ii),

$$\frac{u!}{(u+1)!} \cdot \frac{(2n - (u+1)(2\ell(S) - 1))_{u+1}}{(2n - u(2\ell(S) - 1))_u} = \frac{1}{u+1} \cdot \frac{(2n)^{u+1}}{(2n)^u} \exp(O(1)) \geq c \frac{n}{u} \tag{4.33}$$

for some $c > 0$ and all large n and $u \leq \sqrt{n}$. The result follows. □

Proof of Proposition 4.6. Using Lemma 4.5(ii), we have for all $M \geq 0$:

$$\sum_{u \leq r-M} \sum_{E \in A_u^r} \mathbb{E} \left[\prod_{i \in E} Y_i \right] \leq \sum_{u=1}^{r-M} \binom{r-1}{u-1} (2\ell(S))^{r-u} F_u. \tag{4.34}$$

Letting

$$B_{r,u} := \binom{r-1}{u-1} (2\ell(S))^{r-u} F_u, \tag{4.35}$$

we get from Lemma 4.8 that, for $r \leq \sqrt{n}$ and any $u \leq r - 1$:

$$\frac{B_{r,u+1}}{B_{r,u}} = \frac{1}{2\ell(S)} \frac{r-u}{u} \frac{F_{u+1}}{F_u} \geq \frac{Q}{2\ell(S)} \frac{n(r-u)}{u^2} \geq \frac{Q}{2\ell(S)} \frac{n}{u^2}. \tag{4.36}$$

Hence, there exists $\eta > 0$ small enough such that, for all $u < r \leq \eta\sqrt{n}$, we have $B_{r,u+1} \geq 2B_{r,u}$, and thus by backward induction,

$$B_{r,u} \leq 2^{-(r-u)} B_{r,r}. \tag{4.37}$$

Then, for $r \leq \eta\sqrt{n}$, (4.34) yields

$$\sum_{u \leq r-M} \sum_{E \in A_{r-u}^+} \mathbb{E} \left[\prod_{i \in E} Y_i \right] \leq \sum_{u=1}^{r-M} B_{r,u} \leq 2^{1-M} B_{r,r} = 2^{1-M} F_r. \tag{4.38}$$

This yields (4.31) if we choose M such that $2^{1-M} \leq \varepsilon$, since $r! F_r \leq \mathbb{E} [(X_{S,n})_r]$ by Lemma 4.5(i). \square

Proposition 4.6 shows that we only need to understand the asymptotic behaviour of the configurations with a number of blocks $r - M$ for given $M \geq 0$, and Lemma 4.5(iii) that we can focus on configurations with blocks of size 1 or 2. To actually prove our final result, we need to refine Lemma 4.5(iii) and obtain the explicit constants that appear. We define another set of constants, which will account for the cases with blocks of size 2, that is, cases when two components of shape S overlap.

Definition 4.9. Let S be a shape. There is a finite set of integers $i \geq 1$ such that $\mathbb{E} [Y_1 Y_i] > 0$ and $i - 1 < 2\ell(S)$. Let $I(S)$ be this set and i_1, \dots, i_k its elements. For $i \in I(S)$, let $\ell_i, K_i, c_+(i)$, and $c_-(i)$ be the equivalents of $\ell(S), K, c_+, c_-$ in this case of two components C, C' that overlap and start at positions 1 and i (replacing in the definition the component by the union of the two components). In particular, $\ell_i = \ell(S) + (i - 1)/2$ is the total half-length of the block made of two components of shape S started at positions 1 and i . Furthermore, C and C' together with the horizontal axis define two unbounded faces (F_+ in the upper half-plane and F_- in the lower half-plane) and several bounded faces; let $\mathcal{F}(C, C')$ be the set of bounded faces. For each face F , let $\nu(F)$ be the number of integers in $[[L(C), R(C)] \cup [[L(C'), R(C')]] = [[L(C), R(C')]]$ that are incident to F but do not belong to C nor to C' . We set $K_i := \prod_{F \in \mathcal{F}(C, C')} \text{Cat}_{\nu(F)/2}$. Finally, we define $c_{\pm}(i) := \nu(F_{\pm})/2$. Observe again that all these constants only depend on S and i .

Note that $i \in I(S)$ may be even; in this case $2\ell_i, \nu(F_+)$ and $\nu(F_-)$ are odd, and thus ℓ_i and $c_{\pm}(i)$ are half-integers.

Lemma 4.10. Let $r = O(\sqrt{n})$ with $r \rightarrow \infty$. Then, for every fixed $M \geq 0$,

$$\sum_{E \in A_{r-M}^+} \mathbb{E} \left[\prod_{i \in E} Y_i \right] \underset{n \rightarrow \infty}{\sim} F_r \sum_{\substack{g_i \geq 0, i \in I(S) \\ \sum_i g_i = M}} \prod_{i \in I(S)} \frac{(b_i \frac{r^2}{2n})^{g_i}}{g_i!}, \tag{4.39}$$

where

$$b_i := 4^{4\ell(S) - 2\ell_i + c_+(i) - 2c_+ + c_-(i) - 2c_-} \frac{K_i}{K^2}. \tag{4.40}$$

Note that b_i measures (in a specific way) how much two overlapping components of shape S differ from two non-overlapping ones.

Proof. For each $I(S)$ -tuple $G = (g_i)_{i \in I(S)}$ of integers with sum M , let $A_{r-M,G}^r$ be the set of r -tuples $1 \leq i_1 < \dots < i_r \leq 2n$ with $r - 2M$ blocks of size 1 and M blocks of size 2, such that for each $i \in I(S)$, there are g_i blocks of type $\{i_k, i_{k+1} = i_k + i - 1\}$ with $k < r$. Then $A_{r-M,G}^r$ is the union of some classes $A_{T,J}$ from the proof of Lemma 4.5, with all $T_i \in \{1, 2\}$ and a specified number g_i of k such that $J_k = (1, i)$. Hence, we obtain from (4.21), where the multinomial coefficient in (4.41) is the number of (T, J) that is included in $A_{r-M,G}^r$,

$$\sum_{E \in A_{r-M,G}^+} \mathbb{E} \left[\prod_{i \in E} Y_i \right] = \binom{r-M}{g_1, \dots, g_k, r-2M} \binom{2n - 2\tilde{\ell} + (r-M)}{r-M} \tilde{K} \frac{\text{Cat}_{n-d_+} \text{Cat}_{n-d_-}}{\text{Cat}_n^2}, \tag{4.41}$$

where, by (4.22)–(4.25) and the argument yielding them:

$$\tilde{K} = K^{r-2M} \prod_{i \in I(S)} K_i^{g_i}, \tag{4.42}$$

$$\tilde{\ell} = (r - 2M)\ell(S) + \sum_{i \in I(S)} g_i \ell_i, \tag{4.43}$$

$$d_{\pm} = \tilde{\ell} - (r - 2M)c_{\pm} - \sum_{i \in I(S)} g_i c_{\pm}(i). \tag{4.44}$$

We now argue similarly as in the proof of Lemma 4.5, but this time, we compare to F_r . We have

$$\binom{r - M}{g_1, \dots, g_k, r - 2M} \sim r^M \prod_{i \in I(S)} \frac{1}{g_i!}, \tag{4.45}$$

$$\frac{\binom{2n-2\tilde{\ell}+(r-M)}{r-M}}{\binom{2n-2r\ell(S)+r}{r}} = \frac{r!}{(r - M)!} \cdot \frac{(2n - 2\tilde{\ell} + (r - M))_{r-M}}{(2n - 2r\ell(S) + r)_r} \tag{4.46}$$

$$\sim r^M (2n)^{-M} \exp\left(-\frac{1}{4n}((r - M)(4\tilde{\ell} - r + M) - r(4r\ell(S) - r))\right) \sim r^M (2n)^{-M},$$

$$\frac{\tilde{K}}{K^r} = K^{-2M} \prod_{i \in I(S)} K_i^{g_i}, \tag{4.47}$$

$$\frac{\text{Cat}_{n-d_{\pm}}}{\text{Cat}_{n-r\ell(S)+rc_{\pm}}} \sim 4^{-d_{\pm}+r(\ell(S)-c_{\pm})} = 4^{2M(\ell(S)-c_{\pm})-\sum_{i \in I(S)} (\ell_i - c_{\pm}(i))g_i}. \tag{4.48}$$

and thus, from (4.41) and (4.12), recalling that $\sum_{i \in I(S)} g_i = M$,

$$\frac{\sum_{E \in A_{r-M,G}^r} \mathbb{E} \left[\prod_{i \in E} Y_i \right]}{F_r} \tag{4.49}$$

$$\underset{n \rightarrow \infty}{\sim} r^{2M} (2nK^2)^{-M} 4^{2(\ell(S)-c_+)M - \sum_{i \in I(S)} (\ell_i - c_+(i))g_i} 4^{2(\ell(S)-c_-)M - \sum_{i \in I(S)} (\ell_i - c_-(i))g_i} \prod_{i \in I(S)} \frac{1}{g_i!} K_i^{g_i}$$

$$= \left(B \frac{r^2}{2n} \right)^M \prod_{i \in I(S)} \frac{q_i^{g_i}}{g_i!} = \prod_{i \in I(S)} \frac{(Bq_i \frac{r^2}{2n})^{g_i}}{g_i!},$$

where

$$B := \frac{4^{4\ell(S)-2c_+-2c_-}}{K^2}, \tag{4.50}$$

$$q_i := 4^{-2\ell_i+c_+(i)+c_-(i)} K_i. \tag{4.51}$$

The set $A_{r-M}^r(1, 2)$ defined in Lemma 4.5(iii) is the union of $A_{r-M,G}^r$ over all G with sum M . Hence, (4.49) implies, noting that there is only a finite number of such G ,

$$\sum_{E \in A_{r-M}^r(1,2)} \mathbb{E} \left[\prod_{i \in E} Y_i \right] \underset{n \rightarrow \infty}{\sim} F_r \sum_{g_i \geq 0, i \in I(S)} \prod_{i \in I(S)} \frac{(Bq_i \frac{r^2}{2n})^{g_i}}{g_i!}. \tag{4.52}$$

$\sum_i g_i = M$

The result (4.39) now follows from (4.52) and (4.16), using $Bq_i = b_i$. □

Proof of Theorem 1.2 for weak shapes. Let $r \rightarrow \infty$ with $r \leq \eta\sqrt{n}$, where η is as in Proposition 4.6.

We may sum (4.39) over all $M \geq 0$ (with $A_{r-M}^r := \emptyset$ for $M > r$), since Proposition 4.6 shows that we may approximate the sum by a finite sum with a fixed number of terms. Consequently, recalling (4.17),

$$\begin{aligned} \mathbb{E} [(X_{S,n})_r] &= r! \sum_{M=0}^{\infty} \sum_{E \in A_{r-M}^r} \mathbb{E} \left[\prod_{i \in E} Y_i \right] \sim r! F_r \sum_{M=0}^{\infty} \sum_{\substack{g_i \geq 0, i \in I(S) \\ \sum_i g_i = M}} \prod_{i \in I(S)} \frac{(b_i \frac{r^2}{2n})^{g_i}}{g_i!} \\ &= r! F_r \prod_{i \in I(S)} \exp\left(b_i \frac{r^2}{2n}\right). \end{aligned} \tag{4.53}$$

By Lemmas 2.2 and 2.3, (4.12) implies (similarly to [4.5])

$$r! F_r \sim \left(\frac{2nK}{4^{2\ell(S)-c_+-c_-}}\right)^r \exp\left(-\frac{r^2}{4n}(4\ell(S)-1)\right). \tag{4.54}$$

Finally, (4.53) and (4.54) yield, for $r \rightarrow \infty$ with $r \leq \eta\sqrt{n}$,

$$\mathbb{E} [(X_{S,n})_r] \underset{n \rightarrow \infty}{\sim} \left(\frac{2nK}{4^{2\ell(S)-c_+-c_-}}\right)^r \exp\left(-\frac{r^2}{4n}(4\ell(S)-1) + \frac{r^2}{2n} \sum_{i \in I(S)} b_i\right). \tag{4.55}$$

This is (2.2), with

$$\mu_n := \frac{2nK}{4^{2\ell(S)-c_+-c_-}}, \tag{4.56}$$

$$s_n := \frac{-(4\ell(S)-1) + 2 \sum_{i \in I(S)} b_i}{2n}. \tag{4.57}$$

In particular, (2.2) thus holds for $r = r(n)$ with $\frac{\eta}{2}\sqrt{n} \leq r \leq \eta\sqrt{n}$; as noted in Section 2.1, it then automatically holds uniformly in this range. Furthermore,

$$\mu_n s_n \geq -\frac{K(4\ell(S)-1)}{4^{2\ell(S)-c_+-c_-}} > -1 \tag{4.58}$$

by Lemma 4.3, and we have again $\mu_n = \Theta(n)$ and $\sigma_n = \Theta(\sqrt{n})$. It follows that Theorem 2.1 applies in this case too, which yields (1.2). \square

We obtain from (4.56)–(4.57)

$$\sigma_S^2 = \frac{2K}{4^{2\ell(S)-c_+-c_-}} \left(1 + \frac{K}{4^{2\ell(S)-c_+-c_-}} \left(1 - 4\ell(S) + 2 \sum_{i \in I(S)} b_i\right)\right), \tag{4.59}$$

with b_i given by (4.40). Note that this formula holds also for strong shapes (when $I(S) = \emptyset$) by (4.11).

5. Open problems

We list here some open problems concerning possible extensions of our results.

1. It seems possible to extend the arguments above to joint factorial moments

$$\mathbb{E} [(X_{S_1,n})_{r_1} \cdots (X_{S_k,n})_{r_k}] \tag{5.1}$$

for several shapes S_1, \dots, S_k and then obtain a multivariate version of Theorem 1.2 using a multivariate version of Gao and Wormald’s theorem [3], [8]. However, we have not

checked the details. Such a multivariate theorem would immediately imply, for example, a central limit theorem for the number of components of a given half-length.

2. Considering shapes that are similar, can we obtain a central limit theorem for the number of components that only cross the horizontal axis twice (i.e. the support has size 2, but the half-length is arbitrary)?
3. Is it true, as Kargin [6] has conjectured, that the total number of components is asymptotically normal?

References

- [1] Di Francesco, P., Golinelli, O. and Guitter, E. (1997) Meanders and the Temperley-Lieb algebra. *Commun. Math. Phys.* **186**(1) 1–59.
- [2] Borga, J., Gwynne, E. and Park, M. (2023) On the geometry of uniform meandric systems, arXiv:2212.00534.
- [3] Ojeda, G. B., Holmgren, C. and Janson, S. (2023) Fringe trees for random trees with given vertex degrees, arXiv:2312.04243.
- [4] Féray, V. and Thévenin, P. (2023) Components in meandric systems and the infinite noodle. *Int. Math. Res. Not. IMRN*, **2023**(14) rnac156.
- [5] Gao, J. and Wormald, N. (2004) Asymptotic normality determined by high moments, and submap counts of random maps. *Probab. Theory Relat. Fields* **130**(3) 368–376.
- [6] Kargin, V. (2020) Cycles in random meander systems. *J. Stat. Phys.* **181**(6) 2322–2345.
- [7] Stanley, R. P. (2015) *Catalan Numbers*. Cambridge University Press.
- [8] Wormald, N., Personal communication.