

Echinoderms from the lower Silurian Brassfield Formation of east-central Kentucky

William I. Ausich,¹ Mark E. Peter,¹ and Frank R. Etensohn²

¹School of Earth Sciences, 155 South Oval Mall, The Ohio State University, Columbus, OH 43210, USA, <ausich.1@osu.edu>; <peter.1@osu.edu>

²Department of Earth and Environmental Sciences, University of Kentucky, Lexington, KY 40506, USA, <f.ettensohn@uky.edu>

Abstract.—A new echinoderm fauna is reported from the Brassfield Formation (Rhuddanian, Silurian) of Bath County, Kentucky. The Brassfield Formation was the first extensive marine unit to be deposited following the end-Ordovician glaciation and extinctions and represents several shallow, open-marine facies. These facies supported a diverse pelmatozoan fauna. This report not only extends the geographic distribution of this fauna, but also the temporal range of the fauna back to Rhuddanian time. Six pelmatozoans are reported, including the crinoids *Browerocrinus arthrikos* n. gen. n. sp., *Temnocrinus americanus* n. sp., *Stereoaster* sp., and *Dendrocrinus* sp.; and the glyptocystitids *Brockocystis nodosarius* Foerste, 1919, and *Anartiocystis whitei* Sumrall, 2002. In addition, the asteroid *Gordonaster brassfieldensis* Blake and Etensohn, 2009, was reported previously from this locality. *Browerocrinus* increases the diverse calceocrinid fauna from the Brassfield Formation; *Temnocrinus* was previously only known from the Homerian (Silurian) of England; and this is the first known occurrence of *Stereoaster* beyond the greater Dayton, Ohio, region. Furthermore, this is the first Brassfield locality known with two glyptocystitid taxa.

Introduction

Crinoids suffered a mass extinction during the end-Katian at the onset of end-Ordovician glaciation, and it was not until the Aeronian (Llandovery, early Silurian) that crinoid generic richness recovered to pre-extinction levels (Peters and Ausich, 2008). This extinction and recovery resulted in a reorganization of crinoid faunas from the Early Paleozoic to the Middle Paleozoic Crinoid Evolutionary Fauna (Baumiller, 1994; Ausich et al., 1994). Until the early 1980s, Llandovery crinoid faunas were very poorly known (Ausich and Deline, 2012). However, a new appreciation for the Llandovery crinoid faunal recovery (Ausich and Deline, 2012) is emerging with the discovery of new Llandovery faunas from Iowa (Witzke and Strimple, 1981), Ohio (Ausich, 1984a, b, 1985, 1986a, b, c, d, 1987a, b; Ausich and Dravage, 1988; Boyarko and Ausich, 2009), Britain and Ireland (e.g., Donovan and Sevastopulo, 1989; Donovan, 1993; Donovan and Harper, 2003; Donovan and Lewis, 2005; Fearnhead and Donovan, 2007a, b, c; Fearnhead and Harper, 2007; Donovan et al., 2008, 2009, 2010, 2012), and New York and Canada (e.g., Eckert 1984, 1990; Eckert and Brett, 2001; Ausich and Copper, 2010).

The work by Ausich and colleagues in Ohio described faunas from the upper parts of the Brassfield Formation (Aeronian) and was an extension of the pioneering work of Foerste (1919). With the exception of Ausich and Dravage (1988) and Boyarko and Ausich (2009), Brassfield crinoid faunas are primarily known from the greater Dayton, Ohio, area. Ausich and Dravage (1988) and Boyarko and Ausich (2009) described only two species and recognized two taxa in open

nomenclature from Adams County, Ohio, and northern Kentucky. Thus, the new, low-diversity fauna described herein from Bath County, Kentucky, is particularly significant for understanding the geographic variability of crinoid faunas from the Brassfield Formation and for its even earlier Rhuddanian age. Here, we describe a pelmatozoan fauna with six taxa, including two rhombiferans and four crinoids. Crinoid taxa include one new genus, two new species, and two taxa left in open nomenclature. New taxa include the calceocrinid crinoid, *Browerocrinus arthrikos* n. gen., n. sp. and a flexible crinoid, *Temnocrinus americanus* n. sp. *Temnocrinus* was previously recognized only from the Much Wenlock Limestone (Homerian) of England.

The richness of this crinoid fauna from Bath County is low in comparison to that from the Dayton, Ohio, area, but crinoids, rhombiferans, and an asteroid co-occur in this relatively limited exposure. The asteroid *Gordonaster brassfieldensis* was described by Blake and Etensohn (2009). With *Anartiocystis whitei* Sumrall, 2002, and *Brockocystis nodosarius* Foerste, 1919, this is the only Brassfield locality known in which two rhombiferan genera co-occur, although Broadhead and Etensohn (1981) described the co-occurrence of two *Brockocystis* species (*B. tecumseth* and *B. nodosaria*) at the same horizon in the lower massive member farther to the north in Fleming County, Kentucky. From this same northern locality, Broadhead and Etensohn (1981) also described an Aeronian species of *Brockocystis* from the Oldham Member of the Crab Orchard Formation, immediately overlying the Brassfield Formation.

The Brassfield Formation represents the first major Silurian episode of sea-level rise during the waning phases of the end-Ordovician glaciation. Although small reefs do occur in the

Brassfield Formation (Sheehy, 1981; Schneider and Ausich, 2002), most of this unit was an extensive grainstone/packstone facies that was deposited with the reestablishment of the Silurian epicontinental sea in eastern North America (McLaughlin et al., 2008; Brett et al., 2012; Etensohn et al., 2013). Based on data presented herein, it is clear that considerable faunal diversity existed within the widespread Brassfield facies, and that this diversity developed somewhat earlier than previously thought.

Geographic, stratigraphic, and paleoenvironmental occurrence

Echinoderms discussed in this paper were collected from Bath County, Kentucky, and are from a roadside borrow pit along the southern side of Interstate 64, ~0.5 mi (0.8 km) west of the intersection of Interstate 64 and Kentucky Highway 36 (38° 7.4' N lat., 83° 45.5' W long., USGS Preston Quadrangle) (Gordon and Etensohn, 1984; Blake and Etensohn, 2009). The echinoderms are from the uppermost part of the lower massive or “cherty” member of the Brassfield Formation (Gordon and Etensohn, 1984; Blake and Etensohn, 2009; Etensohn et al., 2013), which occurs in the *Distomodus kentuckyensis* conodont Assemblage Zone (Rexroad and Kleffner, 1984) and is regarded as early Llandovery (Rhuddanian) in age in east-central and northeastern Kentucky (Etensohn et al., 2013). This unit is present below the biostratigraphically significant Aeronian “bead bed” or “cogwheel bed” of Foerste (1919), Rexroad et al. (1965), and Rexroad (1967), and represents a high-energy, sandbelt environment in a westwardly transgressing, shallow, open-marine sequence (Gordon and Etensohn, 1984; Etensohn et al., 2013). Within that setting, the echinoderms inhabited an irregular hardground surface that was apparently persistent throughout much of northeastern and east-central Kentucky (Broadhead and Etensohn, 1981; Gordon and Etensohn, 1984; Blake and Etensohn, 2009). In fact, this surface may have local biostratigraphic significance (Broadhead and Etensohn, 1981), and may be the same surface from which Foerste (1919) described the type specimens of *Brockocystis nodosarius* in Ohio. Like most of the Kentucky Brassfield Formation, the unit has been pervasively dolomitized by secondary processes related to the formation of the Cincinnati Arch (Gordon and Etensohn, 1984).

At first, the Aeronian age of similar echinoderm faunas from the Dayton area (e.g., Peters and Ausich, 2008) may seem to contradict the Rhuddanian age of the Kentucky fauna; however, this is a product of paleogeography. The exposures of the Brassfield Formation in east-central and northeastern Kentucky occur farther eastward than do the exposures near Dayton, Ohio, from which most Brassfield echinoderms have

been described. This means that the early Silurian transgression first reached east-central and northeastern Kentucky during late Rhuddanian time, but not until the Aeronian did it migrate farther westward into the Dayton area. Hence, although Brassfield Formation units in the Dayton area may be homotaxial with those in east-central and northeastern Kentucky, the units in Kentucky are slightly older, and the contained, post-extinction, Llandovery, crinoid faunal recovery (Ausich and Deline, 2012) most likely had a Rhuddanian origin.

Systematic paleontology

Terminology follows Moore (1962), Ubahgs (1978), Brett (1981), and Ausich et al. (1999); classification follows Ausich (1998). Repository abbreviations are as follows: CMCIP, Cincinnati Museum Center, Cincinnati, Ohio, and USNM, U.S. National Museum, Smithsonian Institution. All measurements are in mm; * indicates that the character is incomplete or compressed.

Order Glyptocystitida Bather, 1899

Family Callocystitidae Bernard, 1895

Remarks.—In southwestern Ohio and northeastern Kentucky, *Brockocystis nodosarius* Foerste, 1919, is a reasonably common glyptocystitid in the limestones and dolostones of the Brassfield Formation and lower Crab Orchard Formation, the occurrence of which is based primarily on the distinctive cone-shaped columnals that occur in the proximal portion of the mesistele (Fig. 1.5, 1.6). However, *Anartiocystis* has only been reported from two Brassfield localities: Montgomery County, Ohio (Ausich and Schumacher, 1984), and the present locality in Bath County, Kentucky (Sumrall, 2002). This is the only known locality in the Brassfield Formation where *Anartiocystis* and *Brockocystis* co-occur.

The systematics of the Callocystitidae is currently under consideration. Consequently, these two genera are not placed in a subfamily.

Subfamily uncertain

Genus *Anartiocystis* Ausich and Schumacher, 1984


Type species.—*Anartiocystis foerstei* Ausich and Schumacher, 1984, by monotypy.

Other species.—*Anartiocystis whitei* Sumrall, 2002.

Anartiocystis whitei Sumrall 2002

Figure 1.3, 1.4

Figure 1. Brassfield Formation rhombiferans and crinoids from Kentucky. (1, 2, 5, 6) *Brockocystis nodosarius* Foerste, 1919; (1) D-ray lateral view of a crushed specimen, USNM 594948, ×3.0; (2) mostly disarticulated periproct region, USNM 594948, ×8.0; (5) lateral view of a specialized proxistele columnal, USNM 594950, ×3.5; (6) proximal view of specialized proxistele columnal, USNM 594950, ×3.5; (3, 4) *Anartiocystis whitei* Sumrall, 2002; (3) articulated periproct region, USNM 594952, ×8.0; (4) D-ray lateral view of partially crushed specimen, USNM 594952, ×3.0; (7–13) *Browerocrinus arthrikos* n. gen. n. sp.; (7) E-ray view of aboral cup and proximal arms, holotype, USNM 594961, ×3.5; (8) small plating covering the hinge between the radial and basal plate circlets, holotype, USNM 594961, ×8.0; (9) E-ray view of crown, holotype, USNM 594961, ×3.5; (10) posterior view of crown, holotype, USNM 594961, ×3.5; (11) lateral view of juvenile crown, paratype, USNM 594963, ×4.0; (12) lateral view of crown, holotype, USNM 594961, ×3.5; (13) E-ray view of aboral cup with proximal arms, note small plates covering the hinge between the radial and basal circlets, paratype, USNM 594962, ×3.5; (14) *Dendrocrinus* sp., lateral view of a compressed anal sac, USNM 595072, ×3.5; (15) partial holdfast of *Stereocrinus* sp., USNM 594959, ×3.5.



Material.—Holotype: CMCIP 50601; new material: USNM 594952–594958.

Occurrence.—Brassfield Formation, Rhuddanian (Llandoverly, Silurian), Bath County, Kentucky.

Remarks.—With ambulacra confined to the radial plates, these seven new specimens are assigned to *A. whitei*. The largest (19 mm, USNM 594953) of these new specimens (Fig. 1.4) is larger than those reported by Sumrall (2002). Specimens are either mostly inflated (USNM 594953, USNM 594954, USNM 594956) or compacted (USNM 594952, USNM 594955, USNM 594957). The periproct on USNM 594952 is very well preserved (Fig. 1.3). The periphery of the periproct plating is comprised of 14 or 15 variously sized tetragonal, pentagonal, and hexagonal plates. Articulated to this outer ring is a series of elongate, broadly triangular plates that taper toward the center of the periproct. Although mostly disarticulated, a discontinuous ring of smaller, elongate plates occurs at the summit of the periproct plating.

Genus *Brockocystis* Foerste, 1919

Type species.—*Brockocystis nodosarius* Foerste, 1919, by monotypy.

Brockocystis nodosarius Foerste, 1919
Figure 1.1, 1.2, 1.5, 1.6

Material.—Syntypes: USNM 85346A to USNM 85346E; new material: USNM 594948 to USNM 594951.

Occurrence.—Brassfield Formation, Rhuddanian (Llandoverly, Silurian), Bath County, Kentucky.

Remarks.—Two partially compacted thecae (e.g., Fig. 1.1), distinctive proximal mesistele columnals (Fig. 1.5, 1.6), and numerous isolated plates were collected from the Brassfield Formation of Bath County, Kentucky. These are relatively large specimens (up to ~20 mm). Although not nearly as complete as in *A. whitei* (Fig. 1.3), some periproct plating is preserved on both thecae of *B. nodosarius* (Fig. 1.2). This periproct plating is comprised of much smaller and more numerous plates than the periproct of *A. whitei*. Although these are large specimens, they are still relatively small compared to the unnamed species described by Broadhead and Etensohn (1981) from the overlying Crab Orchard Formation. Plates from *B. nodosarius* have also been reported from the upper massive member (Aeronian) of the Brassfield in what may be another prominent echinoderm-bearing horizon from northeastern Kentucky (Broadhead and Etensohn, 1981).

Class Crinoidea Miller, 1821
Subclass Camerata Wachsmuth and Springer, 1885
Order Diplobathrida Moore and Laudon, 1943
Suborder Eudiplobathrina Ubahgs, 1953
Superfamily Rhodocrinitoidea Roemer, 1853
Family Rhodocrinitidae Roemer, 1853
Genus *Stereoaster* Foerste, 1919

Type species.—*Stereoaster squamosus* Foerste, 1919.

Stereoaster sp.
Figure 1.15

Material.—USNM 594959, 594960.

Occurrence.—Brassfield Formation, Rhuddanian (Llandoverly, Silurian), Bath County, Kentucky.

Remarks.—One holdfast from this fauna corresponds to the distinctive holdfast of *Stereoaster squamosus* Foerste, 1919 from the Brassfield Formation, which was initially identified as an asteroid. This holdfast has multiple rhizoids, each of which is comprised of numerous, irregularly arranged plates (Fig. 1.15). Ausich (1986c) described a complete specimen in which this holdfast was attached to a rhodocrinitid crinoid crown. *Stereoaster squamosus* is the most abundant crinoid in the area of Dayton, Ohio; however, this is the first occurrence of this genus beyond the Dayton region.

Order Cladida Moore and Laudon, 1943
Suborder Dendrocrinida Bather, 1899

Superfamily Dendrocrinoidea Wachsmuth and Springer, 1886
Family Dendrocrinidae Wachsmuth and Springer, 1886
Genus *Dendrocrinus* Hall, 1852

Type species.—*Dendrocrinus longidactylus* Hall, 1852.

Dendrocrinus sp.
Figure 1.14

Material.—USNM 594972 and USNM 594973.

Occurrence.—Brassfield Formation, Rhuddanian (Llandoverly, Silurian), Bath County, Kentucky.

Remarks.—Two fragments of a cladid anal sac were recovered from this site. This anal sac is comprised of columns of large, plicate plates and is consistent with the anal sac of *Dendrocrinus* (Fig. 1.14). *Dendrocrinus daytonensis* Ausich, 1986b, was reported from the Brassfield Formation in the greater Dayton area. However, a species identification is not possible based solely on a partial anal sac; thus, this taxon is left in open nomenclature.

Subclass Disparida Moore and Laudon, 1943
Order Calceocrinida Ausich, 1998
Family Calceocrinidae Meek and Worthen, 1869
Genus *Browerocrinus* new genus

Type species.—*Browerocrinus arthrikos* n. gen. n. sp.

Diagnosis.—As for species, by monotypy.

Etymology.—In honor of James C. Brower.

Occurrence.—Rhuddanian (Llandoverly), Silurian; North America.

Remarks.—Calceocrinids are among the most diverse families of crinoids from the Brassfield Formation. This new genus is the fourth described. *Calceocrinus insertus* Foerste, 1919, *Stibarocrinus centervillensis* (Foerste, 1919), and *Trypheroocrinus brassfieldensis* Ausich, 1984a, occur in the greater Dayton, Ohio area (Ausich, 1984a), and Boyarko and Ausich (2009) described *Trypheroocrinus adamsensis* from Adams County, Ohio. Boyarko and Ausich (2009) also described an isolated aboral cup from an indeterminate calceocrinid, which is now recognized as *Browerocrinus arthrikos* n. gen. n. sp. Disparid crinoids were characteristic of the Early Paleozoic Crinoid Evolutionary Fauna (CEF), as described by Baumiller (1994) and Ausich et al. (1994); and the Calceocrinidae was the only disparid family that underwent a significant radiation immediately after the late Ordovician (end-Katian) extinction event (Ausich and Deline, 2012).

Among calceocrinids, only *Calceocrinus*, *Cremacrinus*, *Stibarocrinus*, and *Browerocrinus* have four basal plates. These genera can be distinguished because *Calceocrinus* has a calyx that is laterally to abanally-adanally compressed, with a short or long suture between the E-ray infer- and superradials, bilateral heterotomous lateral arm branching, and the crown recumbent on the column; *Cremacrinus* has a calyx that is abanally-adanally compressed, long suture between the E-ray infer- and superradials, bilateral heterotomous lateral arm branching, and the crown recumbent or pendant on the column; *Stibarocrinus* has a calyx that is subcylindrical in shape, short suture between the E-ray infer- and superradials, bilateral heterotomous with robust beta ramules for lateral arm branching, and the crown pendant on the column. In contrast, *Browerocrinus* has a calyx that is abanally-adanally compressed, short suture between the E-ray infer- and superradials, poorly isotomous lateral arm branching, and the crown pendant on the column.

Browerocrinus arthrikos new genus new species
Figures 1.7–1.13 and 2.2–2.5

2009 Calceocrinid indeterminate, Boyarko and Ausich, p. 106–107, figs. 3–4.

Diagnosis.—Crown with perfect bilateral symmetry, crown pendant on column, aboral cup adanally-abanally compressed, ligament pit divided, four basal plates, four basal plates part of distal basal cirlet, four basal plates in column concavity, E-ray infer- and superradial plates with short suture, proximal width of E-ray inferradial approximately 30% of hinge length, B and C inferradials not fused with one another and not fused with A and D radials, E-ray arm branches, main axils with non-axillaries, three arms, lateral arm branching poorly isotomous, beta ramule present, but unknown whether branched or unbranched.

Description.—Crown small, slender, and pendant on column. Aboral cup relatively small, abanally-adanally compressed, constricted mid-height (Figs. 1.9, 2.5). Four basal plates; all four basal plates in column concavity; all four basal plates part of distal margin of basal cirlet with DE and AE basal plates together occupying approximately 80% of this margin;

projections from the radial cirlet proximally and basal cirlet distally divide the ligament pit. Radial cirlet abanally-adanally compressed, rectangular with medial restriction in aboral view. A and D radials majority of radial plate cirlet. E infer-radial and E superradial narrow toward the center of radial cirlet with narrow sutural contact. E inferradial 30% of proximal margin of radial cirlet along hinge. E superradial 100% of aboral cup distal margin, supports E-ray arm. A and D lateral arms from upper lateral facet of A and D radial; two lower lateral facets on A and D radial plates support B and C inferradials.

Numerous, minute plates cover the hinge region (Figs. 1.7, 1.8, 1.13, 2.3, 2.4); plates irregular in shape but commonly elongate and oriented more or less with long axis of plates parallel to the long axis of the hinge region. On USNM 594961, the crown is closed and minute plates more widely spaced (Figs. 1.7, 2.3); on USNM 594962 the crown is open and minute plates more completely fill the hinge area (Figs. 1.13, 2.4). Anal X hexagonal, supported beneath by the subanal (fused B and C superradials); four known anal plates above anal X; at least three proximal anal plates rectangular (Fig. 1.10). Anal sac very long, projecting for length of preserved arms.

Three arms. E-ray arm robust and branched, brachials variable in height. Fourth or fifth primibrachial, third or fourth secundibrachial, and fourth or higher tertibrachial axillary (Fig. 1.9, 2.5). E-ray brachial height:width ratio ranges from 2.0 times wider than high to 1.1 times wider than high.

Lateral arms with weakly developed main axil series and arm branching pattern different from normal calceocrinid bilateral heterotomy (Figs. 1.12, 2.2). Four axillaries in main axil, one non-axillary brachial between each axillary. Alpha ramule on the primaxil arm bifurcates on the second alpha-brachial and supports the betabrachitaxis adanally; the abanal alpha ramule is completely covered by other arms, so no details known. Third betabrachial axillary with a gammabrachitaxis abanally and atomous beta ramule adanally. Third gammabrachial axillary with only the adanal deltabrachitaxis visible. Third deltabrachial axillary, which is the final bifurcation preserved; unbranched epsilonbrachials above as known.

On the secundaxil arm, third alphabrachial axillary and supports the betabrachitaxis adanally and abanal branch mostly covered. Third betabrachial axillary; third gammabrachial axillary; third deltabrachial axillary; and no further branching preserved on this arm.

Tertaxil arm bifurcates on third alphabrachial with betabrachitaxis adanal and the alpha ramule abanal. Distally, abanal branch is atomous. Third betabrachial axillary and supports the gammabrachitaxis abanally and the beta ramule adanally. Third gammabrachial axillary, other details unknown.

Quartaxil arm present but not preserved, so branching condition above the second alphabrachial unknown. Omega ramule present but only two brachials preserved, so branching condition unknown.

Proximal-most columnals very thin, heteromorphic, circular; lumen circular; columnal height increases distally. Mesistele and dististele unknown.

Etymology.—*Arthrikos* (Greek, of a joint) in reference to the distinctive bifurcations on the finger-like arms.

Material.—Holotype USNM 594961; paratypes USNM 594962–594964, CMCIP 51205; other material: USNM 594965 and 594966.

Occurrence.—Brassfield Formation. Rhuddanian (Llandovery, Silurian), Bath County, Kentucky.

Remarks.—Seven specimens of *B. arthrikos* include: (1) USNM 594961, a complete crown of an adult; (2) USNM 594962, an aboral cup with proximal arms preserved; (3) USNM 594963, a complete crown of a juvenile (Fig. 1.11); (4) USNM 594964, an aboral cup; (5) CMCIP 51205, an aboral cup; (6) USNM 594965, a partial aboral cup; and (7) USNM 594966, a set of distal arms.

This new material is significant for understanding calceocrinid morphology, in general, because two specimens preserve irregular, minute plates covering the hinge region. The Calceocrinidae are unique among crinoids because the crown is imperfectly to perfectly bilaterally symmetrical (Moore, 1962; Ubaghs, 1978). Rather than two circlets of five plates positioned around the oral-aboral axis of the organisms, the basal circlet is reduced to a crescent shape, and the radial circlet is typically either rectangular or trapezoidal in shape. These two circlets articulate along a hinge that allowed the crown to close recumbently along the column (which ran along the sea floor) or open perpendicular to the sediment-water interface in a feeding position (Moore, 1962; Ausich, 1986d; Brower, 1988). The nature of the soft tissue on the basal plate-radial plate hinge articulation has been a point of discussion, but no evidence exists for any connective tissue other than ligaments.

USNM 594961 and 594962 are exceptional in possessing a series of minute calcareous plates filling the basal-radial plate junction region (see Bather, 1893, pl. 14, figs. 141, 142; Springer, 1926, pl. 28, figs. 20 and 20a). These plates (“supplementary plates” of Bather and Springer) are minute, irregularly shaped, typically elongate plates that are commonly oriented with the long axis approximately oriented along the width of the articulation. USNM 594962 has the hinge open, and these plates are juxtaposed close to one another and nearly fill the gape between the circlets. Conversely, USNM 594961 has the hinge closed, and the small plates are more widely separated from one another, yielding a much less dense covering of these plates over the gape between circlets. Contrasting preservational modes with contrasting crown positions demonstrate that these plates are not an artifact. Herein, these plates are interpreted to have been held within a leathery integument that covered the hinge region. It is reasonable to assume that other calceocrinids also possessed such an integument, presumably to protect tissue vital to the hinge mechanism. The absence of this small plating in almost all other calceocrinids could be a taphonomic artifact, or it is possible that other taxa had an integument that lacked plating.

USNM 594963 is a juvenile specimen (Fig. 1.11), with USNM 594961 (Fig. 1.9, 1.10, 1.12) and USNM 594962 (Fig. 1.13) each progressively larger. Thus, a general indication of plate-shape change through growth can be deduced for *B. arthrikos*. The height-to-width ratio of the basal circlet is relatively constant, as is the height to proximal width of the radial circlet. In contrast, the height to distal width of the radial circlet is much greater in larger specimens, and the height-to-

width ratio of brachial plates is much smaller (Table 1). These changes in shape through growth contrast somewhat with results of descriptions by Brower (1990), who reported a relative increase in the height-to-width ratio in calceocrinids for both the proximal and distal cup width.

Measurements.—CMCIP 51205, basal circlet height, 1.5; basal circlet width, 5.3; radial circlet height, 5.7; radial circlet width, 6.4; and column length, 0.4* (Table 1).

Subclass Flexibilia von Zittel, 1895
Order Sagenocrinida Springer, 1913
Superfamily Sagenocrinitoidea Roemer, 1854
Family Dactylocrinidae Bather, 1899
Genus *Temnocrinus* Springer, 1902

Type species.—*Cyathocrinites tuberculatus* Miller, 1821, by monotypy.

Occurrence.—Rhuddanian (Llandovery), Silurian, North America; Homerian (Wenlock), Silurian, England. *Temnocrinus* was known previously only from the Much Wenlock Limestone Formation at Dudley, England (Ramsbottom, 1953; Donovan et al., 2010), and *T. tuberculatus* was hitherto the only named species (Webster, 2003).

Fearnhead (2009) described and figured a cf. *Temnocrinus* sp. from the Hughley Shales (Telychian, Llandovery, Silurian) at Devil’s Dingle, north of Buildwas, Shropshire (Fearnhead, 2009; Donovan et al., 2010). However, that specimen appears to lack interrarial plates, and Fearnhead (2009) stated that the two primibrachials are free above the radials. The posterior side is not visible on this specimen, so the anal series cannot be compared with that of *Temnocrinus*. Based on the lack of interrarial plates, Fearnhead’s cf. *Temnocrinus* is not here considered to belong to *Temnocrinus*.

Donovan et al. (2008) also listed an occurrence of *T. tuberculatus* from the basal Woolhope Limestone (Sheinwoodian, Wenlock, Silurian) from quarries at Clencher Mill Lane, Ledbury, Herefordshire. However, Donovan et al. (2010) stated that *T. tuberculatus* is known only from the Much Wenlock Limestone at Dudley. As no museum number is cited for the Clencher Mill Lane occurrence, it needs to be independently confirmed (S. K. Donovan, personal comm. 25 February 2014). The *Temnocrinus* species described herein is the first occurrence of the genus from the Rhuddanian and the first from North America (Webster, 2003).

Temnocrinus americanus new species
Figures 2.1; 3.1–3.6

Diagnosis.—Large species with elongate crown; as many as six or more brachitaxes. Aboral cup medium cone shaped. Infrabasals fully visible in side view. Plate sculpturing mostly smooth on calyx; tubercles and prominent lateral nodes on secundibrachials through quartibrachials. Primibrachials two; secundibrachials typically four; tertibrachials five to eight. Arm branches in second through fourth brachitaxes diverge after the second or third plate above the axillary. Interarea with three to eight plates in two to four ranges, followed by small, unsutured plates of the oral region. Distal-most margins of interrarial plates form upward-directed spines.




Figure 2. Camera lucida drawings of Brassfield Formation crinoids. (1) *Temnocrinus americanus* n. sp., paratype, USNM 594968, CD-interray view of calyx (compare to Fig. 3.6); (2–5) *Browerocrinus arthrikos* n. gen. n. sp.; (2) lateral view of holotype, USNM 594961 (compare to Fig. 1.12); (3) plating partially covering the hinge area, holotype, USNM 594961 (compare to Fig. 1.8); (4) plating partially covering the hinge area, paratype, USNM 594962 (compare to Fig. 1.13); (5) E-ray view of holotype, USNM 594961 (compare to Fig. 1.9). Patterns and plate designations: black, radial or superradial plate; horizontal rule, infraradial plate; cross hatch, radianal plate; stippled, interradial and intrabrachial plates; grey, matrix.

Description.—Crown elongate, ovate, expanding distally to lowermost quartibrachials; thereafter more compact with narrower brachials. Arms coiling inward toward the oral surface beginning with quintibrachials.

Aboral cup (Fig. 3.5, 3.6) medium cone shaped; width-to-height ratio ~1.2 to 1.3. Plates gently convex with smooth plate sculpturing. Infrabasals three, unequal with smaller azygous infrabasal in C ray. Infrabasals wider than high and not covered by column cicatrix; plates entirely visible in side view of calyx. Infrabasal circling ~20% of total aboral cup height.

Basals five, in lateral contact; pentagonal (AB and DE), hexagonal (BC and EA), and heptagonal (CD). Basal plates slightly higher than wide or as wide as high; basals slightly smaller than radials. Basal circling ~40% of total aboral cup

height. CD basal truncated distally where it borders with anal X plate.

Radials five, in narrow contact except for CD interrady; hexagonal (C and D), heptagonal (A and E) and octagonal (B); radials wider than high. Radial circling ~40% of total aboral cup height. Radianal immediately below C radial. Radianal full width of C radial and aligned with radial circling; C radial above radial circling, having form of an extra primibrachial plate (Fig. 2.1).

Anal X hexagonal, slightly higher than wide; sutured below to CD basal and radianal and laterally to C and D radials. Eight or more plates follow in uniseries from right shoulder of anal X; series sutured to C ray. These anal plates with a coalesced transverse row of tubercles on each plate, on the plate margins




Figure 3. *Temnocrinus americanus* n. sp. (1, 2) holotype, USNM 594967, $\times 2.0$; (1) B-ray view of crushed crown; (2) D-ray view of crushed crown; (3, 4) paratype, USNM 594969, juvenile specimen, $\times 4.0$; (3) B-ray view; (4) D-ray view; (5, 6) paratype, USNM 594968, $\times 2.5$; (5) A-ray view of crushed crown; (6) CD-interray view of crushed crown.

Table 1. Measurements in mm on growth series of *Browerocrinus arthrikos*. Height, width, and height-width ratio of basal circolets, E-ray arm first primibrachials, and E-ray arm second primibrachials.

	Basal Circolet		E-Ray Arm First Primibrachial		E-Ray Arm Second Primibrachial		E-Ray Arm First Primibrachial		E-Ray Arm Second Primibrachial	
	Height	Width	Height/Width	Height/Width	Height/Width	Height/Width	Height/Width	Height/Width	Height/Width	
Juvenile	0.68	1.76	0.38	1.08	1.89	0.57	2.03	1.08	1.88	
Adolescent	1.35	3.24	0.42	1.49	3.78	0.39	2.30	2.57	0.89	
Adult	1.62	5.27	0.31	1.76	6.76	0.26	1.69	3.11	0.54	

opposite the C ray. Above the left shoulder of the anal X follows a succession of three plates, with one in the first row and two in the second row. Two of these plates sutured to D ray. Interarea above anal plates filled with small, unsutured oral region plates, some of which appear to be cone-shaped.

Regular interrays with two to four ranges of sutured interrachial plates, rising to level of first or second secundibrachial and effectively fixing entire primibrachitaxis and lower part of secundibrachitaxis. One interrachial plate in first range, two in second, and three each in third and fourth ranges, where present. Some in most distal range of interrachial plates with vertically and medially directed spines. Distalmost one or two ranges of interrachial plates with either single large tubercle or smaller tubercles aligned in vertical axial column within plate.

Intrabrachial plates between both secundibrachials and tertibrachials, with one or more large tubercles. As many as five or more intrabrachials between secundibrachials and one to three intrabrachials between tertibrachials. Intrabrachials above tertibrachitaxis absent.

Oral surface unknown, but interareas between arms and above sutured interrachial plates contain small, unsutured plates that were likely imbedded in a flexible integument not preserved in fossil specimens.

Arms branch as many as five times. First two divisions on fixed brachials isotomous; more distal divisions poorly isotomous, with equal-sized arm branches, but with branching occurring at slightly different levels. Primibrachials consistent at two in all of the three known specimens. Secundibrachials nearly consistent at four in the two specimens where preserved. Tertibrachials five to eight, as known; higher brachitaxes variable in length. In quintibrachitaxis, arms begin to coil inward toward oral region, coiling tightly at distal margins. For second through fourth brachitaxes, first few plates of each series sutured medially; arm branches do not diverge immediately after axillary. Intrabrachials above point of arm branching in secundibrachitaxis and tertibrachitaxis. For quintibrachitaxis and higher, arms diverge immediately following axillary of next lower brachitaxis.

Brachials rectilinear uniserial (sensu Webster and Maples, 2008); brachials wider than high proximally to as wide as high distally; aborally rounded; approximate height-to-width ratio 0.7 for both primibrachials and secundibrachials. Brachial sutures in fixed brachials nearly straight, arcuate, or subtly sinuous; free arms with sinuous sutures; tertibrachials and higher brachials with patelloid processes. Plate sculpturing on lower brachials mostly smooth; free brachials in secundibrachitaxis through quartibrachitaxis with transverse rows of tubercles on lateral margins that coalesce, forming pronounced lateral brachial nodes. Above quartibrachitaxis, lateral nodes disappear as brachials narrow transversely and deepen adorally. Tertiar and quartibrachials covered with small tubercles. Ambulacral surface of brachials obscured by matrix on available specimens; largest specimen with small ambulacral cover plates on one of the distal arms.

Proximal column (Fig. 3.3, 3.4) circular and homeomorphic, with thin discoidal columnals widening proximally to slightly narrower than the width of infrabasal circolet. External sutures crenulate. Outer surface of epifacet rounded, latus

profile slightly convex. Lumen roughly circular to polygonal; lumen diameter at level of most proximal columnal 27% of total columnal diameter as measured on USNM 594968. Mesistele, dististele, and means of attachment unknown.

Etymology.—*Americanus* recognizes the first species of *Temnocrinus* described from North America.

Material.—Holotype USNM 594967; paratypes USNM 594968 and 594969.

Occurrence.—Brassfield Formation, Rhuddanian (Llandovery, Silurian), Bath County, Kentucky.

Remarks.—USNM 594967 is a large, nearly complete crown measuring 54 mm high. USNM 594968 is a slightly smaller individual preserving the calyx, the anal series and lower brachitaxa, and parts of the three most proximal columnals. USNM 594969 is a much smaller, immature specimen preserving a crushed calyx, a few free brachials, and the six most proximal columnals. All three specimens have been compacted to some degree.

Previously, *Temnocrinus* was represented only by the type species. With the addition of *T. americanus*, amendment of the generic concept is required. Specimens described herein closely resemble *T. tuberculatus* in the overall construction of the calyx and the anal series. Characters considered important for placement within *Temnocrinus* include: the radianal as the lower part of a compound radial in the primitive position directly below the C ray, truncated distal margin of the CD basal supporting the anal X, anal series articulating with the C ray, multiple interradial plates, presence of intrabrachials, two primibrachials, elongate form of the crown, general shape and plate sculpture of the brachials, sinuous sutures between brachials, and mode of arm branching.

Temnocrinus americanus is distinguished from *T. tuberculatus* in having an overall more elongated crown with more interradial plates and fixed brachials, more intrabrachials, a somewhat smoother plate sculpture on the calyx and lower brachials, and more pronounced lateral brachial nodes.

The crown of *T. americanus* is more elongate overall owing to several factors. Infrabasals are taller and not covered by the proximal column as in *T. tuberculatus*, giving *T. americanus* more of a cone-shaped aboral cup, versus a low bowl-shaped aboral cup in *T. tuberculatus*. The three known specimens of *T. americanus* have four secundibrachials where known, with one exception on one branch in the A ray of USNM 594968, where there must have been more than four because although only four are preserved, the fourth secundibrachial is non-axillary. *Temnocrinus tuberculatus* typically has only three secundibrachials, although the number can vary from one to seven (Donovan et al., 2010), and specimens with four secundibrachials are known (Springer, 1920; Donovan et al., 2010). The anal series of *T. americanus* is similarly elongated, with the anal X followed by as many as eight or more plates, whereas in *T. tuberculatus*, the anal X is followed by four or five plates (Donovan et al., 2010). The holotype of *T. americanus*, USNM 594967, is a large specimen with a total crown height of 54 mm, whereas in *T. tuberculatus* a large crown height is

40 mm (Springer, 1920), and crown height varies between 17 and 50 mm (Donovan et al., 2010).

Interradial plates are more numerous in *T. americanus* than in the type species, with two to four ranges of plates in the regular interareas. *Temnocrinus tuberculatus* has two ranges, with one large plate in the first range and two smaller plates in the second (Springer, 1920; Donovan et al., 2010), which are rarely in contact with one another (Donovan et al., 2010). Interradial plates extend somewhat higher in *T. americanus*, reaching the first or second secundibrachial; in *T. tuberculatus* interradial plates do not rise above the primibrachitaxis.

Intrabrachial plates are also more numerous in *T. americanus* and extend higher in the crown, with intrabrachials between both secundibrachials and tertibrachials. *Temnocrinus tuberculatus* commonly has a single small intrabrachial between the first and second secundibrachial (Donovan et al., 2010). Intrabrachials in *T. americanus* are between the second secundibrachials and extending distally to the first tertibrachials; more distally, intrabrachials are between the third to fifth tertibrachials. Because the intrabrachial plates do not immediately follow the axillary in *T. americanus*, the arms in the second and third brachitaxes are somewhat more firmly joined together than in *T. tuberculatus*. These extra intrabrachial plates may have provided *T. americanus* with additional rigidity in the arms.

Plate sculpture includes tubercles for both species; however, *T. americanus* has smooth calyx plates, except for the distal one to two ranges of interradial plates and the anal plates above the anal X that are articulated with the C ray, which bear tubercles. *Temnocrinus americanus* also has tubercles on the intrabrachial plates and on the brachials in the secundibrachitaxis through the quartibrachitaxis. *Temnocrinus tuberculatus* has tubercles that also cover calyx plates, and tubercles aligned in horizontal rows on radials and lower brachials, tending to coalesce into vertical ridges on higher brachials (Springer, 1920). Springer (1920) noted considerable variation in the degree of ornamentation of *T. tuberculatus*, with some specimens almost smooth, but he attributed some of this to preservational differences. Preservation does not appear to account for the smoother calyx of *T. americanus*, as all three specimens are consistent, and tubercles are preserved as noted above on the same specimens.

Lateral brachial nodes on *T. americanus* are more pronounced than on *T. tuberculatus*; however, Springer (1920) noted that the lateral nodes, which he likened to “wing-like buttresses,” were present to a greater or lesser degree in most specimens of *T. tuberculatus* and were not always apparent in side view because of the roundedness of the brachials. Springer illustrated a large individual of *T. tuberculatus* with the lateral brachial nodes quite prominent (Springer, 1920, pl. 16, fig. 15). All three specimens of *T. americanus* have these lateral nodes, including the smallest individual.

Although arms of both species of *Temnocrinus* are distinctive, they also appear to be somewhat convergent with those of *Protaxocrinus interbrachiatus* (Angelin, 1878) from the Wenlock of Gotland. This species, which also has arms with tubercles and lateral nodes, was illustrated by Springer (1920, pl. 45, figs. 13–16). *Protaxocrinus interbrachiatus* is easily distinguished from the two species of *Temnocrinus*, however, by its very different CD basal, which has a notch on the distal right side where the anal X is sutured. Only the first three plates

of the anal series following the anal X in *P. interbrachiatus* are articulated with the C ray, whereas in *Temnocrinus*, the entire series is in sutural contact with the C ray.

Although Flexibilia are typically among the rarest of crinoids, with many species represented by only a few specimens (Springer, 1920), *T. tuberculatus* is abundant in the well-collected Much Wenlock Limestone at Dudley (Springer, 1920; Ramsbottom, 1953). *Temnocrinus tuberculatus* was eagerly sought by collectors (Springer, 1920) and at least 254 specimens are present in British museums alone (Donovan et al., 2010).

Crinoid genera in the Silurian were commonly cosmopolitan, with many genera common to England, Sweden, and North America (Holland, 1971; Sevastopulo et al., 1989). The genus *Temnocrinus*, however, was previously known only from the type species, *T. tuberculatus*, which in turn was known only from the type horizon and locality, the Much Wenlock Limestone Formation (Homerian, Wenlock) at Dudley, England (Springer, 1920; Ramsbottom, 1953; Webster, 2003; Donovan et al., 2010). The addition of *T. americanus* extends the geographic range of *Temnocrinus* to include North America. The presence of *T. americanus* in the lower Brassfield Formation extends the chronologic range of the genus from the Rhuddanian to the Homerian.

Crinoidea Incertae Sedis

Remarks.—In addition to the holdfast of *Stereoaster squamosus*, other types of holdfasts are also present in this fauna, although they are disassociated from crown material and cannot be identified. This material is deposited as USNM 595074, and includes simple discoidal holdfasts (both digitate and non-digitate) attached to an encrusting bryozoan or free. Larger terminal holdfasts with branching pseudocirri are also present.

Conclusions

A new low-diversity echinoderm fauna, comprised of six pelmatozoan taxa, three of which are new, is described from the lower Brassfield Formation (Rhuddanian) of east-central Kentucky. The fauna also contains a previously described asteroid. Although the Kentucky fauna contains new taxa, the echinoderms are very similar to previously described faunas from the area of Dayton, Ohio that are Aeronian in age. This new fauna represents one of the earliest examples of the Llandovery crinoid faunal recovery after the echinoderm extinctions that accompanied the latest Ordovician glaciation.

Acknowledgments

We thank Dan Levin for his normal efficiency, and S.K. Donovan and G.D. Sevastopulo whose careful reviews greatly improved this manuscript.

References

- Angelin, N.P., 1878, Iconographia Crinoideorum in stratis Sueciae Siluricis fossilium: Samson and Wallin, Holmiae, 62 p.
- Ausich, W.I., 1984a, Calceocrinids from the early Silurian (Llandoveryan) Brassfield Formation of southwestern Ohio: *Journal of Paleontology*, v. 58, p. 1167–1185.
- Ausich, W.I., 1984b, The genus *Clidochirus* from the early Silurian of Ohio (Crinoidea, Llandoveryan): *Journal of Paleontology*, v. 58, p. 1341–1346.
- Ausich, W.I., 1985, New crinoids and revision of the superfamily Glyptocrinacea (early Silurian, Ohio): *Journal of Paleontology*, v. 59, p. 793–808.
- Ausich, W.I., 1986a, New camerate crinoids of the suborder Glyptocrinina from the lower Silurian Brassfield Formation (southwestern Ohio): *Journal of Paleontology*, v. 60, p. 887–897.
- Ausich, W.I., 1986b, Early Silurian inadunate crinoids (Brassfield Formation, Ohio): *Journal of Paleontology*, v. 60, p. 719–735.
- Ausich, W.I., 1986c, Early Silurian rhodocritacean crinoids (Brassfield Formation, Ohio): *Journal of Paleontology*, v. 60, p. 84–106.
- Ausich, W.I., 1986d, Palaeoecology and history of the Calceocrinidae (Palaeozoic Crinoidea): *Palaeontology*, v. 29, p. 85–99.
- Ausich, W.I., 1987a, Brassfield Compsocrinina (early Silurian crinoids) from Ohio: *Journal of Paleontology*, v. 61, p. 552–562.
- Ausich, W.I., 1987b, Revisions of Rowley's Ordovician (?) and Silurian crinoids from Missouri: *Journal of Paleontology*, v. 61, p. 563–578.
- Ausich, W.I., 1998, Phylogeny of Arenig to Caradoc crinoids (Phylum Echinodermata) and suprageneric classification of the Crinoidea: University of Kansas Paleontological Contributions, new series, no. 9, 39 p.
- Ausich, W.I., and Copper, P., 2010, The Crinoidea of Anticosti Island, Québec (late Ordovician to early Silurian): *Palaeontographica Canadiana*, no. 29, 157 p.
- Ausich, W.I., and Deline, B., 2012, Macroevolutionary transition in crinoids following the late Ordovician extinction event (Ordovician to early Silurian): *Palaeogeography, Palaeoclimatology, Palaeoecology*, v. 361–362, p. 38–48, doi:10.1016/j.palaeo.2012.07.022.
- Ausich, W.I., and Dravage, P., 1988, Crinoids from the Brassfield Formation of Adams County, Ohio: *Journal of Paleontology*, v. 62, p. 285–289.
- Ausich, W.I., and Schumacher, G.A., 1984, New lower Silurian rhombiferan cystoid (Echinodermata, Callocystitidae) from southwestern Ohio: *Journal of Paleontology* v. 58, p. 9–15.
- Ausich, W.I., Kammer, T.W., and Baumiller, T.K., 1994, Demise of the middle Paleozoic crinoid fauna: a single extinction event or rapid faunal turnover?: *Paleobiology*, v. 20, p. 345–361.
- Ausich, W.I., Brett, C.E., Hess, H., and Simms, M.J., 1999, Crinoid form and function, in Hess, H., Ausich, W.I., Brett, C.E., and Simms, M.J. *Fossil Crinoids*: Cambridge, U.K., Cambridge University Press, p. 3–30.
- Bather, F.A., 1893, The Crinoidea of Gotland, Part 1, The Crinoidea Inadunata: *Kongliga Svenska Vetenskaps-Akademiens Handlingar*, v. 25, no. 2, 200 p.
- Bather, F.A., 1899, A phylogenetic classification of the Pelmatozoa: British Association for the Advancement of Sciences, Report, 68th Meeting section D (Bristol, 1898), p. 916–923.
- Baumiller, T.K., 1994, Patterns of dominance and extinction in the record of Paleozoic crinoids, in B. David, A. Guille, J.P. Féral, and M. Roux, eds., *Echinoderms Through Time (Echinoderms Dijon)*: Rotterdam, A.A. Balkema, p. 193–198.
- Bernard, F., 1895, *Éléments de Paléontologie*: Paris, J.B. Baillière and Fils, 612 p.
- Blake, D.B., and Etensohn, F.R., 2009, The complex morphology of a new lower Silurian asteroid (Echinodermata): *Journal of Paleontology*, v. 83, p. 63–69.
- Boyarko, D., and Ausich, W.I., 2009, New calceocrinids from the Brassfield Formation of northern Kentucky and southern Ohio: *Southeastern Geology*, v. 46, p. 103–108.
- Brett, C.E., 1981, Terminology and functional morphology of attachment structures in pelmatozoan echinoderms: *Lethaia*, v. 14, p. 343–370, doi: 10.1111/j.1502-3931.1981.tb01110.x.
- Brett, C.E., Cramer, B.D., McLaughlin, P.I., Kleffner, M.A., Showers, W.J., and Thomka, J.R., 2012, Revised Telychian-Sheinwoodian (Silurian) stratigraphy of the Laurentian midcontinent: Building uniform nomenclature along the Cincinnati Arch: *Bulletin of Geosciences*, v. 87, p. 733–753, doi: 10.3140/bull.geosci.1310.
- Broadhead, T.W., and Etensohn, F.R., 1981, A new *Brockocystis* (Echinodermata: Rhombifera) from the lower Silurian (Llandovery) of Kentucky: *Journal of Paleontology*, v. 55, p. 773–779.
- Brower, J.C., 1988, Ontogeny and phylogeny in primitive calceocrinid crinoids: *Journal of Paleontology*, v. 62, p. 917–934.
- Brower, J. C., 1990, Ontogeny and phylogeny of the dorsal cup in calceocrinid crinoids: *Journal of Paleontology*, v. 64, p. 300–318.
- Donovan, S.K., 1993, A Rhuddanian (Silurian, lower Llandovery) pelmatozoan fauna from south-west Wales: *Geological Journal*, v. 28, p. 1–19, doi: 10.1002/gj.3350280102.
- Donovan, S.K., and Harper, D.A.T., 2003, Llandovery Crinoidea of the British Isles, including description of a new species from the Kilbride Formation

- (Telychian) of western Ireland: Geological Journal, v. 38, p. 85–97, doi: 10.1002/gj.934.
- Donovan, S.K., and Lewis, D.N., 2005, Upper Llandovery (Telychian) crinoids (Echinodermata) of Devil's Dingle, Buildwas, Shropshire: Geological Journal, v. 40, p. 343–350, doi: 10.1002/gj.1011.
- Donovan, S. K., and Sevastopulo, G.D., 1989, Myelodactylid crinoids from the Silurian of the British Isles: Palaeontology, v. 32, p. 689–710.
- Donovan, S.K., Lewis, D.N., Crabb, P., and Widdison, R.E., 2008, A field guide to the Silurian Echinodermata of the British Isles: Part 2 – Crinoidea, minor groups and discussion: Proceedings of the Yorkshire Geological Society, v. 57, p. 29–60.
- Donovan, S. K., Lewis, D.N., Fearnhead, F.E., and Widdison, R.E., 2009, The British Silurian Crinoidea, Part 1, Introduction and Disparida: Monographs of the Palaeontographical Society, London, v. 163, n. 632, 51 p.
- Donovan, S.K., Widdison, R.E., Lewis, D.N., and Fearnhead, F.E., 2010, The British Silurian Crinoidea, Part 2, addendum to part 1 and Cladida: Monographs of the Palaeontographical Society, London, v. 164, p. 47–133.
- Donovan, S.K., Lewis, D.N., Fearnhead, F.E., and Widdison, R.E., 2012, The British Silurian Crinoidea, Part 3, addendum to parts 1 and 2, Camerata and columnals: Monographs of the Palaeontographical Society, London, v. 166, p. 135–259.
- Eckert, J.D., 1984, Early Llandovery crinoids and stelleroids from the Cataract Group (lower Silurian), southern Ontario, Canada: Royal Ontario Museum Life Sciences, Contributions, v. 137, 83 p.
- Eckert, J.D., 1990, The early Silurian myelodactylid crinoid *Eomyelodactylus* Foerste: Journal of Paleontology, v. 64, p. 135–141.
- Eckert, J.D., and Brett, C.E., 2001, Early Silurian (Llandovery) crinoids from the Lower Clinton Group, western New York State: Bulletins of American Paleontology, v. 360, 88 p.
- Ettensohn, F.R., Lierman, R.T., Mason, C.E., Andrews, W.M., Hendricks, R.T., Phelps, D.J., and Gordon, L.A., 2013, The Silurian of central Kentucky, U. S.A.: stratigraphy, palaeoenvironments and palaeoecology: Memoirs of the Association of Australasian Palaeontologists, v. 44, p. 159–189.
- Fearnhead, F.E., 2009, Systematics and palaeobiology of the British Llandovery Crinoidea [Ph.D. thesis]: Birkbeck College, University of London, 307 p.
- Fearnhead, F.E., and Donovan, S.K., 2007a, New crinoids (Echinodermata) from the Llandovery (Lower Silurian) of the British Isles: Palaeontology, v. 50, p. 905–915, doi: 10.1111/j.1475-4983.2007.00681.x.
- Fearnhead, F.E., and Donovan, S.K., 2007b, A cladid crinoid (Echinodermata) from the Llandovery (lower Silurian) of the Girvan district, SW Scotland: Scottish Journal of Geology, v. 43, no. 1, p. 180.
- Fearnhead, F.E., and Donovan, S.K., 2007, A robust crinoid from the Llandovery (lower Silurian) of Norbury, Shropshire: systematics, palaeoecology and taphonomy: Proceedings of the Geologists' Association, v. 118, p. 339–345.
- Fearnhead, F.E., and Harper, D.A.T., 2007, *Petalocrinus* (Echinodermata, Crinoidea) from the Llandovery (lower Silurian; Rhuddanian) of the Girvan District, SW Scotland: Scottish Journal of Geology, v. 43, p. 69–74.
- Foerste, A.F., 1919, Echinodermata of the Brassfield (Silurian) Formation of Ohio: Bulletin of the Scientific Laboratories of Denison University, v. 19, p. 3–31.
- Gordon, L.A., and Ettensohn, F.R., 1984, Stratigraphy, depositional environments, and regional dolomitization of the Brassfield Formation (Llandoveryan) in east-central Kentucky: Southeastern Geology, v. 25, p. 101–115.
- Hall, J., 1852, Palaeontology of New York, Containing descriptions of the organic remains of the lower middle division of the New-York system, Natural History of New York, 6. D: New York, Appleton and Co. and Wiley and Putnam and Boston, Gould, Kendall, & Lincoln, v. 2, 362 p.
- Holland, C.H., 1971, Silurian faunal provinces?, in Middlemiss, F.A., Rawson, P.F., and Newell, G., eds., Faunal Provinces in Space and Time: Geological Journal Special Issue, v. 4, p. 61–76.
- McLaughlin, P.L., Cramer, B.D., Brett, C.E., and Kleffner, M.A., 2008, Silurian high-resolution stratigraphy on the Cincinnati Arch: Progress on recalibrating the layer-cake, in A. H. Maria and R. C. Counts, eds., From the Cincinnati Arch to the Illinois Basin: Geological field excursions along the Ohio River valley: Geological Society of America Field Guide, 12, p. 119–180.
- Meek, F. B., and Worthen, A.H., 1869, Descriptions of new Crinoidea and Echinoidea, from the Carboniferous rocks of the western states, with a note on the genus *Onychaster*: Academy of Natural Sciences of Philadelphia Proceedings, v. 21, p. 67–83.
- Miller, J.S., 1821, A Natural History of the Crinoidea or lily-shaped animals, with observations on the genera *Asteria*, *Euryale*, *Comatula*, and *Marsupites*: Bristol, C. Frost, 150 p.
- Moore, R.C., 1962, Revision of Calceocrinidae: University of Kansas Paleontological Contributions, Echinodermata, Article 4, 40 p.
- Moore, R.C., and Laudon, L.R., 1943, Evolution and classification of Paleozoic crinoids: Geological Society of America Special Papers, v. 46, 153 p.
- Paul, C.R.C., and Donovan, S.K., 2011, A review of the British Silurian cystoids: Geological Journal, v. 46, p. 434–450, 10.1002/gj.1287.
- Peters, S.E., and Ausich, W.I., 2008, A sampling-adjusted macroevolutionary history for Ordovician-Silurian crinoids: Paleobiology, v. 34, no. 1, p. 104–116.
- Ramsbottom, W.H.C., 1953, The British Lower Palaeozoic Crinoidea [Ph.D. thesis]: University of London, 290 p.
- Rexroad, C.B., 1967, Stratigraphy and conodont paleontology of the Brassfield (Silurian) in the Cincinnati Arch area: Indiana Geological Survey Bulletin, v. 33, 64 p.
- Rexroad, C.B., Branson, E.R., Smith, M.O., Summerson, C., and Boucot, A.J., 1965, The Silurian formations of east-central Kentucky and adjacent Ohio: Kentucky Geological Survey Bulletin, v. 2, Series 10, 34 p.
- Rexroad, C.B., and Kleffner, M.A., 1984, The Silurian stratigraphy of east-central Kentucky and adjacent Ohio, in Rast, N. and Hays, H.B., eds., Field trip guides for the Geological Society of America Annual Meeting, Southeastern and North-Central Sections: Department of Geology and Kentucky Geological Survey, University of Kentucky, Lexington, p. 44–65.
- Roemer, C.F., 1852–54, Erste Periode, Kohlen-Gebirge, in Bronn, H.G., Lethaea Geognostica, 1851–1856, (third edition): Stuttgart, E. Schweizerbart, v. 2, p. 210–291.
- Sevastopulo, G.D., Ausich, W.I., and Franzén-Bengtson, C., 1989, Echinoderms, in Holland, C.H., and Bassett, M.G., eds., A Global Standard for the Silurian Series: Cardiff, National Museum of Wales, p. 264–267.
- Schneider, K.A., and Ausich, W.I., 2002, Paleoecology of framebuilders in early Silurian reefs (Brassfield Formation, southwestern Ohio): Palaios, v. 17, p. 237–248, doi: 10.1669/0883-1351(2002)017 <0237:POFIES > 2.0.CO;2.
- Sheehy, T.J., 1981, Carbonate Petrology of Carbonate Buildups in the Brassfield Formation near Fairborn, Ohio [M.S. Thesis]: Dayton, Ohio, Wright State University, 52 p.
- Springer, F., 1902, On the crinoid genera *Sagenocrinus*, *Forbesiocrinus*, and allied forms: American Geologist, v. 30, p. 88–97.
- Springer, F., 1913, Crinoidea, in von Zittel, K.A. Text-book of Paleontology, translated and edited by C. R. Eastman. 2nd edition: London, Macmillan and Co., Ltd., 1, p. 173–243.
- Springer, F., 1920, The Crinoidea Flexibilia: Smithsonian Institution Publication, v. 2501, 486 p.
- Springer, F., 1926, American Silurian Crinoids: Smithsonian Institution Publication, v. 2872, 239 p.
- Sumrall, C.D., 2002, A new species of *Anartiocystis* (Callocystitidae, Glyptocystitida) from the Brassfield Formation of central Kentucky: Journal of Paleontology, v. 76, p. 918–920.
- Ubaghs, G., 1953, Classe des Crinoïdes, in Piveteau, J., ed., Traité de Paléontologie: Paris, Masson and Cie, v. 3, p. 658–773.
- Ubaghs, G., 1978, General morphology, in Moore, R.C., and Teichert, C., eds., Treatise on Invertebrate Paleontology. Pt. T. Echinodermata 2: Boulder and Lawrence, Geological Society of America and University of Kansas; p. T58–T216.
- von Zittel, K.A., 1895, Grundzüge der Palaeontologie (Palaeozoologie), 1st edition, R.: Oldenbourg, München, 971 p.
- Wachsmuth, C., and Springer, F., 1880–1886, Revision of the Palaeocrinoidea. Proceedings of the Academy of Natural Sciences of Philadelphia Pt. I. The families Ichthyocrinidae and Cyathocrinidae (1880), p. 226–378 (separate repaged p. 1–153). Pt. II. Family Sphaeroidocrinidae, with the sub-families Platycrinidae, Rhodocrinidae, and Actinocrinidae (1881), p. 177–411, (separate repaged, p. 1–237). Pt. III, Sec. 1. Discussion of the classification and relations of the brachiote crinoids, and conclusion of the generic descriptions (1885), p. 225–364 (separate repaged, 1–138). Pt. III, Sec. 2. Discussion of the classification and relations of the brachiote crinoids, and conclusion of the generic descriptions (1886), p. 64–226 (separate repaged to continue with section 1, 139–302).
- Webster, G.D., 2003, Bibliography and index of Paleozoic crinoids, coronates, and hemistreptocrinoids: Geological Society of America Special Paper 363.
- Webster, G.D., and Maples, C.G., 2008, Cladid crinoid radial facets, brachials, and arm appendages: A terminology solution for studies of lineage, classification, and paleoenvironment, in W.I. Ausich and G.D. Webster, eds., Echinoderm Paleobiology: Bloomington, Indiana University Press, p. 197–226.
- Witzke, B.J., and Strimple, H.L., 1981, Early Silurian crinoids of eastern Iowa: Proceedings of the Iowa Academy of Science, v. 88, p. 101–137.

Accepted 20 June 2014