

mind that simpler machines have reduced memory requirements. I frequently use the Z88 with a modem (and an additional amateur radio 'packet' modem from the car) without undue trouble, although for heavy use a set of rechargeable pencil batteries has usefully added approximately 10 hours of operation.

When considering compatibility, remember that simple text can be transferred between any type of computer. IBM clone compatibility is measured in percentage terms, often scoring less than 100. The current explosion of faster computer processors gives the software writers a choice, either write programs that ignore the newer, expensive features, or cut out most of your potential customer market, and pass on the costs to those that have the faster machines.

If it is your hobby, buy computer potential. If it is for your work, you should buy only the computer power you need. It will all be archaic soon. A lesser machine that is demonstrated, has local service arrangements and some telephone support can still be a better deal overall.

CARL S. LITTLEJOHNS

North Wales Hospital
Denbigh, Clwyd LL16 5SS

Cigarettes and psychiatric patients

DEAR SIRS

I read with interest Dr Brown's paper on cigarette smoking among psychiatric out-patients (*Psychiatric Bulletin*, July 1991, 15, 413–414) in which he reported a prevalence of "about one and a half times that of the general public" for both sexes. He states that their residency in the community at the time of the study rules out the effects of institutionalisation. How can we judge this for ourselves when he does not tell us if and for how long these people had been in-patients in the past. I found that 24% of psychiatric in-patients started their smoking careers as in-patients. In-patients also have very high consumption of tobacco (Masterson & O'Shea, 1984).

BRIAN O'SHEA

Newcastle Hospital
Greystones, Co Wicklow
Republic of Ireland

Reference

MASTERSON, R. & O'SHEA, B. (1984) Smoking and malignancy in schizophrenia. *British Journal of Psychiatry*, 145, 429–432.

DEAR SIRS

I accept Dr O'Shea's point that current residence in the community does not exclude previous institutional experience as an aetiological factor in cigarette smoking. A number of patients in the study had been in-patients in the past; however all were

living independently in the community at the time of the study. None were from the group who have recently moved into the community as a result of the closure of long stay hospital beds, as these patients came under the care of a different service. I therefore feel that institutionalisation in a narrow sense cannot explain the high levels of smoking.

Dr O'Shea's figure of the number of patients who start smoking as in-patients is surely cause for embarrassment, at least. We have a degree of responsibility for the physical health of our patients, smokers and non-smokers alike, especially those involuntary patients who may be unable to leave a smoky environment. The risks of passive smoking are now recognised. Perhaps self-interest may prompt a re-examination of our attitudes to smoking.

STEVE BROWN

Millbrook
King's Mill Hospital
Sutton-in-Ashfield NG17 4JL

The meaning of 'psychosis'

DEAR SIRS

The term 'psychosis' is in widespread use yet little is known about what it means to health workers. A study was performed among 96 members of various professional groups working within Exeter Health Authority (all 41 psychiatric medical staff, eight District General Hospital (DGH) consultants in acute specialities, 36 psychiatric nurses, five occupational therapists and six approved social workers). Each was sent an unstructured questionnaire asking them to write their understanding of the term psychosis. All replies were anonymous and varied in length from 300–500 words. From each it was possible to identify a definition, often with several items, which could be listed, as well as a variable length of commentary. Many definitions contained items expressed in so similar a fashion that they could be grouped together with "disturbance of contact with reality" the most frequent, appearing in 25 replies and "loss of insight" the next commonest feature (in 15 definitions). Thirteen respondents included "disorders of thinking, experience or perception". Half the respondents (26) had only one item in their definition and a further 19 had only two items.

Historically the term 'psychosis' is imprecise, from its appearance early in the 19th century distinguishing mental disorder from neurosis or a functional disease of nerves to the view of Gelder *et al* (1989) of psychosis as meaning broadly the more severe forms of mental illness. DSM-III-R regards psychotic as gross impairment in reality testing and a creation of a new reality while the ICD-9 definition contains three elements – gross interference with insight, with ability