

TWO NON-VANISHING RESULTS CONCERNING THE ANTI-CANONICAL BUNDLE

NIKLAS MÜLLER 

Abstract. Let (X, Δ) be a klt threefold pair with nef anti-log canonical divisor $-(K_X + \Delta)$. We show that $\kappa(X, -(K_X + \Delta)) \geq 0$. To do so, we prove a more general equivariant non-vanishing result for anti-log canonical bundles, which is valid in any dimension.

§1. Introduction

Let (X, Δ) be a klt pair with nef log-canonical divisor $K_X + \Delta$. Recall that the celebrated *Non-Vanishing conjecture* predicts that $\kappa(X, K_X + \Delta) \geq 0$, i.e. that there exists an integer $m \geq 1$ such that $m(K_X + \Delta)$ is Cartier and

$$H^0\left(X, \mathcal{O}_X(m(K_X + \Delta))\right) \neq 0.$$

Along with the *Abundance conjecture*, which predicts that $K_X + \Delta$ is even semiample, the Non-Vanishing conjecture has attracted much attention. One might be tempted to ponder what happens on the opposite end of the spectrum, namely, when the anti-log canonical divisor $-(K_X + \Delta)$ is nef. One quickly realises that the situation is somewhat different: Already the example of \mathbb{P}^2 blown-up in 9 points shows that $-K_X$ might be nef but fail to be semiample, see [17]. On the other hand, in [1] the authors classified rather explicitly smooth projective threefolds with nef anti-canonical class and observed a posteriori that $\kappa(X, -K_X) \geq 0$, at least when X is rationally connected, see also [31] and [32].

Recently, Lazić–Matsumura–Peternell–Tsakanikas–Xie [21] studied varieties with nef anticanonical bundle with a view towards the recently proposed generalised Non-Vanishing conjecture [20], [14]. In particular, they proved that $\kappa(X, -(K_X + \Delta)) \geq 0$ if (X, Δ) is a rationally connected threefold pair with nef anti-log canonical bundle and they asked whether the same conclusion holds more generally, assuming only that $-(K_X + \Delta)$ is nef. In this paper, we want to give the following affirmative answer to this question:

THEOREM A. *Let (X, Δ) be a projective klt pair and assume that the anti-log canonical divisor $-(K_X + \Delta)$ is nef. Let (F, Δ_F) denote a general fibre of the MRC-fibration of X . If $-(K_F + \Delta_F)$ is semiample, then*

$$\kappa(X, -(K_X + \Delta)) \geq 0.$$

Despite giving only a partial answer, Theorem A applies in many cases. For example, as can be seen by using [15], it applies whenever F is a surface with enough automorphisms. Thus, combining our result with [14] and [21] to deal with the other cases enables us to fully settle the three-dimensional case:

Received March 26, 2024. Revised June 14, 2024. Accepted July 28, 2024.

2020 Mathematics subject classification: Primary 14E30; Secondary 14M25, 14J30.

Keywords: non-vanishing, anti-canonical bundle, GIT.

While working on this project, the author was supported by the DFG Research Training Group 2553 “Symmetries and Classifying Spaces: Analytic, Arithmetic and Derived”



COROLLARY B. *Let (X, Δ) be a projective klt threefold pair such that $-(K_X + \Delta)$ is nef. Then*

$$\kappa(X, -(K_X + \Delta)) \geq 0.$$

Besides the results obtained in [9] and [23], which form the backbone of our strategy, the main ingredient in our proof of Theorem A is the following “equivariant Non-Vanishing theorem,” which allows us to lift sections from F to X as explained in Section 2. It holds in greater generality and might be of independent interest:

THEOREM C (Equivariant Non-Vanishing for Anti-Canonical Divisors). *Let (X, D) be a projective sub-log canonical pair and assume that $-(K_X + D)$ is semiample. Then for any commutative, linear algebraic subgroup $H \subseteq \text{Aut}(X, D)$ there exists an integer $m \geq 1$ such that $-m(K_X + D)$ is Cartier and*

$$H^0\left(X, \mathcal{O}_X(-m(K_X + D))\right)^H \neq 0.$$

We will abbreviate this by writing $\kappa(X, -(K_X + D))^H \geq 0$.

Here, $\text{Aut}(X, D) \subseteq \text{Aut}(X)$ denotes the subgroup of automorphisms $\varphi \in \text{Aut}(X)$ leaving D invariant as a subset of X . Then $\text{Aut}(X, D)$ acts on $H^0(X, \mathcal{O}_X(-m(K_X + D)))$ in a natural way and we denote by $H^0(X, \mathcal{O}_X(-m(K_X + D)))^H$ the space of sections which are fixed under this action. For our terminology concerning pairs we refer the reader to Section 1.2 below.

We would like to emphasise that the conclusion of Theorem C is astonishing to the author in several different ways: First, perhaps somewhat unexpectedly, the crucial assumption in Theorem C is the log canonicity and not the semipositivity of $-(K_X + D)$, as the conclusion of Theorem C can fail even if (X, D) is log smooth and if $-(K_X + D)$ is ample as soon as (X, D) is no longer sub-log canonical, see Example 3.24.

Moreover, it is noteworthy that the action of H on the space of sections of $-m(K_X + D)$ is usually far from being trivial and already in very simple examples the ring of invariant sections of $-(K_X + D)$ might be much smaller than the ring of all sections, cf. Example 3.23. Nevertheless, Theorem C shows that there always do exist invariant sections. In particular, in case $\kappa(X, -(K_X + D)) = 0$ the action *must* be trivial after all.

Finally, note that the conclusion of Theorem C certainly fails if one allows for more general subgroups of $\text{Aut}(X, D)$, cf. Example 3.22.

1.1 Overview of the methods used in the proof of Theorem C

By far the most important special case of Theorem C is the one where (X, D) is log smooth and $H \cong (\mathbb{C}^\times)^r$ is an algebraic torus. Indeed, the general case is readily reduced to this one; the precise argument is contained in Section 4.

Under these assumptions, the main idea of the proof is to convert the global question of whether $\kappa(X, -(K_X + D))^H \geq 0$ into a problem in convex geometry determined by the local action of H near the fixed points in X . This method of computing global invariants of an action is well-known and heavily used in Geometric Invariant Theory and Symplectic Reduction. However, it is classically only developed for ample line bundles, while we will also need it for semiample ones. This case seems to have received little—if any—interest so far. Consequently, we re-prove the statements we need in detail.

In the log Fano case, the result boils down to a combination of some explicit local computations, some classical results about moment polytopes and a little piece of convex geometry. The semiample case then follows via some perturbation techniques. All of this will be explained in detail in Section 3.

1.2 Conventions

Throughout the paper, we work over the field \mathbb{C} of complex numbers.

Since the terminology concerning pairs is slightly ambiguous at times in the literature, let us fix explicitly the notation we will use: For us, a *pair* (X, D) consists of a normal variety X and a \mathbb{Q} -Weil divisor $D = \sum a_i D_i$ on X with the property that $K_X + D$ is \mathbb{Q} -Cartier. Here, D may or may not be effective.

We call (X, D) *log smooth* if X is smooth and if D has SNC support. A log smooth pair (X, D) will be called *sub-log canonical* if $a_i \leq 1$ for all i , and *log canonical* if additionally D is required to be effective, $0 \leq a_i \leq 1$. In general, a not necessarily log smooth pair (X, D) is called (sub-)log canonical if for some/any log resolution $f: \hat{X} \rightarrow X$ the log smooth pair (\hat{X}, \hat{D}) is so, where \hat{D} is determined by the rules $K_{\hat{X}} + \hat{D} \sim_{\mathbb{Q}} f^*(K_X + D)$ and $f_*\hat{D} = D$. Similarly, in the definition of a *sub-klt* pair we do not require D to be effective, while in the definition of a *klt* pair we do. In other words, our definition of a sub-log canonical pair coincides precisely with the definition of a log canonical pair in [18]; similarly for (sub-)klt pairs.

§2. Reduction of Theorem A to the equivariant non-vanishing

In this section, we reduce the proof of Theorem A to the equivariant non-vanishing problem Theorem C. The main technical tool we will use is the following structure theorem which was recently obtained in [23], building on the previous works [27], [28], [33], [22], [6], [8], [7], [9], [11], [5] and [30] among others. We refer to [10] for the definition of *MRC-fibrations* (also known as *rational quotients*).

THEOREM 2.1 (Matsumura–Wang [23]). *Let (X, Δ) be a projective klt pair with nef anti-log canonical divisor $-(K_X + \Delta)$. Then there exists a finite quasi-étale cover $\pi: X' \rightarrow X$ such that X' admits a holomorphic, everywhere defined, MRC-fibration $f: X' \rightarrow Y$. Moreover, the following hold true:*

- (1) *Every component of $\Delta' := \pi^* \Delta$ is dominant over Y , that Δ' is ‘horizontal’.*
- (2) *The pair $(Y, 0)$ is klt and $K_Y \sim_{\mathbb{Q}} 0$.*
- (3) *The general fibre $(F, \Delta_F := \Delta'|_F)$ is a (connected) rationally connected klt pair with nef anti-log canonical divisor $-(K_F + \Delta_F)$.*
- (4) *The morphism f is a locally constant fibration, that is, there exists a group homomorphism $\rho: \pi_1(Y) \rightarrow \text{Aut}(F, \Delta_F)$ and an isomorphism over Y*

$$(X', \Delta') \cong \left(\tilde{Y} \times (F, \Delta_F) \right) / \pi_1(Y),$$

where \tilde{Y} denotes the universal cover of Y and where $\pi_1(Y)$ acts diagonally in the natural way on \tilde{Y} and through ρ on (F, Δ_F) .

In view of Theorem 2.1 we expect $-(K_X + \Delta)$ to have sections if and only if $-(K_F + \Delta_F)$ admits sections which are invariant under the action of ρ . This is made precise below:

THEOREM 2.2. *Let (F, Δ_F) be a rationally connected projective klt pair and assume that the anti-log canonical divisor $-(K_F + \Delta_F)$ is nef. Then the following assertions are equivalent:*

- (1) *For any projective klt pair (X, Δ) with nef anti-log canonical divisor $-(K_X + \Delta)$ and whose MRC-fibration has (F, Δ_F) as its general fibre it holds that*

$$\kappa(X, -(K_X + \Delta)) \geq 0.$$

- (2) *There exists a maximal algebraic torus $T \subseteq \text{Aut}(F, \Delta_F)$ such that*

$$\kappa\left(F, -(K_F + \Delta_F)\right)^T \geq 0.$$

Recall that an *algebraic torus* is an algebraic group which is isomorphic to $(\mathbb{C}^\times)^r$ for some integer r . In order to prove Theorem 2.2 we need the following elementary result; we include a proof for the sake of completeness:

LEMMA 2.3. *Let G be a linear algebraic group acting algebraically on a finite dimensional vector space V . Then the following assertions are equivalent:*

- (1) *For any commutative, algebraic subgroup $H \subseteq G$ it holds that $V^H \neq 0$.*
 (2) *For any algebraic torus $T \subseteq G$ it holds that $V^T \neq 0$.*
 (3) *For some maximal algebraic torus $T \subseteq G$ it holds that $V^T \neq 0$.*

Proof. Clearly (1) \Rightarrow (2). Regarding the converse, let $H \subseteq G$ be any commutative, algebraic subgroup. Then H splits as

$$H = T \times U,$$

where T is an algebraic torus and U is unipotent, cf. [24, Theorem 16.13]. Consequently,

$$V^H = V^{(T \times U)} = (V^T)^U.$$

Here, we used in the last step that the elements of U, T commute so that the action of U preserves V^T . Now, by our assumption $V^T \neq 0$. Since (essentially by definition) any algebraic action of a unipotent group on a non-trivial vector space fixes at least one non-trivial vector we conclude that $V^H \neq 0$.

Finally, the equivalence of (2) and (3) is clear as any torus $T \subseteq G$ is contained in a maximal one and since any two maximal tori in G are conjugate to each other, see [24, Theorem 17.10]. \square

Proof of Theorem 2.2. First, let us prove that (2) \Rightarrow (1). Fix (X, Δ) a projective klt pair which has (F, Δ_F) as general fibre of its MRC-fibration. We want to show that $\kappa(X, -(K_X + \Delta)) \geq 0$. Note that to do so we, may replace (X, Δ) by arbitrary quasi-étale covers, cf. [29, Theorem 5.13]. In particular, by Theorem 2.1, we may assume that the MRC-fibration $f: X \rightarrow Y$ is a locally constant fibration with fibre (F, Δ_F) and such that $K_Y = 0$. Fix a group homomorphism $\rho: \pi_1(Y) \rightarrow \text{Aut}(F, \Delta_F)$ such that

$$(X, \Delta) = \left(\tilde{Y} \times (F, \Delta_F) \right) / \pi_1(Y).$$

CLAIM. We may assume that the Zariski closure $H := \overline{\text{Im} \rho} \subseteq \text{Aut}(F)$ is a connected, commutative, linear algebraic group. \square

Indeed, as X is a projective, the image of $\pi_1(Y) \xrightarrow{\rho} \text{Aut}(F) \rightarrow \text{Aut}(F)/\text{Aut}^0(F)$ is finite, see, for example, [25, Lemma 3.4]. Hence, H has only finitely many connected components. Replacing Y by the finite (!) étale cover corresponding to

$$\ker\left(\pi_1(Y) \rightarrow H/H^0\right) \subseteq \pi_1(Y)$$

and replacing X by $X' := X \times_Y Y'$, we may assume that $H = H^0$ is connected. Moreover, as F is rationally connected, $\text{Pic}^0(F) = 0$ and so $G := \text{Aut}^0(F)$ is linear algebraic, see [2, Corollary 2.18]. Thus, $H \subseteq G$ is also linear.

Finally, according to [12, Theorem B], there exists a finite quasi-étale cover $Y' \rightarrow Y$ such that $Y' = A \times Z$ splits as a product of an abelian variety A and a variety Z with $\mathcal{O}_Z(K_Z) = \mathcal{O}_Z$ and vanishing augmented irregularity. Since $G = \text{Aut}^0(F)$ is linear algebraic, the image of the map $\rho: \pi_1(Z) \rightarrow G = \text{Aut}^0(F)$ is finite by [12, Remark 1.4]. In other words, after another finite étale cover we may assume that X splits as

$$X \cong \left((\tilde{A} \times F) / \pi_1(A) \right) \times Z =: X' \times Z.$$

As $\mathcal{O}_Z(K_Z) = \mathcal{O}_Z$ has sections, it suffices to prove the assertion for $(X', \Delta|_{X'})$ and we are thus reduced to the case that $Y = A$ is an abelian variety. In particular, in this case $\pi_1(Y)$ is an abelian group.

Let us denote by $H \subseteq G$ the Zariski-closure of $\text{Im}\rho$. Then, by continuity,

$$[H, H] = \left[\overline{\text{Im}\rho}, \overline{\text{Im}\rho} \right] \subseteq \overline{[\text{Im}\rho, \text{Im}\rho]} = \{1\},$$

and so $H \subseteq G$ is also commutative. In summary, H is a connected, commutative, linear algebraic group. This concludes the proof of the *Claim*.

Let us now continue with the proof of (2) \Rightarrow (1). According to Lemma 2.3 and the *Claim* we may find an integer $m \geq 1$ such that $-m(K_F + \Delta_F)$ is Cartier and such that

$$\left(H^0\left(F, \mathcal{O}_F\left(-m(K_F + \Delta_F)\right)\right) \right)^H \neq 0. \quad (2.1)$$

Here, we used that $G = \text{Aut}^0(F)$ is linear algebraic. In what follows, we will prove that the elements of the vector space in (2.1) lift to non-zero sections in $H^0(X, \mathcal{O}_X(-m(K_X + \Delta)))$. Indeed, note that

$$H^0\left(X, \mathcal{O}_X(-m(K_X + \Delta))\right) = H^0\left(Y, f_*\mathcal{O}_X(-m(K_X + \Delta))\right). \quad (2.2)$$

Now,

$$\begin{aligned} f_*\mathcal{O}_X(-m(K_X + \Delta)) &= f_*\mathcal{O}_X(-m(K_{X/Y} + \Delta)) \\ &= \left(\tilde{Y} \times H^0\left(F, \mathcal{O}_F(-m(K_F + \Delta_F))\right) \right) / \pi_1(Y) \end{aligned} \quad (2.3)$$

is the holomorphically flat vector bundle with fibre $\mathbb{H} := H^0(F, \mathcal{O}_F(-m(K_F + \Delta_F)))$ and monodromy representation $\pi_1(Y) \xrightarrow{\rho} \text{Aut}(F, \Delta_F) \rightarrow \text{GL}(\mathbb{H})$, c.f. [21, Proposition 6.3(b)]. In particular,

$$\begin{aligned} H^0\left(X, \mathcal{O}_X(-m(K_X + \Delta))\right) &\stackrel{(2.2)}{=} H^0\left(Y, f_*\mathcal{O}_X(-m(K_{X/Y} + \Delta))\right) \\ &\supseteq \left(H^0(F, -m(K_F + \Delta_F))\right)^p \\ &\supseteq \left(H^0(F, -m(K_F + \Delta_F))\right)^H \neq 0. \end{aligned}$$

Here we used in the last line that $\text{Im}\rho \subseteq H$ and (2.1). Thus, (2) \Rightarrow (1) is settled.

Let us now turn to (1) \Rightarrow (2); Given (F, Δ_F) as in the theorem and $T \subseteq \text{Aut}(F, \Delta_F)$ any maximal algebraic torus, our goal is to produce a projective klt pair (X, Δ) which admits a locally constant fibration $f: X \rightarrow Y$ onto a variety Y with $K_Y = 0$ and with fibre (F, Δ_F) such that

$$H^0\left(X, \mathcal{O}_X(-m(K_X + \Delta))\right) = H^0\left(F, \mathcal{O}_F(-m(K_F + \Delta_F))\right)^T.$$

To this end, write $T \cong (\mathbb{C}^\times)^r$. Fix an abelian variety Y of dimension r and $\zeta \in \mathbb{S}^1 \subseteq \mathbb{C}^\times$ any number of absolute value one which is *not* a root of unity. Consider the group homomorphism $\rho: \pi_1(Y) \rightarrow T$ determined by the rule

$$\rho: \pi_1(Y) \rightarrow T, \quad e_{2i-1} \mapsto (1, \dots, 1, \zeta, 1, \dots, 1), \quad e_{2i} \mapsto (1, \dots, 1), \quad \forall i = 1, \dots, r,$$

where e_1, \dots, e_{2r} is some \mathbb{Z} -basis for $\pi_1(Y) \cong \mathbb{Z}^{2r}$. Then $\text{Im}\rho$ is Euclidean dense in the compact group $(\mathbb{S}^1)^r \subseteq (\mathbb{C}^\times)^r \cong T$. In particular, $\text{Im}\rho$ is Zariski dense in T .

Let us consider the complex analytic variety $X := (\tilde{Y} \times F)/\pi_1(Y)$ equipped with the \mathbb{Q} -Weil divisor $\Delta := (\tilde{Y} \times \Delta_F)/\pi_1(Y)$. Then (X, Δ) is a projective klt pair with nef anti-log canonical divisor $-(K_X + \Delta)$, see [25, Theorem 5.1]. By assumption there exists $m \geq 1$ such that

$$0 \neq H^0\left(X, \mathcal{O}_X(-m(K_X + \Delta))\right) = H^0\left(Y, f_*\mathcal{O}_X(-m(K_X + \Delta))\right). \quad (2.4)$$

Now, as in (2.3),

$$f_*\mathcal{O}_X(-m(K_X + \Delta)) = \left(\tilde{Y} \times H^0\left(F, \mathcal{O}_F(-m(K_F + \Delta_F))\right)\right)/\pi_1(Y)$$

is the holomorphically flat vector bundle with fibre $\mathbb{H} := H^0(F, \mathcal{O}_F(-m(K_F + \Delta_F)))$ and monodromy representation $\pi_1(Y) \xrightarrow{\rho} \text{Aut}(F, \Delta_F) \rightarrow \text{GL}(\mathbb{H})$. As $\text{Im}\rho$ is contained in the compact subgroup $(\mathbb{S}^1)^r \subsetneq T = (\mathbb{C}^\times)^r$, this representation is unitary. It follows that all global sections of $f_*\mathcal{O}_X(-m(K_{X/Y} + \Delta))$ are flat, see, for example, [30, Theorem 2.2(b)]. In other words,

$$H^0\left(Y, f_*\left(\mathcal{O}_X(-m(K_{X/Y} + \Delta))\right)\right) = \mathbb{H}^\rho = \mathbb{H}^T.$$

Here, in the last step we used that the action of $\text{Aut}(F)$ on $\mathbb{H} = H^0(F, -m(K_F + \Delta_F))$ is algebraic and that $\text{Im}\rho \subsetneq T$ is Zariski dense by construction. As we know that $H^0(Y, f_*(-m(K_{X/Y} + \Delta))) \neq 0$ by (2.4) we conclude.

§3. Torus-invariant sections on semiample line bundles

In this section, we want to prove Theorem C under the additional assumption that (X, D) is log smooth:

THEOREM 3.1. *Let (X, D) be a projective, log smooth, sub-log canonical pair such that $-(K_X + D)$ is semiample. Then for any algebraic torus $T \subseteq \text{Aut}(X, D)$ it holds that*

$$\kappa(X, -(K_X + D))^T \geq 0.$$

To prove Theorem 3.1, we will employ some discrete methods from toric geometry. Let us start by introducing the following concept:

DEFINITION 3.2. Let X be a normal variety and let $G \subseteq \text{Aut}(X)$ be a subgroup. A coherent sheaf \mathcal{L} on X is said to be G -invariant if $g^*\mathcal{L} \cong \mathcal{L}$ for all $g \in G$. In this case, we say that \mathcal{L} is G -linearizable if the action of G on X may be lifted to an action of G on the total space of \mathcal{L} via sheaf automorphisms. A G -linearization of \mathcal{L} is a choice of such a lift. See [26, Definition 1.6] for a more formal definition of linearizations.

EXAMPLE 3.3. Let X be a normal variety, let $G \subseteq \text{Aut}(X)$ be a subgroup and let $\mathcal{L}, \mathcal{L}_1, \mathcal{L}_2$ be G -linearized coherent sheaves on X . Then also \mathcal{L}^* and $\mathcal{L}_1 \otimes \mathcal{L}_2$ inherit natural G -linearizations.

DEFINITION 3.4. Let X be a normal variety, let $G \subseteq \text{Aut}(X)$ be a subgroup and let \mathcal{L} be a G -linearized, coherent, reflexive \mathcal{O}_X -module of rank one. Then G acts on $H^0(X, (\mathcal{L}^{\otimes m})^{**})$ for any integer m and we will say that that $\kappa(X, \mathcal{L})^G \geq 0$ if and only if

$$\bigoplus_{m=1}^{\infty} H^0\left(X, (\mathcal{L}^{\otimes m})^{**}\right)^G \neq 0.$$

EXAMPLE 3.5. Consider $X = \mathbb{P}^1$ equipped with the line bundle $\mathcal{L} = \mathcal{O}_{\mathbb{P}^1}(-1)$.

- (1) Let $G = \text{Aut}(\mathbb{P}^1) = \text{PGL}_2(\mathbb{C})$. Then \mathcal{L} is clearly G -invariant. However, this action is (rather famously) *not* linearizable, see, for example, [3, Example 4.2.4].
- (2) Let $G = T \subsetneq \text{Aut}(\mathbb{P}^1)$ be the subgroup of (equivalence classes of) diagonal matrices. Then $\mathbb{C}^\times \cong T \subsetneq \text{Aut}(\mathbb{P}^1)$ is a maximal torus acting via $t \cdot [x : y] := [tx : y]$ and this action *is* linearizable. In fact, for any $w \in \mathbb{Z}$ an explicit choice of lifting is provided by $t \cdot (x, y) = (t^{w+1}x, t^w y)$. Here we identify, as per usual, $\mathcal{O}_{\mathbb{P}^1}(-1)|_{[x:y]} = \mathbb{C} \cdot (x, y) \subseteq \mathbb{C}^2$.

EXAMPLE 3.6. Let X be a normal variety and let $G \subseteq \text{Aut}(X)$ be a subgroup. Then the anti-canonical bundle $\mathcal{O}_X(-K_X)$ admits a natural linearization given by push-forward of forms: $g \cdot \tau := g_*(\tau)$. Recall that in local coordinates φ_* is simply defined by the formula

$$\left(\varphi_* \left(f \frac{\partial}{\partial z_1} \wedge \dots \wedge \frac{\partial}{\partial z_n} \right) \right) (\varphi(x)) = f(x) \cdot d\varphi \left(\frac{\partial}{\partial z_1} \right) \wedge \dots \wedge d\varphi \left(\frac{\partial}{\partial z_n} \right),$$

where $\varphi: X \rightarrow X$ is any automorphism.

EXAMPLE 3.7. Let D be a Cartier divisor on X and assume that it is invariant (as a subset of X) under the action of G . Then also $\mathcal{O}_X(D)$ carries a natural G -linearization as a subsheaf of $\mathbb{C}(X)$, which is induced by the pull-back of functions: $g \cdot f := (g^{-1})^* f$. Note that we take the inverse in order to obtain a left-action.

Similarly, Ω_X^1 admits a natural linearization given by $g \cdot df = d(g \cdot f) = d((g^{-1})^* f)$.

In conclusion, whenever (X, D) is a pair, then the line bundle $\mathcal{O}_X(-m(K_X + D))$ carries a natural $\text{Aut}(X, D)$ -linearization for all $m \in \mathbb{Z}$ such that $m(K_X + D)$ is Cartier.

In the following, we will require some standard facts about algebraic tori and their representations which we recall below: Let $T \cong (\mathbb{C}^\times)^r$ be an algebraic torus. Then its

character lattice $M := \text{Hom}(T, \mathbb{C}^\times)$ is a finitely generated free abelian group of rank $\text{rk}M = \dim T$. Explicitly, if $T = (\mathbb{C}^\times)^r$ then

$$\mathbb{Z}^r \xrightarrow{\sim} M = \text{Hom}(T, \mathbb{C}^\times), \quad w = (a_1, \dots, a_r) \mapsto \left(t \mapsto t^w := t_1^{a_1} \cdot \dots \cdot t_r^{a_r} \right).$$

In the sequel, following standard practice, we will consider M as an additive group $(M, +)$. In particular, the neutral element $0 \in M$ corresponds to the trivial homomorphism $t \mapsto 1$. We will also consider the real vector space $M_{\mathbb{R}} := M \otimes_{\mathbb{Z}} \mathbb{R}$.

Now, let $\rho: T \rightarrow \text{GL}(V)$ be an algebraic representation of T on a finite-dimensional complex vector space V (we also say that V is a T -module). Then the action of T can be diagonalised, there exists a \mathbb{C} -basis e_1, \dots, e_m , called a basis of *eigenvectors*, for V such that

$$t \cdot e_i = t^{w_i} \cdot e_i, \quad \forall t \in T$$

for some $w_1, \dots, w_m \in M = \text{Hom}(T, \mathbb{C}^\times)$ called the *weights* of the representation ρ . A representation of T is determined up to isomorphism by its weights.

DEFINITION 3.8. Let X be a normal projective variety, let $T \subseteq \text{Aut}(X)$ be an algebraic torus and let \mathcal{L} be a coherent sheaf on X , linearized for the action of T . Then T acts naturally on the space $H^0(X, \mathcal{L})$. We let $W_\Gamma(X, \mathcal{L}) \subseteq M$ denote the set of weights of this action. Moreover, we denote the convex hull of $W_\Gamma(X, \mathcal{L})$ in $M_{\mathbb{R}}$ by $P_\Gamma(X, \mathcal{L})$ and call this the *section polytope* of \mathcal{L} .

NOTATION 3.9. From now on and for the rest of this section, we let X denote a smooth projective variety and we fix an algebraic torus $T \subseteq \text{Aut}(X)$. Then the set of *fixed points* $X^T := \{x \in X \mid t \cdot x = x, \forall t \in T\}$ is a non-empty, closed, *smooth* subvariety of X , see [24, Theorem 13.1, Proposition 13.20]. We let $X^T = Y_1 \sqcup \dots \sqcup Y_c$ denote the decomposition of X^T into its connected components. Moreover, for any $i = 1, \dots, c$ let us fix a point $y_i \in Y_i$.

DEFINITION 3.10. Let \mathcal{L} be a line bundle on X , linearized for the action of T . Then T acts on the one-dimensional \mathbb{C} -vector space $\mathcal{L}|_{y_i}$ for any $i = 1, \dots, c$, with weight μ_i say. Let us set $W_\mu(X, \mathcal{L}) := \{\mu_1, \dots, \mu_c\} \subseteq M$. The convex hull $P_\mu(X, \mathcal{L})$ of $W_\mu(X, \mathcal{L})$ in $M_{\mathbb{R}}$ is called the *moment polytope*.

REMARK 3.11. Note that μ_i does not depend on the choice of $y_i \in Y_i$. Indeed, let $y_i \in U \subseteq X$ be a T -invariant affine open neighbourhood which exists by [3, Corollary 5.3.6]. Pick a section $\sigma \in \mathcal{L}(U)$ which does not vanish at y_i . Then the weight μ_y of the action of T on $\mathcal{L}|_y$ is given by $t^{\mu_y} = \frac{(t \cdot \sigma)(y)}{\sigma(y)}$ for all $y \in Y_i \cap U$ such that $\sigma(y) \neq 0$. This shows that the map $Y_i \rightarrow M$, $y \mapsto \mu_y$ is continuous, hence constant.

EXAMPLE 3.12. Continuing Example 3.5, let $\mathbb{C}^\times \subseteq \text{Aut}(\mathbb{P}^1)$ act on $\mathcal{O}_{\mathbb{P}^1}(-1)$ via the rule $t \cdot (x, y) = (t^{w+1}x, t^w y)$. Then $W_\mu(\mathbb{P}^1, \mathcal{O}(-1)) = \{w, w+1\} \subseteq M = \mathbb{Z}$ and, consequently, $P_\mu(\mathbb{P}^1, \mathcal{O}_{\mathbb{P}^1}(-1)) = [w+1, w] \subseteq M_{\mathbb{R}} = \mathbb{R}$.

The following result is essentially taken from [4, Lemma 2.4] where it is stated only for ample bundles. The proof generalises to the semiample setting without difficulties.

PROPOSITION 3.13. *Let X be a smooth projective variety, let $T \subseteq \text{Aut}(X)$ be an algebraic torus and let \mathcal{L} be a semiample line bundle on X , linearized for the action of T .*

Then for any integer $m \geq 1$ the following hold true:

- (1) $mP_\Gamma(X, \mathcal{L}) \subseteq P_\Gamma(X, \mathcal{L}^{\otimes m})$ and $mP_\mu(X, \mathcal{L}) = P_\mu(X, \mathcal{L}^{\otimes m})$ as subsets of $M_{\mathbb{R}}$,
- (2) $P_\Gamma(X, \mathcal{L}) \subseteq P_\mu(X, \mathcal{L})$, and
- (3) if \mathcal{L} is basepoint free then $P_\Gamma(X, \mathcal{L}) = P_\mu(X, \mathcal{L})$.

Proof. (1): Clearly $mW_\Gamma(X, \mathcal{L}) \subseteq W_\Gamma(X, \mathcal{L}^{\otimes m})$ for if $w_1, \dots, w_m \in W_\Gamma(X, \mathcal{L})$ are weights for the action of T on $H^0(X, \mathcal{L})$ with corresponding eigenvectors $\sigma_1, \dots, \sigma_m$ then

$$\sigma_1 \otimes \dots \otimes \sigma_m \in H^0(X, \mathcal{L}^{\otimes m})$$

is an eigenvector for the action of T of weight $w_1 + \dots + w_m \in mW_\Gamma(X, \mathcal{L})$. We deduce that $mP_\Gamma(X, \mathcal{L}) \subseteq P_\Gamma(X, \mathcal{L}^{\otimes m})$.

That $mP_\mu(X, \mathcal{L}) = P_\mu(X, \mathcal{L}^{\otimes m})$ is obvious for if $y \in X^T$ and if μ is the weight for the action of T on $\mathcal{L}|_y$ then the weight of the action on $\mathcal{L}^{\otimes m}|_y$ is just $m\mu$.

To prove (2), the argument in [4, Lemma 2.4(2)] applies ad verbatim. Note that to prove Theorem 3.1 we will not make use of this inclusion.

Regarding (3), the proof of [4, Lemma 2.4(3)] again goes through without changes. However, as the argument is so elementary we want to quickly repeat it here: Fix $y \in X^T$ and let us denote by μ the weight of the action of T on $\mathcal{L}|_y$. As \mathcal{L} is generated by global sections we have the short exact sequence of T -modules

$$0 \rightarrow H^0(X, \mathcal{L} \otimes \mathfrak{m}_y) \rightarrow H^0(X, \mathcal{L}) \rightarrow \mathcal{L}|_y \rightarrow 0.$$

As it is well-known that any short exact sequence of T -modules splits, we find an eigenvector $\sigma \in H^0(X, \mathcal{L})$ for the action of T of weight μ . This proves that $W_\mu(X, \mathcal{L}) \subseteq W_\Gamma(X, \mathcal{L})$. We infer that $P_\mu(X, \mathcal{L}) \subseteq P_\Gamma(X, \mathcal{L})$ and so the result follows from (2). \square

COROLLARY 3.14. *Let X be a smooth projective variety, let $T \subseteq \text{Aut}(X)$ be an algebraic torus and let \mathcal{L} be a semiample line bundle on X , linearized for the action of T .*

Then $\kappa(X, \mathcal{L})^T \geq 0$ if and only if the origin $0 \in M$ is contained in $P_\mu(X, \mathcal{L})$.

Proof. Let us start by proving that

$$\kappa(X, \mathcal{L})^T \geq 0 \iff \exists m : 0 \in W_\Gamma(X, \mathcal{L}^{\otimes m}) \iff \exists m : 0 \in P_\Gamma(X, \mathcal{L}^{\otimes m}). \quad (3.1)$$

Indeed, the first equivalence holds true by the very definition of W_Γ and also the second ‘ \Rightarrow ’-implication is obvious. Regarding the converse, assume that $0 \in P_\Gamma(X, \mathcal{L}^{\otimes m})$. Let us denote the weights of the action of T on $H^0(X, \mathcal{L}^{\otimes m})$ by w_1, \dots, w_ℓ and let $\sigma_1, \dots, \sigma_\ell$ be a corresponding basis of eigenvectors. Now $0 \in P_\Gamma(X, \mathcal{L}^{\otimes m})$ simply means that there exist real numbers $0 \leq \lambda_i \leq 1$ such that $\sum \lambda_i = 1$ and

$$\sum_{i=1}^{\ell} \lambda_i w_i = 0.$$

In fact, by Proposition 3.15 below one can assume the λ_i ’s to be rational numbers, say $\lambda_i = \frac{p_i}{q_i}$. Set $k_i := q_1 \cdots q_{i-1} \cdot p_i \cdot q_{i+1} \cdots q_\ell \in \mathbb{Z}$. Then

$$\sum_{i=1}^{\ell} k_i w_i = 0.$$

But this in turn just means that

$$\sigma_1^{\otimes k_1} \otimes \dots \otimes \sigma_\ell^{\otimes k_\ell} \in H^0(X, \mathcal{L}^{\otimes m(k_1 + \dots + k_\ell)})$$

is an eigenvector for the action of T of weight $\sum k_i w_i = 0$. In other words, denoting $k = k_1 + \dots + k_\ell$ we deduce that

$$0 \in kW_\Gamma(X, \mathcal{L}^{\otimes m}) \subseteq kP_\Gamma(X, \mathcal{L}^{\otimes m}) \subseteq P_\Gamma(X, \mathcal{L}^{\otimes km}),$$

where we applied Proposition 3.13. This concludes the proof of (3.1).

But now, a further application of Proposition 3.13 yields

$$P_\Gamma(X, \mathcal{L}^{\otimes m}) = P_\mu(X, \mathcal{L}^{\otimes m}) = mP_\mu(X, \mathcal{L})$$

for sufficiently large $m \geq 1$. In view of (3.1), we deduce that $\kappa(X, \mathcal{L})^T \geq 0$ if and only if $0 \in P_\mu(X, \mathcal{L})$ as proclaimed. \square

The following result is just basic linear algebra. Due to the inability of the author to locate a reference in the literature we include a proof anyway.

PROPOSITION 3.15. *Let M be a finitely generated free abelian group, let $w_1, \dots, w_m \in M_{\mathbb{Q}} := M \otimes \mathbb{Q}$ be a finite subset and let $P \subseteq M_{\mathbb{R}}$ be its convex hull. Then*

$$P \cap M_{\mathbb{Q}} = P_{\mathbb{Q}} := \left\{ \sum_{i=1}^m \lambda_i w_i \mid 0 \leq \lambda_i, \lambda_i \in \mathbb{Q}, \sum_{i=1}^m \lambda_i = 1 \right\}.$$

Proof. Clearly $P_{\mathbb{Q}} \subseteq P \cap M_{\mathbb{Q}}$. Regarding the converse, fix $w \in P \cap M_{\mathbb{Q}}$. After replacing w_i by $w_i - w$ we may assume that $w = 0$. Write

$$w = \sum_{i=1}^m \lambda_i^* w_i$$

for some real numbers $0 \leq \lambda_i^*$ such that $\sum_{i=1}^m \lambda_i^* = 1$. Forgetting some of the w_i if necessary, we may assume that $\lambda_i^* > 0$ for all $i = 1, \dots, m$.

Now, consider the rational system of linear equations

$$\left\{ \left(\begin{array}{c} \lambda_1 \\ \vdots \\ \lambda_m \end{array} \right) \mid \sum_{i=1}^m \lambda_i w_i = 0, \sum_{i=1}^m \lambda_i = 1 \right\}.$$

By assumption it admits the real solution $\lambda^* := (\lambda_1^*, \dots, \lambda_m^*)$ such that $\lambda_i^* > 0$. Let $(\lambda_1, \dots, \lambda_m)$ be any rational solution which is sufficiently close to λ^* . Then also $\lambda_i > 0$ and so $w = \sum_{i=1}^m \lambda_i w_i$ with rational λ_i as required. \square

EXAMPLE 3.16. Continuing Example 3.5, let $\mathbb{C}^\times \subseteq \text{Aut}(\mathbb{P}^1)$ act on $\mathcal{O}(-1)$ via the rule $t \cdot (x, y) = (t^{w+1}x, t^w y)$. By Example 3.12 the moment polytope for the induced action of T on $\mathcal{O}(m)$ is given by

$$P_\Gamma(\mathbb{P}^1, \mathcal{O}(m)) = P_\mu(\mathbb{P}^1, \mathcal{O}(m)) = \left[-m(w+1), -mw \right].$$

We deduce that $\kappa(\mathbb{P}^1, \mathcal{O}(1))^T \geq 0$ if and only if $w = 0$ or $w = -1$.

PROPOSITION 3.17. *Let X be a smooth projective variety and let $T \subseteq \text{Aut}(X)$ be an algebraic torus. Choose a fixed point $y \in X^T$ and denote by ν_1, \dots, ν_n the weights of the natural isotropy action of T on the cotangent space $\Omega_X^1|_y$.*

(1) The weight μ of the action of T on $\mathcal{O}_X(-K_X)|_y$ satisfies the equation

$$\mu = -(\nu_1 + \dots + \nu_n).$$

(2) Let D be a T -invariant prime divisor on X and let us denote by μ the weight of the action of T on $\mathcal{O}_X(-D)|_y$. Then $\mu = \sum_i m_i \cdot \nu_i$ for some non-negative integers $m_i \geq 0$. Moreover:

(a) If $y \notin D$ then $\mu = 0$.

(b) If $y \in D$ is a smooth point of D then the natural isomorphism

$$\mathcal{O}_X(-D)|_y \cong \mathcal{N}_{D/X}^*|_y \subseteq \Omega_X^1|_y$$

is compatible with the action of T . In particular, $\mu = \nu_i$ for some i .

Proof. The first statement is clear as $\mathcal{O}_X(-K_X)|_y = \det(TX|_y)$ as T -modules.

Regarding item (2), we claim that there exists a local defining equation $f \in \mathcal{O}_{X,y}$ of D such that

$$t \cdot f = t^\mu \cdot f, \quad \forall t \in T. \tag{3.2}$$

Indeed, let $f' \in \mathcal{O}_{X,y}$ be any local defining equation. Since the action of T on $\mathcal{O}_{X,y}$ is algebraic, the \mathbb{C} -vector space spanned by $\{t \cdot f' \mid t \in T\}$ is finite-dimensional, see [24, Corollary 4.8]. Let f_1, \dots, f_r be a basis of T -eigenvectors, of weights μ_i say. Since $t \cdot f' = (t^{-1})^* f'$ is a local defining equation of D for any $t \in T$ by Example 3.7, it follows that f_i vanishes along D for any i . On the other hand, since f' is a \mathbb{C} -linear combination of the f_i , there must exist at least one $f = f_{i^*}$ which has multiplicity precisely one along D . Then f is a local defining equation for D , that is, $f|_y = f_{i^*}|_y$ generates $\mathcal{O}_X(-D)|_y$. In particular, the weight of the action of T on $\mathcal{O}_X(-D)|_y$ is μ_{i^*} . Thus, $\mu = \mu_{i^*}$, thereby proving (3.2).

Now, it follows from Luna’s étale slice theorem, cf. for example [24, Lemma 13.36], that

$$\widehat{\mathcal{O}_{X,y}} \cong \mathbb{C}[[z_1, \dots, z_n]]$$

for some formal functions z_i satisfying $t \cdot z_i = t^{\nu_i} \cdot z_i$ for any $i = 1, \dots, n$. Express

$$f = \sum_{J \in (\mathbb{Z}_{\geq 0})^n} a_J \cdot z^J$$

as a formal power series. Since $t \cdot f = t^\mu \cdot f$ by (3.2), it follows that $a_J = 0$ for all but possibly those $J = (m_1, \dots, m_n)$ such that $\sum m_i \cdot \nu_i = \mu$. As $f \neq 0$, we deduce that there exists at least one $J = (m_1, \dots, m_n) \in (\mathbb{Z}_{\geq 0})^n$ such that $\mu = \sum_i m_i \cdot \nu_i$. This proves the assertion.

Finally, if $y \notin D$ we may take $f = 1$, the constant function. As $t \cdot 1 = (t^{-1})^*(1) = 1$ we conclude that T acts trivially on $\mathcal{O}_X(-D)|_y$ as proclaimed.

Moreover, if $y \in D$ is a smooth point of D then $\mathcal{N}_{D/X}^*|_y \subseteq \Omega_X^1|_y$ is a one-dimensional subspace, generated by $df|_y$. We compute

$$t \cdot df|_y \stackrel{\text{Ex. 3.7}}{=} d(t \cdot f)|_y = d(t^\mu \cdot f)|_y = t^\mu df|_y.$$

Thus the weights of the actions of T on $\mathcal{O}_X(-D)|_y$ and $\mathcal{N}_{D/X}^*|_y$ are the same and we are done. \square

The presence of the divisor A in the following Lemma should be understood to be of a purely technical nature. Note that in any case Lemma 3.18 is no more general than Theorem C as also the pair $(X, D - A)$ is sub-log canonical, see [18, Corollary 2.35].

LEMMA 3.18. *Let (X, D) be a projective, log smooth, sub-log canonical pair and fix an algebraic torus $T \subseteq \text{Aut}(X, D)$. Let $A \geq 0$ be an effective, T -invariant \mathbb{Q} -divisor.*

If $-(K_X + D - A)$ is ample, then $0 \in P_\mu(X, -(K_X + D - A))$.

REMARK 3.19. Note that in view of item (1) in Proposition 3.13 it makes sense to define the moment polytope $P_\mu(X, -(K_X + D - A))$ even if A, D are only \mathbb{Q} -divisors by choosing m such that $-m(K_X + D - A)$ is Cartier and setting

$$P_\mu\left(X, -(K_X + D - A)\right) := \frac{1}{m}P_\mu\left(X, \mathcal{O}_X(-m(K_X + D - A))\right) \subseteq M_{\mathbb{R}}.$$

Proof of Lemma 3.18. Choose $y \in X^T \neq \emptyset$ and let us denote by $\mu \in M_{\mathbb{R}}$ the weight of the action of T on $\mathcal{O}_X(-(K_X + D - A))|_y$ ¹. Below, we will prove that

$$(1 - \varepsilon)\mu \in P_\mu\left(X, -(K_X + D - A)\right) \tag{3.3}$$

for all sufficiently small $\varepsilon \geq 0$. Using some completely elementary convex geometry this immediately yields the result, cf. Proposition 3.20 below.

To this end, let us denote the weights of the action of T on $\Omega_X^1|_y$ by ν_1, \dots, ν_n . As D has SNC-support, we may write

$$D = \sum_{i=1}^s \delta_i D_i$$

in some analytic open neighbourhood of $y \in X$ so that the D_i are smooth divisors meeting transversely at y and where $s \leq n$. Then the $\mathcal{N}_{D_i/X}^*|_y \subseteq \Omega_X^1|_y$ are distinct, one-dimensional, T -invariant subspaces. In particular, by Proposition 3.17 and after possibly re-indexing the D_i , we may assume that for any $i = 1, \dots, s$ the weight of the action of T on $\mathcal{N}_{D_i/X}^*|_y \subseteq \Omega_X^1|_y$ is given by ν_i . Note that $\delta_i \leq 1$ as (X, D) is assumed to be sub-log canonical. Let us set $\delta_i = 0$ for $s + 1 \leq i \leq n$. Write $A = \sum a_j A_j$, where, by assumption, $a_j \geq 0$ for all j .

Then by Proposition 3.17 the weight μ of the action of T on $\mathcal{O}_X(-(K_X + D - A))|_y$ satisfies the following equation:

$$\mu = -\sum_{i=1}^n \nu_i + \sum_{i=1}^n \delta_i \nu_i - \sum_j a_j m_{j,i} \nu_i = -\sum_{i=1}^n (1 - \delta_i) \nu_i - \sum_{i,j} a_j m_{j,i} \cdot \nu_i \in M_{\mathbb{R}}, \tag{3.4}$$

for some integers $m_{i,j} \geq 0$.

On the other hand, it is an important fact that if \mathcal{L} is any T -linearized ample line bundle on X and if μ denotes the weight of the action of T on $\mathcal{L}|_y$ then

$$\mu + \varepsilon \nu_i \in P_\mu(X, \mathcal{L}) \tag{3.5}$$

for sufficiently small $\varepsilon \geq 0$, see [4, Corollary 2.14] for a quick and purely algebraic proof².

¹ Here, to be precise we should really choose $m \geq 1$ such that $-m(K_X + D - A)$ is Cartier, denote the weight of the action of T on $\mathcal{O}_X(-m(K_X + D - A))|_y$ by μ' and set $\mu := \frac{\mu'}{m}$.

² In fact, the original statement in [13, Theorem 3] holds much more generally for any symplectic manifold and even shows that locally around μ , $P_\mu(X, \mathcal{L})$ is just the cone spanned by the ν_1, \dots, ν_n , see Figure 1 below. Note that unfortunately [13] conventions regarding moment maps differ from ours by a minus sign.




Figure 1.

Locally around the vertex μ the moment polytope P_μ (drawn in blue) looks like the cone spanned by the weights of the action on the cotangent space (drawn in lilac).

In any case in view of (3.4) and (3.5) we see that

$$(1 - \varepsilon)\mu = \mu + \varepsilon \left(\sum_{i=1}^n (1 - \delta_i)\nu_i + \sum_{i,j} a_j m_{j,i} \cdot \nu_i \right) \in P_\mu \left(X, \mathcal{O}_X(-(K_X + D - A)) \right), \quad (3.6)$$

where we use that $\delta_i \leq 1$ as (X, D) is sub-log canonical and $a_j \geq 0$ as A is effective. This proves (3.3) as required. Finally, as was indicated already above, the claimed Lemma 3.18 now follows immediately from (3.3) and the following elementary consideration: \square

PROPOSITION 3.20. *Let V be a finite dimensional real vector space, let $W \subseteq V$ be a finite subset and let $P \subseteq V$ be its convex hull. Assume that for any $w \in W$ it holds that $(1 - \varepsilon)w \in P$ for all sufficiently small $0 \leq \varepsilon$. Then $0 \in P$.*

Proof. Let us assume that $0 \notin P$. Then there exists a linear functional φ on V such that $\varphi|_P > 0$. As $P \subseteq V$ is compact and convex so is $\varphi(P) \subseteq \mathbb{R}$, that is, it is a (compact) interval, say $\varphi(P) = [a, b]$. Note that as P is the convex hull of W , $\varphi(P)$ is the convex hull of $\varphi(W)$ and so there exists at least one $w \in W$ for which $\varphi(w) = a$. But then our assumption that $(1 - \varepsilon)w \in P$ for all sufficiently small $0 < \varepsilon$ contradicts the fact that

$$\varphi((1 - \varepsilon)w) = (1 - \varepsilon) \cdot \varphi(w) = (1 - \varepsilon) \cdot a \notin [a, b].$$

This completes the proof of the proposition. \square

COROLLARY 3.21. *Let (X, D) be a projective, log smooth, log canonical pair and fix an algebraic torus $T \subseteq \text{Aut}(X, D)$. If $-(K_X + D)$ is nef, then $0 \in P_\mu(X, -(K_X + D))$.*

Proof. Let us start by proving the following:

CLAIM. There exists a T -invariant, effective, ample divisor A on X . \square

Indeed, let \mathcal{L} be a very ample line bundle on X . Then \mathcal{L} is T -invariant and, possibly replacing \mathcal{L} by some multiple, even linearizable, see [3, Theorem 5.2.1]. Let us choose an eigenvector $\sigma \in H^0(X, \mathcal{L})$ for the action of T of weight $w \in M$. Then the zero-set $A := Z(\sigma)$ is a T -invariant, effective, ample divisor on X as required.

Now, write $A = \sum_j a_j A_j$ as the sum of distinct prime divisors. For any rational number $\varepsilon > 0$ we consider the ample (!) divisor

$$-(K_X + D - \varepsilon A) = -(K_X + D) + \varepsilon A.$$

According to Lemma 3.18 it holds that

$$0 \in P_\varepsilon := P_\mu(X, -(K_X + D - \varepsilon A)), \quad \varepsilon > 0.$$

We are now ready to conclude that $0 \in P := P_0 := P_\mu(X, -(K_X + D))$: Let us denote by

$$d(P_\varepsilon, P) := \max \left\{ \sup_{w \in P} d(P_\varepsilon, w), \sup_{v \in P_\varepsilon} d(P, v) \right\}$$

the Hausdorff distance between P_ε and P with respect to some norm on $M_{\mathbb{R}}$. Note that by definition and for any $\varepsilon \geq 0$, the polytope P_ε is the convex hull of the weights $\mu_{\varepsilon,k}$ of the action of T on $\mathcal{O}_X(-(K_X + D - \varepsilon A))|_{y_k}$. Moreover, by (3.4) we have

$$\mu_{0,k} = \mu_{\varepsilon,k} + \varepsilon \left(\sum_{i,j} a_j m_{j,i}^{(k)} \cdot \nu_i \right)$$

for some integers $m_{j,i}^{(k)} \geq 0$, depending only on the weights of the action of T on $\mathcal{O}_X(-A_j)|_{y_k}$ but not on $\varepsilon \geq 0$, cf. Proposition 3.17. We infer that clearly $d(P_\varepsilon, P_0) \rightarrow 0$ as $\varepsilon \rightarrow 0$. In particular, $d(P_0, 0) \leq d(P_\varepsilon, P_0) \rightarrow 0$ and so $0 \in P = P_0$, using that the latter set is closed.

Proof of Theorem 3.1. Follows immediately by simply combining Corollary 3.21 with Proposition 3.13. □

The rest of this section is devoted to giving examples which demonstrate that the assumptions in Theorem 3.1, and hence in Theorem C, are essentially optimal.

EXAMPLE 3.22. The conclusion of Theorem 3.1 fails if we allow for more general subgroups $H \subseteq \text{Aut}(F, \Delta_F)$. In fact, the following example shows that it is already wrong for solvable groups:

Let $X = \mathbb{P}^1$, $D = 0$ and let $\mathbb{C}^\times \cong T \subseteq \text{PGL}_2(\mathbb{C})$ be the group of diagonal matrices as before. Then one easily checks that

$$\bigoplus_{m=0}^{\infty} H^0(X, \mathcal{O}_X(-mK_X))^T = \mathbb{C}[z\partial_z]$$

is generated by a single section. However, the vector field $z\partial_z$ is *not* invariant under the action of the subgroup $B \subseteq \text{PGL}_2(\mathbb{C})$ of upper triangular matrices.

EXAMPLE 3.23. The above example shows that usually $\kappa(X, -(K_X + D))^T$ is strictly smaller than $\kappa(X, -(K_X + D))$ even when the assumptions in Theorem 3.1 are satisfied. In fact, it is straightforward to extend Example 3.22 to show that

$$0 = \kappa(\mathbb{P}^n, -K_{\mathbb{P}^n})^T < \kappa(\mathbb{P}^n, -K_{\mathbb{P}^n}) = n,$$

for any $n \geq 1$. Here, $T \subseteq \text{Aut}(\mathbb{P}^n) = \text{PGL}_{n+1}(\mathbb{C})$ denotes the torus of diagonal matrices.

EXAMPLE 3.24. The assumption that (X, D) is sub-log canonical is optimal as the following example demonstrates: Consider $X = \mathbb{P}^1$ equipped with $D = a \cdot \{0\}$ for some $a \in \mathbb{Q}$. Then the standard torus of diagonal matrices $T \cong \mathbb{C}^\times \subseteq \text{Aut}(\mathbb{P}^1)$ acts on \mathbb{P}^1 via $t \cdot [x : y] = [tx : y]$. Note that the weight of the action of T on $\mathcal{T}_{\mathbb{P}^1}|_0$ is given by $1 \in \mathbb{Z} \cong M$ and the weight of the action of T on $\mathcal{T}_{\mathbb{P}^1}|_\infty$ is given by $-1 \in M$. Consequently, the weights of the action of T on $\Omega_{\mathbb{P}^1}^1|_0$ and $\Omega_{\mathbb{P}^1}^1|_\infty$ are given by -1 and 1 , respectively. Thus,

$P_\mu(\mathbb{P}^1, \mathcal{O}(-K_{\mathbb{P}^1})) = [-1, 1]$ and, hence, using Example 3.17

$$P_\mu\left(\mathbb{P}^1, -(K_{CP^1} + D)\right) = [-1 + a, 1].$$

We conclude that $-(K_{\mathbb{P}^1} + D)$ has T -invariant sections if and only if $a \leq 1$ which is the case if and only if (X, D) is sub-log canonical. At the same time, notice that $-(K_{\mathbb{P}^1} + D)$ is ample as long as $a \leq 2$.

§4. Proof of the main results

4.1 Proof of Theorem C

Let (X, D) be a projective, sub-log canonical pair and assume that $-(K_X + D)$ is semiample. Fix a connected, commutative, linear algebraic subgroup $H \subseteq \text{Aut}(X, D)$. We need to show that

$$\kappa(X, -(K_X + D))^H \geq 0.$$

If (X, D) is log smooth, then this was already shown in Theorem 3.1, cf. also Lemma 2.3. In general, we choose an H -equivariant log resolution $f: \hat{X} \rightarrow X$ of (X, D) . Such a map always exists by [19, Proposition 3.9.1]. Then, in particular $\text{Exc}(f)$ will be H -invariant.

Let us write

$$K_{\hat{X}} + \hat{D} \sim_{\mathbb{Q}} f^*(K_X + D), \quad (4.1)$$

where \hat{D} is supported on the strict transform of D and the exceptional locus of f . Then (\hat{X}, \hat{D}) is a log smooth pair which is sub-log canonical as (X, D) is so, see [18, Lemma 2.30]. Also, by construction,

$$-(K_{\hat{X}} + \hat{D}) \sim_{\mathbb{Q}} -f^*(K_X + D),$$

is semiample. Thus the conditions in Theorem 3.1 are satisfied and we deduce that

$$\kappa\left(\hat{X}, -(K_{\hat{X}} + \hat{D})\right)^H \geq 0.$$

Let us fix an integer $m \geq 1$ such that $-m(K_{\hat{X}} + \hat{D})$ is Cartier and such that there exists an H -invariant form

$$\begin{aligned} \hat{\tau} &\in H^0\left(\hat{X}, \mathcal{O}_{\hat{X}}\left(-m(K_{\hat{X}} + \hat{D})\right)\right) \\ &\cong H^0\left(\hat{X}, \mathcal{O}_{\hat{X}}\left(f^*\left(-m(K_X + D)\right)\right)\right) \\ &= H^0\left(X, \mathcal{O}_X\left(-m(K_X + D)\right)\right). \end{aligned}$$

Here, for the isomorphism in the second line we used (4.1). Let us denote by $\tau \in H^0(X, \mathcal{O}_X(-m(K_X + D)))$ the image of $\hat{\tau}$ under this isomorphism. We claim that τ is H -invariant, concluding the proof of the Lemma. Note that this is not entirely clear as the natural H -actions on both sides may a priori differ.

However, as $f: \hat{X} \rightarrow X$ is birational and H -equivariant, the set $U := \hat{X} \setminus (\hat{D} \cup \text{Exc}(f))$ is a dense open H -invariant subset of \hat{X} on which f is an isomorphism. In particular, we see that

$$\hat{\tau}|_U = \tau|_U \in H^0(U, \mathcal{O}_X(-K_U)).$$

As $\hat{\tau}$ is H -invariant we infer that so is $\hat{\tau}|_U = \tau|_U$ and, hence, (by continuity) τ . Thus, we have produced a non-trivial H -invariant form $\tau \in H^0(X, \mathcal{O}_X(-m(K_X + D)))$ and we are done.

4.2 Proof of Theorem A

Theorem A is an immediate consequence of Theorem C and Theorem 2.2.

4.3 Proof of Corollary B

Let (X, Δ) a projective klt threefold pair with nef anti-log canonical divisor $-(K_X + \Delta)$. We want to prove that

$$\kappa(X, -(K_X + \Delta)) \geq 0. \quad (4.2)$$

Let us start with some preliminary considerations. Let us denote by (F, Δ_F) a general fibre of the MRC fibration of (X, Δ) as in Theorem 2.1. Then F is a rationally connected projective variety, (F, Δ_F) is a klt pair and $-(K_F + \Delta_F)$ is nef. We distinguish four cases according to whether $\dim F = 0, 1, 2, 3$.

If $\dim F = 3$, then $X = F$ and (4.2) follows from [21, Theorem A]. In the other cases, according to Theorem 2.2 we need to prove that for any algebraic torus $T \subseteq \text{Aut}(F, \Delta_F)$ it holds that

$$\kappa(F, -(K_F + \Delta_F))^T \geq 0. \quad (4.3)$$

In case $\dim F = 0$, if $F = \{*\}$ is a point, there is nothing to prove, cf. Theorem 2.1. Also the case $\dim F = 1$ is easy to settle, for example because in this case clearly $F = \mathbb{P}^1$. Then since $-(K_F + \Delta_F)$ is nef it is even semiample and (4.2) is a direct consequence of Theorem A. Thus, it remains to deal with the case when F is a surface.

So let us then assume that F is a surface. If $T = \{1\}$ is trivial, then we simply need to verify that

$$\kappa(F, -(K_F + \Delta_F)) \geq 0,$$

which was proved in [14, Theorem 1.5]. Alternatively, this statement is also contained in [21, Theorem A]. In any case, note that this is precisely the situation when $X = Y \times F$ is a product.

It remains to deal with the case that $\dim T \geq 1$. Proceeding exactly as in the proof of Theorem C we may replace F by a log resolution. Note that this may force us to allow for (F, Δ_F) to be only *sub*-klt but this will not be an issue henceforth. In any case, we may assume F to be smooth.

Now, as $\dim T \geq 1$, the surface F is certainly a *T-variety of complexity one*, a variety equipped with the action of an algebraic torus such that the maximal dimensional orbits have codimension one in F . But then F is an *Mori Dream Space* in the sense of [16], so that $-(K_F + \Delta_F)$ is not only nef but also semiample. Hence, (4.2) follows from Theorem C.

The assertion that F is a Mori Dream Space under the above hypothesis seems to be well-known, for the convenience of the reader we nevertheless provide a detailed explanation below: As F is a smooth, rationally connected surface, it is in particular rational, birational to \mathbb{P}^2 . It follows that the class group $\text{Cl}(F)$ is finitely generated. Thus, to prove that F is a Mori Dream Space it only remains to show that the Cox ring of F is finitely generated, see [16, Proposition 2.9]. But indeed, Hausen and Süß in [15, Theorem 1.3] can even determine

an explicit finite list of generators and relations in terms of combinatorial data associated to the T -action. The theorem is proved.

Acknowledgements. First and foremost, the author would like to express his sincere gratitude towards Jarosław Wiśniewski for sketching to the author the proof of Theorem C in the case of a smooth Fanos without boundary. This paper would not exist without him. The author would also like to take this opportunity to warmly thank Roland Půček and Andriy Regeta for organising the “Regional Workshop in Algebraic Geometry” at Jena where the author and Jarosław Wiśniewski met.

Second, the author is extremely grateful towards the authors of [21] for bringing up the question whether Theorem A is true and, in particular, Shin-Ichi Matsumura and Thomas Peternell for several inspiring and fruitful initial discussions on the issue. Moreover, he is indebted to Vladimir Lazić and to Nikolaos Tsakanikas for many helpful comments on an early draft of this text and especially to Nikolaos Tsakanikas and Lingyao Xie for pointing out a serious mistake in an earlier version of this text.

He would also like to thank Daniel Greb for his advice and for always trying to make the author see the positive side of things, and Jochen Heinloth for several helpful discussions on torus-actions and for always being glad to answer any question of anyone stumbling by his office. Finally, he would like to warmly thank the anonymous referee for their detailed review and their valuable suggestions.

The author gladly wants to acknowledge that while writing this article he was supported from the DFG Research Training Group 2553 ”Symmetries and Classifying Spaces: Analytic, Arithmetic, and Derived.

REFERENCES

- [1] T. Bauer and T. Peternell, *Nef reduction and anticanonical bundles*, Asian J. Math. **8** (2004), no. 2, 315–352.
- [2] M. Brion, *Notes on automorphism groups of projective varieties*, unpublished, 2018. Available from the author’s website https://www-fourier.univ-grenoble-alpes.fr/mbrion/autos_final.pdf.
- [3] M. Brion, *Linearisations of algebraic group actions*, unpublished, 2018. Available from the author’s website <https://www-fourier.ujf-grenoble.fr/mbrion/lin>.
- [4] J. Buczyński and J. Wiśniewski, *Algebraic torus actions on contact manifolds*, J. Differ. Geom. **121** (2022), no. 2, 227–289.
- [5] F. Campana, J. Cao and S.-I. Matsumura, *Projective klt pairs with nef anti-canonical divisor*, Algebr. Geom. **8** (2021), no. 4, 430–464.
- [6] J. Cao, *Vanishing theorems and structure theorems of compact Kähler manifolds*. Thesis, Institut Fourier, France, 2013. <https://tel.archives-ouvertes.fr/tel-00919536>.
- [7] J. Cao, *Albanese maps of projective manifolds with nef anticanonical bundles*, Ann. Sci. Éc. Norm. Supér. (4) **52** (2019), no. 5, 1137–1154.
- [8] J. Cao and A. Höring, *Manifolds with nef anticanonical bundle*, J. Reine Angew. Math. **724** (2017), 203–244.
- [9] J. Cao and A. Höring, *A decomposition theorem for projective manifolds with nef anticanonical bundle*, J. Algebraic Geom. **28** (2019), no. 3, 567–597.
- [10] O. Debarre, *Higher-dimensional algebraic geometry*, Universitext, Springer-Verlag, New York, 2001.
- [11] S. Ejiri and Y. Gongyo, *Nef anti-canonical divisors and rationally connected fibrations*, Compos. Math. **155** (2019), no. 7, 1444–1456.
- [12] D. Greb, H. Guenancia and S. Kebekus, *Klt varieties with trivial canonical class: holonomy*, Geom. Topol. **23** (2019), no. 4, 2051–2124.
- [13] V. Guillemin and S. Sternberg, *Convexity properties of the moment mapping*, Invent. Math. **67** (1982), no. 3, 491–513.
- [14] J. Han and W. Liu, *On numerical nonvanishing for generalized log canonical pairs*, Doc. Math. **25** (2020), 93–123.

- [15] J. Hausen and H. Süß, *The Cox ring of an algebraic variety with torus action*, Adv Math. **225** (2010), no. 2, 977–1012.
- [16] Y. Hu and S. Keel, *Mori dream spaces and GIT*, Michigan Math J. **48** (2000), no. 1, 331–348.
- [17] T. Koike, *Ueda theory for compact curves with nodes*, Indiana Univ. Math. J. **66** (2017), no. 3, 845–876.
- [18] J. Kollár and S. Mori, *Birational geometry of algebraic varieties*, Cambridge Tracts in Mathematics, Vol. 134, Cambridge University Press, Cambridge, 1998.
- [19] J. Kollár, *Lectures on resolution of singularities*, Annals of Mathematics Studies, Vol. 166, Princeton University Press, Princeton, NJ, 2007.
- [20] V. Lazić and T. Peternell, *On generalised abundance, I*, Publ. Res. Inst. Math. Sci. **56** (2020), no. 2, 353–389.
- [21] V. Lazić, S.-I. Matsumura, T. Peternell, N. Tsakanikas and Z. Xie, *The nonvanishing problem for varieties with nef anticanonical bundle*, Doc. Math. **28** (2023), no. 6, 1393–1440.
- [22] S. Lu, Y. Tu, Q. Zhang and Q. Zheng, *On semistability of Albanese maps*, Manuscripta Math. **131** (2010), no. 3–4, 531–535.
- [23] S.-I. Matsumura and J. Wang, *Structure theorem for projective klt pairs with nef anti-canonical divisor*, preprint, 2021, [arXiv.2105.14308](https://arxiv.org/abs/2105.14308).
- [24] J. Milne, *Algebraic groups: The theory of group schemes of finite type over a field*, Cambridge Studies in Advanced Mathematics, Cambridge University Press, Cambridge, 2017.
- [25] N. Müller, *Locally constant fibrations and positivity of curvature*, preprint, 2022, [arXiv:2212.11530v1](https://arxiv.org/abs/2212.11530v1).
- [26] D. Mumford, D. Fogarty and F. Kirwan, *Geometric invariant theory*, Ergebnisse der Mathematik und ihrer Grenzgebiete (2), Vol. 34, Springer-Verlag, Berlin, 1994.
- [27] M. Păun, *Sur le groupe fondamental des variétés kählériennes compactes à classe de Ricci numériquement effective*, C. R. Acad. Sci. Paris Sér. I Math. **324** (1997), no. 11, 1249–1254.
- [28] M. Păun, *On the Albanese map of compact Kähler manifolds with numerically effective Ricci curvature*, Comm. Anal. Geom. **9** (2001), no. 1, 35–60.
- [29] K. Ueno, *Classification theory of algebraic varieties and compact complex spaces*, Lecture Notes in Mathematics, Vol. 439, Springer-Verlag, Heidelberg, 1975.
- [30] J. Wang, *Structure of projective varieties with nef anticanonical divisor: the case of log terminal singularities*, Math. Ann. **384** (2022), no. 1–2, 47–100.
- [31] Z. Xie, *Rationally connected threefolds with nef and bad anticanonical divisor*, Ann. Inst. Fourier **74** (2024), no. 5, 1819–1850.
- [32] Z. Xie, *Rationally connected threefolds with nef and bad anticanonical divisor, II*, preprint, 2023, [arXiv.2301.09466](https://arxiv.org/abs/2301.09466).
- [33] Q. Zhang, *On projective varieties with nef anticanonical divisors*, Math. Ann. **332** (2005), no. 3, 697–703.

Niklas Müller
Department of Mathematics
Universität Duisburg-Essen
Thea-Leymann-Strasse 9
45127 Essen
Germany
niklas.mueller@uni-duisburg-essen.de