

## Original Research

**Cite this article:** Dadario NB, Bellido S, Restivo A, Kulkarni M, Singh M, Yoon A, Shapiro J, Quintero F, Tagami T, Yang CJ, Jafri FN (2022) Using a logic model to enable and evaluate long-term outcomes of a mass casualty training program: a single center case study. *Disaster Med Public Health Prep* 16: 1116–1122. doi: <https://doi.org/10.1017/dmp.2021.66>.

First published online: 28 May 2021


### Keywords:

mass casualty incidents; medical simulation; emergency preparedness; emergency medicine; logic model

### Corresponding author:

Farrukh N Jafri,  
Email: [fjafri@wphospital.org](mailto:fjafri@wphospital.org).

# Using a Logic Model to Enable and Evaluate Long-Term Outcomes of a Mass Casualty Training Program: A Single Center Case Study

Nicholas B Dadario BS<sup>1</sup>, Simon Bellido BA<sup>2</sup>, Andrew Restivo MD<sup>3,4</sup>, Miriam Kulkarni MD<sup>5</sup>, Maninder Singh MD<sup>4,6</sup> , Andrew Yoon MD<sup>4,7</sup>, Jared Shapiro MPH<sup>8</sup>, Frank Quintero DO<sup>2</sup>, Tianna Tagami MEd<sup>9,10</sup>, Christina J Yang MD<sup>4,11</sup> and Farrukh N Jafri MD, MS-HPed<sup>2,4</sup>

<sup>1</sup>Robert Wood Johnson School of Medicine, Rutgers University, New Brunswick, NJ, USA; <sup>2</sup>White Plains Hospital Center, Emergency Medicine, White Plains, NY, USA; <sup>3</sup>Department of Emergency Medicine, Montefiore Medical Center, Weiler Division, Bronx, NY, USA; <sup>4</sup>Albert Einstein College of Medicine, Bronx, NY, USA; <sup>5</sup>Department of Emergency Medicine, St. John's Riverside Hospital, Yonkers, NY, USA; <sup>6</sup>Department of Emergency Medicine, Jacobi Medical Center, Bronx, NY, USA; <sup>7</sup>Department of Emergency Medicine, Montefiore Medical Center, Moses Division, Bronx, NY, USA; <sup>8</sup>Environmental Health and Safety, Montefiore Health System, Bronx, NY, USA; <sup>9</sup>MGH Institute of Health Professions, Boston, MA, USA; <sup>10</sup>Pearson Education, Boston, MA, USA and <sup>11</sup>Department of Otorhinolaryngology, Montefiore Medical Center, Bronx, NY, USA

## Abstract

**Purpose:** Global health disasters are on the rise and can occur at any time with little advance warning, necessitating preparation. The authors created a comprehensive evidence-based Emergency Preparedness Training Program focused on long-term retention and sustained learner engagement.

**Method:** A prospective observational study was conducted of a simulation-based mass casualty event training program designed using an outcomes-based logic model. A total of 25 frontline healthcare workers from multiple hospital sites in the New York metropolitan area participated in an 8-hour immersive workshop. Data was collected from assessments, and surveys provided to participants 3 weeks prior to the workshop, immediately following the workshop, and 3 months after completion of the workshop.

**Results:** The mean percentage of total knowledge scores improved across pre-workshop, post-workshop and retention (3 months post-workshop) assessments (53.2% vs. 64.8% vs. 67.6%,  $P < 0.05$ ). Average comfort scores in the core MCI competencies increased across pre-workshop, post-workshop and retention self-assessments ( $P < 0.01$ ). Of the participants assessed at 3 months retention ( $n = 14$ , 56%), 50.0% ( $n = 7$ ) assisted in updating their hospital's emergency operations plan and 50.0% ( $n = 7$ ) pursued further self-directed learning in disaster preparedness medicine.

**Conclusions:** The use of the logic model provided a transparent framework for the design, implementation, and evaluation of a competency-based EPT program at a single academic center.

## Introduction

Current literature has frequently identified deficiencies in emergency preparedness as well as performance in previous disasters, raising concerns for a broad lack of system preparedness for future mass casualty incidents (MCI).<sup>1–3</sup> Annual Emergency Preparedness Training (EPT) programs allow clinicians to learn and refine disaster management approaches to protect both patients and healthcare professionals in the event of an MCI, and have been associated with improved performance in actual MCI events.<sup>4–6</sup> There are scant competency-based EPT programs in the United States with non-uniform curriculum and a paucity of objective data suitable for actionable evaluation.<sup>7–9</sup> Few programs are deliberately designed to evaluate long-term individual disaster knowledge retention, continuing engagement with training or contributions to local hospital emergency operations plans.<sup>10,11</sup> With an increased incidence of global health disasters,<sup>12,13</sup> it is imperative that disaster preparedness educators apply evidence-based methodologies to guide deliberate design and evaluation of training programs, to ensure they result in significant, durable learning, and increased levels of engagement and readiness.

Logic models have been successfully used to evaluate programs in the fields of public health and medical education.<sup>14–16</sup> Well-designed logic models provide a transparent program framework to enable design collaboration and articulate extractable performance outcomes to guide program implementation, evaluation and continuous quality improvement.<sup>17–19</sup> The use of a logic model projects outcomes beyond immediate, direct learning, to measure intermediate, and long-term impact.<sup>20</sup>

The objective of the MCI Foundations program was to examine the use of a logic model to align a training program's activities and assumed inputs with its intended effects over time, to determine whether the use of the model facilitated significant, durable learning, and increased engagement and impact.

## Methods

### Study Design

This was an open-label, observational study following the reporting model of Strengthening the Reporting of Observational Studies in Epidemiology (STROBE) checklist. MCI Foundations was a 1-day comprehensive 8-hour experiential workshop, held on November 13, 2019 at the Montefiore Einstein Center for Innovation in Simulation (MECIS). The training was developed by the MCI Foundations Committee, which consisted of disaster medicine experts, simulation educators and clinical educators. The study was reviewed and determined exempt by the Montefiore Medical Center Institutional Review Board (IRB), #2019-10614.

This program's metrics for success were sustained improvements in disaster medicine knowledge and comfort with core disaster competencies, pursuit of further study in emergency preparedness, and involvement of participants in their individual hospital's disaster management.

### Participants

Participants were recruited by electronic mail sent to chief medical officers, directors of emergency medicine, and vice presidents of nursing across all affiliated hospitals in the Montefiore Network by the director of emergency management at Montefiore as well as by members of the MCI Foundations Committee. Inclusion criteria for invitation to the program included full-time employment in the Montefiore Health System (academic and non-academic centers) with interest in disaster medicine, but little to no expertise. Priority was offered to employees from the departments of emergency medicine, surgery and intensive care. Residents, students and part-time employees were excluded from invitation. 30 slots were available in a "first-come, first served" system. Participation was voluntary and oral informed consent was obtained with each participant having the ability to withdraw at any time.

### Implementation of the Logic Model

The use of a logic model for program design begins with the definition of intended short-term, intermediate and long-term program outcomes; these are important for EPT programs to demonstrate impact. The next stage in applying a logic model is to identify the activities and outputs that will enable the identified outcomes. For the MCI Foundations program, this stage involved the actual curriculum design and development. Once activities and outputs are defined, then inputs, or required materials, and resources can be identified. The inputs, activities, outputs and outcomes identified for this training are shown in [Table 1](#), along with the tool designed to measure each outcome.

### Learning Design

In order to enable significant, durable learning, the MCI Foundations Committee aimed to ground the training in relevant, profession-related activities by articulating requirements as

entrustable professional activities (EPAs) and support skills and knowledge acquisition with meaningful personal connections by using Fink's Taxonomy of Significant Learning to articulate learning outcomes and define learning experiences.

EPAs are units of professional practice, defined as tasks or responsibilities, to be entrusted to an unsupervised trainee for execution once they have attained sufficient specific competence.<sup>21</sup> The praxis focus of EPAs ensures the curriculum design drives toward operationalized skills and knowledge.

The MCI Foundations training was designed to enable the following EPAs:

- 1) Identify a potential critical event, appropriate safety precautions for that event type, and perform the appropriate simulated notification and actions for mobilization.
- 2) Identify the Incident Command System (ICS) defined individual task and scope of responsibility.
- 3) Correctly utilize the Team Strategies and Tools to Enhance Performance and Patient Safety (TeamSTEPPS) principles of "Brief, Huddle, and Debrief" during a mass casualty exercise.
- 4) Apply knowledge and skills concerning MCI triage systems to rapidly assign victims to appropriate triage categories.

The training focused on the following scope of content to enable these EPAs:

- Critical event safety principles ("Safety")
- The hospital incident command system and the participant's role in it ("HICS")
- Effective critical event communications ("TeamSTEPPS")
- Knowledge of the MCI Triage system ("MCI Triage")

[Table 2](#) illustrates how a single MCI Foundations learning outcome correlates to an EPA and a core competency,<sup>9</sup> and is aligned to a selected-response assessment item. This program contained a complete list of targeted learning outcomes aligned to EPAs and healthcare worker disaster training competencies. Further information on our curriculum is outlined in the Supplementary material.

Fink's Taxonomy of Significant Learning was used to articulate the targeted learning outcomes and design appropriate learning and assessment activities. Fink's Taxonomy is designed to encourage curriculum developers to go beyond a content-driven approach to course design and consider how to affect significant durable change in the learner.<sup>22</sup> Fink's Taxonomy is composed of dynamically interacting dimensions that consider the learner's relationship with the content, themselves and their own learning and others. The dimensions of Fink's Taxonomy are: (1) Foundational Knowledge (2) Application (3) Integration (4) Human Dimension (5) Caring (6) Learning how to learn. Organizing the outcomes and learning activities around Fink's Taxonomy kept the MCI Foundations team focused on making meaningful connections to the learners' lives and emphasizing the personal and professional value of becoming a resource for their communities in preparation for a mass casualty event. [Table 3](#) contains the activities that comprised the training, organized by Fink's dimensions.

### Inputs

MCI Foundations was funded internally by the Office of Emergency Management at Montefiore Health. Funding covered the following: facility space, consumable, and non-consumable

**Table 1.** Logic model of MCI foundations emergency preparedness training program

| Inputs   |   | Activities  | Output  |
|--|---|---|---|
| <ul style="list-style-type: none"> <li>• Facility space</li> <li>• Faculty time</li> <li>• Trainee time</li> <li>• Consumable equipment               <ul style="list-style-type: none"> <li>○ Tape, markers, signs, papers, moulage</li> </ul> </li> <li>• Non-Consumable               <ul style="list-style-type: none"> <li>○ Simulation, audiovisual, costumes</li> </ul> </li> <li>• Funding (catering, event space, actors)</li> <li>• Administration approval</li> </ul> |   | <ul style="list-style-type: none"> <li>• Online module to introduce core concepts and competencies</li> <li>• Rapid-fire panel discussion on core concepts by disaster experts</li> <li>• Tabletop exercises focused on disaster triage and comprehensive MCI evaluations</li> <li>• Multiple immersive simulations focused on MCI</li> </ul> | <ul style="list-style-type: none"> <li>• Completion of the program with pre and post event assessments</li> <li>• Number of participants (by discipline/by hospital)</li> <li>• Number of facilitators (by discipline)</li> </ul> |
| Outcome Term   | Outcome   | Measurement Tool  |   |
| Short-Term   | Disaster preparedness knowledge                 | Selected-response assessment (post-test data compared to pre-test)  |   |
|  | Comfort with core competencies                  | Survey using the 5-point linear numeric response format   |   |
| Intermediate   | Continuing interest in disaster preparedness    | Web traffic   |   |
| Long-Term  | Disaster preparedness knowledge                 | Selected-response assessment (retention data compared to post-test and pre-test data)   |   |
|  | Familiarity with core competencies              | Survey using the 5-point linear numeric response format   |   |
|  | Pursuit of further study                        | Survey  |   |
|  | Involvement with hospital preparedness planning | Survey  |   |

**Table 2.** An MCI foundations learning outcome with correlated EPA and core competency

|   |   |
|---|---|
| <b>EPA:</b> Manage critical event actions using the Disaster Triage System              | <b>Assessment item:</b>   |
| <b>Competency:</b> Demonstrate disaster triage knowledge and skills                     | Comparing the START and JumpSTART algorithms, identify a key difference.  |
| <b>Learning Outcome:</b> Identify differences between the START and JumpSTART algorithm | a) If opening the airway of a patient does not restart breathing, tag as black in both algorithms<br>b) If an apneic child has a radial pulse, 5 rescue breaths can be given in the JumpSTART algorithm<br>c) If opening the airway of a patient does not restart breathing, tag as black only in JumpSTART<br>d) If an apneic adult has a radial pulse, 5 rescue breaths can be given in the START algorithm |

**Abbreviations:** EPA, Entrustable Professional Activity.

**Table 3.** MCI foundations activities according to Fink's taxonomy of significant learning

| Fink's Dimension                              | Training Activities  |
|---|--|
| Foundational knowledge                        | First day final<br>Online modules<br>Rapid fire/panel              |
| Application<br>Integration<br>Human dimension | Tabletop triage<br>Tabletop comprehensive<br>Immersive simulations |
| Caring  | 3-minute message   |
| Learning how to learn                         | Learning goal list<br>Stamped postcards                            |

equipment, as well as actors, and moulage. Faculty time was considered on a volunteer basis.

### Activities

Exactly 3 weeks prior to the workshop, participants completed a pre-test to establish their baseline disaster medicine knowledge and familiarity, followed by completing online modules. The in-person workshop was an 8-hour immersive program held at the Montefiore Einstein Center for Innovation in Simulation (MECIS), that covered

panel discussions, table-top exercises, a personal reflection, and 2 large scale disaster simulation exercises involving multiple high and low fidelity as well as professional actors and volunteer ambulance crews.

### Outcome Measures

#### Short-term Knowledge

Disaster medicine knowledge was measured using a 16-question selected-response assessment, administered before the training, immediately following the training, and 3 months later. The test answers were not reviewed during, or after the program to limit bias on future assessments. The test was developed by the Principal Investigator (FJ) and First Author (ND), internally validated by 4 internationally recognized disaster medicine experts with a combined 90 years of training and with experience in actual MCI events, and then reviewed by the MCI Foundations Committee. The assessment covered 4 main categories of disaster medicine knowledge: Hospital incident command System (HICS), MCI triage, Safety assessment, and Communication (using the TeamSTEPPS model). HICS questions were designed following the Federal Emergency Management Agency (FEMA) hospital incident command system (ICS) guidelines; MCI triage questions were developed following the Simple Triage and Rapid Treatment (START) and JumpSTART, the pediatric disaster triage algorithm, and protocols; TeamSTEPPS questions were created following guidelines set forth

by the Agency for Healthcare Research and Quality in collaboration with the U.S. Department of Defense; safety assessment questions were created based on principles put forth by FEMA in its Incident Command System courses.<sup>23</sup> While TeamSTEPPS does not traditionally play a role in Disaster Management programs, the MCI Foundations Committee felt teaching Crisis Resource Management (CRM) in such a setting would be an integral part of the program.

The disaster medicine knowledge pre-test and post-test scores were compared with a paired-t test. The effect size for mean percentage of pre-test and post-test was measured using a Cohens D. This study expected training participants to score at least 67% on the post-test, which was administered immediately following the training. This goal was determined by consensus from the MCI Foundations team using the strategy of averaging scores of the pilot test obtained by 4 independent disaster medicine experts (83%) to 6 independent emergency department physicians without prior disaster management training (51%). The MCI Foundations committee determined if the program could improve the average score to 67% (an increase by 31%), the group would consider the foundational program successful in improving baseline knowledge.

#### Short Term Comfort

Comfort with core disaster competencies was assessed using a questionnaire with a 5-point linear numeric response format (1 = Very Uncomfortable, 5 = Very Comfortable) relating to fundamental competencies in disaster preparedness, HICS, TeamSTEPPS, emergency preparedness, and triage. The group's Pre and Post mean scores in these categories were all compared using Wilcoxon signed rank tests and the effect size was measured with the Wilcoxon test effect size ( $r$ ).

#### Intermediate Outcome

To measure intermediate progress toward the MCI Foundation goal of encouraging participants to pursue self-learning activities and engage in their hospital's emergency operations plans, traffic to a website created specifically for the program was monitored during 3 months, from the workshop to retention testing. The website contained resources for MCI and information for all regional and local upcoming meetings in our hospital network, upcoming course offerings, seminars, interactive learning games and online disaster educational opportunities. The link was sent out immediately upon conclusion of the program and again 3 months later.

#### Long-term Outcomes

In order to assess the acquisition and maintenance of any benefits from the program on specific individuals over time, the group's knowledge and comfort scores were compared with paired subject analyses at pre, post and retention. Knowledge and comfort assessments included the same questions as those used in the post-test. Total knowledge scores were compared using one-way repeated measures ANOVA to evaluate pre-test, post-test, and retention scores on paired subjects. Post Hoc tests were conducted using pairwise t-test comparisons to analyze the mean total score differences.

Paired subjects for the comfort self assessments at pre, post and retention were compared using a Friedman's test. Post Hoc data was conducted using Wilcoxon signed ranks tests to determine the difference between individual effects and identify any skill decay after the program. The Friedman test effect size for mean percentage of pre/post/retention was measured using Kendall's

W, which follows the interpretation guidelines of Cohen's D effect size test.

## Results

All statistical analyses were performed in R 4.0.0 (R Foundation for Statistical Computing, Vienna, Austria). Short-term, intermediate and long-term outcomes were evaluated in order to assess the efficacy of the program (Table 1). The effect of this program on knowledge and comfort were assessed using paired samples between the pre-assessments and post-assessments. For the participants who were retained in all 3 of the testing phases, paired samples were compared across pre-test, post-test and retention.

#### Outputs

A total of 25 frontline staff participated in this program from 8 different hospitals within the Montefiore Hospital Network in the southern NY region that included community (64%) and academic (36%) centers. These 25 participants had varying clinical experience and included 10 physicians (40%), 7 registered nurses (28%), 3 advanced practice providers (12%), 2 nursing technicians (8%), 2 administration roles (8%) and 1 respiratory therapist (4%). 5 participants (20%) reported previous training in mass casualty incidents. Participants predominantly worked in the Department of Emergency Medicine (23; 92%), followed by the Department of Surgery (1; 4%) and the Intensive Care Unit (1; 4%). There were 25 requests for 30 available slots, so all those who responded with interest were admitted.

The 15 instructors who participated in the workshop included 5 disaster management experts covering 2 hospital systems as well as a representative from the New York State Division of Homeland Security and Emergency Services, 10 instructors with backgrounds in Medical Simulation and Healthcare Education, alongside 8 non-medical volunteers assisted in the program.

#### Short Term Knowledge


The mean percentage of the group's total knowledge-based scores improved from pre-test to post-test (51.3% vs. 63.8%,  $P < 0.001$ ; 95% CI: 0.075-0.174;  $d = 1.034$ ). The breakdown of the total scores demonstrated improvements of knowledge in the mean score of each category: TeamSTEPPS (35.1% to 50.7%), HICS (47.3% to 62.0%), MCI Triage (59.8% to 68.8%), and Safety (59.0% to 70.0%).

#### Short Term Comfort

2 participants were excluded from the pre/post familiarity score analysis because they did not complete the comfort self-assessment before or directly after the course (Figure 2). The mean scores for all categories of comfort in core MCI competencies significantly increased from pre-test to post-test: disaster preparedness (2.6 vs. 3.5,  $P < 0.001$ ; 95% CI: 0.999-1.500,  $r = 0.77$ ), Triage (2.8 vs. 3.9,  $P < 0.001$ ; 95% CI: 0.999-1.50,  $r = 0.80$ ), the HICS framework (2.3 vs. 3.3,  $P < 0.001$ ; 95% CI: 0.999-1.999,  $r = 0.75$ ), Team STEPPS (2.2 vs 3.7,  $P < 0.001$ ; 95% CI: 1.00-2.00,  $r = 0.84$ ), and emergency preparedness (2.7 vs. 3.6,  $P < 0.001$ ; 95% CI: 1.50-2.00,  $r = 0.76$ ).

#### Intermediate Outcome

In the 3 months following the workshop, our website measured a total of 64 views amongst 53 different devices. The website was



**Figure 1.** Assessment of disaster knowledge in the total group (left) and in only the group that was retained throughout the whole study period (right). Testing periods relate to pre-workshop, post-workshop, and retention 3 months after workshop. Higher scores relate to better performance. Bold lines refer to median scores, horizontal lines refer to the 1st and 3rd quartiles in the distribution, whiskers represent the lowest and highest datum within 1.5 interquartile range of the lower and upper quartiles, and dots represent outliers. \* $P < 0.05$  \*\* $P < 0.01$  \*\*\* $P < 0.001$  \*\*\*\* $P < 0.0001$ .

emailed both after the program and in the 3-month follow up to all participants and program instructors.

### Long-term Outcomes

Among the 14 participants (56.0%) who completed the 3-month survey, 7 participants (50%) reported engaging in further self-directed learning. A total of 7 participants (50.0%) assisted in updating the disaster response plan for their hospital, 6 participants (42.9%) attended a local disaster meeting, and 7 people (50.0%) took an extra disaster preparedness course or workshop. Of these participants, 5 (35.7%) said they would not have attended further disaster workshops if it was not for this course and 9 (64.3%) reported further pursuing their self-reported learning objectives that they wrote in this program with “time” being the most reported barrier to their disaster goals ( $n = 2$ , 14.3%). When assessing only the participants who completed all 3 of the testing phases, this program had a significant effect on *knowledge* and *comfort* over the course of 3 months.

### Long-term Knowledge

During all 3 time periods, 2 participants were excluded from the disaster medicine knowledge analysis for not completing the assessments. The mean percentage of the retention group’s ( $n = 12$ ) total knowledge-based scores improved across pre, post and retention testing (53.2% vs. 64.8% vs 67.6%,  $P < 0.05$ ). These scores exceeded this programs 67% score predetermined threshold for post and retention testing. Post hoc comparisons demonstrated that the mean total score difference was significant between pre-test and post-test ( $P < 0.05$ ,  $d = 0.99$ ) and pre-test and retention ( $P < 0.05$ ,  $d = 0.80$ ), but not between post-test and retention

( $P > 0.05$ ,  $d = 0.17$ ). Together, these data suggest that this program has a large effect on total MCI knowledge after the program with a continued moderate effect at 3 months retention (Figure 1).

### Long-term Comfort

A total of 14 participants (56%) completed the comfort self-assessment after 3 months. However, 1 participant was excluded from the analysis at all 3 time periods due to an incomplete pre-course comfort assessment. The mean comfort scores in core MCI

competencies in disaster, triage and HICS increased across pre, post and retention self-assessments relating to: disaster preparedness (2.69 vs. 3.62 vs. 3.85,  $P < 0.01$ ,  $W = 0.62$ ), Triage (2.92 vs. 3.85 vs. 4.00,  $P < 0.01$ ,  $W = 0.54$ ) and HICS (2.46 vs 3.46 vs 3.85,  $P < 0.01$ ,  $W = 0.72$ ). Post hoc comparisons demonstrated that the score differences in each of these categories were significant between pre-test and post-test ( $P < 0.05$ ) and pretest and retention ( $P < 0.05$ ), but not between post-test and retention ( $P > 0.05$ ). The mean comfort score in overall emergency preparedness and TeamSTEPPS both increased after the workshop, but the scores remained the same from post testing to retention for overall emergency preparedness, and decreased for TeamSTEPPS (Overall Emergency Preparedness: 2.77 vs. 3.77 vs. 3.77,  $P < 0.01$ ,  $W = 0.56$ ; TeamSTEPPS: 2.62 vs. 4.08 vs. 4.00,  $P < 0.01$ ,  $W = 0.73$ ). Post hoc comparisons demonstrated that the pre to post improvements for both of these categories significantly differed ( $P < 0.05$ ), and the decline from post to retention testing for TeamSTEPPS was not significant ( $P > 0.05$ ).


### Discussion

The MCI Foundations program successfully implemented an outcomes-based logic model in the inception, implementation, and assessment of emergency preparedness training (EPT). The study’s successful application of the model to the design and evaluation of this program enables targeted improvements with long-term impact as the program continues to expand towards a more prepared hospital network. It also provides an early example for future emergency preparedness program designers to utilize, and improve upon to create high-quality, transparent, outcomes-based programs.

The authors observed a short-term improvement in disaster knowledge immediately after the program compared to baseline knowledge (51.3% vs. 63.8%); however, this data did not meet the predetermined 67% success mark, indicating a need for program improvement in this area. As all aspects of the curriculum and assessment design were aligned to specific learning objectives and knowledge outcomes, this design allowed for targeted improvements for future programs.

In addition to the novelty of the logic model, this study is unique in measuring long-term retention of knowledge and comfort with core competencies following EPT programs. Subgroup analysis on the individuals who were retained throughout the entire program (pre-test, post-test, and retention) demonstrated that, while their post test average was 65%, their retention average met the 67% threshold for success, suggesting further self-study, and improvement in disaster preparedness in this group. It is noteworthy that 50% of the retention group engaged in disaster response planning at their hospital.

This retention group had higher scores on average compared to the entire program group in knowledge and in all categories of comfort after the program. These scores improved on post testing without significant decay (Figure 2). These findings suggest the MCI Foundations program, which is designed to support durable learning, played a role in the reduction of decay of disaster knowledge and comfort with core competencies in this subgroup. This is significant given that an MCI can occur anytime and makes a compelling argument that future studies of disaster preparedness training should include long-term outcomes’ metrics. Importantly, this program utilized a curriculum deliberately designed to foster significant, durable learning in participants through personally meaningful and professionally relevant activities and assessments



**Figure 2.** Assessment of comfort in individual MCI competencies. Testing periods relate to pre-workshop, post-workshop, and retention 3 months after workshop. (Scale: 1= not comfortable, 5= very comfortable). \* $P < 0.05$  \*\* $P < 0.01$  \*\*\* $P < 0.001$ .

developed using EPAs and Fink's Taxonomy of Significant Learning. The application of these frameworks to guide curriculum development warrants further research in the context of clinical disaster education.

### Limitations

This program encountered a number of limitations and challenges in its design, implementation, and evaluation. Using a logic model as an evaluation tool requires alignment of specific metrics to outcomes. This alignment was problematic in the case of intermediate outcomes related to individual learner commitments to continuing education. By monitoring traffic on a website sent to participants after the program, it was discovered that participants were accessing resources for course offerings and disaster planning meetings; however, the authors noted sparse attendance at these meetings and only 7 participants (50%) in the retention group reported engaging in further learning or helping their hospital develop a disaster response plan. Still, the authors observed 1 participant in the study rewrote much of her hospital's critical event annex in the emergency operations plan and has since enrolled in a Masters in Emergency Management, demonstrating a need for more individualized observations. While the MCI Foundations program cannot take credit for this career trajectory, this individual observation is precisely the reason Fink's taxonomy was utilized as the backbone structure for the program. The current study was unable to measure the impact participants had in their home institutions. The low retention rate prevented this study from understanding the full impact of the intervention utilized. Only 14 participants responded to the retention survey, some of which were incomplete. Given that the pool of self-identified disaster novices in this program's hospital network is likely over 10000 employees,

and this study only enrolled 25, this program is subject to both selection bias and attrition bias. Also, the logic model was not designed to directly measure impact, but rather had a quantitative function for involvement, and attendance-only, that was proposed to lead to impact. This led the MCI Foundations Committee to consider future directions of focusing on individual hospitals at a time using a similar model to further understand potential impact.

### Conclusion

The first iteration of the MCI Foundations model provided the committee with a transparent framework for the design, implementation, evaluation and continuous quality improvement of a competency-based EPT program. While successful in evaluating targeted improvements with participant long-term disaster knowledge and comfort, future models will consider the impact of individual learners on their hospital's disaster preparedness rather than simply the number who engage. Future work should further implement this model within individual hospitals for more customized, system-based training and evaluation to understand the multitude of benefits offered from the MCI Foundations model.

**Acknowledgments.** We would like to acknowledge additional faculty for MCI Foundations: Ed Tangredi CEM, Erik Larsen MD, and Nadine Macura MBA CEM as content specialists, and support for the program and panelists; Bernadette Amicucci DNS, MBA, FNP-BC, CNE, Marc J Gibber MD, and David DiMattia MS as faculty instructors; Marvin Fried MD and Emily Kaplan MPA for facilities and technical support; and Stefan Ravello and Steven Haimowitz MD for online module support from RealCME, Inc.

**Ethics statement.** The study was reviewed and determined exempt by the Montefiore Medical Center Institutional Review Board (IRB).

**Supplementary material.** To view supplementary material for this article, please visit <https://doi.org/10.1017/dmp.2021.66>

## References

1. **Lyznicki J, Subbarao I, Benjamin GC, James JJ.** Developing a consensus framework for an effective and efficient disaster response health system: a national call to action. *Disaster Med Public Health Prep.* 2007;1(1 Suppl): S51-S54.
2. **Epstein SK, Burstein JL, Case RB, et al.** The National Report Card on the state of emergency medicine: Evaluating the emergency care environment state by state 2009 edition. *Ann Emerg Med.* 2009;53(1):4-148.
3. **Report Card Task Force Members; ACEP Staff.** America's emergency care environment, a state-by-state Report Card: 2014 edition. *Ann Emerg Med.* 2014;63(2):97-242.
4. **Resnick L.** It Takes a Team - The 2013 Boston Marathon: Preparing for and recovering from a mass-casualty event. Needham, MA: *JBJS and JOSPT*; 2014.
5. **Tobert D, von Keudell A, Rodriguez EK.** Lessons from the Boston Marathon Bombing: An orthopedic perspective on preparing for high-volume trauma in an urban academic center. *J Orthop Trauma.* 2015;29 Suppl 10:S7-S10.
6. **Walls RM, Zinner MJ.** The Boston Marathon response: Why did it work so well?. *JAMA.* 2013;309(23):2441-2442.
7. **Subbarao I, Lyznicki JM, Hsu EB, et al.** A consensus-based educational framework and competency set for the discipline of disaster medicine and public health preparedness. *Disaster Med Public Health Prep.* 2008;2(1):57-68.
8. **Scott LA, Smith C, Jones EM, Manaker LW, Seymore AC, Fulkerson J.** Regional approach to competency-based patient care provider disaster training: the Center for Health Professional Training and Emergency Response. *South Med J.* 2013;106(1):43-48.
9. **Hsu EB, Thomas TL, Bass EB, Whyne D, Kelen GD, Green GB.** Healthcare worker competencies for disaster training. *BMC Med Educ.* 2006;6:19.
10. **Scott LA, Swartzentruber DA, Davis CA, Maddux PT, Schnellman J, Wahlquist AE.** Competency in chaos: Lifesaving performance of care providers utilizing a competency-based, multi-actor emergency preparedness training curriculum. *Prehosp Disaster Med.* 2013;28(4):322-333.
11. **Scott LA, Carson DS, Greenwell IB.** Disaster 101: A novel approach to disaster medicine training for health professionals. *J Emerg Med.* 2010;39(2):220-226.
12. **Rossati A.** Global warming and its health impact. *Int J Occup Environ Med.* 2017;8(1):7-20.
13. **Tekola B, Myers L, Lubroth J, Plee L, Calistri P, Pinto J.** International health threats and global early warning and response mechanisms. *Rev Sci Tech.* 2017;36(2):657-670.
14. **Evers U, Jones SC, Iverson D, Caputi P.** 'Get Your Life Back': Process and impact evaluation of an asthma social marketing campaign targeting older adults. *BMC Public Health.* 2013;13:759.
15. **Kaplan SA, Garrett KE.** The use of logic models by community-based initiatives. *Eval Program Plann.* 2005;28(2):167-172.
16. **Sitaker M, Jernigan J, Ladd S, Patanian M.** Adapting logic models over time: The Washington State heart disease and stroke prevention program experience. *Prev Chronic Dis.* 2008;5(2):A60.
17. **Loberti AM, Dewsbury BM.** Using a logic model to direct backward design of curriculum. *J Microbiol Biol Educ.* 2018;19(3):19.3.105.
18. **Goeschel CA, Weiss WM, Pronovost PJ.** Using a logic model to design and evaluate quality and patient safety improvement programs. *Int J Qual Health Care.* 2012;24(4):330-337.
19. **Armstrong EG, Barsion SJ.** Using an outcomes-logic-model approach to evaluate a faculty development program for medical educators. *Acad Med.* 2006;81(5):483-488.
20. **WK Kellogg Foundation.** Using logic models to bring together planning, evaluation and action: Logic model development guide; 2004.
21. **Ten Cate O.** Nuts and bolts of entrustable professional activities. *J Grad Med Educ.* 2013;5(1):157-158.
22. **Fink, LD.** 1: Evaluating teaching: A new approach to an old problem. *To Improve the Academy.* 2008;26(1):3-21.
23. **Federal Emergency Management Agency.** *National Incident Management System (NIMS).* <https://training.fema.gov/nims/>. Accessed June 12, 2019.