



On the complete separation of unique ℓ_1 spreading models and the Lebesgue property of Banach spaces

Harrison Gaebler, Pavlos Motakis , and Bünyamin Sarı

Abstract. We construct a reflexive Banach space $X_{\mathcal{D}}$ with an unconditional basis such that all spreading models admitted by normalized block sequences in $X_{\mathcal{D}}$ are uniformly equivalent to the unit vector basis of ℓ_1 , yet every infinite-dimensional closed subspace of $X_{\mathcal{D}}$ fails the Lebesgue property. This is a new result in a program initiated by Odell in 2002 concerning the strong separation of asymptotic properties in Banach spaces.

1 Introduction

In Banach space theory, an asymptotic notion refers to a concept describing the behavior of the norm of linear combinations of vectors sampled in a specified asymptotic fashion from a certain type of structure. For example, the seminal notion of a spreading model of a bounded sequence $(x_n)_{n=1}^{\infty}$ is defined as the unit vector basis $(e_i)_{i=1}^{\infty}$ of c_{00} with the seminorm given by

$$\left| \sum_{i=1}^n a_i e_i \right| = \lim_{k_1 \rightarrow \infty} \lim_{k_2 \rightarrow \infty} \cdots \lim_{k_n \rightarrow \infty} \left\| \sum_{i=1}^n a_i x_{k_i} \right\|,$$

provided that this iterated limit exists for all choices of coefficients a_1, \dots, a_n . Here, the structure in question is a bounded sequence of vectors in a Banach space, and vectors are sampled over a sparse set of sufficiently large indices. Using Ramsey's Theorem, Brunel and Sucheston proved in [9] that every bounded sequence in a Banach space has a subsequence generating some spreading model. Similar notions about different types of structures have been introduced and studied over the years. One such example is the notion of asymptotic models, introduced by Odell and Halbeisen in [13], concerning the norm of linear combinations of vectors sampled from an array of sequences. A different type of such a notion is asymptotic spaces, introduced by Maurey, Milman, and Tomczak-Jaegerman in [17], concerning the norm of linear combinations of vectors picked in a finite-round two-player game between a vector chooser and a finite-codimensional subspace chooser. The different types of asymptotic notions are relevant to the study of many subjects in Banach

Received by the editors February 22, 2024; accepted August 19, 2024.

The second author was supported by NSERC Grant RGPIN-2021-03639.

AMS subject classification: 46B06, 46B20, 46B25, 46G12.

Keywords: Lebesgue property of Banach spaces, spreading models, complete separation of asymptotic structures.



space theory, such as the distortion problem in [18] using asymptotic spaces, the invariant subspace problem in reflexive Banach spaces in [4] using spreading models, the uniform approximation properties of bounded linear operators in [2] using joint spreading models, the coarse geometry of Banach spaces in [7] using asymptotic models, unique maximal ideals in the operators of stopping time-type Banach spaces in [16] using spreading models, and quotients algebras of operator spaces in [20] using asymptotic versions. The wide applicability of these tools has led to an independent study of their uniqueness properties and a desire to better understand their interconnections. Among the most outspoken promoters of this program was Odell, who in this context posed several questions of the following type (see [22], [23], and [15]): Let P, Q be properties of Banach spaces that are stable under passing to closed subspaces and assume that P implies Q . If a Banach space X satisfies Q , must it have an infinite-dimensional closed subspace satisfying P ? For such P and Q , if there exists a Banach space X satisfying Q and all infinite-dimensional subspaces of which fail P , then we say that P is completely separated from Q . For a detailed discussion on positive and negative results in this context, we refer the reader to [5]. We prove such a complete separation theorem for P the Lebesgue property and Q the property of admitting a unique ℓ_1 spreading model.

Let X be a Banach space. A function $f : [0, 1] \rightarrow X$ is Riemann-integrable if it is bounded and continuous up to a set of Lebesgue measure zero (see, for example, [12, Theorem 18] or [10, Theorem. 2.1.3.]). However, the converse statement is false in general. Indeed, it is easy to check that the function which maps the rationals in $[0, 1]$ to the unit vector basis of c_0 is Riemann-integrable. A Banach space X is therefore said to have the Lebesgue property if every Riemann-integrable function $f : [0, 1] \rightarrow X$ is continuous up to a set of Lebesgue measure zero. The first and third named authors recently in [11] characterized the Lebesgue property in terms of a new asymptotic notion. In particular, a Banach space X has the Lebesgue property if and only if every normalized basic sequence in X is Haar- ℓ_1^+ (see [11] or Section 2 of this paper for the precise definition of Haar- ℓ_1^+). It is also proved in [11] that for the properties

- (P1) every asymptotic model of X is equivalent to the unit vector basis of ℓ_1 ,
- (P2) X satisfies the Lebesgue property,
- (P3) every spreading model of X is equivalent to the unit vector basis of ℓ_1

the implications (P1) \implies (P2) \implies (P3) are true. However, (P2) $\not\implies$ (P1) because the Tsirelson sum of Tsirelson spaces, $(T \oplus T \oplus \dots)_T$, has the Lebesgue property but no unique asymptotic model (see, for example, [11]) and (P3) $\not\implies$ (P2) because there exist Schur spaces that do not have the Lebesgue property (e.g. [14],[21]) and Schur spaces satisfy (P3). The complete separation of (P1) from (P2) was proved in [11] by showing that the Banach space X_{i_w} constructed in [5] has the Lebesgue property, and yet, every subspace of X_{i_w} contains a c_0 -asymptotic model. The complete separation of (P2) from (P3) was not attempted in [11], and it is the purpose of the present article. By carrying out a substantial modification of the construction in [5], we are able to achieve this new result. We construct a Banach space $X_{\mathcal{D}}$ with an unconditional basis such that every spreading model of $X_{\mathcal{D}}$ is uniformly equivalent to the unit vector basis of ℓ_1 and such that every subspace of $X_{\mathcal{D}}$ fails the Lebesgue property.

Let us explain the main properties of $X_{\mathcal{D}}$. Denote by \mathcal{D} the binary tree $\cup_{n=0}^{\infty} \{0, 1\}^n$. Following standard convention, $\{0, 1\}^0$ is a singleton containing the empty sequence. For $n \in \mathbb{N} \cup \{0\}$ and $\lambda \in \{0, 1\}^n$, we write $|\lambda| = n$ and say λ has height n . We consider \mathcal{D} with the usual partial order with minimum the empty sequence, and, otherwise, $\lambda \leq \mu$ means $|\lambda| \leq |\mu|$ and $\lambda(i) = \mu(i)$, for $1 \leq i \leq |\lambda|$. The characterization of the Lebesgue property in terms of Haar- ℓ_1^+ sequences from [11] easily yields the following (see Remark 2.2): a Banach space X has the Lebesgue property if and only if, for every collection $(x_\lambda)_{\lambda \in \mathcal{D}}$ of norm-one vectors in X , there exists a constant $\theta > 0$ such that, for every $n \in \mathbb{N}$, there exists a subset $V \subset \{0, 1\}^n$ such that

$$\theta \leq \frac{1}{2^n} \left\| \sum_{\lambda \in V} x_{\mu_\lambda} \right\|$$

for some choice of nodes $(\mu_\lambda)_{\lambda \in V}$ with $\mu_\lambda \geq \lambda$. In the space $X_{\mathcal{D}}$, every infinite-dimensional subspace contains a lexicographically block collection $(x_\lambda)_{\lambda \in \mathcal{D}}$ of norm-one vectors such that for any $n \in \mathbb{N}$ and collection $(\mu_\lambda)_{\lambda \in \{0,1\}^n}$ in \mathcal{D} such that $\mu_\lambda \geq \lambda$, $\lambda \in \{0, 1\}^n$,

$$(1.1) \quad \max_{\lambda \in \{0,1\}^n} |a_\lambda| \leq \left\| \sum_{\lambda \in \{0,1\}^n} a_\lambda x_{\mu_\lambda} \right\| \leq 3 \max_{\lambda \in \{0,1\}^n} |a_\lambda|,$$

for any scalars $(a_\lambda)_{\lambda \in \{0,1\}^n}$. Therefore, $X_{\mathcal{D}}$ fails the Lebesgue property. At the same time, every normalized weakly null sequence in $X_{\mathcal{D}}$ has a subsequence generating a spreading model that is 102-equivalent to the unit vector basis of ℓ_1 . In particular, it follows from James’s characterization of reflexivity for spaces with an unconditional basis that $X_{\mathcal{D}}$ is reflexive.

Our construction relies on the method of saturation under constraints with increasing weights, also employed in [5]. The major difference lies in the involvement of an additional metric constraint. A distantly related version of this constraint was also present in [19], but its application and role were significantly different as it produced an operator-algebraic outcome (see [19, Section 2]). The earliest form of saturation under constraints is due to Odell and Schlumprecht (see [24] and [25]). It has since been extensively developed – for example, in the papers [4], [8], [5], and many others. Let us elaborate on the main ingredients of our construction. The norm of the space $X_{\mathcal{D}}$ is defined via a norming set; that is, there is a subset W of c_{00} containing the unit vector basis $(e_i^*)_{i=1}^{\infty}$ such that, for $x \in c_{00}$, $\|x\| = \sup\{|\langle f, x \rangle| : f \in W\}$, where $\langle \cdot, \cdot \rangle$ denotes the standard duality pairing of $c_{00} \times c_{00}$. The space $X_{\mathcal{D}}$ is the completion of c_{00} under this norm. We fix a pair of lacunary sequences $m_j, n_j, j \in \mathbb{N}$, of natural numbers. The first sequence is used to define a collection of weights $\{\prod_{i=1}^{\ell} m_{j_i} : j_1, \dots, j_{\ell} \in \mathbb{N}\}$, which is closed under multiplication. To each weight $\prod_{i=1}^{\ell} m_{j_i}$, we associate a Schreier family $\mathcal{S}_{n_{j_1} + n_{j_2} + \dots + n_{j_{\ell}}}$. This is a compact collection of finite subsets of \mathbb{N} recalled in Section 2. Every f in W satisfies $\|f\|_{\infty} \leq 1$, and it is either of the form $\pm e_i^*$ or

$$(1.2) \quad f = \frac{1}{w(f)} \sum_{q=1}^d f_q,$$

where $w(f) = \prod_{i=1}^{\ell} m_{j_i}$ and f_1, \dots, f_d are successively supported vectors such that the set $\{\min \text{supp}(f_q) : 1 \leq q \leq d\}$ is in the collection $\mathcal{S}_{n_{j_1} + n_{j_2} + \dots + n_{j_d}}$. A Banach space defined by a norming set on which only the aforementioned restrictions are imposed is asymptotic- ℓ_1 (see, for example, [1]), an asymptotic property implying having a unique ℓ_1 asymptotic model and thus also the Lebesgue property. To avoid this without compromising the uniform uniqueness of ℓ_1 spreading models, just as in [5], we additionally demand that for f in W as in (1.2), the sequence $w(f_1), \dots, w(f_d)$ is increasing sufficiently rapidly in relation to the supports of f_1, \dots, f_d . Imposing these additional restrictions on W yields a Banach space with a uniformly unique ℓ_1 spreading model and no asymptotic- ℓ_1 subspace; however, as it was proved in [11], it still satisfies the Lebesgue property. The intuitive explanation for this is the following: despite the imposed restrictions on W , there are sufficiently many functionals yielding the required ℓ_1^+ estimates on normalized collections $(x_\lambda)_{\lambda \in \mathcal{D}}$ in the defined Banach space.

To address this, we impose an additional metric constraint on W . Recall that \mathcal{D} is a totally bounded metric space with the metric $d(\lambda, \mu) = 2^{-n}$, where, if $\lambda \neq \mu$, $n \in \mathbb{N} \cup \{0\}$ is the maximum height for which there exists $v \in \{0, 1\}^n$ with $v \leq \lambda$ and $v \leq \mu$. We fix a bijection $\phi : \{m_j : j \in \mathbb{N}\} \rightarrow \mathcal{D}$, and to every $f \in W$ of weight $\prod_{i=1}^{\ell} m_{j_i}$, we associate the node $\lambda = \phi(m(f))$, where $m(f)$ is the maximum of $m_{j_1}, \dots, m_{j_\ell}$. We then impose on W the restriction that for every f as in (1.2), the nodes $\phi(m(f_1)), \dots, \phi(m(f_d))$ approximate some λ in the completion of \mathcal{D} , and the quantification of the approximation depends on the tuple $(n_{j_1}, \dots, n_{j_\ell})$ and the supports of the f_q , $1 \leq q \leq d$. Modulo some subtle but significant details, the construction of the space $X_{\mathcal{D}}$ is complete.

Using a standard technique, in every block-subspace Y of $X_{\mathcal{D}}$, we build a collection of vectors $(x_{m_j})_{j \in \mathbb{N}}$ such that, for every $j \in \mathbb{N}$, the norm of x_{m_j} can only be effectively estimated by functionals of weight $w(f) = m_j$; that is, x_{m_j} is an exact vector. We then consider the \mathcal{D} -indexed collection $(x_{\phi^{-1}(\lambda)})_{\lambda \in \mathcal{D}}$. This witnesses the failure of the Lebesgue property in Y . Indeed, for $n \in \mathbb{N}$, denote $\{0, 1\}^n = \{\lambda_1, \dots, \lambda_{2^n}\}$ and let $m_{j_1}, \dots, m_{j_{2^n}}$ such that $\phi(m_{j_1}) \geq \lambda_1, \dots, \phi(m_{j_{2^n}}) \geq \lambda_{2^n}$. Then the nodes $\phi(m_{j_1}), \dots, \phi(m_{j_{2^n}})$ form a 2^{-n} separated set in \mathcal{D} . For every $f \in W$ as in (1.2), $\phi(m(f_1)), \dots, \phi(m(f_d))$ approximate a λ in the completion of \mathcal{D} , and thus, roughly speaking, the components of f only contribute to the norm of very few of the $x_{m_{j_i}}$, $1 \leq i \leq 2^n$, yielding the ℓ_∞ -estimate (1.1).

Our paper is organized as follows. In Section 2, we recall the required terminology surrounding Schreier families and special convex combinations. We also recall the definition of a Haar- ℓ_1^+ sequence and remark on the formulation of the Lebesgue property in terms of normalized collections $(x_\lambda)_{\lambda \in \mathcal{D}}$. In Section 3, we define the space $X_{\mathcal{D}}$ and prove the uniform uniqueness of ℓ_1 spreading models. In Section 4, we define an auxiliary Banach space $X_{\mathcal{D}}^{\text{aux}, \delta}$ and prove ℓ_∞ -type estimates for linear combinations of “simple” vectors in this space. In Section 5, we define and study rapidly increasing sequences (RISs) and the basic inequality. The former comprise a special class of sequences in $X_{\mathcal{D}}$, and the latter provides bounds for their linear combinations in terms of linear combinations of basis vectors in the auxiliary space. In Section 6, we combine all ingredients prepared in Sections 4 and 5 to construct in every infinite-dimensional

closed subspace of $X_{\mathcal{D}}$ a normalized collection $(x_\lambda)_{\lambda \in \mathcal{D}}$ witnessing the failure of the Lebesgue property.

2 Preliminary information

Let $A, B \subset \mathbb{N}$. If $\max(A) < \min(B)$, then we write $A < B$. We also write $n \leq A$ if $n \in \mathbb{N}$ is at most $\min(A)$ and, by convention, $\emptyset < A$ and $A < \emptyset$ for every $A \subset \mathbb{N}$. We denote by c_{00} the vector space that consists of sequences of real numbers that have at most finitely-many nonzero terms, and we let (e_i) be the unit vector basis of c_{00} . Note that the members of the norming set W are themselves members of c_{00} and, in particular, that $\{\pm e_i\}_{i=1}^\infty \subset W$. For this reason, we write $e_i^* \in c_{00}$ when referring to e_i as a norming functional. The support of $x = (c_i) \in c_{00}$ is the set $\text{supp}(x) = \{i \in \mathbb{N} \mid c_i \neq 0\}$, and we say that vectors $x_1, \dots, x_n \in c_{00}$ are successive if $\text{supp}(x_{i-1}) < \text{supp}(x_i)$ for $2 \leq i \leq n$. In this case, we use the shorthand notation $x_1 < \dots < x_n$, and we also write $n \leq x$ if $n \leq \min \text{supp}(x)$. A finite or infinite sequence of successive vectors in c_{00} is said to be a block sequence of (e_i) .

The Schreier sets are an increasing sequence of sets of finite subsets of \mathbb{N} . These sets are defined inductively as follows:

$$S_0 = \{\{i\} \mid i \in \mathbb{N}\} \text{ and } S_1 = \{F \subset \mathbb{N} \mid \text{card}(F) \leq 2\},$$

and we write that

$$S_{n+1} = \{F \subset \mathbb{N} \mid F = \cup_{i=1}^d F_i \text{ where } d \leq F_1 < \dots < F_d \in S_n\}$$

once S_n has been defined. We also define the set $\mathcal{A}_3 = \{F \subset \mathbb{N} \mid \text{card}(F) \leq 3\}$, and we will be interested in sets of the form

$$S_n * \mathcal{A}_3 = \{F \subset \mathbb{N} \mid F = \cup_{i=1}^d F_i \text{ where } F_1 < \dots < F_d \in \mathcal{A}_3 \text{ and } \{\min(F_i)\}_{i=1}^d \in S_n\}.$$

It is easy to see that $F \in S_n * \mathcal{A}_3$ is the union of at most three sets in S_n . A block sequence $x_1 < \dots < x_k \in c_{00}$ is said to be S_n -admissible (resp. $S_n * \mathcal{A}_3$ -admissible) if $\{\min \text{supp}(x_i)\}_{i=1}^k \in S_n$ (resp. $\{\min \text{supp}(x_i)\}_{i=1}^k \in S_n * \mathcal{A}_3$).

Let $n \in \mathbb{N}$ and let $\varepsilon > 0$. Then, the vector $x = \sum_{i \in F} c_i e_i \in c_{00}$ is said to be an (n, ε) -basic special convex combination ((n, ε) -basic scc) if

- (1) $F \in S_n$, $c_i \geq 0$ for each $i \in F$, and $\sum_{i \in F} c_i = 1$,
- (2) For every $G \subset F$ with $G \in S_{n-1}$, we have that $\sum_{i \in G} c_i < \varepsilon$,

and we refer the reader to [1], [3], or [6] for more information. In particular, note that $\sum_{i \in G} c_i < 3\varepsilon$ if $G \subset F$ with $G \in S_k * \mathcal{A}_3$ for $k < n$, and there is also the following useful result which is Proposition 2.3 of [6].

Proposition 2.1 *Let $n \in \mathbb{N}$. Then, for every infinite subset $M \subset \mathbb{N}$ and for every $F \subset M$ such that $F \in S_n$ is maximal with respect to inclusion, there exist real numbers $(c_i)_{i \in F}$ so that the vector $x = \sum_{i \in F} c_i e_i \in c_{00}$ is an $(n, 3/\min(F))$ -basic scc.*

Similarly, if $x_1 < \dots < x_d \in c_{00}$ and if $s_i = \min \text{supp}(x_i)$ for each $1 \leq i \leq d$, then the vector $x = \sum_{i=1}^d c_i x_i$ is said to be an (n, ε) -special convex combination ((n, ε) -scc) if

$\sum_{i=1}^d c_i e_{s_i}$ is an (n, ε) -basic scc. It follows in this case that $\sum_{i=1}^d c_i e_{t_i}$ is an $(n, 2\varepsilon)$ -basic scc as well where $t_i = \max \text{supp}(x_i)$ for each $1 \leq i \leq d$ because

$$\sum_{i \in G} c_i = \sum_{i \in G_1} c_i + \sum_{i \in G_2} c_i < \varepsilon + \varepsilon = 2\varepsilon$$

if $G \subset \{t_i\}_{i=1}^d$ is S_{n-1} -admissible and is therefore a translation to the right of at most two sets $G_1, G_2 \subset \{s_i\}_{i=1}^d$ with $G_1, G_2 \in S_{n-1}$.

Finally, two specific asymptotic structures appear in this paper. The first is that of a spreading model from [9], recalled in page 2. It is a consequence of Ramsey’s theorem that every normalized basic sequence (x_i) in a Banach space X has a subsequence that generates a spreading model of X . We now define the notion of a Haar- ℓ_1^+ sequence, and we need first the definition of a Haar system of partitions of \mathbb{N} .

Definition 2.1 A collection $(A_j^n)_{j=0, n \in \mathbb{N}}^{2^n-1}$ of infinite subsets of \mathbb{N} is said to be a Haar system of partitions of \mathbb{N} if the following two conditions are satisfied:

- (1) For every $n \in \mathbb{N}$, we have that $\cup_{j=0}^{2^n-1} A_j^n = \mathbb{N}$ and $A_j^n \cap A_{j'}^n = \emptyset$ if $j \neq j'$.
- (2) For every $n \in \mathbb{N}$ and for every $0 \leq j \leq 2^n - 1$, we have that $A_j^n = A_{2j}^{n+1} \cup A_{2j+1}^{n+1}$.

The connection between the Haar- ℓ_1^+ condition (defined below) and Riemann integrability is due to the fact that the infinite set A_j^n of the Haar system of partitions of \mathbb{N} corresponds to the dyadic rational numbers in the dyadic interval $(\frac{j}{2^n}, \frac{j+1}{2^n})$ (see, for example, [11, Theorem 3.1]).

Definition 2.2 A normalized basic sequence (x_i) in X is said to be Haar- ℓ_1^+ if, for every Haar system of partitions $(A_j^n)_{j=0, n \in \mathbb{N}}^{2^n-1}$ of \mathbb{N} , there exists a constant $C > 0$ such that

$$C \leq \limsup_{n \rightarrow \infty} \left\{ \frac{1}{2^m} \left\| \sum_{j=0}^{2^m-1} x_{i_j} \right\| \mid m \geq n \text{ and } 2^m \leq i_j \in A_j^m \right\}.$$

Remark 2.2 For a Banach space X , the following are equivalent.

- (1) X has the Lebesgue property.
- (2) Every normalized basic sequence in X is Haar- ℓ_1^+ .
- (3) For every collection $(x_\lambda)_{\lambda \in \mathcal{D}}$ of norm-one vectors in X , there exists a constant $\theta > 0$ such that, for every $n \in \mathbb{N}$, there exists a subset $V \subset \{0, 1\}^n$ such that

$$\theta \leq \frac{1}{2^n} \left\| \sum_{\lambda \in V} x_{\mu_\lambda} \right\|$$

for some choice of nodes $(\mu_\lambda)_{\lambda \in V}$ with $\mu_\lambda \geq \lambda$.

Indeed, the assertion (1) \iff (2) is established in [11]. In addition, it is easy to see that (1) \implies (3) as collection witnessing the negation of (3) can be easily used to construct a Dirichlet-type function with Riemann integral zero. Lastly, for (3) \implies (2), if there exists a normalized basic sequence (x_i) in X that is not Haar- ℓ_1^+ , then it is easy to construct a collection $(x_\lambda)_{\lambda \in \mathcal{D}}$ of norm-one vectors for which the claim of (3) does not hold as follows. Fix an order-preserving bijection $\mathcal{D} \rightarrow \{A_j^n \mid n \in \mathbb{N}, 0 \leq j \leq 2^n - 1\}$ such that, for every $n \in \mathbb{N}$, $\lambda \mapsto A_{j_\lambda}^{|\lambda|}$, for some $0 \leq j_\lambda \leq 2^{|\lambda|} - 1$.

Then, for every $\lambda \in \mathcal{D}$, choose $i_\lambda \in A_{j_\lambda}^{|\lambda|}$ such that $i_\lambda \geq 2^{|\lambda|}$ and put $x_\lambda = x_{i_\lambda}$. Note that $(x_\lambda)_{\lambda \in \mathcal{D}}$ can be chosen Schauder basic in the lexicographical order of \mathcal{D} .

3 Construction of $X_{\mathcal{D}}$ and uniformly unique ℓ_1 spreading model

In this section, we define the norming set W inducing the Banach space $X_{\mathcal{D}}$. It is defined as an increasing union $W = \cup_{k=0}^\infty W_k$, where $W_0 = \{\pm e_i^*\}_{i=1}^\infty$, and, in each step, having defined W_k , we formulate the operations and constraints that are used to produce W_{k+1} . The quantification of the metric constraints discussed in the introduction is defined inductively and carries a certain complexity; this is necessary to balance the uniform uniqueness of ℓ_1 spreading models with the ℓ_∞ estimates on tree-indexed collection. At the end of this section, we prove that every normalized weakly null sequence in $X_{\mathcal{D}}$ has a subsequence generating a spreading model that is 102-equivalent to the unit vector basis of ℓ_1 .

Fix a pair of strictly increasing sequences (m_j) and (n_j) of positive integers such that $m_1 = 100, n_1 = 1$,

- (1) $m_{j+1} \geq j^2 m_j^2$, for every $j \in \mathbb{N}$,
- (2) $n_{j+1} > \frac{4}{\log(100)} n_j \log(m_{j+1})$.

Let $\mathcal{D} = \cup_{n=0}^\infty \{0, 1\}^n$ with the partial order and metric d defined in the introduction. We note that the completion $\overline{\mathcal{D}}$ of \mathcal{D} is a compact metric space. Denote $\mathcal{M} = \{\infty\} \cup \{m_j : j \in \mathbb{N}\}$ and let $\phi : \mathcal{M} \rightarrow \mathcal{D}$ be a bijection so that the nodes of \mathcal{D} are ordered lexicographically with respect to \mathcal{M} ; that is, $\phi(\infty) = \emptyset, \phi(m_1) = (0), \phi(m_2) = (1), \phi(m_3) = (0, 0)$, and so on. Lastly, as a matter of notational convenience, we always write $d(m_j, \cdot)$ instead of the formally correct $d(\phi(m_j), \cdot)$.

Now, let $W_0 = \{\pm e_i^*\}_{i=1}^\infty \subset c_{00}$ and write $w(\pm e_i^*) = m(\pm e_i^*) = \infty$ for each $i \in \mathbb{N}$. Then, define the set

$$W_1 = W_0 \cup \left\{ \frac{1}{m_{j_1} \cdots m_{j_\ell}} \sum_{k=1}^\ell e_{i_k}^* \mid \begin{array}{l} e_{i_1}^* < \dots < e_{i_\ell}^* \text{ are} \\ S_{n_{j_1} + \dots + n_{j_\ell}}\text{-admissible} \end{array} \right\},$$

where $w(f) = m_{j_1} \cdots m_{j_\ell}, \bar{w}(f) = (m_{j_1}, \dots, m_{j_\ell})$, and $m(f) = \max_{1 \leq z \leq \ell} m_{j_z}$ for $f \in W_1 \setminus W_0$. For f in $W_1 \setminus W_0$, $\bar{w}(f)$ may not be uniquely defined, and thus neither are $w(f)$ and $m(f)$. The notation $\bar{w}(f) = (m_{j_1}, \dots, m_{j_\ell})$ means that f admits a representation as above, and it is fixed while working with f . This will also apply to $f \in W_k, k > 1$. This is common in this type of construction and does not pose a problem. In fact, insisting on uniquely defining the weight may interfere with constructing functionals that are used to prove the properties of the space. We next define the following notion of proximity with respect to $\overline{\mathcal{D}}$ for functionals in W_1 .

Definition 3.1 The functionals $f_1 < \dots < f_d$ in W_1 are said to be (n, λ) -packed for some $\lambda \in \overline{\mathcal{D}}$ if the following conditions are satisfied:

- (1) $f_1 < \dots < f_d$ are S_n -admissible,
- (2) $w(f_i) \geq 2^{\max \text{supp}(f_{i-1})}$ for $2 \leq i \leq d$,
- (3) $d(m(f_i), \lambda) \leq 2^{-\max \text{supp}(f_{i-1})}$ for $2 \leq i \leq d$,

and if, in addition, we have that $\max_{1 \leq i \leq d} d(m(f_i), \lambda) \leq \eta$. Then $f_1 < \dots < f_d$ are said to be (n, λ, η) -packed.

This definition allows for the following inductive definition as well.

Definition 3.2 The functionals $f_1 < \dots < f_d$ in W_1 are said to be $(n_{j_1}, \dots, n_{j_d}, \lambda)$ -packed for some $\lambda \in \overline{\mathcal{D}}$ if the following claims hold:

- (1) $w(f_i) \geq 2^{\max \text{supp}(f_{i-1})}$ for $2 \leq i \leq d$,
- (2) There exists a partition $F_1 < \dots < F_M$ of $\{1, \dots, d\}$ such that the set $\{f_{\min(F_k)}\}_{k=1}^M$ is $S_{n_{j_i}}$ -admissible,
- (3) There exist nodes $\lambda_1, \dots, \lambda_M \in \overline{\mathcal{D}}$ such that
 - (a) $(f_i)_{i \in F_1}$ is $(n_{j_1}, \dots, n_{j_{l-1}}, \lambda_1)$ -packed,
 - (b) $(f_i)_{i \in F_k}$ is $(n_{j_1}, \dots, n_{j_{l-1}}, \lambda_k, 2^{-\max \text{supp}(f_{\max(F_{k-1})})})$ -packed for $2 \leq k \leq M$,
 - (c) $d(\lambda_k, \lambda) \leq 2^{-\max \text{supp}(f_{\max(F_{k-1})})}$ for $2 \leq k \leq M$,

and if, in addition, we have that $\max_{1 \leq i \leq d} d(m(f_i), \lambda) \leq \eta$, then we say that $f_1 < \dots < f_d$ are $(n_{j_1}, \dots, n_{j_d}, \lambda, \eta)$ -packed.

We now define the next set of weighted functionals

$$W_2 = W_1 \cup \left\{ \frac{1}{m_{j_1} \cdots m_{j_d}} \sum_{i=1}^d f_i \mid \begin{array}{l} f_1 < \dots < f_d \text{ are in } W_1 \text{ and are} \\ (n_{j_1}, \dots, n_{j_d}, \lambda)\text{-packed for some } \lambda \in \overline{\mathcal{D}} \end{array} \right\}$$

with $w(f) = m_{j_1} \cdots m_{j_d}$, $\vec{w}(f) = (m_{j_1}, \dots, m_{j_d})$, and $m(f) = \max_{1 \leq z \leq l} m_{j_z}$. Now, once we have defined the sets W_0, W_1, \dots, W_k , we replace W_1 in Definitions 3.1 and 3.2 with W_k , and we write that

$$W_{k+1} = W_k \cup \left\{ \frac{1}{m_{j_1} \cdots m_{j_d}} \sum_{i=1}^d f_i \mid \begin{array}{l} f_1 < \dots < f_d \text{ are in } W_k \text{ and are} \\ (n_{j_1}, \dots, n_{j_d}, \lambda)\text{-packed for some } \lambda \in \overline{\mathcal{D}} \end{array} \right\}$$

for each $k \in \mathbb{N}$. We then define our norming set $W = \cup_{k=0}^\infty W_k$, and we let the Banach space $X_{\mathcal{D}}$ be the completion of c_{00} with respect to the norm that is given by $\|x\| = \sup\{f(x) \mid f \in W\}$ for $x \in c_{00}$.

Remark 3.1 By induction on k , where $W = \cup_{k=0}^\infty W_k$, it is easy to prove the following.

- (1) For every $(n_{j_1}, \dots, n_{j_\ell}, \lambda)$ -packed sequence $f_1 < \dots < f_d$ in W ,

$$f = \frac{1}{m_{j_1} \cdots m_{j_\ell}} \sum_{i=1}^d f_i$$

is in W .

Incorporating a nested induction on ℓ , the following items can be shown.

- (2) A subsequence of a $(n_{j_1}, \dots, n_{j_\ell}, \lambda)$ -packed sequence $f_1 < \dots < f_d$ in W is also $(n_{j_1}, \dots, n_{j_\ell}, \lambda)$ -packed.
- (3) If $f_1 < \dots < f_d$ is a $(n_{j_1}, \dots, n_{j_\ell}, \lambda)$ -packed sequence in W and $g_1 < \dots < g_d$ are in c_{00} such that, for $1 \leq i \leq d$ and $s \in \text{supp}(g_i)$, we have $|g_i(e_s)| = |f_i(e_s)|$, then g_1, \dots, g_d are also in W , and they are $(n_{j_1}, \dots, n_{j_\ell}, \lambda)$ -packed.

In particular, (2) and (3) yield that W is closed under restricting the supports of its members to subsets and changing the signs of their scalar coefficients, and thus, (e_i) is a 1-unconditional basis for $X_{\mathcal{D}}$.

We now prove that every spreading model of $X_{\mathcal{D}}$ is uniformly equivalent to the unit vector basis of ℓ_1 . In fact, the precise statement of the following theorem is slightly stronger than this assertion; it will be required in its full generality later, in the proof of Proposition 5.2.

Theorem 3.2 *Let (x_r) be a normalized block sequence of (e_i) . Then, there exists an infinite subset $\Omega \subset \mathbb{N}$ such that, for every $j_0 \in \mathbb{N}$, there is the estimate*

$$\left\| \sum_{r \in F} c_r x_r \right\| \geq 0.99 \frac{1}{m_{j_0}} \sum_{r \in \Gamma} |c_r|$$

for every subset $\Gamma \subset \Omega$ such that $(x_r)_{r \in \Gamma}$ are $S_{n_{j_0}}$ -admissible and for every choice $(c_r)_{r \in \Gamma}$ of scalars.

Proof Choose for every $r \in \mathbb{N}$ a norming functional $f_r \in W$ so that $f_r(x_r) = 1$ and so that $\text{supp}(f_r) \subset \text{supp}(x_r)$. Then, there are two cases to consider.

(i) $\limsup_{r \rightarrow \infty} w(f_r) = L < \infty$

Here, we may pass to an infinite subset $\Omega_1 \subset \mathbb{N}$ so that $w(f_r) = m_{j_1} \cdots m_{j_l}$ for each $r \in \Omega_1$. Note that each f_r with $r \in \Omega_1$ is then $(n_{j_1}, \dots, n_{j_l}, \lambda_r)$ -packed for some $\lambda_r \in \overline{\mathcal{D}}$. We may then pass to a further infinite subset $\Omega_2 \subset \Omega_1$ so that $d(\lambda_r, \lambda) \leq \frac{1}{2} \cdot 2^{-\max \text{supp}(x_{r-1})}$ for each $r \in \Omega_2 \setminus \{\min(\Omega_2)\}$ and for some $\lambda \in \overline{\mathcal{D}}$. Now, let $\Omega = \Omega_2$ and consider the subsequence $(x_r)_{r \in \Omega_2}$. Let $\Gamma \subset \Omega$ be a subset such that the block vectors $(x_r)_{r \in \Gamma}$ are $S_{n_{j_0}}$ -admissible and let

$$f_r = \frac{1}{m_{j_1} \cdots m_{j_l}} \sum_{q=1}^{d_r} f_q^r \quad r \in \Gamma$$

be the associated norming functionals. For each $r \in \Gamma$, let $F_1^r < \dots < F_{M_r}^r$ be the partition of $\{1, \dots, d_r\}$ and let $\lambda_1^r, \dots, \lambda_{M_r}^r \in \overline{\mathcal{D}}$ be the corresponding nodes so that Definition 3.2 is satisfied because the functionals $f_1^r < \dots < f_{d_r}^r$ are $(n_{j_1}, \dots, n_{j_l}, \lambda_r)$ -packed. We will now show that the concatenation

$$f_{\min(F_2^{\min(\Gamma)})}^{\min(\Gamma)} < \dots < f_{d_{\min(\Gamma)}}^{\min(\Gamma)} < \dots < f_{\min(F_2^{\max(\Gamma)})}^{\max(\Gamma)} < \dots < f_{d_{\max(\Gamma)}}^{\max(\Gamma)}$$

is in W . First, note that there is some $\kappa \in \mathbb{N}$ so that each functional in this concatenation is a member of W_{κ} . It therefore remains to show that these functionals are $(n_{j_1}, \dots, n_{j_l}, n_{j_0}, \lambda)$ -packed. Indeed, it follows immediately that they are $S_{n_{j_1} + \dots + n_{j_l} + n_{j_0}}$ -admissible and are very fast-growing in the sense of Definition 3.2. Then, let $F_{\min(\Gamma)} < \dots < F_{\max(\Gamma)}$ be the partition of the number of functionals in the above concatenation such that $(f_i)_{i \in F_r} = \{f_{\min(F_2^r)}^r, \dots, f_{d_r}^r\}$ and note that $\{f_{\min(F_r)}\}_{r \in \Gamma} = \{f_{\min(F_2^r)}^r\}_{r \in \Gamma}$ is $S_{n_{j_0}}$ -admissible. Moreover,

- (1) $(f_i)_{i \in F_1} = \{f_{\min(F_2^1)}^1, \dots, f_{d_1}^1\}$ is $(n_{j_1}, \dots, n_{j_l}, \lambda_1)$ -packed.
- (2) $(f_i)_{i \in F_r} = \{f_{\min(F_2^r)}^r, \dots, f_{d_r}^r\}$ is $(n_{j_1}, \dots, n_{j_l}, \lambda_r)$ -packed for each $r \in \Gamma \setminus \{\min(\Gamma)\}$ and, in addition, for each f_i with $i \in F_k^r$ (where $2 \leq k \leq M_r$), there is

$$d(m(f_i), \lambda_r) \leq d(m(f_i), \lambda_k^r) + d(\lambda_k^r, \lambda_r) \leq 2 \cdot 2^{-\max \text{supp}(f_{\max(F_{k-1}^r)})} \leq 2^{-\max \text{supp}(f_{\max(F_{r-1})})}$$

so that, in fact, $(f_i)_{i \in F_r}$ is $(n_{j_1}, \dots, n_{j_l}, \lambda_r, 2^{-\max \text{supp}(f_{\max(F_{r-1})})})$ -packed.

(3) $d(\lambda_r, \lambda) \leq \frac{1}{2} \cdot 2^{-\max \text{supp}(x_{r-1})} \leq 2^{-\max \text{supp}(f_{\max(F_{r-1})})}$ for each $r \in \Gamma \setminus \{\min(\Gamma)\}$.

Putting these pieces together, it follows that the functional

$$\mathcal{F} = \frac{1}{m_{j_1} \cdots m_{j_l} m_{j_0}} \sum_{r \in \Gamma} \sum_{q=\min(F_2^r)}^{d_r} f_q^r$$

is $(n_{j_1}, \dots, n_{j_l}, n_{j_0}, \lambda)$ -packed and is therefore a member of $W_{\kappa+1} \subset W$. Now, because $(x_r)_{r \in \Omega}$ is itself 1-unconditional, there is the estimate

$$\begin{aligned} \left\| \sum_{r \in \Gamma} c_r x_r \right\| &= \left\| \sum_{r \in \Gamma} |c_r| x_r \right\| \geq \mathcal{F} \left(\sum_{r \in \Gamma} |c_r| x_r \right) \\ &= \frac{1}{m_{j_1} \cdots m_{j_l} m_{j_0}} \sum_{r \in \Gamma} \sum_{q=\min(F_2^r)}^{d_r} f_q^r \left(\sum_{p \in \Gamma} |c_p| x_p \right) \\ &= \frac{1}{m_{j_0}} \sum_{r \in \Gamma} \frac{|c_r|}{m_{j_1} \cdots m_{j_l}} \sum_{q=\min(F_2^r)}^{d_r} f_q^r(x_r) \\ &= \frac{1}{m_{j_0}} \sum_{r \in \Gamma} |c_r| \left(f_r(x_r) - \frac{1}{m_{j_1} \cdots m_{j_l}} \sum_{q \in F_1^r} f_q^r(x_r) \right) \\ &\geq \frac{1}{m_{j_0}} \sum_{r \in \Gamma} |c_r| \left(1 - \frac{1}{m_{j_1}} \cdot \frac{1}{m_{j_1} \cdots m_{j_{l-1}}} \sum_{q \in F_1^r} f_q^r(x_r) \right) \\ &\geq \frac{1}{m_{j_0}} \sum_{r \in \Gamma} |c_r| \left(1 - \frac{1}{m_1} \right) = \frac{m_1 - 1}{m_1 m_{j_0}} \sum_{r \in \Gamma} |c_r|, \end{aligned}$$

where we have used the fact that $\frac{1}{m_{j_1} \cdots m_{j_{l-1}}} \sum_{q \in F_1^r} f_q^r \in W$, and this completes the proof for the first case.

(ii) $\limsup_{r \rightarrow \infty} w(f_r) = \infty$

In this case, there exists an infinite subset $\Omega_1 \subset \mathbb{N}$ such that $w(f_r)$ for $r \in \Omega_1 \setminus \{\min(\Omega_1)\}$ is very fast-growing in the sense of Definition 3.1. Then, let λ_r be the node in $\overline{\mathcal{D}}$ that corresponds to $m(f_r)$ for each $r \in \Omega_1$. We may then pass to a further infinite subset $\Omega_2 \subset \Omega_1$ so that $d(\lambda_r, \lambda) \leq \frac{1}{2} \cdot 2^{-\max \text{supp}(x_{r-1})}$ for some $\lambda \in \overline{\mathcal{D}}$ and for each $r \in \Omega_2 \setminus \{\min(\Omega_2)\}$. Now, let $\Omega = \Omega_2$ and consider the subsequence $(x_r)_{r \in \Omega}$. Let $\Gamma \subset \Omega$ be a subset such that the block vectors $(x_r)_{r \in \Gamma}$ are $S_{n_{j_0}}$ -admissible and let $f_{\min(\Gamma)} < \dots < f_{\max(\Gamma)}$ be the corresponding norming functionals. Note that there is some $\kappa \in \mathbb{N}$ so that $f_r \in W_\kappa$ for each $r \in \Gamma$. Note also that, by construction, these norming functionals are $S_{n_{j_0}}$ -admissible and are very fast-growing in the sense of Definition 3.1. Moreover, there is the estimate

$$d(\lambda_r, \lambda) \leq 2^{-\max \text{supp}(x_{r-1})} \leq 2^{-\max \text{supp}(f_{r-1})}$$

for each $r \in \Gamma \setminus \{\min(\Gamma)\}$ so that these functionals are (n_{j_0}, λ) -packed. Then, we have that $\frac{1}{m_{j_0}} \sum_{r \in \Gamma} f_r \in W_{\kappa+1} \subset W$ so that

$$\begin{aligned} \left\| \sum_{r \in \Gamma} c_r x_r \right\| &= \left\| \sum_{r \in \Gamma} |c_r| x_r \right\| \geq \frac{1}{m_{j_0}} \sum_{r \in \Gamma} f_r \left(\sum_{p \in \Gamma} |c_p| x_p \right) \\ &= \frac{1}{m_{j_0}} \sum_{r \in \Gamma} |c_r| f_r(x_r) = \frac{1}{m_{j_0}} \sum_{r \in \Gamma} |c_r| \geq \frac{m_1 - 1}{m_1 m_{j_0}} \sum_{r \in \Gamma} |c_r|, \end{aligned}$$

and this completes the proof of the theorem. ■

4 Auxiliary Banach spaces and c_0 -type estimate

So far, we have proved that all spreading models admitted by normalized weakly null sequences in $X_{\mathcal{D}}$ are uniformly equivalent to the unit vector basis of ℓ_1 . The remaining goal is to define in each subspace of $X_{\mathcal{D}}$ a normalized collection $(x_\lambda)_{\lambda \in \mathcal{D}}$ satisfying the ℓ_∞ -estimate (1.1), and this will require some effort. As a first step, we define an auxiliary norming set $W^{\text{aux}, \delta}$ inducing a norm on c_{00} and an auxiliary Banach space $X_{\mathcal{D}}^{\text{aux}, \delta}$. We then prove estimates for the norm of linear combinations of “simple” vectors, specifically weighted (n, ε) basic special convex combinations, in the auxiliary space. These will be relevant later when a tool called the basic inequality will yield high-precision estimates of linear combinations of special sequences of vectors, called rapidly increasing sequences, in $X_{\mathcal{D}}$, in terms of sequences of basis vectors in $X_{\mathcal{D}}^{\text{aux}, \delta}$.

We derive in this section a c_0 -type estimate for the action of auxiliary functionals on weighted (n, ε) -basic scc vectors. The idea is that the functional evaluates to something small if its weight is in some sense a mismatch to the (n, ε) -basic scc vector weight. Let us first define the auxiliary functionals and auxiliary Banach spaces.

We will define an auxiliary Banach space for each $\delta > 0$. First, fix $\delta > 0$ and let $W_0^{\text{aux}, \delta} = \{\pm e_i^*\}_{i=1}^\infty \subset c_{00}$ as before. Then, also as before, define $w(\pm e_i^*) = m(\pm e_i^*) = \infty$ and define the set of auxiliary weighted functionals

$$W_1^{\text{aux}, \delta} = W_0^{\text{aux}, \delta} \cup \left\{ \frac{1}{m_{j_1} \cdots m_{j_l}} \sum_{k=1}^d e_{i_k}^* \mid \begin{array}{l} e_{i_1}^* < \dots < e_{i_d}^* \text{ are} \\ S_{n_{j_1} + \dots + n_{j_l}} * \mathcal{A}_3\text{-admissible} \end{array} \right\},$$

where $w(f) = m_{j_1} \cdots m_{j_l}$, $m(f) = \max_{1 \leq z \leq l} m_{j_z}$, and $\vec{w}(f) = (m_{j_1}, \dots, m_{j_l})$, for $f \in W_1^{\text{aux}, \delta} \setminus W_0^{\text{aux}, \delta}$. We next define the following notion of proximity with respect to $\overline{\mathcal{D}}$ for functionals in $W_1^{\text{aux}, \delta}$.

Definition 4.1 The functionals $f_1 < \dots < f_d$ in $W_1^{\text{aux}, \delta}$ are said to be $(n, \lambda)_\delta$ -packed for some $\lambda \in \overline{\mathcal{D}}$ if the following conditions are satisfied:

- (1) $f_1 < \dots < f_d$ are $S_n * \mathcal{A}_3$ -admissible,
- (2) $w(f_i) \geq 2^{\max \text{supp}(f_{i-2})}$ for $3 \leq i \leq d$,
- (3) $d(m(f_i), \lambda) \leq \delta$ for $f_i \notin W_0^{\text{aux}, \delta}$ and for $2 \leq i \leq d$.

This definition is the base case for the following inductive definition.

Definition 4.2 The functionals $f_1 < \dots < f_d$ in $W_1^{\text{aux},\delta}$ are said to be $(n_{j_1}, \dots, n_{j_l}, \lambda)_\delta$ -packed for some $\lambda \in \overline{\mathcal{D}}$ if the following conditions hold:

- (1) $w(f_i) \geq 2^{\max \text{supp}(f_{i-2})}$ for $3 \leq i \leq d$,
- (2) There exists a partition $F_1 < \dots < F_M$ of $\{1, \dots, d\}$ such that the set $\{f_{\min(F_k)}\}_{k=1}^M$ is $S_{n_{j_l}}$ -admissible and such that the functionals $(f_i)_{i \in F_k}$ are $S_{n_{j_1} + \dots + n_{j_{l-1}}} * \mathcal{A}_3$ -admissible for $1 \leq k \leq M$,
- (3) There exists a node $\lambda_1 \in \overline{\mathcal{D}}$ such that $(f_i)_{i \in F_1}$ is $(n_{j_1}, \dots, n_{j_{l-1}}, \lambda_1)_\delta$ -packed,
- (4) $d(m(f_i), \lambda) \leq \delta$ for $f_i \in (\cup_{k=2}^M F_k) \setminus W_0^{\text{aux},\delta}$.

It is worth noting that a subset of $(n_{j_1}, \dots, n_{j_l}, \lambda)_\delta$ -packed functionals in $W_1^{\text{aux},\delta}$ is still $(n_{j_1}, \dots, n_{j_l}, \lambda)_\delta$ -packed, and we now define the next set of auxiliary weighted functionals

$$W_2^{\text{aux},\delta} = W_1^{\text{aux},\delta} \cup \left\{ \frac{1}{m_{j_1} \dots m_{j_l}} \sum_{i=1}^d f_i \mid \begin{array}{l} f_1 < \dots < f_d \text{ are in } W_1^{\text{aux},\delta} \text{ and are} \\ (n_{j_1}, \dots, n_{j_l}, \lambda)_\delta \text{-packed for some } \lambda \in \overline{\mathcal{D}} \end{array} \right\}$$

with $w(f) = m_{j_1} \dots m_{j_l}$, $m(f) = \max_{1 \leq z \leq l} m_{j_z}$, and $\vec{w}(f) = (m_{j_1}, \dots, m_{j_l})$. Now, once we have defined the sets $W_0^{\text{aux},\delta}, W_1^{\text{aux},\delta}, \dots, W_k^{\text{aux},\delta}$, we replace $W_1^{\text{aux},\delta}$ in Definitions 4.1 and 4.2 with $W_k^{\text{aux},\delta}$ and, we write that

$$W_{k+1}^{\text{aux},\delta} = W_k^{\text{aux},\delta} \cup \left\{ \frac{1}{m_{j_1} \dots m_{j_l}} \sum_{i=1}^d f_i \mid \begin{array}{l} f_1 < \dots < f_d \text{ are in } W_k^{\text{aux},\delta} \text{ and are} \\ (n_{j_1}, \dots, n_{j_l}, \lambda)_\delta \text{-packed for some } \lambda \in \overline{\mathcal{D}} \end{array} \right\}$$

for each $k \in \mathbb{N}$. We then define our norming set $W^{\text{aux},\delta} = \cup_{k=0}^\infty W_k^{\text{aux},\delta}$, and we let the Banach space $X_{\overline{\mathcal{D}}}^{\text{aux},\delta}$ be the completion of c_{00} with respect to the norm that is given by $\|x\|_{W^{\text{aux},\delta}} = \sup\{f(x) \mid f \in W^{\text{aux},\delta}\}$ for $x \in c_{00}$. We will need the following lemma before we can prove the main result of this section.

Lemma 4.1 Let $f = \frac{1}{m_{j_1} \dots m_{j_l}} \sum_{i=1}^d f_i \in W^{\text{aux},\delta}$. Then, $\{i \mid f_i \notin W_0^{\text{aux},\delta}\} = E_0 \cup E_1 \cup \dots \cup E_l$, where $\text{card}(E_0) \leq 1$ and, for each $1 \leq z \leq l$, $(f_i)_{i \in E_z}$ is $S_{n_{j_1} + \dots + n_{j_z}} * \mathcal{A}_3$ -admissible and $\max_{i \in E_z} d_T(m(f_i), \lambda_z) \leq \delta$ for some $\lambda_z \in \overline{\mathcal{D}}$.

Proof We proceed by induction on the length l of the tuple $(m_{j_1}, \dots, m_{j_l})$. If $l = 1$, then $f = \frac{1}{m_{j_1}} \sum_{i=1}^d f_i \in W_k^{\text{aux},\delta}$ for some $k \geq 1$, and we put

$$E_0 = \{1\} \setminus \{i \mid f_i \in W_0^{\text{aux},\delta}\}$$

$$E_1 = \{2, \dots, d\} \setminus \{i \mid f_i \in W_0^{\text{aux},\delta}\}$$

and note that, if $k = 1$, then $E_0 = E_1 = \emptyset$ so that the result holds vacuously. Assume then that $k > 1$ and note that $f_1 < \dots < f_d$ are in $W_{k-1}^{\text{aux},\delta}$ and are $(n_{j_1}, \lambda)_\delta$ -packed for some $\lambda \in \overline{\mathcal{D}}$. It therefore follows by definition that $(f_i)_{i \in E_1}$ are $S_{n_{j_1}} * \mathcal{A}_3$ -admissible and that $\max_{i \in E_1} d(m(f_i), \lambda) \leq \delta$.

Assume now that the result holds for functionals of the form $f = \frac{1}{m_{j_1} \dots m_{j_l}} \sum_{i=1}^d f_i \in W^{\text{aux},\delta}$ and let $g = \frac{1}{m_{j_1} \dots m_{j_l} m_{j_{l+1}}} \sum_{i=1}^d g_i \in W^{\text{aux},\delta}$. As above, the result holds vacuously if $g \in W_1^{\text{aux},\delta}$, so assume that $g \in W_k^{\text{aux},\delta}$ for some $k > 1$. In this

case, we have that $g_1 < \dots < g_d$ are in $W_{k-1}^{\text{aux}, \delta}$ and are $(n_{j_1}, \dots, n_{j_l}, n_{j_{l+1}}, \lambda)_\delta$ -packed for some $\lambda \in \overline{\mathcal{D}}$. It follows that there is a partition $F_1 < \dots < F_M$ of $\{1, \dots, d\}$ as in Definition 4.2. Note that in particular that $(g_i)_{i \in F_1}$ are $(n_{j_1}, \dots, n_{j_l}, \lambda)_\delta$ -packed for some $\lambda_1 \in \overline{\mathcal{D}}$ so that the functional $\tilde{g} = \frac{1}{m_{j_1} \dots m_{j_l}} \sum_{i \in F_1} g_i \in W^{\text{aux}, \delta}$. Then, by the inductive hypothesis, there are sets $E_0 \cup E_1 \cup \dots \cup E_l \subset F_1$ so that the required claim holds for \tilde{g} . Now, let $E_{l+1} = (\bigcup_{r=2}^M F_r) \setminus \{i \mid g_i \in W_0^{\text{aux}, \delta}\}$, and it follows that the required claim holds for g itself so that the proof is complete. ■

We now derive the aforementioned c_0 -type estimate for auxiliary weighted functionals whose weights in some sense mismatch the weights of weighted (n, ε) -basic scc vectors. First, we make a remark about the relation between the weight of a functional and the admissibility of its components.

Remark 4.2 Let $j \in \mathbb{N}$ and $f \in W$ such that $m(f) < m_j$ and $w(f) < m_j^4$. Then, if $\bar{w}(f) = (m_{j_1}, \dots, m_{j_\ell})$, $n_{j_1} + \dots + n_{j_\ell} < n_j$. Indeed, $m_j^l \leq w(f) < m_j^4$, and thus, $l < 4 \log(m_j) / \log(100)$. By condition (2) on page 8,

$$n_{j_1} + \dots + n_{j_\ell} \leq l n_{j-1} < n_{j-1} 4^{\frac{\log(m_j)}{\log(100)}} \leq n_j.$$

Proposition 4.3 Let $N \in \mathbb{N} \cup \{0\}$ and choose $(t_i)_{i=1}^{2^N}$ such that $d(m_{t_i}, m_{t_{i'}}) \geq 2^{-N}$ for each $i \neq i'$. Then, choose $\varepsilon_i \in (0, \frac{1}{3m_{t_i} 2^N})$ and define a (n_{t_i}, ε_i) -basic scc by

$$x_i = m_{t_i} \sum_{r \in F_i} c_r^i e_r$$

for each $1 \leq i \leq 2^N$. It follows that $|f(\sum_{i \in \Omega} a_i x_i)| \leq (1 + \frac{4}{\sqrt{w(f)}}) \max_{i \in \Omega} |a_i|$ for every subset $\Omega \subset \{1, \dots, 2^N\}$, for every choice $(a_i)_{i \in \Omega}$ of scalars, and for every auxiliary weighted functional $f = \frac{1}{m_{j_1} \dots m_{j_l}} \sum_{q=1}^d f_q \in W^{\text{aux}, \delta}$ where $\delta < 2^{-(N+1)}$.

Proof Let $\Omega \subset \{1, \dots, 2^N\}$ be given. We will proceed by induction on the level k norming sets $W_k^{\text{aux}, \delta}$, and the inductive hypothesis will include that $|f(\sum_{i \in A_f(\Omega)} a_i x_i)| < \frac{4\gamma}{\sqrt{w(f)}}$ if $f \notin W_0^{\text{aux}, \delta}$, where $\gamma = \max_{i \in \Omega} |a_i|$ and where $A_f(\Omega) := \{i \in \Omega \mid m_{t_i} \neq m(f)\}$. The base case is the level $k = 1$ norming set, and we first let $f \in W_0^{\text{aux}, \delta}$ so that

$$\left| f\left(\sum_{i \in \Omega} a_i x_i\right) \right| = \left| e_{i_0}^* \left(\sum_{i \in \Omega} a_i m_{t_i} \sum_{r \in F_i} c_r^i e_r \right) \right| \leq \max_{i \in \Omega} \varepsilon_i m_{t_i} |a_i| < \max_{i \in \Omega} |a_i|$$

for some $i_0 \in \mathbb{N}$. It therefore remains to check functionals of the form

$$f = \frac{1}{m_{j_1} \dots m_{j_l}} \sum_{q=1}^d e_{i_q}^* \in W_1^{\text{aux}, \delta} \setminus W_0^{\text{aux}, \delta},$$

and we write $A_f(\Omega) = \{i \in \Omega \mid m_{t_i} \neq m(f)\} = A_1 \cup A_2 \cup A_3$, where

$$\begin{aligned} A_1 &= \{i \in \Omega \mid m_{t_i} < m(f)\} \\ A_2 &= \{i \in \Omega \mid m(f) < m_{t_i} \text{ and } n_{j_1} + \dots + n_{j_l} < n_{t_i}\} \\ A_3 &= \{i \in \Omega \mid m(f) < m_{t_i} \text{ and } n_{j_1} + \dots + n_{j_l} \geq n_{t_i}\} \end{aligned}$$

so that $\Omega \setminus A_f(\Omega)$ is at most a singleton $\{i_0\}$ with $m_{t_{i_0}} = m(f)$. Then, we have that

$$\left| f \left(\sum_{i \in \Omega} a_i x_i \right) \right| \leq \sum_{p=1}^3 \left| f \left(\sum_{i \in A_p} a_i x_i \right) \right| + |f(a_{i_0} x_{i_0})| \leq \sum_{p=1}^3 \left| f \left(\sum_{i \in A_p} a_i x_i \right) \right| + \max_{i \in \Omega} |a_i|$$

because $|f(x_{i_0})| \leq \frac{m_{t_{i_0}}}{w(f)} \leq 1$. Now, recall that $\gamma = \max_{i \in \Omega} |a_i|$ and note that, for A_1 , there is the estimate

$$\begin{aligned} \left| f \left(\sum_{i \in A_1} a_i x_i \right) \right| &\leq \sum_{i \in A_1} |f(a_i x_i)| \leq \frac{\gamma}{w(f)} \sum_{i \in A_1} m_{t_i} \\ &\leq \frac{\gamma}{w(f)} \text{card}(A_1) m_{t_{\max(A_1)}} \leq \frac{\gamma}{w(f)} t_{\max(A_1)} m_{t_{\max(A_1)}} \\ &\leq \frac{\gamma}{w(f)} (\text{ind}(m(f)) - 1) m_{\text{ind}(m(f)) - 1} \leq \frac{\gamma m_{\text{ind}(m(f))}}{w(f) \sqrt{m_{\text{ind}(m(f))}}} \leq \frac{\gamma}{\sqrt{w(f)}}, \end{aligned}$$

where we have used the first condition on the sequence (m_j) and the fact that $t_i < \text{ind}(m(f)) := \max_{1 \leq z \leq l} j_z$ for each $i \in A_1$. Next, for A_2 , note that

$$\begin{aligned} \left| f \left(\sum_{i \in A_2} a_i x_i \right) \right| &\leq \frac{\gamma}{w(f)} \sum_{i \in A_2} m_{t_i} \left| \sum_{q=1}^d e_{i_q}^* \left(\sum_{r \in F_i} c_r^i e_r \right) \right| \\ &\leq \frac{\gamma}{w(f)} \sum_{i \in A_2} 3m_{t_i} \varepsilon_i \leq \frac{\gamma}{w(f)} \sum_{i \in A_2} \frac{1}{2^N} \leq \frac{\gamma}{w(f)} \leq \frac{\gamma}{\sqrt{w(f)}}, \end{aligned}$$

where we have used the observation that immediately precedes Proposition 2.1 and the fact that $e_{i_1}^* < \dots < e_{i_d}^*$ are $S_{n_{j_1} + \dots + n_{j_l}} * \mathcal{A}_3$ -admissible with $n_{j_1} + \dots + n_{j_l} < n_{t_i}$ for each $i \in A_2$. Lastly, for A_3 , note that

$$\begin{aligned} \left| f \left(\sum_{i \in A_3} a_i x_i \right) \right| &\leq \frac{\gamma}{w(f)} \sum_{i \in A_3} m_{t_i} \leq \frac{\gamma}{w(f)} |A_3| m_{t_{\max(A_3)}} \\ &\leq \frac{\gamma m_{t_{\max(A_3)}}^2}{w(f)} \leq \frac{\gamma \sqrt{w(f)}}{w(f)} = \frac{\gamma}{\sqrt{w(f)}} \end{aligned}$$

because $m_{t_i}^2 \leq \sqrt{w(f)}$ for each $i \in A_3$, by Remark 4.2. It now follows by putting these pieces together that

$$\left| f \left(\sum_{i \in \Omega} a_i x_i \right) \right| \leq \gamma + \frac{3\gamma}{\sqrt{w(f)}}$$

for each $f \in W_1^{\text{aux}, \delta}$, and, in particular, we have that $\left| f \left(\sum_{i \in A_f(\Omega)} a_i x_i \right) \right| < \frac{4\gamma}{\sqrt{w(f)}}$, as is required as part of the inductive hypothesis.

Assume now that the statement of Proposition 4.3 holds for every functional $f \in W_k^{\text{aux},\delta}$, let $\Omega \subset \{1, \dots, 2^N\}$ be given, and let

$$f = \frac{1}{m_{j_1} \cdots m_{j_d}} \sum_{q=1}^d f_q \in W_{k+1}^{\text{aux},\delta} \setminus W_k^{\text{aux},\delta},$$

where $f_q \in W_k^{\text{aux},\delta}$ for each $1 \leq q \leq d$. Define the sets $B = \{q \mid f_q \in W_0^{\text{aux},\delta}\}$ and $C = \{3, \dots, d\} \setminus B$. We may then write $f = \sum_{\alpha=1}^4 g_\alpha$, where

$$g_1 = \frac{1}{w(f)} f_1, \quad g_2 = \frac{1}{w(f)} f_2, \quad g_3 = \frac{1}{w(f)} \sum_{q \in B} f_q, \quad \text{and} \quad g_4 = \frac{1}{w(f)} \sum_{q \in C} f_q,$$

and it follows that

$$\left| f \left(\sum_{i \in \Omega} a_i x_i \right) \right| \leq \left| f \left(\sum_{i \in A_f(\Omega)} a_i x_i \right) \right| + |f(a_{i_0} x_{i_0})| \leq \max_{i \in \Omega} |a_i| + \left| f \left(\sum_{i \in A_f(\Omega)} a_i x_i \right) \right|,$$

where, as before, $A_f(\Omega) = \{i \in \Omega \mid m_{t_i} \neq m(f)\}$ so that $\Omega \setminus A_f(\Omega)$ is at most a singleton $\{i_0\}$ with $m_{t_{i_0}} = m(f)$, in which case, $|f(x_{i_0})| \leq \frac{m_{t_{i_0}}}{w(f)} \leq 1$. Let $\gamma = \max_{i \in \Omega} |a_i|$ as before, and it now remains to estimate

$$\begin{aligned} \left| f \left(\sum_{i \in A_f(\Omega)} a_i x_i \right) \right| &\leq \sum_{\alpha=1}^4 \left| g_\alpha \left(\sum_{i \in A_f(\Omega)} a_i x_i \right) \right| \\ &\leq \frac{1}{w(f)} \left| f_1 \left(\sum_{i \in A_f(\Omega)} a_i x_i \right) \right| + \frac{1}{w(f)} \left| f_2 \left(\sum_{i \in A_f(\Omega)} a_i x_i \right) \right| \\ &\quad + \frac{1}{w(f)} \left| \sum_{q \in B} f_q \left(\sum_{i \in A_f(\Omega)} a_i x_i \right) \right| + \frac{1}{w(f)} \left| \sum_{q \in C} f_q \left(\sum_{i \in A_f(\Omega)} a_i x_i \right) \right|, \end{aligned}$$

where, after applying the inductive hypothesis to the g_1 and g_2 terms and using the same argument as is used in the base case for the g_3 term, this quantity is bounded above by

$$\begin{aligned} &\frac{\gamma}{w(f)} \left(1 + \frac{3}{\sqrt{w(f_1)}} \right) + \frac{\gamma}{w(f)} \left(1 + \frac{3}{\sqrt{w(f_2)}} \right) + \frac{3\gamma}{\sqrt{w(f)}} + \frac{1}{w(f)} \left| \sum_{q \in C} f_q \left(\sum_{i \in A_f(\Omega)} a_i x_i \right) \right| \\ &\leq \frac{13}{5} \frac{\gamma}{w(f)} + \frac{3\gamma}{\sqrt{w(f)}} + \frac{1}{w(f)} \left| \sum_{q \in C} f_q \left(\sum_{i \in A_f(\Omega)} a_i x_i \right) \right| \\ &\leq \frac{(3 + \frac{13}{50})\gamma}{\sqrt{w(f)}} + \frac{1}{w(f)} \left| \sum_{q \in C} f_q \left(\sum_{i \in A_f(\Omega)} a_i x_i \right) \right| \end{aligned}$$

because $m_1 \geq 100$. Now, note that there are sets $E_0 \cup E_1 \cup \dots \cup E_l$ from Lemma 4.1 such that $\{q \mid f_q \in \mathcal{C}\} = E_0 \cup E_1 \cup \dots \cup E_l$ and such that, for each $1 \leq z \leq l$, we have that $(f_q)_{q \in E_z}$ is $S_{n_{j_1} + \dots + n_{j_z}} * \mathcal{A}_3$ -admissible and $\max_{q \in E_z} d(m(f_q), \lambda_z) \leq \delta$ for some

$\lambda_z \in \overline{\mathcal{D}}$. Next, let $\Gamma_z = \{i \in A_f(\Omega) \mid m_{t_i} = m(f_q) \text{ for some } q \in E_z\}$ for each $1 \leq z \leq l$ and note that if $i_1, i_2 \in \Gamma_z$, then we have that

$$\begin{aligned} 2^{-N} &\leq d(m_{t_{i_1}}, m_{t_{i_2}}) \\ &= d(m(f_{q_1}), m(f_{q_2})) \leq d(m(f_{q_1}), \lambda_z) + d(\lambda_z, m(f_{q_2})) \leq 2\delta \end{aligned}$$

for the corresponding $q_1, q_2 \in E_z$, and this cannot happen. It follows that Γ_z is at most a singleton, say i_z , for each $1 \leq z \leq l$ so that

$$\begin{aligned} \left| \sum_{q \in \mathbb{C}} f_q \left(\sum_{i \in A_f(\Omega)} a_i x_i \right) \right| &\leq \sum_{z=0}^l \sum_{q \in E_z} \left| f_q \left(\sum_{i \in A_f(\Omega) \setminus \{i_z\}} a_i x_i \right) \right| + \sum_{z=0}^l \sum_{q \in E_z} |f_q(a_{i_z} x_{i_z})| \\ &\leq \sum_{z=0}^l \sum_{q \in E_z} \frac{4\gamma}{\sqrt{w(f_q)}} + \gamma \sum_{z=0}^l \sum_{q \in E_z} \frac{m_{t_{i_z}}}{w(f_q)} \\ &\leq (l+1)\gamma + \sum_{z=0}^l \sum_{q \in E_z} \frac{4\gamma}{\sqrt{w(f_q)}} \leq (l+5)\gamma, \end{aligned}$$

where again, we have used the inductive hypothesis, and, by putting all of these pieces together, there is the estimate

$$\begin{aligned} \left| f \left(\sum_{i \in \Omega} a_i x_i \right) \right| &\leq \gamma + \frac{(3 + \frac{13}{50})\gamma}{\sqrt{w(f)}} + \frac{(l+5)\gamma}{w(f)} \\ &\leq \gamma + \frac{(3 + \frac{13}{50})\gamma}{\sqrt{w(f)}} + \frac{(l+5)\gamma}{10^l \sqrt{w(f)}} \leq \gamma + \frac{(3 + \frac{13}{50} + \frac{3}{5})\gamma}{\sqrt{w(f)}} \leq \gamma + \frac{4\gamma}{\sqrt{w(f)}}, \end{aligned}$$

where, in particular, $\left| f \left(\sum_{i \in A_f(\Omega)} a_i x_i \right) \right| \leq \frac{4\gamma}{\sqrt{w(f)}}$ so that the proof is complete. ■

5 Rapidly increasing sequences and basic inequality

Rapidly increasing sequences (RISs) and the basic inequality are standard concepts used in most Banach spaces constructions using saturated norms, such as $X_{\mathcal{D}}$. The main utility of such sequences is that they can be found in every infinite-dimensional subspace of $X_{\mathcal{D}}$, and, via the basic inequality, the norm of their linear combinations is bounded above by the same linear combinations of basis vectors in the auxiliary space $X_{\mathcal{D}}^{\text{aux}, \delta}$. RISs first appeared in [26], and the basic inequality has its roots in [1]. These tools will be essential in Section 6 when we construct, in an arbitrary infinite-dimensional block subspace of $X_{\mathcal{D}}$, a tree of normalized vectors that witness the failure of the Lebesgue property.

We begin with a short technical lemma, and we recall that $\text{supp}(0) = \emptyset$ and that $\emptyset < A$ and $A < \emptyset$ for every subset $A \subset \mathbb{N}$.

Lemma 5.1 *Fix $\delta > 0$ and assume that $f_1 < \dots < f_d \in W$ with $2^{-\max \text{supp}(f_i)} \leq \frac{\delta}{2}$ are $(n_{j_1}, \dots, n_{j_l}, \lambda)$ -packed for some $\lambda \in \overline{\mathcal{D}}$. Next, suppose that there exist sets of functionals $\mathcal{F}_1, \dots, \mathcal{F}_d$ such that, for each $1 \leq q \leq d$, $\mathcal{F}_q = \{g_1^q, g_2^q, g_3^q\}$ with $g_r^q \in \{0\} \cup W^{\text{aux}, \delta}$, and*

- (1) $\cup_{r=1}^3 \text{supp}(g_r^q) \subset \text{supp}(f_q)$ and $\text{supp}(g_1^q) < \text{supp}(g_2^q) < \text{supp}(g_3^q)$,
- (2) *there is at most one $g_r^q \in \mathcal{F}_q$ such that g_r^q is not a unit vector,*

- (3) there is at most one $g_r^q \in \mathcal{F}_q$ whose support precedes the support of g_r^q , and
- (4) for $g_r^q \in \mathcal{F}_q$ that is not a unit vector, $\tilde{w}(g_r^q) = \tilde{w}(f_q)$.

Then, the functionals $(g_r^q)_{r=1, q=1}^{3, d}$ are $(n_{j_1}, \dots, n_{j_l}, \lambda)_\delta$ -packed.

Proof We proceed by induction on the length l of the tuple $(m_{j_1}, \dots, m_{j_l})$. Let $l = 1$ and assume that $f_1 < \dots < f_d \in W$ with $2^{-\max \text{supp}(f_i)} \leq \delta$ are (n_{j_1}, λ) -packed for some $\lambda \in \overline{\mathcal{D}}$. Then, $(g_r^q)_{r=1, q=1}^{3, d} \subset W_k^{\text{aux}, \delta}$ for some $k \in \mathbb{N}$ are $S_{n_{j_1}} * \mathcal{A}_3$ -admissible, and we have that

$$w(g_r^q) = w(f_q) \geq 2^{\max \text{supp}(f_{q-1})} \geq 2^{\max \text{supp}(\varphi)}$$

for each $g_r^q \in \mathcal{F}_q$ with at least two preceding nonzero functionals, where φ is the nonzero functional that is two before g_r^q . It follows that $(g_r^q)_{r=1, q=1}^{3, d}$ are skipped very fast-growing in the sense of Definition 4.1. Moreover, we have that

$$d(m(g_r^q), \lambda) = d(m(f_q), \lambda) \leq 2^{-\max \text{supp}(f_{q-1})} \leq 2^{-\max \text{supp}(f_1)} \leq \frac{\delta}{2} < \delta$$

for $2 \leq q \leq d$ so that $(g_r^q)_{q=1, r=1}^{d, 3}$ are $(n_{j_1}, \lambda)_\delta$ -packed.

Assume now that the result holds for tuples of length l and let $f_1 < \dots < f_d \in W$ be $(n_{j_1}, \dots, n_{j_l}, n_{j_{l+1}}, \lambda)$ -packed for some $\lambda \in \overline{\mathcal{D}}$. Let $\mathcal{F}_1, \dots, \mathcal{F}_d$ be sets of at most three auxiliary weighted functionals as in the statement of the lemma and let $F_1 < \dots < F_M$ be a partition of $\{1, \dots, d\}$ as in Definition 3.2. Note that $(f_q)_{q \in F_1}$ are $(n_{j_1}, \dots, n_{j_l}, \lambda_1)$ -packed for some $\lambda_1 \in \overline{\mathcal{D}}$. It therefore follows from the inductive hypothesis that $(g_r^q)_{r=1, q \in F_1}^3$ are $(n_{j_1}, \dots, n_{j_l}, \lambda_1)_\delta$ -packed. Moreover, for each g_r^q with $q \in F_p$ for $p > 1$, we have that

$$\begin{aligned} d(m(g_r^q), \lambda) &= d(m(f_q), \lambda) \leq d(m(f_q), \lambda_p) + d(\lambda_p, \lambda) \\ &\leq 2 \cdot 2^{-\max \text{supp}(f_{\max(F_{p-1})})} \leq 2 \cdot 2^{-\max \text{supp}(f_1)} \leq \delta, \end{aligned}$$

and because $(g_r^q)_{q=1, r=1}^{d, 3}$ are skipped very fast-growing in the sense of Definition 4.2 by the same reasoning that is used in the base case, it follows that these functionals are $(n_{j_1}, \dots, n_{j_l}, n_{j_{l+1}}, \lambda)_\delta$ -packed so that the proof is complete. ■

We now recall from [5] the notion of a rapidly increasing sequence (RIS).

Definition 5.1 Let $C \geq 1$, let $I \subset \mathbb{N}$ be an interval, and let $(j_i)_{i \in I}$ be a strictly increasing sequence of positive integers. Then, a block sequence $(x_i)_{i \in I}$ is said to be a $(C, (j_i)_{i \in I})$ -RIS if the following three statements hold:

- (1) $\|x_i\| \leq C$ for each $i \in \mathbb{N}$.
- (2) $\max \text{supp}(x_{i-1}) < \sqrt{m_{j_i}}$ for each $i \in I \setminus \{\min(I)\}$.
- (3) $|f(x_i)| \leq \frac{C}{w(f)}$ for each $i \in I$ and for every $f \in W$ with $w(f) < m_{j_i}$.

We recall the steps of a standard argument that shows that there exists a RIS in every block subspace of $X_{\mathcal{D}}$. For some of the details, we refer to specific parts of [5]. The proposition below is proved word-for-word as [5, Proposition 6.3], and it requires

only two things: Theorem 3.2 (in place of [5, Proposition 4.1]) and that, for every $C > 1$, $m_j^{-1}C^{n_j} \rightarrow \infty$. The latter is true because, by condition (2) on page 8,

$$\log\left(\frac{C^{n_j}}{m_j}\right) > \frac{4\log(C)}{\log(100)}n_{j-1}\log(m_j) - \log(m_j) = \log(m_j)\left(\frac{4\log(C)n_{j-1}}{\log(100)} - 1\right).$$

Proposition 5.2 *Let Y be a block subspace of $X_{\mathcal{D}}$. Then, for every $n \in \mathbb{N}$, $\varepsilon > 0$, and $\delta > 0$, there exists a (n, ε) special convex combination $x = \sum_{i=1}^m c_i x_i$ such that $\|x\| > 1/(1 + \delta)$ and x_1, \dots, x_m are block vectors in Y of norm at most one.*

The next proposition is also proved verbatim as [5, Proposition 6.4], and it uses Remark 4.2.

Proposition 5.3 *Let $j \in \mathbb{N}$, $\varepsilon > 0$, and $x = \sum_{i=1}^m c_i x_i$ be an (n_j, ε) -special convex combination such that $\|x_i\| \leq 1$, $1 \leq i \leq m$. Then, for every $f \in W$ such that $w(f) < m_j$,*

$$|f(x)| \leq \frac{1 + 2w(f)\varepsilon}{w(f)}.$$

The desired conclusion follows immediately from Proposition 5.2 and Proposition 5.3.

Corollary 5.4 *For every $C > 1$ and in every block subspace Y of $X_{\mathcal{D}}$, there exists a $(C, (j_i)_{i \in \mathbb{N}})$ -RIS (x_i) , with $\|x_i\| \geq 1$, for $i \in \mathbb{N}$.*

We now prove the basic inequality that transfers the action of a weighted functional $f \in W$ to the action of auxiliary weighted functionals on appropriate unit vectors.

Proposition 5.5 *Let $(x_i)_{i \in I}$ be a $(C, (j_i)_{i \in I})$ -RIS with $2^{-\min \text{supp}(x_{\min(I)})} \leq \delta$. Then, for every functional $f = \frac{1}{m_{j_1} \cdots m_{j_l}} \sum_{q=1}^d f_q \in W$, there exist functionals $h \in \{0\} \cup W_0^{\text{aux}, \delta}$ and $g \in W^{\text{aux}, \delta}$ with $\tilde{w}(g) = \tilde{w}(f)$ such that*

$$\left| f\left(\sum_{i \in I} a_i x_i\right) \right| \leq C \left(1 + \frac{1}{\sqrt{m_{j_{\min(I)}}}}\right) \left| (h + g)\left(\sum_{i \in I} a_i e_{t_i}\right) \right|$$

for every choice $(a_i)_{i \in I}$ of scalars where $t_i = \max \text{supp}(x_i)$ for each $i \in I$.

Proof We proceed by induction on the level k norming sets W_k , and the base case is the level $k = 0$ set $W_0 = \{\pm e_i^* \mid i \in \mathbb{N}\}$. Indeed, if $f = e_{i_0}^* \in W_0$, then

$$\left| e_{i_0}^*\left(\sum_{i \in I} a_i x_i\right) \right| \leq |a_{i_0} e_{i_0}^*(x_{i_0})| \leq C|a_{i_0}| = C|e_{t_{i_0}}^*(a_{t_{i_0}} e_{t_{i_0}})|,$$

where $i_0 \in \text{supp}(x_{i_0})$ so that the desired estimate holds with $h = \{0\}$ and with $g = e_{t_{i_0}}^*$. Note that, as part of the inductive hypothesis, we have that $\text{supp}(h), \text{supp}(g) \subset \{t_i\}_{i \in I}$ and that either $h = 0$ or $h \in W_0^{\text{aux}, \delta}$, $\text{supp}(h) < \min \text{supp}(g)$.

Assume now that the desired estimate holds for functionals in the level k norming set W_k and consider the functional

$$f = \frac{1}{m_{j_1} \cdots m_{j_l}} \sum_{q=1}^d f_q \in W_{k+1} \setminus W_k,$$

which is to say that $f_1 < \dots < f_d$ are in W_k and are $(n_{j_1}, \dots, n_{j_l}, \lambda)$ -packed for some $\lambda \in \overline{\mathcal{D}}$. We omit if necessary the functionals f_q that are below the support of $x_{\min(I)}$. It then follows that

$$\frac{1}{\delta} \leq 2^{\min \text{supp}(x_{\min(I)})} \leq 2^{\max \text{supp}(f_i)} \implies 2^{-\max \text{supp}(f_i)} \leq \delta,$$

and we will use this observation later. First, define the quantity i_0 by

$$i_0 = \max\{i \in I \mid m_{j_1} \cdots m_{j_l} \geq m_{j_i}\}$$

if it exists. If there does exist such an i_0 , then choose $i_1 \in [\min(I), i_0]$ so that $|a_{i_1}| = \max\{|a_i|\}_{i=\min(I)}^{i_1}$ and define the functional $h = \text{sign}(a_{i_1})e_{t_{i_1}}^* \in W_0^{\text{aux}, \delta}$. Note that, if $\min(I) = i_0 = i_1$, then

$$\left| f\left(\sum_{i \leq i_0} a_i x_i\right) \right| = |f(a_{i_1} x_{i_1})| \leq C|a_{i_1}| = Ch(a_{i_1} e_{t_{i_1}}),$$

and, if $\min(I) < i_0$, then

$$\begin{aligned} \left| f\left(\sum_{i \leq i_0} a_i x_i\right) \right| &\leq |a_{i_0} f(x_{i_0})| + \sum_{i < i_0} |a_i f(x_i)| \leq C|a_{i_0}| + \sum_{i < i_0} \left| \frac{a_i}{w(f)} \sum_{q=1}^d f_q(x_i) \right| \\ &\leq C|a_{i_1}| + \frac{|a_{i_1}|}{m_{j_{i_0}}} \sum_{i < i_0} \sum_{q=1}^d \left| f_q\left(\sum_{r \in \text{supp}(x_i)} b_r e_r\right) \right| \\ &\leq C|a_{i_1}| + \frac{|a_{i_1}|}{m_{j_{i_0}}} \sum_{i < i_0} \sum_{r \in \text{supp}(x_i)} |b_r| \leq C|a_{i_1}| + \frac{C|a_{i_1}|}{m_{j_{i_0}}} \max \text{supp}(x_{i_0-1}) \\ &\leq C\left(1 + \frac{1}{\sqrt{m_{j_{i_0-1}}}}\right) |a_{i_1}| \leq C\left(1 + \frac{1}{\sqrt{m_{j_{\min(I)}}}}\right) h\left(\sum_{i \leq i_0} a_i e_{t_i}\right). \end{aligned}$$

If, however, there is no such number i_0 , then the sum $\sum_{i \leq i_0} a_i x_i$ is empty, so we take $h = 0$.

Next, let $\tilde{I} = \{i \in I \mid i > i_0\}$ if such an i_0 exists and put $\tilde{I} = I$ otherwise. We then define the subsets

$$\begin{aligned} A &= \{i \in \tilde{I} \mid \text{supp}(x_i) \cap \text{supp}(f_q) \neq \emptyset \text{ for at most one } q\} \\ I_q &= \{i \in A \mid \text{supp}(x_i) \cap \text{supp}(f_q) \neq \emptyset\} \\ D &= \{q \mid I_q \neq \emptyset\} \\ B &= \tilde{I} \setminus A \end{aligned}$$

so that the intervals I_q are pairwise disjoint and $\text{supp}(x_i)$ intersects the support of at least two f_q if $i \in B$. It follows by the inductive hypothesis that, for each $q \in D$, there exist functionals $h_q \in \{0\} \cup W_0^{\text{aux}, \delta}$ and $g_q \in W^{\text{aux}, \delta}$ with $\vec{w}(g_q) = \vec{w}(f_q)$ so that

$$\left| f_q\left(\sum_{i \in I_q} a_i x_i\right) \right| \leq C\left(1 + \frac{1}{\sqrt{m_{j_{\min(I)}}}}\right) \left| (h_q + g_q)\left(\sum_{i \in I_q} a_i e_{t_i}\right) \right|,$$

where $\text{supp}(h_q), \text{supp}(g_q) \subset \text{supp}(f_q)$ and we have that either $h_q = 0$ or $h_q \in \{e_{t_i}\}_{i \in I_q}$ and $t_i < \min \text{supp}(g_q)$. Now, replace $h_q + g_q$ by $\text{sign}[(h_q + g_q)(\sum_{i \in I_q} a_i e_{t_i})]$ $(h_q + g_q)$ if necessary and note that

$$\begin{aligned} \left| f\left(\sum_{i \in \bar{I}} a_i x_i\right) \right| &\leq \sum_{i \in B} |f(x_i)| + \frac{1}{w(f)} \sum_{q \in D} \left| f_q\left(\sum_{i \in I_q} a_i x_i\right) \right| \\ &\leq \frac{C}{w(f)} \sum_{i \in B} |a_i| + C \left(1 + \frac{1}{\sqrt{m_{j_{\min(I)}}}}\right) \frac{1}{w(f)} \sum_{q \in D} (h_q + g_q) \left(\sum_{i \in I_q} a_i e_{t_i}\right) \\ &\leq C \left(1 + \frac{1}{\sqrt{m_{j_{\min(I)}}}}\right) \frac{1}{w(f)} \left[\sum_{i \in B} \tilde{h}_i + \sum_{q \in D} (h_q + g_q) \right] \left(\sum_{i \in \bar{I}} a_i e_{t_i}\right), \end{aligned}$$

where we have used the third condition from Definition 5.1 and where $\tilde{h}_i = \text{sign}(a_i) e_{t_i}^*$ for each $i \in B$. It therefore follows that

$$\left| f\left(\sum_{i \in I} a_i x_i\right) \right| \leq C \left(1 + \frac{1}{\sqrt{m_{j_{\min(I)}}}}\right) \left[(h + g) \left(\sum_{i \in I} a_i e_{t_i}\right) \right],$$

where either $h = 0$ or $h \in \{e_{t_i}\}_{i=\min(I)}^{i_0}$ and $\text{supp}(h) = t_{i_1} < \min \text{supp}(g)$. Indeed, we have that

$$g = \frac{1}{w(f)} \left(\sum_{i \in B} \tilde{h}_i + \sum_{q \in D} (h_q + g_q) \right)$$

so that $\tilde{w}(f) = \tilde{w}(g)$. It remains to prove that $g \in W^{\text{aux}, \delta}$. To that end, define for each $i \in B$ the quantity $q_i = \max\{1 \leq q \leq d \mid \min \text{supp}(f_q) \leq \max \text{supp}(x_i) = t_i\}$ and note that the correspondence $i \mapsto q_i$ is strictly increasing. If $q = q_i$ for some $i \in B$, then write $\mathcal{F}_q = \{h_q, g_q, \tilde{h}_i\}$. If $q \neq q_i$ for any $i \in B$, then write $\mathcal{F}_q = \{h_q, g_q\}$. Note that \mathcal{F}_q may consist only of \tilde{h}_i or may be empty if $q \notin D$. It therefore follows from Lemma 5.1 (applied to $(f_q)_{q \in D}$) that $g \in W^{\text{aux}, \delta}$, and this completes the proof. ■

6 The failure of the Lebesgue property in subspaces of $X_{\mathcal{D}}$

In this section, we put together the results of the previous two sections in order to prove that every subspace of $X_{\mathcal{D}}$ contains a normalized collection $(y_{\lambda})_{\lambda \in \mathcal{D}}$ that satisfies the estimates (1.1), and thus witnesses the failure of the Lebesgue property.

Theorem 6.1 *Let Y be an infinite-dimensional block subspace of $X_{\mathcal{D}}$. Then, there exists a normalized collection $(y_{\lambda})_{\lambda \in \mathcal{D}}$ in Y that is block with the lexicographical order of \mathcal{D} such that the following holds: for every $N \in \mathbb{N}$ and $(\mu_{\lambda})_{\lambda \in \{0,1\}^N}$ in \mathcal{D} such that, for $\lambda \in \{0,1\}^N$, $\mu_{\lambda} \geq \lambda$ and for any scalars $(a_{\lambda})_{\lambda \in \{0,1\}^N}$,*

$$\max_{\lambda \in \{0,1\}^N} |a_{\lambda}| \leq \left\| \sum_{\lambda \in \{0,1\}^N} a_{\lambda} y_{\mu_{\lambda}} \right\| \leq 3 \max_{\lambda \in \{0,1\}^N} |a_{\lambda}|.$$

In particular, the Lebesgue property fails in every infinite-dimensional closed subspace of $X_{\mathcal{D}}$.

Proof Let $Y \subset X_{\mathcal{D}}$ and let $Z \subset Y$ be a block subspace. Then, fix $C > 1$ and let (z_i) be an infinite $(C, (j_i)_{i \in \mathbb{N}})$ -RIS in Z with $1 \leq \|z_i\|$ for each $i \in \mathbb{N}$, which exists by Corollary 5.4. By passing to a subsequence, we may assume that the conclusion of Theorem 3.2 is satisfied. Recall that the nodes of \mathcal{D} are in lexicographical order with respect to \mathcal{M} and choose for each weight m_j a number $\varepsilon_j \in \left(0, \frac{1}{3m_j 2^{N_j}}\right)$, where $N_j = |\phi(m_j)|$.

Now, let $F_1 < F_2 < \dots$ be subsets of \mathbb{N} so that $\{\min \text{supp}(z_i) \mid i \in F_j\}$ is a maximal set in S_{n_j} with respect to inclusion for each $j \in \mathbb{N}$. It follows from Proposition 2.1 that there exist coefficients $(c_i^j)_{i \in F_j}$ so that the vector $\sum_{i \in F_j} c_i^j e_i$ is an $(n_j, \frac{\varepsilon_j}{2})$ -basic scc. In particular, we have that $x_{m_j} = \sum_{i \in F_j} c_i^j z_i$ is an $(n_j, \frac{\varepsilon_j}{2})$ -scc, and we note also that

$$\|x_{m_j}\| = \left\| \sum_{i \in F_j} c_i^j z_i \right\| \geq \frac{m_1 - 1}{m_1 m_j} \sum_{i \in F_j} c_i^j = \frac{m_1 - 1}{m_1 m_j}$$

from Theorem 3.2. Let $\tilde{x}_{m_j} = m_j x_j$ and define, for $\lambda \in \mathcal{D}$, $y_\lambda = \tilde{x}_{\phi^{-1}(m_j)} / \|\tilde{x}_{\phi^{-1}(m_j)}\|$ so that (y_λ) is a normalized sequence in Z that is block in the lexicographical order of \mathcal{D} .

Now, fix $N \in \mathbb{N}$ and, for $\lambda \in \{0, 1\}^N$, let $\mu_\lambda \in \mathcal{D}$ such that $\mu_\lambda \geq \lambda$. Because $(y_\lambda)_{\lambda \in \mathcal{D}}$ is lexicographically block, there are $2^N \leq j_1 < \dots < j_{2^N}$ such that $\phi(\{m_{j_1}, \dots, m_{j_{2^N}}\}) = \{\mu_\lambda : \lambda \in \{0, 1\}^N\}$. In particular, $\min \text{supp}(y_{\mu_\lambda}) \geq N$, $\lambda \in \{0, 1\}^N$. Furthermore, for $\lambda \in \{0, 1\}^N$, $\mu_\lambda \geq \lambda$, and thus, for $1 \leq k \neq k' \leq 2^N$, $d(m_{j_k}, m_{j_{k'}}) \geq 2^{-N}$. Then, it follows that

$$\left\| \sum_{\lambda \in \{0, 1\}^N} y_{\mu_\lambda} \right\| = \mathcal{F} \left(\sum_{\lambda \in \{0, 1\}^N} y_{\mu_\lambda} \right) = \mathcal{F} \left(\sum_{k=0}^{2^N-1} \frac{1}{\|\tilde{x}_{m_{j_k}}\|} m_{j_k} \sum_{i \in F_{j_k}} c_i^{j_k} z_i \right) \quad (\#)$$

for some norming functional $\mathcal{F} \in W$, and, by Proposition 5.5, this quantity is bounded above by

$$C \left| (h + g) \left(\sum_{k=0}^{2^N-1} \frac{1}{\|\tilde{x}_{m_{j_k}}\|} m_{j_k} \sum_{i \in F_{j_k}} c_i^{j_k} e_{\rho(i)} \right) \right| \quad (\#\#),$$

where $h \in W_0^{\text{aux}, \delta} \cup \{0\}$, $g \in W^{\text{aux}, \delta}$ with $\vec{w}(g) = \vec{w}(\mathcal{F})$, and where $\rho(i) = \max \text{supp}(z_i)$ for each $i \in \mathbb{N}$. Note that $\sum_{i \in F_{j_k}} c_i^{j_k} e_{\rho(i)}$ is for each $0 \leq k \leq 2^N - 1$ an $(n_{j_k}, \varepsilon_{j_k})$ -basic scc by the observation that immediately follows Proposition 2.1, and note also that

$$\frac{m_{j_k}}{\|\tilde{x}_{m_{j_k}}\|} = \frac{1}{\|x_{m_{j_k}}\|} \leq \frac{m_1 m_{j_k}}{m_1 - 1}.$$

It therefore follows that $(\#\#)$ is bounded above by

$$C \max_{0 \leq k \leq 2^N-1} \frac{m_{j_k} \varepsilon_{j_k}}{\|\tilde{x}_{j_k}\|} + C \left| g \left(\sum_{k=0}^{2^N-1} \frac{1}{\|\tilde{x}_{j_k}\|} m_{j_k} \sum_{i \in F_{j_k}} c_i^{j_k} e_{\rho(i)} \right) \right| \quad (\#\#\#),$$

and lastly, by Proposition 4.3, we note that (###) is bounded above by

$$C \frac{m_1 m_{j_k}}{m_1 - 1} \frac{1}{3 m_{j_k} 2^{N_{j_k}}} + C \left(1 + \frac{4}{\sqrt{w(g)}} \right) \max_{0 \leq k \leq 2^N - 1} \frac{1}{\|\tilde{x}_{j_k}\|} \leq C \frac{100}{99} \left(1 + \left(1 + \frac{2}{5} \right) \right) \leq 3,$$

for $C > 1$ sufficiently close to one. The estimate for arbitrary scalars coefficients follows from the 1-unconditionality of the basis of $X_{\mathcal{D}}$. ■

References

- [1] S. A. Argyros and I. Deliayanni, *Examples of asymptotic l_1 Banach spaces*. Trans. Amer. Math. Soc. 349(1997), no. 3, 973–995.
- [2] S. A. Argyros, A. Georgiou, A.-R. Lagos and P. Motakis, *Joint spreading models and uniform approximation of bounded operators*. Studia Math. 253(2020), no. 1, 57–107.
- [3] S. A. Argyros, G. Godefroy and H. P. Rosenthal, *Descriptive set theory and Banach spaces*. In *Handbook of the geometry of Banach spaces*, Vol. 2, North-Holland, Amsterdam, 2003, 1007–1069.
- [4] S. A. Argyros and P. Motakis, *A reflexive hereditarily indecomposable space with the hereditary invariant subspace property*. Proc. Lond. Math. Soc. (3) 108(2014), no. 6, 1381–1416.
- [5] S. A. Argyros and P. Motakis, *On the complete separation of asymptotic structures in Banach spaces*. Adv. Math. 362(2020), 106962, 51.
- [6] S. A. Argyros and A. Toliás, *Methods in the Theory of Hereditarily Indecomposable Banach spaces*. Mem. Amer. Math. Soc. 170(2004), no. 806, vi+114.
- [7] F. Baudier, G. Lancien, P. Motakis and Th. Schlumprecht, *A new coarsely rigid class of Banach spaces*. J. Inst. Math. Jussieu 20(2021), no. 5, 1729–1747.
- [8] K. Beanland, D. Freeman and P. Motakis, *The stabilized set of p 's in Krivine's theorem can be disconnected*. Adv. Math. 281(2015), 553–577.
- [9] A. Brunel and L. Sucheston, *On B -convex Banach spaces*. Math. Systems Theory 7(1974), no. 4, 294–299.
- [10] H. Gaebler, *Towards a characterization of the property of Lebesgue*. Real Anal. Exchange 46(2021), no. 2, 319–344.
- [11] H. Gaebler and B. Sari, *Banach spaces with the Lebesgue property of Riemann integrability*. J. Funct. Anal. 287(2024), no. 2, 110462, 23.
- [12] R. Gordon, *Riemann integration in Banach spaces*. Rocky Mountain J. Math. 21(1991), no. 3, 923–949.
- [13] L. Halbeisen and E. Odell, *On asymptotic models in Banach spaces*. Israel J. Math. 139(2004), 253–291.
- [14] R. Haydon, *Darboux integrability and separability of types in stable Banach spaces*. In Seminar on functional analysis, 1983/1984, Publ. Math. Univ. Paris VII, Vol. 20, Univ. Paris VII, Paris, 1984, 95–115.
- [15] M. Junge, D. Kutzarova and E. Odell, *On asymptotically symmetric Banach spaces*. Studia Math. 173(2006), no. 3, 203–231.
- [16] T. Kania and R. Lechner, *Factorisation in stopping-time Banach spaces: identifying unique maximal ideals*. Adv. Math. 409(2022), Paper No. 108643, 35.
- [17] B. Maurey, V. D. Milman and N. Tomczak-Jaegermann, *Asymptotic infinite-dimensional theory of Banach spaces*. In Geometric aspects of functional analysis (Israel, 1992–1994), Oper. Theory Adv. Appl., Vol. 77, Birkhäuser, Basel, 1995, 149–175.
- [18] V. D. Milman and N. Tomczak-Jaegermann, *Asymptotic l_p spaces and bounded distortions*. In Banach spaces (Mérida, 1992), Contemp. Math., Vol. 144, Amer. Math. Soc., Providence, RI, 1993, 173–195.
- [19] P. Motakis, *Separable spaces of continuous functions as Calkin algebras*. J. Amer. Math. Soc. 37(2024), no. 1, 1–37.
- [20] P. Motakis and A. Pelczar-Barwacz, *Reflexive Calkin algebras*. Preprint, 2024, [arXiv:2401.18037](https://arxiv.org/abs/2401.18037).
- [21] K. M. Naralenskoy, *Asymptotic structure of Banach spaces and Riemann integration*. Real Anal. Exchange 33(2008), no. 1, 111–124.
- [22] E. Odell, *Stability in Banach spaces*. In IV Course on Banach Spaces and Operators (Spanish) (Laredo, 2001), Vol. 17, 2002, 385–425.
- [23] E. Odell, *On the structure of separable infinite dimensional Banach spaces*, Chern Institute of Mathematics, Nankai University, Tianjin, China, 2007.

- [24] E. Odell and Th. Schlumprecht, *On the richness of the set of p 's in Krivine's theorem*. In Geometric aspects of functional analysis (Israel, 1992–1994), Oper. Theory Adv. Appl., Vol. 77, Birkhäuser, Basel, 1995, 177–198.
- [25] E. Odell and Th. Schlumprecht, *A Banach space block finitely universal for monotone bases*. Trans. Amer. Math. Soc. 352(2000), no. 4, 1859–1888.
- [26] Th. Schlumprecht, *An arbitrarily distortable Banach space*. Israel J. Math. 76(1991), no. 1–2, 81–95.

University of North Texas, Denton, USA

e-mail: harrison.gaebler@unt.edu

York University, Toronto, Canada

e-mail: pmotakis@yorku.ca

University of North Texas, Denton, USA

e-mail: bunyamin.sari@unt.edu