

CRYSTAL FLEX BASES AND THE RUM SPECTRUM

G. BADRI¹, D. KITSON²  AND S. C. POWER³ 

¹ *Department of Mathematical Sciences, Umm Al-Qura University, Mecca Saudi Arabia*
(gmbadri@uqu.edu.sa)

² *Department of Mathematics and Computer Studies, Mary Immaculate College,*
Thurles, Co. Tipperary, Ireland (derek.kitson@mic.ul.ie)

³ *Department of Mathematics and Statistics, Lancaster University, Lancaster LA1 4YF,*
UK (s.power@lancaster.ac.uk)

(Received 19 October 2019; first published online 24 November 2021)

Abstract A theory of infinite spanning sets and bases is developed for the first-order flex space of an infinite bar-joint framework, together with space group symmetric versions for a crystallographic bar-joint framework \mathcal{C} . The existence of a crystal flex basis for \mathcal{C} is shown to be closely related to the spectral analysis of the rigid unit mode (RUM) spectrum of \mathcal{C} and an associated *geometric flex spectrum*. Additionally, infinite spanning sets and bases are computed for a range of fundamental crystallographic bar-joint frameworks, including the honeycomb (graphene) framework, the octahedron (perovskite) framework and the 2D and 3D kagome frameworks.

Keywords: rigid unit modes; periodic frameworks; flexibility

2020 *Mathematics subject classification:* Primary 52C25; 74N05;
Secondary 74N10

1. Introduction

The analysis of the rigidity and flexibility of periodic infinite bond-node structures is an ongoing endeavour in materials science and pure mathematics (Connelly, Ivić and Whiteley [2], Guest, Fowler and Power [7]). In this connection, it is well known in crystallography and engineering that many crystals and periodic structures which are critically coordinated exhibit a rich set of localized zero energy crystal vibrations (phonons) (Giddy et al. [5], Dove et al. [3, 4], Wegner [17]) and localized infinitesimal mechanisms (Hutchinson and Fleck [8]). Inspired by this, we wish to understand for which crystal frameworks there exists a finite or countable set of localized modes u_1, u_2, \dots which tells the whole story, so to speak, by providing a *complete set* in the sense that every first-order

(i.e. infinitesimal) flex may be expressed as an infinite linear combination

$$u = \sum_{n=1}^{\infty} \alpha_n u_n, \quad \alpha_n \in \mathbb{R}.$$

In the dynamical theory of a crystal framework \mathcal{C} , it is common to limit attention to first-order motions with some form of periodicity, or periodicity up to a phase factor for a Bloch wave vector. However, one of our motivations is to determine the structure of the space $\mathcal{F}(\mathcal{C}; \mathbb{R})$ of *all* first-order flexes, including unbounded ones, as well as the structure of the space $\mathcal{F}_{\infty}(\mathcal{C}; \mathbb{R})$ of all bounded first-order flexes.

To approach these problems, we formalize, in §2, the notions of a *free spanning set* and a *free basis* for a space of velocity fields for an arbitrary countable bar-joint framework, with no periodicity assumptions, while in §4, we define their space group symmetric counterparts in the case of crystal frameworks. Also, in §5, we determine such spanning sets and bases for a range of fundamental examples, including the honeycomb (graphene) framework, the octahedron (perovskite) framework and the 2D and 3D kagome frameworks. Our main result, Theorem 4.4, shows that if the infinitesimal flex space of a crystallographic bar-joint framework is infinite dimensional then in any space group symmetric free spanning set, there necessarily exists a *band-limited* flex, that is, one whose support lies uniformly close to a proper linear subspace. We show that this requirement implies that there can be obstacles to the existence of crystal flex bases which arise from nonlinearity in the RUM spectrum $\Omega(\mathcal{C})$ or nonlinearity in a more general *geometric flex spectrum* $\Gamma(\mathcal{C})$ which we introduce here. These obstacles are expressed in Theorems 4.9 and 4.11.

The rigid unit modes, or RUMs, of a material crystal are the zero energy oscillation modes observed in the long-wavelength limit [4, 5, 17]. These vibration modes are bounded and periodic modulo a multiphase factor and in fact, they correspond precisely to non-zero infinitesimal flexes with the corresponding boundedness and periodicity properties (see [1, 14, 15]). The spectrum of multiphase factors for such modes is a subset of the d -torus known as the RUM spectrum and it corresponds to the points of rank degeneracy of a matrix function $\Phi_{\mathcal{C}}(z)$ computable from a building block unit for \mathcal{C} . In this way, the study of RUM modes is reduced to the spectral analysis of function matrices and their associated eigenspaces.

The determination of the space of *all* real or complex infinitesimal flexes requires an understanding of unbounded flexes, possibly localized to a hyperplane, and in this connection, we introduce the *transfer function* $\Psi_{\mathcal{C}}(z)$ associated with \mathcal{C} and a building block unit. This is an analytic matrix-valued function on the d -fold product \mathbb{C}_*^d of the punctured complex plane $\mathbb{C}_* = \mathbb{C} \setminus \{0\}$ defined as the unique extension of $\Phi_{\mathcal{C}}(z)$. The points $z = \omega$ of rank degeneracy of the transfer function indicate the presence of non-zero *geometrically periodic flexes*. Such periodicity is characterized by a set of equations of the form

$$u(p_k) = \omega^k u(p_0) = \omega_1^{k_1} \cdots \omega_d^{k_d} u(p_0)$$

which relate the velocity $u(p_0)$ of a node p_0 in the base unit cell to the velocity $u(p_k)$ of the corresponding node p_k in the cell with label $k \in \mathbb{Z}^d$. In particular, if the multifactor $\omega = (\omega_1, \dots, \omega_d)$ has $|\omega_i| > 1$ for some i then the local velocities $u(p_k)$ are unbounded

(or zero) as $k_i \rightarrow +\infty$, with k_j fixed for $j \neq i$, and are geometrically decaying as $k_i \rightarrow -\infty$. The geometrically periodic flexes are also referred to as *factor periodic* flexes with multifactor ω . The novel geometric spectrum $\Gamma(\mathcal{C})$ referred to above is the set of such multifactors, that is, the set of points of rank degeneracy of the transfer function.

The development is organized as follows. In §2, we show, using an abstract non-constructive argument, that for any countable bar-joint framework, periodic or not, every infinite dimensional linear subspace of the space of velocity fields has a free basis in the sense of Definition 2.3. In §3, we consider group actions on free spanning sets, proving several key results which are then applied in §4 to crystal flex spanning sets for periodic frameworks. Our first main result, Theorem 4.4, shows that a crystal flex spanning set for an infinite dimensional space of flexes necessarily contains localized velocity fields which moreover have geometric periodicity relative to their support. We also show that the existence of a natural crystal flex basis can lead to a simple description of the bounded flexes as those whose infinite expansion in the basis have bounded coefficients. Following this, the RUM spectrum and the geometric flex spectrum, associated with a periodic structure for \mathcal{C} , are introduced and we obtain necessary spectral conditions for the existence of various crystal spanning sets. We also pose here the intriguing problem of determining sufficient conditions, including spectral conditions, which ensure the existence of a crystal flex basis. In the final section, which is largely independent of earlier results, we compute crystal flex bases and spanning sets for several fundamental examples.

2. Free spanning sets and free bases

Let A be a non-empty set and let X be a finite-dimensional vector space over a field \mathbb{K} , where $\mathbb{K} = \mathbb{R}$ or \mathbb{C} . Endow X with a norm and the norm topology. Let X^A denote the topological vector space of maps $f: A \rightarrow X$ with the usual pointwise vector space operations and the topology of pointwise convergence (i.e. the product topology). The support of a map f is denoted $\text{supp}(f)$.

Definition 2.1. A sequence (f_n) in $X^A \setminus \{0\}$ *tends to zero strictly* if, for each $a \in A$, the sequence $(f_n(a))$ in X has at most finitely many non-zero terms.

Lemma 2.2. Let (f_n) be a sequence in $X^A \setminus \{0\}$. The following conditions are equivalent.

(i) (f_n) tends to zero strictly.

(ii) $\sum_{n=1}^{\infty} \alpha_n f_n(a)$ converges in X for every sequence of scalars (α_n) in \mathbb{K} and every $a \in A$.

Proof. If (i) holds and (α_n) is an arbitrary sequence of scalars then, for each $a \in A$, the series $\sum_{n=1}^{\infty} \alpha_n f_n(a)$ has finitely many non-zero terms and hence converges in X . This establishes (ii). If (i) does not hold, then there exists $a \in A$ and a subsequence (f_{n_j}) such

that $f_{n_j}(a)$ is non-zero for each j . Choose a norm $\|\cdot\|$ on X and for each $n \in \mathbb{N}$ define,

$$\alpha_n = \begin{cases} \frac{1}{\|f_{n_j}(a)\|} & \text{if } n = n_j \text{ for some } j \in \mathbb{N}, \\ 0 & \text{otherwise.} \end{cases}$$

Then the series $\sum_{n=1}^{\infty} \alpha_n f_n(a)$ fails to converge in X and so (ii) is not satisfied. \square

If a sequence (f_n) in $X^A \setminus \{0\}$ tends to zero strictly then, by Lemma 2.2, for every sequence of scalars (α_n) in \mathbb{K} the sum $\sum_{n=1}^{\infty} \alpha_n f_n$ represents an element of X^A . Note that the partial sums, $s_N = \sum_{n=1}^N \alpha_n f_n$, converge to $s = \sum_{n=1}^{\infty} \alpha_n f_n$ in the strict sense that the sequence $(s - s_N)$ tends to zero strictly in X^A . Also note that, setting $S = \{f_n : n \in \mathbb{N}\}$, the set $\mathcal{M}(S) = \{\sum_{n=1}^{\infty} \alpha_n f_n : \alpha_n \in \mathbb{K}\}$ is a vector subspace of X^A .

In the next definition, the term *free* for a countable set of vectors is used in the sense that there are no coefficient restrictions needed to ensure that an infinite linear combination of the vectors is pointwise convergent.

Definition 2.3. Let W be an infinite-dimensional vector subspace of X^A .

- (a) A *free spanning set* for W is a countable subset $S = \{f_1, f_2, \dots\}$ of $W \setminus \{0\}$ for which the sequence (f_n) tends to zero strictly and satisfies $W \subseteq \mathcal{M}(S)$.
- (b) A *free basis* for W is a free spanning set $S = \{f_1, f_2, \dots\}$ for W such that each $w \in W$ has a unique representation,

$$w = \sum_{n=1}^{\infty} \alpha_n f_n, \quad \alpha_n \in \mathbb{K}.$$

Example 2.4. Let $A = \{a_k : k \in \mathbb{N}\}$ be a countable set and let x_1, \dots, x_d be a basis for X . For each $n \in \mathbb{N}$ and each $\sigma \in \{1, \dots, d\}$ define,

$$e_{n,\sigma}(a_k) = \begin{cases} x_\sigma & \text{if } n = k, \\ 0 & \text{otherwise.} \end{cases}$$

Then $\{e_{n,\sigma} : n \in \mathbb{N}, \sigma \in \{1, \dots, d\}\}$ is a free basis for X^A .

2.1. Existence of free bases

Let A be a countably infinite set and let $\alpha = (A_i)$ be an increasing sequence of finite subsets of A which cover A (i.e. $A_i \subset A_{i+1}$ for all $i \in \mathbb{N}$ and $A = \cup_{i \in \mathbb{N}} A_i$). For each $i \in \mathbb{N}$, let $\pi_i : X^A \rightarrow X^{A_i}$ denote the natural restriction map. Let W be an infinite dimensional vector subspace of X^A and let $W_i = \pi_i(W)$ for each $i \in \mathbb{N}$. For all $i, j \in \mathbb{N}$ with $j \geq i$, let $\pi_{j,i} : W_j \rightarrow W_i$ be the natural restriction maps. Note that the pair $((W_i), (\pi_{j,i}))$ is an *inverse sequence* of finite dimensional vector spaces.

Lemma 2.5. *There exists a sequence (B_j) of disjoint finite sets in W with the following properties.*

- (i) $\pi_i(B_j) = \{0\}$ for all $i < j$.
- (ii) $\pi_j(B_1 \cup \dots \cup B_j)$ is a basis for W_j .
- (iii) $|B_1 \cup \dots \cup B_j| = \dim W_j$ for each $j \in \mathbb{N}$.

Proof. Choose a minimal finite subset $B_1 \subset W$ with the property that $\pi_1(B_1)$ is a basis for W_1 . Note that $\pi_1(w) = \pi_{2,1} \circ \pi_2(w)$ for all $w \in W$ and so it follows that $\pi_2(B_1)$ is a linearly independent set in W_2 . Let Y_2 be the linear span of $\pi_2(B_1)$ in W_2 . Since the restriction map $\pi_{2,1}: W_2 \rightarrow W_1$ is linear and surjective we have $W_2 = Y_2 \oplus \ker \pi_{2,1}$ and $\dim Y_2 = \dim W_1$. Now choose a minimal finite subset $B_2 \subset W$ with the property that $\pi_2(B_2)$ is a basis for $\ker \pi_{2,1}$. Note that B_1 and B_2 are disjoint. Then $\pi_1(B_2) = \pi_{2,1} \circ \pi_2(B_2) = 0$, $\pi_2(B_1 \cup B_2)$ is a basis for W_2 and $|B_1 \cup B_2| = \dim W_2$. Repeating these arguments, the sequence (B_j) can now be constructed inductively. \square

Theorem 2.6. *If A is a countably infinite set then every infinite dimensional vector subspace of X^A has a free basis.*

Proof. Let W be a subspace of X^A and let $\alpha = (A_i)$ be an increasing sequence of finite subsets which cover A . There exists a sequence (B_j) of disjoint finite sets in W with properties (i)–(iii) as in the statement of Lemma 2.5. It remains only to show that the countable set $\cup_{j \in \mathbb{N}} B_j = \{f_n : n \in \mathbb{N}\}$ is a free basis for W . By property (i), the sequence (f_n) tends to zero strictly. Suppose $w \in W$. By properties (i)–(iii), there exists a unique sequence (h_j) in W such that $h_j \in \text{span}(B_j)$ and $\pi_j(w) = \pi_j(h_1 + \dots + h_j)$ for each $j \in \mathbb{N}$. Define $h = \sum_{j=1}^\infty h_j$. Then $w = h \in \mathcal{M}(S)$, where $S = \{f_n : n \in \mathbb{N}\}$, and so $W \subseteq \mathcal{M}(S)$. \square

Remark 2.7. Note that the proof of Theorem 2.6 is non-constructive in the sense that the resulting free basis depends both on a choice of covering sequence $\alpha = (A_i)$ and a process of selection for the sequence (B_j) .

Given W and α as above, denote by $\varprojlim W_i$ the inverse limit of the inverse sequence $((W_i), (\pi_{j,i}))$. Thus $\varprojlim W_i$ is the vector space of all (f_i) in the Cartesian product $\prod_{i \in \mathbb{N}} W_i$ with the property that $\pi_{j,i}(f_j) = f_i$ whenever $j \geq i$. In the following, we consider the linear map $\Phi_\alpha : X^A \rightarrow \prod_{i \in \mathbb{N}} W_i, f \mapsto (\pi_i(f))$.

Lemma 2.8. *The following statements are equivalent.*

- (i) W is closed, with respect to the product topology on X^A .
- (ii) $\Phi_\alpha(W) = \varprojlim W_i$.

Proof. To show (i) \rightarrow (ii), suppose W is closed in X^A . Let $(\pi_i(w_i)) \in \varprojlim W_i$, where $w_i \in W$ for each $i \in \mathbb{N}$, and define $w \in X^A$ by setting $w(a) = w_j(a)$ for all $a \in A_j$, and all $j \in \mathbb{N}$. Note that the sequence (w_j) converges pointwise to w . Thus, $w \in W$ and $\Phi_\alpha(w) = (\pi_i(w_i))$.

To show $(ii) \rightarrow (i)$, suppose $\Phi_\alpha(W) = \varprojlim W_i$ and let (w_n) be a sequence in W which converges pointwise to $h \in X^A$. For each $i \in \mathbb{N}$, the sequence $(\pi_i(w_n))$ in W_i converges pointwise to $\pi_i(h) \in X^{A_i}$. Since X^{A_i} is finite dimensional, W_i is closed and so $\pi_i(h) \in W_i$. Thus, the sequence $(\pi_i(h))$ lies in the inverse limit $\varprojlim W_i$ and so there exists $w \in W$ such that $\Phi_\alpha(w) = (\pi_i(h))$. Note that $\pi_i(h) = \pi_i(w)$ for all $i \in \mathbb{N}$ and so $h = w \in W$. \square

Theorem 2.9. *If A is a countably infinite set then every infinite dimensional closed vector subspace W of X^A has a free basis $S = \{f_n : n \in \mathbb{N}\}$ with $W = \mathcal{M}(S)$.*

Proof. Let W be a closed subspace of X^A and let $S = \{f_n : n \in \mathbb{N}\}$ be the free basis for W obtained from a sequence (B_j) as in the proof of Theorem 2.6. Then $W \subseteq \mathcal{M}(S)$ and so it only remains to show the reverse inclusion holds. Let $h \in \mathcal{M}(S)$. By properties (i) and (ii) of the sequence (B_j) , $\pi_i(h) \in W_i$ for each $i \in \mathbb{N}$ and so $(\pi_i(h)) \in \varprojlim W_i$. Since W is closed, by Lemma 2.8 there exists $w \in W$ such that $\Phi_\alpha(w) = (\pi_i(h))$. Now $h = w \in W$ and so $\mathcal{M}(S) \subseteq W$. \square

Given $f \in X^A$ we write $\|f\|_\infty = \sup_{a \in A} \|f(a)\|$. A mapping $f \in X^A$ is *bounded* if $\|f\|_\infty < \infty$. Otherwise, f is said to be *unbounded*.

Corollary 2.10. *Let A be a countably infinite set. If W is an infinite dimensional closed subspace of X^A then W contains a countable linearly independent set of unbounded elements.*

Proof. By Theorem 2.9, there exists a free basis S for W with the property that $W = \mathcal{M}(S)$. Since S is countably infinite, we may choose a sequence (h_k) in S and a sequence (a_k) in A so that

$$h_k(a_k) \neq 0 \text{ and } h_k(a_j) = 0, \quad \text{for } 1 \leq j \leq k-1.$$

Replacing h_k by a scalar multiple of h_k , we may assume for convenience that $\|h_k(a_k)\| = 1$ for all k . Choose non-zero scalars $(\alpha_{1,n})$ successively such that $\alpha_{1,1} = 1$ and $|\alpha_{1,n}| \geq n + \|\sum_{k=1}^{n-1} \alpha_{1,k} h_k(a_n)\|$ for each $n \geq 2$. Let $g_1 = \sum_{k=1}^\infty \alpha_{1,k} h_k \in W$ and note that for each $n \in \mathbb{N}$,

$$\|g_1(a_n)\| = \left\| \sum_{k=1}^n \alpha_{1,k} h_k(a_n) \right\| \geq \|\alpha_{1,n} h_n(a_n)\| - \left\| \sum_{k=1}^{n-1} \alpha_{1,k} h_k(a_n) \right\| \geq n.$$

Thus, g_1 is unbounded. For $m \geq 2$, similarly define $g_m = \sum_{k=m}^\infty \alpha_{m,k} h_k$ with non-zero scalars $(\alpha_{m,k})$ chosen so that g_m is unbounded. To see that the set $\{g_m : m \in \mathbb{N}\}$ is linearly independent note that if $\sum_{m=1}^d \lambda_m g_m = 0$ for some $\lambda_1, \dots, \lambda_d \in \mathbb{K}$ then $\lambda_1 \alpha_{1,1} h_1(a_1) = \sum_{m=2}^d \lambda_m g_m(a_1) = 0$ and so $\lambda_1 = 0$. By similar arguments it follows that $\lambda_j = 0$ for each $j = 2, \dots, d$. \square

2.2. Application to bar-joint frameworks

The results of the previous section apply to the infinitesimal flex spaces of countable bar-joint frameworks (and in particular, to crystallographic bar-joint frameworks). Let




Figure 1. A motif for the bar-joint framework $\mathcal{C}_{\text{kite}}$ together with four velocity vectors from the unbounded infinitesimal flex a_{kite} .

Fig. 1 - B/W online, B/W in print

$G = (V, E)$ be a simple graph with vertex set V and edge set E . A *bar-joint framework* for G in \mathbb{R}^d is a pair $\mathcal{G} = (G, p)$ where $p : V \rightarrow \mathbb{R}^d, v \mapsto p_v$, is an injective map. The points p_v are referred to as *joints* of \mathcal{G} and the line segments (p_v, p_w) , where $vw \in E$, are referred to as *bars*. Let $\mathcal{V}(\mathcal{G}; \mathbb{K}) = X^A$ where $A = p(V)$ and $X = \mathbb{K}^d$ (for $\mathbb{K} = \mathbb{R}$ or \mathbb{C}). The elements of $\mathcal{V}(\mathcal{G}; \mathbb{K})$ are referred to as *velocity fields* for \mathcal{G} over \mathbb{K} . If $u \in \mathcal{V}(\mathcal{G}; \mathbb{K})$ then $u(p_v)$ is referred to as the *velocity vector* for u at p_v . An *infinitesimal flex* (or *first order flex*) for \mathcal{G} over \mathbb{K} is a velocity field $u \in \mathcal{V}(\mathcal{G}; \mathbb{K})$ with the property that $\langle p_v - p_w, u(p_v) \rangle = \langle p_v - p_w, u(p_w) \rangle$ for each edge $vw \in E$. The set of infinitesimal flexes for \mathcal{G} is a closed vector subspace of $\mathcal{V}(\mathcal{G}; \mathbb{K})$, denoted $\mathcal{F}(\mathcal{G}; \mathbb{K})$. The vector subspace of bounded infinitesimal flexes for \mathcal{G} is denoted $\mathcal{F}_\infty(\mathcal{G}; \mathbb{K})$.

Example 2.11. Consider the crystallographic bar-joint framework $\mathcal{C}_{\text{kite}}$ in the Euclidean plane which is defined by the kite-shaped motif indicated in Figure 1 consisting of two joints and five bars. We assume that the vertical crossbar is closer to the apex joint, positioned to the left, than it is to the tail joint on the right. The figure also indicates a pair of periodicity vectors $\{a_1, a_2\}$ and four velocity vectors at the four joints of a kite subframework of $\mathcal{C}_{\text{kite}}$. These velocity vectors are indicative of an infinitesimal rotational flex of this subframework about the centre of the crossbar.

Let \vec{x}, \vec{y} be a choice of non-zero infinitesimal translation flexes of $\mathcal{C}_{\text{kite}}$, for the x and y directions, respectively, and let \vec{r} be a non-zero infinitesimal rotation flex of $\mathcal{C}_{\text{kite}}$. Additionally, note that there is an unbounded infinitesimal flex a_{kite} in $\mathcal{F}(\text{nn}\mathcal{C}_{\text{kite}}; \mathbb{R})$ which restricts to alternating rotations of the 5-bar kite framework and its translates, each of these restriction flexes being rotational about the centre of the crossbar. The magnitude of these rotations tends to infinity exponentially in the positive x -direction and is constant in magnitude in the y -direction.

It is straightforward to show that the finite set $\{\vec{x}, \vec{y}, \vec{r}, a_{\text{kite}}\}$ is a vector space basis for $\mathcal{F}(\mathcal{C}_{\text{kite}}; \mathbb{R})$. Indeed, let \mathcal{G} be a finite subframework of $\mathcal{C}_{\text{kite}}$ formed by 2 kite subframeworks which meet at a common joint. The infinitesimal flex space of \mathcal{G} is four-dimensional and is spanned by the restrictions of the quartet $\vec{x}, \vec{y}, \vec{r}, a_{\text{kite}}$. Thus, if u is an infinitesimal flex of $\mathcal{C}_{\text{kite}}$ then we may subtract from u a linear combination of this quartet to obtain a flex u_1 whose restriction to \mathcal{G} is 0. One can see readily from the geometry of $\mathcal{C}_{\text{kite}}$ and the rigidity of the kite subframeworks that this implies that $u_1 = 0$.

Let $\mathcal{G} = (G, p)$ be a finite or countable bar-joint framework whose joints do not lie in a hyperplane. The space $\mathcal{F}_{\text{rig}}(\mathcal{G}; \mathbb{R})$ of trivial infinitesimal flexes, or *rigid motion infinitesimal flexes*, is the vector subspace of $\mathcal{F}(\mathcal{G}; \mathbb{R})$ consisting of real infinitesimal flexes of the complete graph bar-joint framework (K_V, p) . Here K_V is the complete graph on the vertex set V . The bar-joint framework \mathcal{G} is *infinitesimally rigid* if $\mathcal{F}(\mathcal{G}; \mathbb{R}) = \mathcal{F}_{\text{rig}}(\mathcal{G}; \mathbb{R})$ and *boundedly infinitesimally rigid* if $\mathcal{F}_{\infty}(\mathcal{G}; \mathbb{R}) \subset \mathcal{F}_{\text{rig}}(\mathcal{G}; \mathbb{R})$.

Example 2.12. The three velocity fields $\vec{x}, \vec{y}, \vec{r}$ in Example 2.11 span $\mathcal{F}_{\text{rig}}(\mathcal{C}_{\text{kite}}; \mathbb{R})$. Also a_{kite} is not a bounded infinitesimal flex. It follows that $\mathcal{C}_{\text{kite}}$ is boundedly infinitesimally rigid, but not infinitesimally rigid.

In contrast to this, consider the semi-crystallographic framework $\mathcal{C}_{\text{kite}}^{x \leq 0}$ which is formed by removing from $\mathcal{C}_{\text{kite}}$ all the 5-bar kite frameworks not lying in the half-plane $x \leq 0$. This is not boundedly infinitesimally rigid. Indeed it can be shown, as in the consideration of $\mathcal{C}_{\text{kite}}$, that the flex space is four-dimensional with basis given by the restrictions to $\mathcal{C}_{\text{kite}}^{x \leq 0}$ of the quartet $\vec{x}, \vec{y}, \vec{r}, a_{\text{kite}}$. Since the restriction of a_{kite} to $\mathcal{C}_{\text{kite}}^{x \leq 0}$ is bounded, the semi-crystallographic framework is not boundedly rigid. One may view this restriction infinitesimal flex (and its associated zero energy mechanical mode) as a *bounded surface flex* (or surface mode), of the bulk crystal $\mathcal{C}_{\text{kite}}$, which is associated with the domain wall $x = 0$.

Theorem 2.13. Let $\mathcal{G} = (G, p)$ be a bar-joint framework in \mathbb{R}^d with a countable set of joints. If the space of infinitesimal flexes $\mathcal{F}(\mathcal{G}; \mathbb{K})$ is infinite dimensional then,

- (i) $\mathcal{F}(\mathcal{G}; \mathbb{K})$ has a free basis, and,
- (ii) $\mathcal{F}(\mathcal{G}; \mathbb{K})$ contains a countable linearly independent set of unbounded infinitesimal flexes.

Proof. The statements follow from Theorem 2.9 and Corollary 2.10 since $\mathcal{F}(\mathcal{G}, \mathbb{K})$ is closed in $\mathcal{V}(\mathcal{G}; \mathbb{K})$. □

Several examples of free bases for the infinitesimal flex spaces of crystallographic bar-joint frameworks are presented in §5.

3. Group actions on free spanning sets

Once again let A be a countably infinite set and X a finite-dimensional normed space over \mathbb{K} . Throughout this section, Γ denotes a multiplicative free abelian group with identity element 1, $\theta: \Gamma \times A \rightarrow A$ is a free group action on A and $\pi: \Gamma \times X^A \rightarrow X^A$ is the induced faithful group action on X^A given by $\pi(\gamma, f)(a) = f(\gamma^{-1}a)$ for all $a \in A$. To simplify notation, $\theta(\gamma, a)$ will be written as γa for all $\gamma \in \Gamma$ and $a \in A$. Similarly, $\pi(\gamma, f)$ will be written as γf for all $\gamma \in \Gamma$ and $f \in X^A$. The *orbit* of an element $a \in A$ under the group action θ is the set $\Gamma a = \{\gamma a: \gamma \in \Gamma\}$ and the *quotient set* $A/\Gamma = \{\Gamma a: a \in A\}$ is the set of all such orbits.

Definition 3.1. A *geometric direction* for $f \in X^A \setminus \{0\}$ is an element $\gamma \in \Gamma$ such that γf is a non-zero scalar multiple of f .

The set of all geometric directions for f is denoted Γ_f .

Lemma 3.2. Let $f \in X^A \setminus \{0\}$.

- (i) Γ_f is a subgroup of Γ .
- (ii) $\Gamma_f = \Gamma_{\lambda(\gamma f)}$ for all non-zero scalars $\lambda \in \mathbb{K}$ and all $\gamma \in \Gamma$.
- (iii) If f has finite support then $\Gamma_f = \{1\}$.

Proof. The proofs of parts (i) and (ii) are elementary and left to the reader. To show (iii), suppose $\gamma \in \Gamma$ is a geometric direction for f and choose $a \in \text{supp}(f)$. Then $\gamma^k a \in \text{supp}(f)$ for each $k \in \mathbb{Z}$. Since θ is a free group action, the elements $\gamma^k a$, $k \in \mathbb{Z}$, are distinct unless $\gamma = 1$. □

Definition 3.3. Let $S \subset X^A \setminus \{0\}$. A group action $\pi: \Gamma \times X^A \rightarrow X^A$ acts on S (up to scalar multiples) if for each $f \in S$ and each $\gamma \in \Gamma$ there exists a scalar $\lambda \in \mathbb{K}$ and $g \in S$ such that $\gamma f = \lambda g$.

Lemma 3.4. Let S be a free spanning set for an infinite-dimensional vector subspace of X^A and let $f \in S$. If the quotient set A/Γ under the action θ is finite and the induced action π acts on S up to scalar multiples then the following statements are equivalent.

- (i) f has finite support.
- (ii) $\Gamma_f = \{1\}$.

Proof. The implication (i) \rightarrow (ii) is Lemma 3.2(iii). To show (ii) \rightarrow (i), suppose $\Gamma_f = \{1\}$. Let $S = \{f_n : n \in \mathbb{N}\}$ and let a_1, \dots, a_r be a set of representatives for the finitely many orbits in A/Γ . For $i \in \{1, \dots, r\}$, define $C_i = \{\gamma f : \gamma \in \Gamma, (\gamma f)(a_i) \neq 0\}$. Note that, since the sequence (f_n) tends to zero strictly, the set $S_i = \{g \in S : g(a_i) \neq 0\}$ is finite and, since π acts on S , each element of C_i is a scalar multiple of an element of S_i . If C_i is not finite, for some $i \in \{1, \dots, r\}$, then there exist distinct elements $\gamma_1 f, \gamma_2 f \in C_i$ and a non-zero scalar $\lambda \in \mathbb{K}$ such that $\gamma_1 f = \lambda(\gamma_2 f)$. Thus $(\gamma_1 \gamma_2^{-1})f = \lambda f$ and so $\gamma_1 \gamma_2^{-1}$ is a geometric direction for f , which contradicts $\Gamma_f = \{1\}$. Thus C_i must be finite for each $i \in \{1, \dots, r\}$. Now for each C_i , either $\gamma_1 f = \gamma_2 f$ for some distinct $\gamma_1, \gamma_2 \in \Gamma$ or there are only finitely many $\gamma \in \Gamma$ with $(\gamma f)(a_i) \neq 0$. If the former case holds for some C_i , then $f = (\gamma_1 \gamma_2^{-1})f$ and so $\gamma_1 \gamma_2^{-1}$ is a geometric direction for f , which contradicts $\Gamma_f = \{1\}$. Thus, the latter case holds for each C_i . Since $f(\gamma^{-1} a_i) = (\gamma f)(a_i)$ for all $\gamma \in \Gamma$ and $i \in \{1, \dots, r\}$, and since $A = \{\gamma a_i : \gamma \in \Gamma, i \in \{1, \dots, r\}\}$, it follows that f has finite support. □

Definition 3.5. A set $S \subset X^A \setminus \{0\}$ is *finitely generated* by a group action $\pi: \Gamma \times X^A \rightarrow X^A$ (up to scalar multiples) if π acts on S , and, there is a finite subset $S_0 \subset S$ such that for every $f \in S$ there exists $\gamma \in \Gamma$, $f_0 \in S_0$, and a scalar $\lambda \in \mathbb{K}$, with $f = \lambda(\gamma f_0)$.

Lemma 3.6. Let S be a free spanning set for an infinite-dimensional vector subspace of X^A . If S is finitely generated by a group action π then there exists $f \in S$ such that the quotient group Γ/Γ_f has infinite order.

Proof. Let $\mathcal{I} = \{\mathbb{K}f : f \in S\}$ where $\mathbb{K}f$ denotes the one-dimensional subspace of X^A spanned by f . If \mathcal{I} is a finite set, then there exists a sequence (f_n) in S for which the one-dimensional spaces $\mathbb{K}f_n$ are equal. In particular, the support sets $\text{supp}(f_n)$ are equal for all $n \in \mathbb{N}$. This is a contradiction since (f_n) tends to zero strictly and so \mathcal{I} is countably infinite. Since S is finitely generated by π , there exists a finite subset $S_0 \subset S$ such that $\mathcal{I} = \{\mathbb{K}(\gamma f) : \gamma \in \Gamma, f \in S_0\}$. Note that, for each $f_0 \in S_0$, the cardinality of $\{\mathbb{K}(\gamma f_0) : \gamma \in \Gamma\}$ is bounded by the order of Γ/Γ_{f_0} . Thus, Γ/Γ_{f_0} has infinite order for some $f_0 \in S_0$. \square

3.1. Lattice group actions

Let $L = \{\sum_{i=1}^m n_i b_i : n_i \in \mathbb{Z}\}$ be a rank m lattice in \mathbb{R}^d determined by linearly independent vectors $b_1, \dots, b_m \in \mathbb{R}^d$. For each $b \in L$, let $T_b : \mathbb{R}^d \rightarrow \mathbb{R}^d$ denote the translation $T_b(x) = x + b$. The translation group associated with L is the multiplicative abelian group $\mathcal{T} = \{T_b : b \in L\}$. A *multilattice* on L is a set $A = A_1 \cup \dots \cup A_s$ where A_1, \dots, A_s are pairwise disjoint translates of L . Define a free group action $\theta_L : \mathcal{T} \times A \rightarrow A$ by setting $\theta_L(T, a_i) = T(a_i)$ for $a_i \in A_i, 1 \leq i \leq s$. The *lattice group action* on X^A induced by θ_L is the group action $\pi_L : \mathcal{T} \times X^A \rightarrow X^A$ with $\pi_L(T, f) = f \circ T^{-1}$. Note that the quotient set A/\mathcal{T} is finite and a translation $T_b \in \mathcal{T}$ is a geometric direction for $f \in X^A$ if and only if there exists a non-zero scalar λ such that $f(a - b) = \lambda f(a)$ for all $a \in A$. If $\Gamma_f \neq \{1\}$ then the sublattice $L_f = \{b \in L : T_b \in \Gamma_f\}$ is referred to as the lattice of geometric directions for f .

Definition 3.7. Let L' be a rank n sublattice of L with basis $b_1, \dots, b_n \in \mathbb{R}^d$. A mapping $f \in X^A$ is *factor periodic* on L' if there exists $\omega = (\omega_1, \dots, \omega_n) \in \mathbb{K}_*^n$ such that $T_b f = \omega_1^{k_1} \dots \omega_n^{k_n} f$ for all $b = \sum_{j=1}^n k_j b_j \in L'$. In this case, ω is referred to as the periodicity multifactor for f determined by the basis $b_1, \dots, b_n \in \mathbb{R}^d$.

Lemma 3.8. Let A be a multilattice on L and let $f \in X^A \setminus \{0\}$.

- (i) If $\Gamma_f \neq \{1\}$ then f is factor periodic on L_f .
- (ii) If f is bounded and factor periodic, with periodicity multifactor ω , then the components of ω are unimodular.

Proof. (i) If $\Gamma_f \neq \{1\}$ then L_f is a lattice of rank n say. Let $b_1, \dots, b_n \in \mathbb{R}^d$ be a set of generators for the lattice L_f . For each $i = 1, \dots, n, T_{b_i}$ is a geometric direction for f and so $T_{b_i} f = \omega_i f$ for some non-zero scalar $\omega_i \in \mathbb{K}$. Thus, $T_b f = T_{b_1}^{k_1} \dots T_{b_n}^{k_n} f = \omega_1^{k_1} \dots \omega_n^{k_n} f$ for all $b = \sum_{j=1}^n k_j b_j \in L_f$.

(ii) Suppose f is factor periodic on the lattice L' and let $\omega = (\omega_1, \dots, \omega_n) \in \mathbb{K}_*^n$ be the periodicity factor for f determined by a basis b_1, \dots, b_n for L' . Let $a \in \text{supp}(f)$. Note that, for each $j = 1, \dots, n, \|f(a - kb_j)\| = \|T_{b_j}^k f(a)\| = |\omega_j|^k \|f(a)\|$ for all $k \in \mathbb{Z}$. Thus, if $|\omega_j| \neq 1$ for some j then f is clearly unbounded. \square

If $S \subset X^A$ then, for each $a \in A$, let $S_a = \{f \in S : f(a) \neq 0\}$.

Lemma 3.9. Let A be a multilattice on L and let S be a free spanning set for an infinite-dimensional subspace of X^A . If the lattice group action π_L acts on S then each $f \in S$ satisfies one of the following conditions.

- (i) f is finitely supported.
- (ii) f is factor periodic on L_f and the support of f is a multilattice on L_f .

Proof. Suppose $f \in S$ is not finitely supported. By Lemma 3.4, $\Gamma_f \neq \{1\}$. Thus, by Lemma 3.8, f is factor periodic on L_f . Choose $a_1, \dots, a_s \in A$ such that $A = \cup_{i=1}^s (a_i + L)$. Since S is a free spanning set, S_{a_i} is finite and so $S_{a_i} = \{g_{a_i,1}, \dots, g_{a_i,s_i}\}$ say. By Lemma 3.4, the subgroup L_f is a non-zero sublattice of L . Let $L_{a_i,k} = \{b \in L : T_b^{-1}f = \lambda g_{a_i,k}, \text{ for some } \lambda \neq 0\}$ for $k=1, \dots, s_i$. Note that $(T_b^{-1}f)(a_i) = f(a) \neq 0$ for each $a \in \text{supp}(f)$ with $a = a_i + b$ and $b \in L$. Thus, since π_L acts on S , it follows that $\text{supp}(f) = \cup_{i=1}^s \cup_{k=1}^{s_i} (a_i + L_{a_i,k})$. If $b, b' \in L_{a_i,k}$ then $T_b f = \lambda(T_{b'} f)$ for some non-zero scalar $\lambda \in \mathbb{K}$ and so $T_{b-b'}$ is a geometric direction for f . In particular, $b - b' \in L_f$. Thus, if $b_{a_i,k} \in L_{a_i,k}$ then it follows that $L_{a_i,k} = b_{a_i,k} + L_f$, and so $\text{supp}(f)$ is a finite union of translates of the lattice L_f . □

In the following, \mathbb{R}^d is endowed with a norm and $\text{dist}(a, E) = \inf_{x \in E} \|a - x\|$ denotes the distance between a point $a \in \mathbb{R}^d$ and a subset $E \subset \mathbb{R}^d$.

Definition 3.10. Let A be a multilattice in \mathbb{R}^d . A mapping $f \in X^A$ is band-limited with respect to a proper linear subspace \mathcal{K} of \mathbb{R}^d if there exists $C > 0$ such that the support of f is contained in the ‘band’ $\{x \in \mathbb{R}^d : \text{dist}(x, \mathcal{K}) \leq C\}$.

Theorem 3.11. Let A be a multilattice and let S be a free spanning set for an infinite-dimensional vector subspace of X^A . If S is finitely generated by the lattice group action π_L , and contains no finitely supported elements, then there exists $f \in S$ such that the linear span \mathcal{K} of L_f is a proper subspace of \mathbb{R}^d and f is band-limited with respect to \mathcal{K} .

Proof. Suppose no element of S has finite support. By Lemma 3.6, there exists $f \in S$ such that the quotient group Γ/Γ_f has infinite order. Recall that a quotient of two free abelian groups with equal rank has finite order. Thus, Γ_f has rank k strictly less than d . Let \mathcal{K} be the proper linear subspace in \mathbb{R}^d spanned by L_f . By Lemma 3.9, the support of f is a multilattice on L_f and hence is contained in a band $\{x \in \mathbb{R}^d : d(x, \mathcal{K}) \leq C\}$ for some $C > 0$. □

A subset S of X^A is bounded if each element of S is bounded and $\sup_{f \in S} \|f\|_\infty < \infty$.

Lemma 3.12. Let A be a multilattice on L and let $S = \{f_n : n \in \mathbb{N}\}$ be a bounded free spanning set for an infinite-dimensional subspace of X^A . If the lattice group action π_L acts on S and $(\alpha_n) \in \ell^\infty(\mathbb{N})$ is a bounded sequence of scalars then $\sum_{n=1}^\infty \alpha_n f_n$ is bounded.

Proof. Suppose $f = \sum_{n=1}^\infty \alpha_n f_n$ for some $\alpha = (\alpha_n) \in \ell^\infty(\mathbb{N})$. Since (f_n) tends to zero strictly, S_a is a finite set for each $a \in A$. Note that for each $b \in L$, $f(a + b) \neq 0$ if and only if $(T_b^{-1}f)(a) \neq 0$. Thus, since π_L acts on S , the set S_{a+b} has the same cardinality as S_a . Choose $a_1, \dots, a_s \in A$ such that $A = \cup_{i=1}^s (a_i + L)$. Let $N = \sup_{a \in A} |S_a| = \max_{i=1, \dots, s} |S_{a_i}|$ and let $M = \sup_n \|f_n\|_\infty$. Then $\|f\|_\infty = \sup_{a \in A} \|f(a)\| \leq NM \|\alpha\|_\infty$ and so f is bounded. □

In the following, a set $S \subset X^A \setminus \{0\}$ has the *local basis property* if, for each $a \in A$, the set $\{f(a): f \in S_a\}$ is a basis for X . Also, W_∞ denotes the set of bounded elements in a subspace W of X^A .

Theorem 3.13. *Let A be a multilattice on L and let $S = \{f_n : n \in \mathbb{N}\}$ be a free spanning set for an infinite-dimensional subspace W of X^A with the following properties.*

- (i) *The lattice group action π_L acts on S .*
- (ii) *S is bounded and has the local basis property.*
- (iii) *No element of S is finitely supported.*
- (iv) $\inf\{\|g(a)\| : a \in A, g \in S_a\} > 0$.

Then,

$$W_\infty = \left\{ f \in W : f = \sum_{n=1}^{\infty} \alpha_n f_n \text{ and } (\alpha_n) \in \ell^\infty(\mathbb{N}) \right\}.$$

Proof. Let $f \in W_\infty$. Then $f = \sum_{n=1}^{\infty} \alpha_n f_n$ for some sequence of scalars (α_n) and $\|f\|_\infty < \infty$. Since S has the local basis property, given any $a \in A$, the set $\{g(a): g \in S_a\}$ is a basis for X and so we may consider the norm on X given by $\|x\|_a = \max_{g \in S_a} |\lambda_g|$ where $x = \sum_{g \in S_a} \lambda_g g(a)$. Moreover, since X is finite-dimensional, there exists $c_a > 0$ such that $c_a \|x\|_a \leq \|x\|$ for all $x \in X$. Suppose $\alpha_m \neq 0$ and choose $a \in A$ with $f_m(a) \neq 0$. Since the lattice group action π_L acts on S and no element of S is finitely supported, it follows from Lemmas 3.4 and 3.8(i) that each $g \in S_a$ is factor periodic on L_g . Moreover, since each element of S is bounded, it follows from Lemma 3.8(ii) that the components of the periodicity factors for each $g \in S_a$ are unimodular. Choose $a_1, \dots, a_s \in A$ such that $A = \cup_{i=1}^s (a_i + L)$. Then $a = a_j + b$ for some j and some $b \in L$. Thus, since the lattice group action π_L acts on S , there exist scalars μ_g such that $S_a = \{\mu_g g : g \in S_{a_j}\}$. It follows that $\|x\|_a \leq \frac{1}{M} \|x\|_{a_j}$ where $M > 0$ is a lower bound for $\{\|g(a)\| : a \in A, g \in S_a\}$. Thus, $c_{a_j} \|x\|_a \leq c_{a_j} (\frac{1}{M}) \|x\|_{a_j} \leq \frac{1}{M} \|x\|$ for each $x \in X$. Note that $|\alpha_m| \leq \|f(a)\|_a \leq \frac{1}{cM} \|f(a)\| \leq \frac{1}{cM} \|f\|_\infty$ where $c = \min_{i=1, \dots, s} c_{a_i}$. Thus $(\alpha_n) \in \ell^\infty(\mathbb{N})$.

For the reverse inclusion, apply Lemma 3.12. □

4. Crystal flex bases and the RUM spectrum

An *automorphism* of a graph $G = (V, E)$ is a bijection $\beta: V \rightarrow V$ with the property that $vw \in E$ if and only if $\beta(v)\beta(w) \in E$. The *space group* for a bar-joint framework $\mathcal{G} = (G, p)$ in \mathbb{R}^d is the group $\mathfrak{S}(\mathcal{G})$ of Euclidean isometries $T: \mathbb{R}^d \rightarrow \mathbb{R}^d$ with the property that $T(p(V)) = p(V)$ and the induced map $\beta: V \rightarrow V, v \mapsto p^{-1}(T(p_v))$ is an automorphism of G . The subgroup of $\mathfrak{S}(\mathcal{G})$ consisting of translations is denoted $\mathcal{T}(\mathcal{G})$.

Definition 4.1. A bar-joint framework $\mathcal{C} = (G, p)$ in \mathbb{R}^d is referred to as a *crystal framework* if there exists a rank d lattice L in \mathbb{R}^d with the following properties.

- (a) The translation group $\mathcal{T} = \{T_b : b \in L\}$ is a subgroup of $\mathfrak{S}(\mathcal{C})$.
- (b) \mathcal{C} has only finitely many distinct vertex orbits and edge orbits under \mathcal{T} .

In this case, the lattice L (or equivalently, the translation group \mathcal{T}) is referred to as a *periodic structure* for \mathcal{C} .

Let \mathcal{C} be a crystal framework in \mathbb{R}^d . The *space group action* on $\mathcal{V}(\mathcal{C}; \mathbb{R})$ is the faithful group action $\pi_{\mathfrak{S}(\mathcal{C})} : \mathfrak{S}(\mathcal{C}) \times \mathcal{V}(\mathcal{C}; \mathbb{R}) \rightarrow \mathcal{V}(\mathcal{C}; \mathbb{R})$ with $\pi_{\mathfrak{S}(\mathcal{C})}(T, f) = T_2 \circ f \circ T^{-1}$ where $T = T_1 T_2$ is the unique factorization of the Euclidean isometry T with T_1 a translation and T_2 an orthogonal linear transformation. Also, we define the *space group action* on $\mathcal{V}(\mathcal{C}; \mathbb{C})$ as the natural induced action. Alternatively, for $f \in \mathcal{V}(\mathcal{C}; \mathbb{C})$ we may define $\pi_{\mathfrak{S}(\mathcal{C})}(T, f) = \tilde{T}_2 \circ f \circ T^{-1}$ where \tilde{T}_2 has the diagonal action $T_2 \oplus T_2$ on the direct sum $\mathbb{C}^d = \mathbb{R}^d \oplus i\mathbb{R}^d$. The *lattice group action* on $\mathcal{V}(\mathcal{C}; \mathbb{K})$ determined by a periodic structure L with translation group $\mathcal{T} = \{T_b : b \in L\}$ is the group action $\pi_{\mathcal{T}} : \mathcal{T} \times \mathcal{V}(\mathcal{C}; \mathbb{K}) \rightarrow \mathcal{V}(\mathcal{C}; \mathbb{K})$, $\pi_{\mathcal{T}}(T, f) = f \circ T^{-1}$.

Lemma 4.2. *Let $\mathcal{C} = (G, p)$ be a crystal framework in \mathbb{R}^d and let $S \subset \mathcal{V}(\mathcal{C}; \mathbb{K})$. If S is finitely generated by the space group action $\pi_{\mathfrak{S}(\mathcal{C})}$ then S is finitely generated by the lattice group action $\pi_{\mathcal{T}}$ for any choice of periodic structure \mathcal{T} .*

Proof. Suppose S is finitely generated by $\pi_{\mathfrak{S}(\mathcal{C})}$ and let \mathcal{T} be a periodic structure for \mathcal{C} . Then given any $f \in S$ and any $T \in \mathfrak{S}(\mathcal{C})$ there exists $\lambda \in \mathbb{K}$ and $g \in S$ such that $Tf = \lambda g$. In particular, this holds for all $T \in \mathcal{T}$ and so $\pi_{\mathcal{T}}$ acts on S . Also, there exists a finite subset $S_0 \subset S$ such that every member of S can be expressed as a scalar multiple of Tf for some $T \in \mathfrak{S}(\mathcal{C})$ and some $f \in S_0$. Note that the set of joints of \mathcal{C} is a multilattice and the quotient group $\mathfrak{S}(\mathcal{C})/\mathcal{T}$ acts faithfully on the finite set of translation orbits. It follows that the quotient group $\mathfrak{S}(\mathcal{C})/\mathcal{T}$ is finite. Let R_1, \dots, R_m be a set of representatives for the finitely many elements in $\mathfrak{S}(\mathcal{C})/\mathcal{T}$ and let $S_0' = \{R_j f : f \in S_0, j = 1, \dots, m\}$. Then every member of S is a scalar multiple of Tf for some $T \in \mathcal{T}$ and some $f \in S_0'$ and so S is finitely generated over \mathcal{T} . □

A *geometric flex* for a crystal framework \mathcal{C} is an infinitesimal flex which is factor periodic with respect to a sublattice of a periodic structure for \mathcal{C} . A *local geometric flex* for \mathcal{C} is an infinitesimal flex which is factor periodic with respect to a lower rank sublattice of a periodic structure for \mathcal{C} . A *band-limited flex* for \mathcal{C} is an infinitesimal flex which is also a band-limited vector field in $\mathcal{V}(\mathcal{C}; \mathbb{K})$. For crystal frameworks, it is of interest to determine free bases for $\mathcal{F}(\mathcal{C}, \mathbb{K})$ which incorporate localized or band-limited flexes, should such flexes exist, and which, moreover, incorporate the crystallographic symmetry group. Accordingly, we make the following definitions.

Definition 4.3. Let \mathcal{C} be a crystal framework and let W be an infinite-dimensional vector subspace of $\mathcal{V}(\mathcal{C}; \mathbb{K})$.

- (a) A *crystal spanning set* (respectively, *crystal basis*) for W is a free spanning set (respectively, free basis) which is finitely generated by the space group action $\pi_{\mathfrak{S}(\mathcal{C})}$.

- (a) A crystal spanning set (respectively crystal basis) for $W = \mathcal{F}(\mathcal{C}; \mathbb{K})$ is also referred to as a *crystal flex spanning set* (respectively *crystal flex basis*) for \mathcal{C} .

In the next section, we identify a number of crystal frameworks, such as the grid frameworks and the kagome frameworks which possess a crystal flex basis in the sense above. In several cases, these bases consist entirely of local geometric flexes. Also, we see that the flex space of the octahedron framework has a free basis which is an *essential crystal flex basis* in the sense that there is a subset which is a crystal flex basis for a vector subspace of $\mathcal{F}(\mathcal{C}; \mathbb{R})$ which is complementary to the three-dimensional space of infinitesimal rotation flexes.

Theorem 4.4. *Let $\mathcal{C} = (G, p)$ be a crystal framework in \mathbb{R}^d and suppose $\mathcal{F}(\mathcal{C}; \mathbb{K})$ is infinite dimensional. Let S be a crystal flex spanning set for \mathcal{C} .*

- (i) *If $f \in S$ then f is either finitely supported or a geometric flex for \mathcal{C} .*
- (ii) *S must contain a band-limited flex. Moreover, $f \in S$ is a band-limited flex if and only if it is either finitely supported, or, a local geometric flex which, for any periodic structure L , is band-limited with respect to the linear span of L_f .*
- (iii) *If S is bounded, has the local basis property, has no finitely supported elements, and $\inf\{\|f(p(v))\| : v \in V, f \in S_{p(v)}\} > 0$, then,*

$$\mathcal{F}_\infty(\mathcal{C}; \mathbb{K}) = \{f \in \mathcal{F}(\mathcal{C}; \mathbb{K}) : f = \sum_{n=1}^\infty \alpha_n f_n \text{ and } (\alpha_n) \in \ell^\infty(\mathbb{N})\}.$$

Proof. By Lemma 4.2, S is finitely generated by the lattice group action $\pi_{\mathcal{T}}$ for any choice of periodic structure \mathcal{T} . The set of joints of \mathcal{C} is a multilattice in \mathbb{R}^d and so the results of §3 may be applied with $A = p(V)$ and $X = \mathbb{K}^d$. Statement (i) follows from Lemmas 3.4, 3.8 and 3.9. In particular, given any periodic structure L for \mathcal{C} , if $f \in S$ and f is not finitely supported then f is factor periodic for the sublattice L_f of L . Statement (ii) now follows since, by Lemma 3.9, the support of f is a multilattice on L_f . Thus, if f is band-limited then L_f must have rank t , where $t < d$, and so f is band-limited with respect to the linear span of L_f . Statement (iii) follows from Theorem 3.13. □

We now define a *transfer function* $\Psi_{\mathcal{C}}(z)$ associated with a crystal framework \mathcal{C} , a choice of periodic structure L with translation group \mathcal{T} and a choice of basis b_1, \dots, b_d for the lattice L . Let $F_v = \{p_\kappa : 1 \leq \kappa \leq |F_v|\}$ be a finite set of joints representing the \mathcal{T} -translation classes of the joints of \mathcal{C} . Then we may conveniently label the joints of \mathcal{C} as,

$$\{p_{\kappa,k} : 1 \leq \kappa \leq |F_v|, k \in \mathbb{Z}^d\},$$

where $p_{\kappa,k} = T_k p_\kappa$. Also let F_e be a finite set of representative bars for the translation classes of the bars. The pair (F_v, F_e) , referred to as a *motif* for \mathcal{C} , together with the periodic structure \mathcal{T} carries the essential geometric information which defines \mathcal{C} [14, 15].

Definition 4.5. Let \mathcal{C} be a crystal framework in \mathbb{R}^d with motif (F_v, F_e) and for $e = vw \in F_e$ let $p(e) = p(v) - p(w)$. The *transfer function* $\Psi_{\mathcal{C}}(z)$ is a matrix-valued function on \mathbb{C}_*^d whose rows are labelled by the edges of F_e and whose columns are labelled

by the vertex-coordinate pairs in $F_v \times \{1, \dots, d\}$. The row for an edge $e = (v, k)(w, l)$ with $v \neq w$ takes the form,

$$e \left[\begin{array}{ccccccc} 0 & \cdots & 0 & \overset{v}{p(e)z^{-k}} & \cdots & 0 & \overset{w}{-p(e)z^{-l}} & \cdots & 0 \end{array} \right],$$

while if $v = w$ it takes the form,

$$e \left[\begin{array}{ccccccc} 0 & \cdots & 0 & \overset{v}{p(e)(z^{-k} - z^{-l})} & \cdots & 0 & \end{array} \right].$$

The restriction of $\Psi_{\mathcal{C}}(z)$ to the d -torus \mathbb{T}^d gives the symbol function $\Phi_{\mathcal{C}}(z)$ considered in [1, 14, 15]. With the labelling of the joints of \mathcal{C} given above, note that a complex velocity field $u \in \mathcal{Y}(\mathcal{C}, \mathbb{C})$ is a map $u : F_v \times \mathbb{Z}^d \rightarrow \mathbb{C}^d$ where $u(p_{\kappa}, k)$ is the velocity vector assigned to the joint $p_{\kappa, k}$. Note that u is factor periodic, with periodicity multifactor $\omega \in \mathbb{C}_*^d$, if $u(p_{\kappa}, k) = \omega^k u(p_{\kappa}, 0)$ for all $p_{\kappa} \in F_v, k \in \mathbb{Z}^d$. Here ω^k is the product $\omega_1^{k_1} \dots \omega_d^{k_d}$. We write $u = b \otimes e_{\omega}$ for this vector field, where b is the vector $(u(p_{\kappa}, 0))_{\kappa \in F_v}$ in $\mathbb{C}^{d|F_v|}$ and e_{ω} is the multi-sequence $(\omega^k)_{k \in \mathbb{Z}^d}$.

Theorem 4.6. *Let \mathcal{C} be a crystal framework in \mathbb{R}^d with motif (F_v, F_e) and let $u = b \otimes e_{\omega}$ where $\omega \in \mathbb{C}_*^d$ and $b \in \mathbb{C}^{d|F_v|}$. The following conditions are equivalent.*

- (i) $u \in \mathcal{F}(\mathcal{C}, \mathbb{C})$.
- (ii) $\Psi(\omega^{-1})b = 0$.

Proof. The proof is similar to the unimodular case given in Power [15]. □

Definition 4.7. Let \mathcal{C} be a crystal framework in \mathbb{R}^d with a transfer function $\Psi_{\mathcal{C}}(z)$.

- (a) The *geometric flex spectrum* $\Gamma(\mathcal{C})$ of \mathcal{C} is defined to be the set,

$$\Gamma(\mathcal{C}) := \{\omega \in \mathbb{C}_*^d : \ker \Psi_{\mathcal{C}}(\omega^{-1}) \neq \{0\}\}.$$

- (b) The *rigid unit mode spectrum*, or RUM spectrum, of \mathcal{C} is the set,

$$\Omega(\mathcal{C}) := \{\omega \in \mathbb{T}^d : \ker \Phi_{\mathcal{C}}(\bar{\omega}) \neq \{0\}\},$$

where $\Phi_{\mathcal{C}}(z)$ is the restriction of $\Psi_{\mathcal{C}}(z)$ to the d -torus.

Note that $\Gamma(\mathcal{C})$ is the set of points $\omega \in \mathbb{C}_*^d$ for which there exists a non-zero factor periodic infinitesimal flex with periodicity multifactor ω . This set depends both on the choice of periodic structure L for \mathcal{C} and the choice of basis b_1, \dots, b_d for L . The RUM spectrum is the subset $\Omega(\mathcal{C}) = \Gamma(\mathcal{C}) \cap \mathbb{T}^d$ of multifactors ω with unimodular coordinates. Such multifactors are also referred to as *multi-phases*, or simply *phases*. Note also that for a critically coordinated crystal framework, in the elementary sense that $|F_e| = d|F_v|$, the transfer function is a square matrix-valued analytic function on its domain. It follows in this case that the determinant provides a multivariable analytic function and that the geometric spectrum is given by its set of zeros.

A small gallery of crystal frameworks and their transfer functions is given in Badri, Kitson and Power [1] and their RUM spectra are determined. See also Power [15] where the connection with mode vibrations in materials science is discussed. In § 5, we examine the geometric flex spectrum and the RUM spectrum of a novel bipyramid framework. In close analogy with the semi-crystallographic kite framework of Example 2.12, we see that there are geometrically decaying surface flexes associated with directions with no half-turn symmetry.

Definition 4.8. Let \mathcal{C} be a crystal framework in \mathbb{R}^d with a transfer function $\Psi_{\mathcal{C}}(z)$. The RUM spectrum $\Omega(\mathcal{C})$ is said to *contain linear structure* if the logarithmic representation of $\Omega(\mathcal{C})$ in $[0, 2\pi)^d$ contains a t -dimensional set of the form $[0, 2\pi)^d \cap H$ where H is an affine subspace of \mathbb{R}^d and $1 \leq t \leq d$.

If the RUM spectrum for a particular periodic structure contains linear structure then the same is true for the RUM spectrum for any periodic structure. This follows from the fact that the RUM spectrum for a periodic structure arises as the image of the primitive RUM spectrum under a natural surjective map [15].

Theorem 4.9. *Let \mathcal{C} be a crystal framework in \mathbb{R}^d and let S be a crystal flex spanning set for \mathcal{C} . If S contains a bounded band-limited flex then $\Omega(\mathcal{C})$ contains the linear structure.*

Proof. Let $u \in S$ be a bounded band-limited flex for \mathcal{C} . If u is finitely supported then, by [15, Theorem 5.6], $\Omega(\mathcal{C}) = \mathbb{T}^d$ and so $\Omega(\mathcal{C})$ contains the linear structure. Suppose u is not finitely supported. By Theorem 4.4(ii) and its proof, given a periodic structure L for \mathcal{C} , the sublattice L_u has rank t , where $1 \leq t \leq d - 1$, and u is factor periodic with respect to L_u . Moreover, u is band-limited with respect to the proper subspace \mathcal{K} of \mathbb{R}^d spanned by L_u .

Let $\mathcal{T}_u = \{T_b \in \mathcal{T} : b \in L_u\}$ be the subgroup of the translation group \mathcal{T} determined by this sublattice. Also, let $\omega_0 = (\omega_1, \dots, \omega_t)$ be the periodicity multifactor. By Lemma 3.8, since u is bounded, $\omega_0 \in \mathbb{T}^t$. Assume that \mathcal{T}_u has generators $T_{g(1)}, \dots, T_{g(t)}$ and choose translations $T_{g(t+1)}, \dots, T_{g(d)}$ such that $T_{g(1)}, \dots, T_{g(d)}$ is a set of generators for a full rank subgroup \mathcal{T}' of \mathcal{T} . Let $\omega_* = (\omega_{t+1}, \dots, \omega_d)$ be an arbitrary point in \mathbb{T}^{d-t} and define

$$w = \sum_{k' \in \mathbb{Z}^{d-t}} \omega_*^{k'} X_{k'} u,$$

where $X_{k'} = k'_1 T_{g(t+1)} + \dots + k'_{d-t} T_{g(d)}$. This velocity field is well defined, since u is band-limited relative to \mathcal{K} and is an infinitesimal flex since $\mathcal{F}(\mathcal{C}; \mathbb{K})$ is invariant under the lattice group action $\pi_{\mathcal{T}}$. Also, w is factor periodic for $\omega = (\omega_0, \omega_*)$ and the periodic structure \mathcal{T}' . It follows that $\Omega(\mathcal{C})$ contains (ω_0, ω_*) for every point ω_* and so the RUM spectrum contains the linear structure of dimension $d - t$. □

Corollary 4.10. *Let \mathcal{C} be a crystal framework in \mathbb{R}^2 whose RUM spectrum in $[0, 2\pi)^2$ is a proper infinite subset which contains no line segments. Then \mathcal{C} does not possess a crystal flex spanning set which includes a bounded band-limited flex.*

This corollary applies in particular to the 2D zeolite framework \mathcal{C}_{oct} whose motif has bars belonging to a regular octagonal ring of equilateral triangles. Indeed the RUM spectrum of \mathcal{C}_{oct} has been shown to be a curve containing no line segments [1, 15].

The same argument in the proof of Theorem 4.9 gives a parallel corollary for the geometric flex spectrum. Indeed, let us say that the geometric flex spectrum $\Gamma(\mathcal{C})$ contains linear structure if there exists a point $\omega = (r_1 e^{i\eta_1}, \dots, r_d e^{i\eta_d})$ in $\Gamma(\mathcal{C})$ such that the intersection of $\Gamma(\mathcal{C})$ with the torus

$$\mathbb{T}_\omega^d := \{(r_1 e^{i\theta_1}, \dots, r_d e^{i\theta_d}) : \theta_1, \dots, \theta_d \in [0, 2\pi)\}$$

contains t -dimensional linear structure in the sense above for \mathbb{T}^d . Then if there is a crystal spanning set for \mathcal{C} which contains a band-limited flex it follows, as in the proof of Theorem 4.9, that $\Gamma(\mathcal{C})$ contains the linear structure.

We can use this observation together with Theorem 4.4 to obtain the following necessary condition for the existence of a crystal flex spanning set or basis.

Theorem 4.11. *Let \mathcal{C} be a crystal framework in \mathbb{R}^d whose geometric flex spectrum is a proper infinite subset which contains no linear structure. Then \mathcal{C} does not possess a crystal flex spanning set.*

Proof. A finite set of geometric flexes u^1, \dots, u^r with distinct periodicity multifactors $\omega^1, \dots, \omega^r$ is linearly independent and so, by the hypotheses, the infinitesimal flex space of \mathcal{C} is infinite dimensional. By Theorem 4.4, any crystal flex spanning set for \mathcal{C} contains a band-limited flex. By the discussion above, it follows that the geometric flex spectrum contains linear structure, a contradiction. \square

Remark 4.12. We recall that early experimental and computational studies of rigid unit modes in silicates typically revealed linear structure in the RUM spectrum for their high-temperature phases. See, for example, Dove et al. [3] where studies of curved RUM surfaces, occurring in tridymite, for example, are also indicated.

It is natural to pose the following questions regarding converse implications to the statements in the results above. If the geometric flex spectrum (respectively RUM spectrum) has linear structure does it follow that there exists a band-limited flex (respectively bounded band-limited flex)? More generally, it would be of interest to obtain sufficient conditions, including linear structure conditions on the geometric flex spectrum, for the existence of a crystal flex basis or an essential crystal flex basis. Such sufficient conditions would explain more fully the nature of the experimental phenomenon in terms of the existence and non-existence of flexes which are band-limited with respect to lines and planes in the crystal structure.

Remark 4.13. We remark that while unbounded flexes do not correspond to physical modes for the bulk crystal their consideration is nevertheless important for the identification of *surface modes* associated with a boundary wall or free surface. Indeed, such modes can be identified with bounded restrictions of unbounded flexes of the bulk crystal. In particular, in three dimensions, a point $\omega = (\omega_1, \omega_2, \omega_3)$ in $\Gamma(\mathcal{C})$ with $|\omega_i| \neq 1$ for a single value of i indicates the existence of a first-order surface mode associated with a hyperplane boundary wall which is normal to the period vector of the bulk crystal

corresponding to i . Thus, a further motivation for the introduction of the geometric flex spectrum and the identification of crystal flex spanning sets lies in their connections with surface modes and isolated Weyl modes (Rocklin et al. [16]), and their analogies with boundary modes for topological insulators (Graf and Porter [6], Lubensky et al. [13]).

5. Examples

We now determine crystal flex spanning sets and bases for a number of elementary crystal frameworks. In these examples, the spanning sets are finitely generated by a small number of band-limited flexes. To quantify this, we introduce the following associated measure of flex complexity for any crystal framework \mathcal{C} .

Definition 5.1. The *flex complexity* $cpx(\mathcal{C})$ is the minimum of the cardinalities of the generating sets for a crystal spanning set for $\mathcal{F}(\mathcal{C}; \mathbb{R})$. Moreover, $cpx(\mathcal{C}) = \infty$ if there is no such generating set.

5.1. The frameworks $\mathcal{C}_{\mathbb{Z}^2}$ and \mathcal{C}_{hex}

Let $\mathcal{C}_{\mathbb{Z}^2}$ be the grid framework in two dimensions, that is, the framework with joints located on the \mathbb{Z}^2 lattice and bars between nearest neighbours. Let $\mathcal{S}_{\text{grid}} = \{u_n, v_n : n \in \mathbb{Z}\}$ be a set of velocity fields with u_n (respectively v_m) supported by the joints on the line $y = n$ (respectively $x = m$) and with unit velocities in the direction of the support line. Then it is elementary to check, by an exhaustion argument in the style of the proofs below, that $\mathcal{S}_{\text{grid}}$ is a crystal basis for the space $\mathcal{F}(\mathcal{C}_{\mathbb{Z}^2}; \mathbb{R})$.

Let \mathcal{C}_{hex} be the honeycomb framework associated with the regular hexagonal tiling of the plane. Note that any hexagon ring subframework, \mathcal{H} say, is the support of a (normalised) local infinitesimal flex, $u_{\mathcal{H}}$ say, which acts as infinitesimal rotation of \mathcal{H} . Let \mathcal{S}_{hex} be a set of identical non-zero flexes of this type for all the honeycomb cells. We claim this is a crystal flex spanning set. We give the argument for this since it is typical of the simple exhaustion argument needed to show that a given set is a free spanning set or a free basis. Note also that $\sum_{\mathcal{H}} u_{\mathcal{H}}$ is the zero infinitesimal flex of \mathcal{C}_{hex} so \mathcal{S}_{hex} is not a free basis. On the other hand, as the proof below shows, it has a curious minimal redundancy property, in the sense that the removal of any flex from \mathcal{S}_{hex} gives a free basis, although not a crystal basis.

From these observations and Proposition 5.2, it follows that $cpx(\mathcal{C}_{\mathbb{Z}^2}) = cpx(\mathcal{C}_{\text{hex}}) = 1$.

Proposition 5.2. \mathcal{S}_{hex} is a crystal flex spanning set for the space $\mathcal{F}(\mathcal{C}_{\text{hex}}; \mathbb{R})$ of real infinitesimal flexes of \mathcal{C}_{hex} .

Proof. Let z be an infinitesimal flex of \mathcal{C}_{hex} . Subtracting a linear combination of the local flexes for the cells A and B indicated in Figure 2, we may assume that the velocity field for z at the origin is 0. Consider the infinite path of framework points to the right of the origin, lying on cells 1, 2, 3, ..., as indicated in Figure 2. We may subtract a multiple of the local flex for cell 1 to 'fix' the first joint, that is, to create a flex with zero velocities at O and at the first joint of the path. We may continue similarly to see that there is an infinite linear combination w of the local flexes for cells 1, 2, 3, ... such that $z - w$ has zero velocity at each joint of the infinite path.




Figure 2. Part of \mathcal{C}_{hex} .

In the same manner, we can subtract an infinite linear combination of the local flexes to the left of cells A and B to obtain a new flex which fixes all the framework points on the two-way infinite horizontal path, π say, through O . This may be achieved without making use of the local flex u_C for the other hexagon incident to the origin. We may assume then that z has zero velocities on the joints on π . The line of joints on $y = 1$ may be now be ‘fixed’ by subtraction of a unique infinite linear combination of the local flexes for the next horizontal line of cells, a, b, c, \dots etc. At this point, the next horizontal line of joints is necessarily fixed by z , in view of the flex condition. Continuing this process with the horizontal hexagonal strips above and below π we see that there is an infinite linear combination of the local flexes which is equal to the original flex. Also, the coefficients of this infinite linear combination are determined uniquely, with the proviso that the local flex u_C is not used in the representation. It follows that \mathcal{S}_{hex} is a crystal flex spanning set. \square

5.2. The 2D kagome framework

We next consider the kagome framework in two dimensions, part of which is indicated in Figure 3. Let a, b, c be the vertices of a triangular subframework of \mathcal{C}_{kag} , with horizontal base edge $[p_a, p_b]$, and let \mathcal{L}_u^0 be the (infinite) linear subframework which contains this edge. Note that there is an evident one-dimensional subspace of infinitesimal flexes of \mathcal{C}_{kag} that are supported on this linear subframework. Consider the non-zero element u_0 in this space which acts on alternate joints of \mathcal{L}_u^0 with unit norm velocity fields, with

$$u_0(p_a) = (\cos \pi/6, -\sin \pi/6), \quad u_0(p_b) = (\cos \pi/6, \sin \pi/6).$$

Let u_n , where $n \in \mathbb{Z}$, be the parallel translates of u , naturally labelled, with u_1 supported by the first linear subframework \mathcal{L}_u^1 above \mathcal{L}_u^0 . Also, let $\{v_n : n \in \mathbb{Z}\}$ (respectively $\{w_n : n \in \mathbb{Z}\}$) be obtained from $\{u_n : n \in \mathbb{Z}\}$ by rotation about the centroid of the triangle abc by $2\pi/3$ (respectively by $4\pi/3$).

The next theorem is due to A. Sait [9] from which it follows that $\text{cpx}(\mathcal{C}_{\text{kag}}) = 1$.

Theorem 5.3. *The set $\mathcal{B}_{\text{kag}} = \{u_n, v_n, w_n : n \in \mathbb{Z}\}$ is a crystal flex basis for $\mathcal{F}(\mathcal{C}_{\text{kag}}; \mathbb{R})$.*

Proof. Since the space group acts on the set it will be sufficient to show that \mathcal{B}_{kag} is a free basis. Write \mathcal{L}_u^n (respectively $\mathcal{L}_v^n, \mathcal{L}_w^n$), for $n \in \mathbb{Z}$, for the supporting linear subframeworks for u_n (respectively v_n, w_n). Let z be an infinitesimal flex of \mathcal{C}_{kag} . By subtracting an appropriate linear combination of the three infinitesimal flexes u_0, v_0, w_0 , we may assume that the three velocity vectors z_a, z_b, z_c for the joints p_a, p_b, p_c of a central

Fig. 2 - B/W online, B/W in print

Fig. 3 - Colour online, B/W in print




Figure 3. The horizontal band-limited flexes u_n for $n \in \mathbb{Z}$.

triangle subframework are zero. Subtracting an appropriate multiple of w_1 we may then arrange $z_d=0$, where d is the next vertex in the direction from a to b . Following this, subtracting an appropriate multiple of v_{-1} , we may arrange $z_e=0$ for the next vertex. Continuing in this way, we obtain an infinite linear combination

$$z' = \sum_{n \in \mathbb{Z}} \beta_n v_n + \gamma_n w_n$$

such that the infinitesimal flex $z'' = z - z'$ imparts only zero velocities to the joints of \mathcal{L}_u^0 . From the flex condition and the rigidity of triangles, we deduce that the flex velocities are also zero on the apex vertices for the triangle subframeworks, such as def , which are horizontal translates of abc . Now subtract an appropriate multiple of u_1 so that the resulting flex is zero on \mathcal{L}_u^1 . Continuing upwards in this manner, and similarly downwards, we obtain an infinite sum representation for z'' in terms of the infinitesimal flexes $u_n, n \in \mathbb{Z}$. Thus, the original flex z is an infinite sum of the basis vectors. Also, the representation is unique and so the proof is complete. \square

Combining this result with Theorem 4.4(iii), we obtain the following description of the space of bounded infinitesimal flexes.

Corollary 5.4. With $\mathcal{B}_{kag} = \{u_n, v_n, w_n : n \in \mathbb{Z}\}$ as above,

$$\mathcal{F}_\infty(\mathcal{C}_{kag}; \mathbb{R}) = \left\{ u \in \mathcal{F}(\mathcal{C}_{kag}; \mathbb{R}) : u = \sum_{n \in \mathbb{Z}} (\alpha_n u_n + \beta_n v_n + \gamma_n w_n) : (\alpha_n), (\beta_n), (\gamma_n) \in \ell^\infty(\mathbb{Z}) \right\}.$$




Figure 4. A motif of 3 joints and 12 bars for the framework \mathcal{C}_{Oct} .

Fig. 4 - B/W online, B/W in print

5.3. The octahedron crystal framework \mathcal{C}_{Oct}

We now consider some crystal frameworks in three dimensions.

Write \mathcal{C}_{Oct} for the crystal framework of corner-connected regular octahedra, in the symmetric placement for which there is one translation class for them. The construction of a crystal flex basis for \mathcal{C}_{Oct} may be determined by viewing \mathcal{C}_{Oct} as the union of countably many copies of the 2D grid framework whose orientations and joint connections are consistent with Figure 4.

To be precise, assume that the period vectors for \mathcal{C}_{Oct} are $(2, 0, 0)$, $(0, 2, 0)$, $(0, 0, 2)$ and let \mathcal{C}_z denote the grid framework in the xy -plane which contains the adjacent joints $p_1 = (0, -1, 0)$, $p_2 = (1, 0, 0)$, $p_3 = (0, 1, 0)$ and $p_4 = (-1, 0, 0)$. These joints lie on a 4-cycle of bars in the xy -plane. Similarly, let \mathcal{C}_x denote the 2D grid framework in the yz -plane which contains the adjacent joints p_1 , $p_5 = (0, 0, -1)$, p_3 and $p_6 = (0, 0, 1)$, and let \mathcal{C}_y denote the 2D grid framework parallel to the xz -plane which contains the adjacent joints p_2 , p_5 , p_4 and p_6 .

Let \mathcal{C}_x^n be the translated bar-joint frameworks $\mathcal{C}_x + (2n, 0, 0)$, for $n \in \mathbb{Z}$, and similarly define \mathcal{C}_y^n and \mathcal{C}_z^n . Then \mathcal{C}_{Oct} is the union of all of these frameworks, that is, it is the bar-joint framework whose joint set is the union of the joints (without multiplicity) and whose set of bars is the union of all the bars.

Let us also define $\mathcal{C}_{\text{Oct}}^+$ as the augmented framework in which each regular octahedron is augmented by three bars parallel to the coordinate axes. In view of the infinitesimal rigidity of a convex octahedron it follows that the vector spaces $\mathcal{F}(\mathcal{C}_{\text{Oct}}; \mathbb{R})$ and $\mathcal{F}(\mathcal{C}_{\text{Oct}}^+; \mathbb{R})$ are equal.

Let us write \mathcal{C}_{sq} for a 2D bar-joint framework composed of corner-connected rigid squares. This is obtained from the 2D grid framework by adding an edge to each alternate square. Let $\mathcal{C}_{\text{sq}}^+$ be the related framework which has both cross diagonals added to the rigid squares. We may thus view $\mathcal{C}_{\text{Oct}}^+$ as the union of copies of $\mathcal{C}_{\text{sq}}^+$, where these copies

are augmentation frameworks, \tilde{C}_x^n , \tilde{C}_y^n and \tilde{C}_z^n say, of the frameworks C_x^n , C_y^n and C_z^n . It follows immediately that the alternation flex a of \tilde{C}_x^n extends to a flex a_n^x of C_{Oct}^+ which has zero velocities on all the other vertices. Let us similarly define the ‘local alternation flexes’ a_n^y and a_n^z for $n \in \mathbb{Z}$.

Let r^x , r^y , r^z be infinitesimal flexes for axial rotations of C_{Oct} about the principal axes of the central octahedron. These infinitesimal flexes act on the entire framework and are unbounded flexes. Also, we assume the normalization such that for $\sigma = x, y$ or z the restrictions of a_n^σ to the octahedron meeting the σ -axis agrees with the restriction of r^σ .

Finally, let \vec{x} be the velocity field in $\mathcal{F}(C_{Oct}; \mathbb{R})$ with joint velocities $(1, 0, 0)$ and let \vec{y} and \vec{z} be analogous velocity fields for the y and z directions.

In the next proof, we use the following elementary flex projection principle. If the bar $[p_a, p_b]$ lies in a plane \mathcal{P} of \mathbb{R}^3 and if the joint velocities v_a, v_b in \mathbb{R}^3 give an infinitesimal flex of the bar $[p_a, p_b]$ then the \mathcal{P} components v_a', v_b' of v_a, v_b also give an infinitesimal flex of the bar. We say that such a flex is an *in-plane* flex when the plane in question is understood.

Theorem 5.5. *The set \mathcal{S} of velocity fields*

$$\{r^x, r^y, r^z\} \cup \{\vec{x}, \vec{y}, \vec{z}\} \cup \{a_n^x, a_n^y, a_n^z : n \in \mathbb{Z}\}$$

is an essential crystal flex basis for $\mathcal{F}(C_{Oct}; \mathbb{R})$.

Proof. The subset $\mathcal{S}_0 \subseteq \mathcal{S}$ of non-rotational flexes has the crystallographic group action property and so it will suffice to show that \mathcal{S} is a free basis for C_{Oct} . Equivalently, we show that \mathcal{S} is a free basis for C_{Oct}^+ .

Let z be an infinitesimal flex in $\mathcal{F}(C_{Oct}^+; \mathbb{R})$. There is a linear combination z_{rig} of $\vec{x}, \vec{y}, \vec{z}, r^x, r^y, r^z$ which agrees with z on the joints p_1, \dots, p_6 . Replacing z by $z - z_{rig}$, for some rigid motion infinitesimal flex z_{rig} , we may assume that these velocities for z are zero.

We now make use of the flex projection principle. Note that the velocity field z_{xy} given by the xy -plane projection of the joint velocities $z(p)$, for joints in \tilde{C}_z^0 , is an infinitesimal flex of \tilde{C}_z^0 . Since z has zero velocity vectors on the central octahedron it follows that the xy -plane projection of z also has zero velocities on the central square of \tilde{C}_z^0 . It follows that this in-plane flex is *equal* to the restriction of a scalar multiple of $a_0^z - r^z$. In this way, we obtain scalar multiples $\alpha_0(a_0^z - r^z)$, $\beta_0(a_0^x - r^x)$ and $\gamma_0(a_0^y - r^y)$ which provide the in-plane flexes of z for the planes $z = 0$, $x = 0$ and $y = 0$.

Consider now the tower subframework given by the tower of octahedra whose connecting joints lie on the z -axis. Since z is zero on the central octahedron supported by p_1, \dots, p_6 , denoted $O_{(0,0,0)}$, it follows that the z component of the joint velocity for a joint on this line is zero. It also follows that there is an infinitesimal flex $\beta_0(a_0^x - r^x) + \gamma_0(a_0^y - r^y)$ with joint velocities agreeing with those of z for the joints on the axial line. It follows similarly that there is a flex of the form

$$w = \alpha(a_0^z - r^z) + \beta_0(a_0^x - r^x) + \gamma_0(a_0^y - r^y)$$

with this agreement property for the three axial lines through $O_{(0,0,0)}$.

Replacing z by $z - w$, we may assume that z is zero on $O_{(0,0,0)}$ and on the joints of the three axial lines of $O_{(0,0,0)}$. Note that the restriction of such a flex z to any other octahedron O with an axis on the coordinate axes must be an infinitesimal rotation flex of the octahedron about this axis. Also each such flex of an individual octahedron O , on the σ -axis say, agrees with the restriction of a scalar multiple of the local alternation flex a_n^σ , for some $n \neq 0$. Evidently, these infinitesimal flexes act on distinct octahedra on the axial lines.

It follows that there is an infinite linear combination of these flexes, w_2 say, whose restriction to any octahedron on a coordinate axis is equal to the restriction of z . Replacing z by $z - w_2$, we may assume that z is zero on this triple tower, \mathcal{T}_{Oct} say. We now observe that any infinitesimal flex which is zero on \mathcal{T}_{Oct} is the zero flex. This follows from the fact that the entire framework may be built up from \mathcal{T}_{Oct} by attaching octahedra in groups of four such that at each stage every flex which is zero on \mathcal{T}_{Oct} is the zero flex. It follows that z must be identically zero. Thus, every velocity field z in $\mathcal{F}(\mathcal{C}_{Oct}; \mathbb{R})$ is an infinite linear combination of the vectors in the set \mathcal{S} .

Note that \mathcal{S} is a countable set of velocity fields which tend to zero strictly and is a free spanning set for a vector space of infinitesimal flexes. Also, the scalar coefficients in the identifications above are determined uniquely by the joint velocities of the flex z . Thus, \mathcal{S} is a free infinitesimal flex basis, as required. \square

Remark 5.6. From a simple adaptation of the above proof, it follows that the space $\mathcal{F}_\infty(\mathcal{C}_{Oct}; \mathbb{R})$ of bounded infinitesimal flexes is the space of infinitesimal flexes of the form

$$u = \alpha \vec{x} + \beta \vec{y} + \gamma \vec{z} + \sum_{n \in \mathbb{Z}} \alpha_n a_n^x + \beta_n a_n^y + \gamma_n a_n^z, \quad (\alpha_n), (\beta_n), (\gamma_n) \in \ell^\infty(\mathbb{Z})$$

The octahedral framework serves to model the rigid unit atomic structure of the crystal perovskite. Its zero mode (or RUM mode) phonon spectrum was among early examples to be computed by experiment and simulation. Other examples were quartz and cristobalite. See Giddy et al. [5]. The RUM spectrum for cristobalite has complete linear structure in $[0, 2\pi]^3$, being the union of the three line segments $(t, 0, 0)$, $(0, t, 0)$, $(0, 0, t)$, for $0 \leq t < 2\pi$. The existence of this linear structure in the RUM spectrum also follows from the band-limited flexes in \mathcal{S} .

5.4. The 3D kagome framework

The kagome framework in three dimensions, \mathcal{C}_{Kag} , has the structure of the kagome net, a network of regular tetrahedral units connected pairwise at their vertices. It may be constructed from the 2D kagome net by first completing its triangles to tetrahedra with alternating up and down orientations to create a horizontal layer framework, and then joining translational copies of such layers. The period vectors determine a parallelepiped unit which is occupied by a single tetrahedron and one can determine a motif for \mathcal{C}_{Kag} with a corresponding 12-by-12 matrix-valued symbol function on the 3-torus [14]. However, the simple layer structure description, together with the symmetry of \mathcal{C}_{Kag} , are sufficient to determine a crystal basis by an exhaustion argument, as before, as we now sketch.

For $m \in \mathbb{Z}$, let $\{u_n^m, v_n^m, w_n^m : n \in \mathbb{Z}\}$ be the crystal flex bases for the in-plane infinitesimal flex space of the m^{th} layer and note that the infinitesimal flexes in these sets extend

Fig. 5 - B/W online, B/W in print




Figure 5. A motif of 2 joints and 9 bars for the bipyramid framework.

(by zero velocity specifications) to infinitesimal flexes of \mathcal{C}_{Kag} . If w is an arbitrary infinitesimal flex of \mathcal{C}_{Kag} , we may subtract an appropriate infinite linear combination of these flexes to replace w by a flex w_1 with the property that the velocity vectors at each joint in the horizontal layers are in the vertical direction. Let us make a distinction between the ‘thick’ horizontal layer frameworks and their ‘thin’ kagome subframeworks (which support the added tetrahedra). For layer $m = 0$, fix a triangle subframework in the thin layer to be a base triangle. Its tetrahedron has three additional bars. Each of these bars indicates the direction of a linearly localized flex, each of which is a space group element image of u_0^0 . We may choose a linear combination of these flexes and subtract them from w_1 to obtain a flex w_2 so that the tetrahedron has zero velocities (under w_2) at all four of its joints.

Continuing in this way we see that, as in the 2D kagome, there is a crystal flex basis with a single generator and it consists of the space group element images of u_0^0 . In particular, $\text{cpx}(\mathcal{C}_{\text{Kag}}) = 1$.

5.5. The bipyramid framework $\mathcal{C}_{\text{Bipyrr}}$

We next define the bipyramid crystal framework $\mathcal{C}_{\text{Bipyrr}}$ and compute the transfer function associated with the natural primitive periodic structure. An appropriate basis of period vectors takes the form

$$(1, 0, 0), \quad (1/2, \sqrt{3}/2, 0), \quad (0, 0, 2h)$$

and we consider a motif (F_v, F_e) where F_v consists of the two joints $p_1 = (0, 0, 0)$, $p_2 = (1/2, \alpha, -h)$ and F_e consists of the nine bars $p_i p_j$ associated with the nine directed bars indicated in Figure 5. Here $h = \sqrt{2/3}$ is the height of a regular tetrahedron of unit sidelength and $\alpha = \sqrt{3}/6$ is the distance from B to C , that is, the distance of the centroid of a unit sidelength equilateral triangle to the triangle boundary.

The following data for the motif edges will suffice for the computation of the associated transfer function $\Psi_{\text{Bipyrr}}(z)$.

A factor-periodic infinitesimal flex, with factor $\omega = (\omega_1, \omega_2, \omega_3) \in \mathbb{C}_*^3$, has the form

$$u = (u_{\kappa, k}) = (\omega^k a) = (\omega_1^{k_1} \omega_2^{k_2} \omega_3^{k_3} a), \quad k \in \mathbb{Z}^3,$$

i	$p(e_i)$	$k = (k_1, k_2, k_3)$	$l = (l_1, l_2, l_3)$
1	$(-1, 0, 0)$	$(0, 0, 0)$	$(1, 0, 0)$
2	$(1/2, -\sqrt{3}/2, 0)$	$(1, 0, 0)$	$(0, 1, 0)$
3	$(-1/2, -\sqrt{3}/2, 0)$	$(0, 0, 0)$	$(0, 1, 0)$
4	$(-1/2, -\alpha, h)$	$(0, 0, 0)$	$(0, 0, 0)$
5	$(1/2, -\alpha, h)$	$(1, 0, 0)$	$(0, 0, 0)$
6	$(0, 2\alpha, h)$	$(0, 1, 0)$	$(0, 0, 0)$
7	$(-1/2, -\alpha, -h)$	$(0, 0, 0)$	$(0, 0, 1)$
8	$(1/2, -\alpha, -h)$	$(1, 0, 0)$	$(0, 0, 1)$
9	$(0, 2\alpha, -h)$	$(0, 1, 0)$	$(0, 0, 1)$

where $a = (u_{1,x}, u_{1,y}, u_{1,z}, u_{2,x}, u_{2,y}, u_{2,z}) \in \mathbb{R}^6$ is the velocity field for the motif joints, being the list of the coordinate velocities of u at the joints p_1 and p_2 . We have,

$$\Psi_{\text{Bipyr}}(z^{-1}) = \begin{matrix} & \begin{matrix} v_{1,x} & v_{1,y} & v_{1,z} \end{matrix} \\ \begin{matrix} e_1 \\ e_2 \\ e_3 \\ e_4 \\ e_5 \\ e_6 \\ e_7 \\ e_8 \\ e_9 \end{matrix} & \left[\begin{array}{ccc|ccc} - (1 - z_1) & 0 & 0 & 0 & 0 & 0 \\ (z_1 - z_2)/2 & -(z_1 - z_2)\sqrt{3}/2 & 0 & 0 & 0 & 0 \\ (1 - z_2)/2 & -(1 - z_2)\sqrt{3}/2 & 0 & 0 & 0 & 0 \\ -1/2 & -\alpha & h & 1/2 & \alpha & -h \\ z_1/2 & -\alpha z_1 & z_1 h & -1/2 & \alpha & -h \\ 0 & 2z_2\alpha & z_2 h & 0 & -2\alpha & -h \\ -1/2 & -\alpha & -h & z_3/2 & z_3\alpha & z_3 h \\ z_1/2 & -z_1\alpha & -z_1 h & -z_3/2 & z_3\alpha & z_3 h \\ 0 & 2z_2\alpha & -z_2 h & 0 & -2z_3\alpha & z_3 h \end{array} \right] \end{matrix}$$

The submatrix of $\Psi_{\text{Bipyr}}(z^{-1})$ for columns 3, 4, 5 is a 9×3 function matrix. We note that the first three rows are zero. Indeed, the three edges $e_1 = (v_1, (0, 0, 0))(v_1, (1, 0, 0))$, $e_2 = (v_1, (1, 0, 0))(v_1, (0, 1, 0))$ and $e_3 = (v_1, (0, 0, 0))(v_1, (0, 0, 1))$ are of $v = w$ type and so the entries for columns 4, 5, 6 are zero. Also, the entries for column 3 are zero since the z -component is zero for the vectors $p(e_i)$, for $i = 1, 2, 3$.

We next determine the set of factors $\omega \in \mathbb{C}_*^3$ for which $\Psi_{\text{Bipyr}}(\omega^{-1})a = 0$ for some non-zero vector of the form $a = (0, 0, u_{1,z}, u_{2,x}, u_{2,y}, 0)$. In doing so, we shall determine the factor periodic infinitesimal flexes u with the property that their velocity fields impart only vertical velocities to the joints that lie in the horizontal copies of the fully triangulated framework \mathcal{C}_{tri} and impart only horizontal velocities to the other joints, the polar joints of the constituent bipyramids. In view of the latter condition, we refer to these infinitesimal flexes simply as *sheering flexes*. The required solutions for ω are the values

of $z = (z_1, z_2, z_3)$ for which the submatrix $\Psi_{\text{sub}}(z^{-1})$ has non-zero kernel, where

$$\Psi_{\text{sub}}(z^{-1}) = \begin{matrix} & \begin{matrix} v_{1,z} \\ v_{2,x} \\ v_{2,y} \end{matrix} \\ \begin{matrix} e_4 \\ e_5 \\ e_6 \\ e_7 \\ e_8 \\ e_9 \end{matrix} & \left[\begin{array}{c|cc} h & 1/2 & \alpha \\ z_1 h & -1/2 & \alpha \\ z_2 h & 0 & -2\alpha \\ -h & z_3/2 & z_3\alpha \\ -z_1 h & -z_3/2 & z_3\alpha \\ -z_2 h & 0 & -2z_3\alpha \end{array} \right] \end{matrix}$$

Noting the similarity between the rows for e_6 and e_9 , it follows that if z_3 is not equal to -1 then the kernel is trivial. On the other hand if $z_3 = -1$ then the rows for e_7, e_8, e_9 are the negatives of the rows for e_4, e_5, e_6 . We conclude that $(\omega_1, \omega_2, \omega_3)$ is a periodicity multifactor for a sheering flex u if and only if $\omega_3 = -1$ and the determinant of the 3×3 submatrix for the first 3 rows is zero at ω_1, ω_2 . This determinant is $\alpha h(1 + z_1 + z_2)$. We conclude that when z_1 and z_2 are unimodular then there are exactly two solutions and so the RUM spectrum $\Omega(\mathcal{C}_{\text{BipyR}})$ contains the set

$$\{(1, 1, 1), (e^{2\pi i/3}, e^{4\pi i/3}, e^{\pi i}), (e^{4\pi i/3}, e^{2\pi i/3}, e^{\pi i})\}.$$

We also note that there are unbounded geometric flexes associated with the solutions (ω_1, ω_2) of the equation of $1 + z_1 + z_2 = 0$, where $\omega_1 = r, \omega_2 = -(1 + r)$, and $0 < r < 1$. These infinitesimal flexes bear some analogy with the unbounded flex of the kite framework considered in [Example 2.11](#). Since there are uncountably many linearly independent unbounded flexes of this type the infinitesimal flex space $\mathcal{F}(\mathcal{C}_{\text{BipyR}}; \mathbb{R})$ is infinite dimensional.

5.6. Two crystal extensions of $\mathcal{C}_{\text{BipyR}}$

Consider the crystal framework $\mathcal{C}_{\text{BipyR}}^e$ which is obtained from $\mathcal{C}_{\text{BipyR}}$ by adding bonds of length 1 between polar joints whenever this is possible. It is straightforward to see that $\mathcal{C}_{\text{BipyR}}^e$ is sequentially infinitesimally rigid [[12, 14](#)] and so is infinitesimally rigid in the strongest possible sense. Note that every 2D subframework parallel to the xy -plane is a copy of the fully triangulated framework \mathcal{C}_{tri} . Let $\mathcal{C}_{\text{BipyR}}^+$ be obtained from $\mathcal{C}_{\text{BipyR}}$ by adding edges and vertices so that every triangle in the horizontal copies of \mathcal{C}_{tri} in $\mathcal{C}_{\text{BipyR}}$ is the equator of a bipyramid. This framework may be described as having horizontal layers of maximally packed bipyramids. The joints are once again of two types, with polar joints having degree 6, as before, and equatorial joints having degree 18. One can observe that in fact the two sheering RUMs of $\mathcal{C}_{\text{BipyR}}$ extend to this framework and so, despite the edge rich structure, $\mathcal{C}_{\text{BipyR}}^+$ is not infinitesimally rigid.

Added in proof. The geometric spectrum has been used by the third author and E. Kastis [[10, 11](#)] to obtain necessary and sufficient conditions for the infinitesimal rigidity of a crystallographic bar-joint framework.

Acknowledgements. This work was supported by the Engineering and Physical Sciences Research Council [grant number EP/P01108X/1].

References

1. G. BADRI, D. KITSON AND S. C. POWER, The almost periodic rigidity of crystallographic bar-joint frameworks, *Symmetry* **6** (2014), 308–328.
2. R. CONNELLY, W. A. IVIĆ AND W. WHITELEY, *Rigidity and symmetry*, Fields Institute Communications, Volume **70** (Springer, 2014).
3. M. T. DOVE, V. HEINE AND K. D. HAMMONDS, Rigid unit modes in framework silicates, *Mineral. Mag.* **59** (1995), 629–639.
4. M. T. DOVE, A. K. A. PRYDE, V. HEINE AND K. D. HAMMONDS, Exotic distributions of rigid unit modes in the reciprocal spaces of framework aluminosilicates, *J. Phys. Condens. Matter* **19** (2007), 275209. doi:10.1088/0953-8984/19/27/275209
5. A. P. GIDDY, M. T. DOVE, G. S. PAWLEY AND V. HEINE, The determination of rigid unit modes as potential soft modes for displacive phase transitions in framework crystal structures, *Acta Crystallogr.* **A49** (1993), 697–703.
6. G. M. GRAF AND M. PORTA, Bulk-edge correspondence for two-dimensional topological insulators, *Comm. Math. Phys.* **324** (2013), 851–895.
7. S. D. GUEST, P. W. FOWLER AND S. C. POWER, (Editors), Rigidity of periodic and symmetric structures in nature and engineering, *Phil. Trans. R. Soc. A* **372** (2014), 0358.
8. R. G. HUTCHINSON AND N. A. FLECK, The structural performance of the periodic truss, *J. Mech. Phys. Solids* **54** (2006), 756–782.
9. A. KASIM SAIT, Rigidity of Infinite Frameworks, MPhil thesis, Lancaster University (2011).
10. E. KASTIS AND S. C. POWER, Algebraic spectral synthesis and crystal rigidity, *J. Pure Appl. Algebra* **223** (2019), 4954–4965.
11. E. KASTIS AND S. C. POWER, The first-order flexibility of a crystal framework, *J. Math. Anal. Appl.* **504** (2021), 125404. doi:10.1016/j.jmaa.2021.125404
12. D. KITSON AND S. C. POWER, The rigidity of infinite graphs, *Discrete Comput. Geom.* **60** (2018), 531–557.
13. T. C. LUBENSKY, C. L. KANE, X. MAO, A. SOUSLOV AND K. SUN, Phonons and elasticity in critically coordinated lattices, *Rep. Progress Phys.* **78**(7) (2015), 073901.
14. J. C. OWEN AND S. C. POWER, Infinite bar-joint frameworks, crystals and operator theory, *New York J. Math.* **17** (2011), 445–490.
15. S. C. POWER, Polynomials for crystal frameworks and the rigid unit mode spectrum, *Phil. Trans. R. Soc. Lond. Ser. A Math. Phys. Eng. Sci.* **372** (2014), 20120030.
16. D. Z. ROCKLIN, B. GIN-GE CHEN, M. FALK, V. VITELLI AND T. C. LUBENSKY, Mechanical Weyl modes in topological Maxwell lattices, *Phys. Rev. Lett.* **116** (2016), 135503.
17. F. WEGNER, Rigid-unit modes in tetrahedral crystals, *J. Phys. Condens. Matter* **19** (2007), 406218. doi:10.1088/0953-8984/19/40/406218