


Regular Article

Exposure to early childhood maltreatment and its effect over time on social cognition

Katherine M. Crawford¹, Karmel Choi^{1,2,3}, Kathryn A. Davis¹, Yiwen Zhu¹, Thomas W. Soare^{1,2}, Andrew D.A.C. Smith⁴, Laura Germine^{3,5} and Erin C. Dunn^{1,2,5} 

¹Center for Genomic Medicine, Massachusetts General Hospital, Boston, MA, USA; ²Stanley Center for Psychiatric Research, The Broad Institute of Harvard and MIT, Cambridge, MA, USA; ³Institute for Technology in Psychiatry, McLean Hospital, Belmont, MA, USA; ⁴Applied Statistics Group, University of the West of England, Bristol, UK and ⁵Department of Psychiatry, Harvard Medical School, Boston, MA, USA

Abstract

Social cognitive deficits can have many negative consequences, spanning social withdrawal to psychopathology. Prior work has shown that child maltreatment may associate with poorer social cognitive skills in later life. However, no studies have examined this association from early childhood into adolescence. Using data from the Avon Longitudinal Study of Parents and Children (ALSPAC; $n = 4,438$), we examined the association between maltreatment (caregiver physical or emotional abuse; sexual or physical abuse), assessed repeatedly (every 1–3 years) from birth to age 9, and social cognitive skills at ages 7.5, 10.5, and 14 years. We evaluated the role of both the developmental timing (defined by age at exposure) and accumulation of maltreatment (defined as the number of occasions exposed) using a least angle regression variable selection procedure, followed by structural equation modeling. Among females, accumulation of maltreatment explained the most variation in social cognitive skills. For males, no significant associations were found. These findings underscore the importance of early intervention to minimize the accumulation of maltreatment and showcase the importance of prospective studies to understand the development of social cognition over time.

Keywords: adversity, ALSPAC, sensitive periods, social cognition, structural equation modeling

Introduction

Epidemiological studies suggest that approximately one out of every six young people in the United States and other developed countries worldwide has experienced some type of childhood maltreatment, such as physical abuse, sexual abuse, emotional abuse, or neglect (Finkelhor, Turner, Shattuck, & Hamby, 2013; Gilbert et al., 2009; Stoltenborgh et al., 2015). The ubiquity of maltreatment exposure is concerning due to the wide ranging negative health outcomes linked to childhood maltreatment, including depression, posttraumatic stress disorder (PTSD), other psychiatric problems, as well as brain-based structural and functional changes (McLaughlin et al., 2010; Teicher et al., 2003).

Recent evidence also suggests that exposure to childhood maltreatment may lead to difficulties in social cognition abilities. Social cognition refers to the information processing mechanisms underlying the perception, interpretation, and response to social information that drive social interactions (Crick & Dodge, 1994; Green, Olivier, Crawley, Penn, & Silverstein, 2005; Piskulic & Addington, 2011; Sergi, Rassovsky, Nuechterlein, & Green,

2006; Vauth, Rüscher, Wirtz, & Corrigan, 2004). It is often conceptualized as the psychological processes that allow individuals to benefit from being part of a social group, such as human society (Frith, 2007). Such processes include skills to recognize other's emotions and infer their mental states via body language and other social cues. Social cognitive deficits, or difficulties in interpreting social cues and situations, can have many negative consequences. For example, prior studies have linked deficits in social cognition to social misperceptions and even social withdrawal (Green et al., 2005; Piskulic & Addington, 2011), as well as poor vocational outcomes stemming from deficits in work-related social skills (Vauth et al., 2004). Of consequence, children who have experienced abuse or maltreatment have been found to have a wide range of social cognition deficits as compared to their nonmaltreated peers, including difficulties comprehending complex social situations, understanding emotions, identifying facial emotions, and imitating multiple roles in social interactions (Barahal, Waterman, & Martin, 1981; Luke & Banerjee, 2013). For example, children exposed to institutional caregiving environments were found to have worse social cognition in all domains, but particularly reciprocal social interactions (Levin et al., 2015). Many theorize that social cognitive deficits can occur following exposure to maltreatment, due to maltreatment being linked with insecure attachments, lowered threshold for limbic system reactivity, and other neurobiological changes (Dvir, Ford, Hill, & Frazier, 2014).

Social cognition may play also an important role in mental illness. To date, deficits in social cognition abilities have been linked

Author for Correspondence: Erin C. Dunn, Psychiatric and Neurodevelopmental Genetics Unit, Center for Genomic Medicine, Massachusetts General Hospital, 185 Cambridge Street, Simches Research Building 6th Floor, Boston, MA 02114; E-mail: edunn2@mg.harvard.edu.

Cite this article: Crawford KM, Choi K, Davis KA, Zhu Y, Soare TW, Smith ADAC, Germine L, Dunn EC (2022). Exposure to early childhood maltreatment and its effect over time on social cognition. *Development and Psychopathology* 34, 409–419. <https://doi.org/10.1017/S095457942000139X>

to the etiology, course, and treatment of a wide range of psychiatric disorders, including schizophrenia, autism, posttraumatic stress disorder, depression, and early-onset conduct problems (Ladegaard, Larsen, Videbech, & Lysaker, 2014; Oliver, Barker, Mandy, Skuse, & Maughan, 2011; Piskulic & Addington, 2011; Sasson, Nowlin, & Pinkham, 2013; Sergi *et al.*, 2006). For example, prior research has shown that among people with schizophrenia, social cognition abilities can predict functional outcome (i.e., ability to work independently and social problem solving) and, when considered with broader cognitive skills (i.e., verbal memory and attention), can explain nearly 80% of the variance in social functioning (Addington, Girard, Christensen, & Addington, 2010; Sergi *et al.*, 2006). In addition, mentalizing impairment – a subset of social cognition defined as a person's capacity to understand one's own and others' behavior in regards to a mental state – has been shown to partially mediate or explain the association between childhood abuse and negative symptoms in nonaffective psychotic disorders (Weijers *et al.*, 2018). As social cognition is widely implicated in psychiatric disorders, research into the features that predict social cognition problems, including childhood maltreatment, are needed.

However, research on the relationship between childhood maltreatment and social cognition has been limited in three important ways. First, studies of maltreated children have often assessed social cognition deficits in adulthood – and not throughout childhood and adolescence, when social cognitive skills are developing (Happé & Frith, 2014). In particular, few studies have examined social cognition during middle childhood, when important gains in social perspective taking are made (Bosacki, 2000; Van Der Graaff *et al.*, 2014). This is a shortcoming as social cognition is not a fixed state during early life, but rather the result of an ongoing neurodevelopmental process that continues throughout childhood and adolescence. The capacity to mentalize – or understand the mental state of others – develops in the first 5 years of life (Frith & Frith, 2007). By age 4, most children have developed the understanding that others may hold beliefs that are different from their own and that other's beliefs can be untrue (Barresi & Moore, 1996; Kilford, Garrett, & Blakemore, 2016). During adolescence, brain structures important for social cognition, including grey matter density in the superior temporal lobe, undergo rapid development and change (Burnett, Sebastian, Cohen Kadosh, & Blakemore, 2011; Mills, Lalonde, Clasen, Giedd, & Blakemore, 2014; Vetter, Leipold, Kliegel, Phillips, & Altgassen, 2013). Previous work has indicated that social cognition may be highly plastic and malleable depending on the social network and social experiences of the child across time and development (Cabrera, Fagan, Wight, & Schadler, 2011; Ford, Clark, & Stansfeld, 2011; Happé & Frith, 2014). Although there are a number of longitudinal studies of social cognition, few of these start with children; of these, none to our knowledge examine repeated measures of social cognition throughout childhood (Magiati, Tay, & Howlin, 2014). These shortcomings exist despite evidence suggesting that the foundations of social cognitive skills are gained in very early life (Happé & Frith, 2014; Mills *et al.*, 2014). To address these gaps, prospective and longitudinal studies are needed that include repeated, standardized measures to capture the development of social cognition beginning in infancy throughout childhood and adolescence.

Second, although childhood maltreatment experiences have been linked to social cognition deficits, including emotional regulation problems (Dvir *et al.*, 2014), there is limited knowledge of

how certain features of these maltreatment experiences predict social cognition. For example, little is known whether the developmental timing of occurrence and the number of times a child is exposed to maltreatment shapes their social cognition, or whether being exposed to maltreatment, regardless of these timing aspects, is more important. Thus, efforts are needed to statistically evaluate hypotheses described by life course theory (Ben-Shlomo & Kuh, 2002; Kuh, Ben-Shlomo, Lynch, Hallqvist, & Power, 2003), including those focused on sensitive periods, accumulation, ever-exposed models, respectively. Studies on a host of other psychosocial outcomes, such as depression risk, psychosis risk, lower ego resiliency, and neurocognitive functioning (Barahal *et al.*, 1981; Dale *et al.*, 2010; Dunn *et al.*, 2018; Luke & Banerjee, 2013; Manly, Kim, Rogosch, & Cicchetti, 2001) have found that the effects of childhood maltreatment, and of adversity more broadly, can vary considerably depending on the timing and accumulation of abuse exposures. There is evidence to suggest that there may be sensitive periods when the developing brain is particularly susceptible to these adverse social experiences (Dunn, McLaughlin, Slopen, Rosand, & Smoller, 2013; Dunn, Nishimi, Powers, & Bradley, 2017). Further, prior work has also demonstrated the importance of exposure timing in predicting neurophysiology, neurobiological structure, and broader cognitive functioning (Bosch *et al.*, 2012; Cicchetti, Cowell, Rogosch, & Toth, 2015). Given this growing body of evidence to support time-dependent effects of maltreatment, there is a need for research that accounts for not only changes in social cognition over time, but also the temporal features of maltreatment exposure across childhood. Such investigations will allow researchers to evaluate the effects of different theories and determine which theoretical models alone or in combination best explain child development outcomes.

Finally, there is a distinct lack of population-based social cognition research, with the majority of prior studies being conducted in clinical cohorts and other highly selected or homogeneous samples (e.g., college undergraduates). While social cognition has been an excellent predictor of functional outcomes (e.g., vocational functioning) in previous studies using clinical samples (e.g., people diagnosed with schizophrenia), these findings have not been generalized to typically developing populations. Efforts to expand work on the links between maltreatment exposure and social cognition to more representative samples can improve reproducibility and generate results that are more widely generalizable. Moreover, a focus on triangulation, where multiple approaches are used to address the same research question – including studying different population groups – can yield results that are less likely to be driven by methodological artifacts and are closer to the underlying truth (Munafò & Davey Smith, 2018).

The current study aimed to address these gaps and expand upon prior literature by prospectively examining the relationship between exposure to childhood maltreatment and social cognition deficits. Data came from a large, population-based sample of children followed from infancy to adolescence called the Avon Longitudinal Study of Parents and Children (ALSPAC). Our aims were to: (a) determine how exposure to different types of maltreatment between birth and age 9 years was associated with the development of social cognition skill deficits from ages 7.5 through 14; and (b) evaluate the effects of the different life course models in relation to these later social cognition skill deficits. Specifically, the three life course models tested were: (a) a sensitive period model (Ben-Shlomo & Kuh, 2002), in which the effect of maltreatment depends on the developmental time period of the

exposure; (b) an accumulation model (Evans, Li, & Whipple, 2013), in which the effect of maltreatment increases with the number of occasions exposed, regardless of timing; and (c) an ever-exposed model, in which the effect of maltreatment does not depend on the timing or amount of exposure, but rather its presence or absence.

Method

Sample and procedures

ALSPAC is a prospective, longitudinal birth-cohort of children born to pregnant mothers living in the county of Avon England, which is 120 miles west of London (Boyd et al., 2013; Fraser et al., 2013). ALSPAC was designed to generate new knowledge on the pathways to health across the life span, with a focus on genetic and environmental determinants. Women residing in Avon, UK who gave birth between 1 April 1991 to 31 December 1992 were recruited through media advertisements and visits by research staff at multiple sites within the community. The study was also advertised at routine antenatal and maternity health services appointments. Approximately 85% of eligible pregnant women agreed to participate ($N = 14,541$), and 76% of eligible live births ($N = 14,062$) who were alive at 12 months of age ($N = 13,988$ children) were enrolled. An additional 913 children who would have been eligible, but whose mothers did not choose to participate during pregnancy, were enrolled after age 7 years. Response rates to data collection have been good (75% have completed at least one follow-up). Compared to the general population in the UK based on the 1991 Census, mothers enrolled in the ALSPAC were more likely to be White, married, and home owners. However, despite having generally higher socioeconomic positions than non-ALSPAC enrolled women, ALSPAC-enrolled women were more likely to live in overcrowded conditions (Fraser et al., 2013). Ethical approval for the study was obtained from the ALSPAC Ethics and Law Committee and the Local Research Ethics Committee. The ALSPAC website contains details of all the data that are available through a fully searchable data dictionary and variable search tool: <http://www.bristol.ac.uk/alspac/researchers/our-data/>.

The current analysis was based on an analytic sample of 4,438 children (out of a possible 9,677 children with one measure of social cognition) who met three inclusion criteria. First, we restricted our analytic sample to singleton births to prevent confounding associated with the unique social structure and support that multiple-birth children can have in the home (Lang, Cox, & Flores, 2013; Prino et al., 2016). Second, the measurement of social cognition came from mailed questionnaires, and we further restricted the sample to only those children whose caregivers had completed these questionnaires for all of the first three timepoints (ages 7.5, 10.5, and 14 years) when social cognition was measured. Finally, to ensure a consistent reporter of child social cognition across time, we restricted the sample to only those children who had mothers and maternal figures as the sole reporters of their social cognition skills over the three timepoints of assessment. As expected, given attrition patterns in the ALSPAC, children in the included sample ($n = 4,438$) were slightly more socio-demographically advantaged as compared to children in the excluded sample who had at least one social cognition measure ($n = 5,239$) (Supplemental Table 1). In addition, we compared the distribution of covariates and outcome scores between the analytic sample ($n = 4,438$) and the subset of excluded

participants with complete social cognition outcome data at all three time points, but who lacked consistent maternal reports ($n = 688$). These two samples had largely similar sociodemographic characteristics and social cognition scores at all three time points; however, the excluded sample (without consistent maternal reports) were more likely to be born to mothers with slightly higher education (Supplemental Table 2).

Measures

Exposure to child maltreatment

We examined two types of child maltreatment, measured using mailed questionnaires. Each maltreatment type was measured on seven occasions before age 10; the time frame covered by each assessment varied, with an average duration of 19 months (Table 1 and Supplemental Table 3).

Caregiver physical or emotional abuse. Children were coded as having been exposed to physical or emotional abuse if the mother, partner, or both responded affirmatively to any of the following items: (a) "Your partner was physically cruel to your children"; (b) "You were physically cruel to your children"; (c) "Your partner was emotionally cruel to your children"; (d) "You were emotionally cruel to your children". Physical abuse and emotional abuse items were examined together, rather than separately, given their moderately strong correlation overall (average correlation = 0.55) and to ensure an adequate sample size for these analyses given the rarity of abuse exposure in this population-based sample (Supplemental Table 4).

Sexual or physical abuse. Exposure to sexual or physical abuse was determined through an inventory asking the mother to indicate whether or not the child had been either "sexually abused" or "physically hurt by someone." If the mother indicated the child had been exposed to either physical or sexual abuse, the child was coded as having experienced abuse. Owing to low prevalence of sexual abuse in the sample, under 10 cases per timepoint, sexual and physical abuse were considered in the same measure; the tetrachoric correlation between any exposure to physical abuse before age 10 and any exposure to sexual abuse before age 10 was moderate ($r_{\text{tetrachoric}} = .39$).

Variable encoding. For each type of maltreatment, we generated three sets of encoded variables: (a) a single variable denoting the total number of time periods of exposure to each type of maltreatment, to test the accumulation hypothesis (coded as 0–6); (b) a set of variables indicating presence versus absence of each type of maltreatment at a specific developmental stage, to test the sensitive period hypothesis; and (c) a single variable denoting whether the child had ever experienced exposure to a specific type of maltreatment regardless of timing or number of exposures, to test the exposure hypothesis.

Social cognition

Social cognition skills were measured using the Social Communication Disorder Checklist at ages 7.5, 10.5, and 14 years. This 12-item measure was designed to capture the main features of individual social cognition ability, as reported by caregivers, and was originally developed and created for children with Turner's syndrome, which is a disorder characterized by social adjustment problems (Skuse et al., 1997). Sample items included "Child did not realize when others were upset/angry," "Child was

Table 1. Exposure to childhood maltreatment in the analytic sample, stratified by type, sex, and age at exposure ($n=4,438$)

	Sexual or physical abuse (by anyone)				Physical or emotional abuse			
	Female		Male		Female		Male	
	<i>N</i>	(%)	<i>N</i>	(%)	<i>N</i>	(%)	<i>N</i>	(%)
Unexposed	1,964	89.03	1,884	84.41	1,812	82.14	1,852	82.97
Exposed	242	10.97	348	15.59	394	17.86	380	17.03
Age at exposure								
Infancy								
Age 8 mo.	---	---	---	---	82	3.81	70	3.24
Age 1.5/1.75	48	2.22	52	2.39	89	4.18	72	3.36
Age 2.5/2.75	59	2.81	83	3.88	110	5.22	123	5.79
Preschool								
Age 3.5	51	2.39	71	3.29	---	---	---	---
Age 4/4.75	48	2.27	84	3.91	115	5.42	85	3.95
Age 5/5.75	40	1.92	63	2.97	138	6.60	136	6.40
Middle childhood								
Age 6/6.75	40	1.89	74	3.47	123	5.89	93	4.36
Age 8/9	42	2.00	95	4.45	74	3.51	84	3.95

Percentages for each age represent proportions of children exposed out of the total analytic sample. --- indicates that the variable was not assessed at the corresponding time point.

very demanding of other people's time," and "Child did not pick up on body language." Response options to each item on this questionnaire were: 1 = not true, 2 = quite true or sometimes true, and 3 = very or often true and were asked in relation to the child's behavior over the past 6 months. These responses were summed to create a total score, with higher scores indicating more social cognition difficulties. The Social Communication Disorder Checklist has demonstrated excellent internal consistency reliability overall $\alpha = 0.98$ (Skuse et al., 1997), and in our analytic sample (age 7.5 $\alpha = 0.79$; age 10.5 $\alpha = 0.73$; age 14 $\alpha = 0.80$).

Covariates

All multivariable analyses controlled for the following covariates, measured at the time of the child's birth: child race/ethnicity; maternal marital status; highest level of maternal education; maternal age; homeownership status; parent social class; number of previous pregnancies; and levels of maternal depression, as assessed by the Edinburgh Postnatal Depression Scale (EPDS) (Cox, Holden, & Sagovsky, 1987) (see Supplemental Materials). Covariates were selected for inclusion because they were found to be potential confounders in our sample (see Supplemental Table 1), or because they have been included routinely in longitudinal birth-cohorts when studying child outcomes (Hibbeln et al., 2007; Oliver et al., 2011).

Analyses

We began by running univariate and bivariate analyses to examine the distribution of covariates and exposure to maltreatment in the total analytic sample. We then used a two-stage structured life course modeling approach (SLCMA) (Dunn et al., 2018; Smith et al., 2015, 2016) to evaluate the extent to which the temporal

characteristics of maltreatment were associated with deficits in social cognition. For these analyses, we tested – for each type of maltreatment – three life course theoretical models to determine which one explained the most outcome variability (i.e., r^2) (Ben-Shlomo & Kuh, 2002). The major advantage of the SLCMA relative to other methods, including standard multiple regression, is that it provides an unbiased way to compare multiple competing theoretical models simultaneously and identify the most parsimonious explanation for the observed outcome variation (see Supplemental Materials and Supplemental Figure 1 for more details). These analyses were performed so that each life course theoretical model was tested for both types of maltreatment at each time point when social cognition was measured (age 7.5, 10.5, and 14).

Given that the final maltreatment measurement time point (at age 8/9) occurred after the first social cognition measurement time point (at age 7.5), the theoretical models tested to explain social cognition at age 7.5 did not include maltreatment exposure at age 9 as a predictor. Maltreatment exposure at age 9 was included in the analyses examining social cognition at 10.5 years and 14 years, allowing us to examine the links between maltreatment (between ages 0–9) on social cognition over almost a decade-long period (Figure 1).

We conducted all analyses using a multiply imputed dataset to reduce potential bias and minimize loss of power due to attrition (see Supplemental Materials). All analyses were stratified by sex, given previous literature documenting sex differences in exposure to childhood maltreatment (Briere & Elliott, 2003), and social cognition development (Gur et al., 2012).

After selecting the life course theoretical models in the first stage of the SLCMA that explained the most outcome variability, we then performed a linear regression of the theoretical model chosen in the second stage of the SLCMA within each of the 20




Figure 1. The time periods of exposure and outcome measurement.

multiply imputed datasets and calculated pooled effect estimates (regression coefficients) across datasets using Rubin's rules (Rubin, 1987; van Buuren & Groothuis-Oudshoorn, 2011). We used the p value from the covariance test to calculate unbiased confidence intervals for the effect estimates (Lockhart, Taylor, Tibshirani, & Tibshirani, 2014; Smith et al., 2015).

Finally, we evaluated how well the theoretical models selected in the SLCMA analyses fit the social cognition data across multiple timepoints and related to one other. To do this, we used structural equation modeling (SEM), which allowed us to further explore within a single analysis how maltreatment exposure was associated with social cognition across timepoints. In SEM, one or more measurement models – describing the relationships between latent factors and observed indicators – are joined together in a structural model, where associations between latent variables, covariates, and observed variables are estimated (Kline, 2010). We modeled the effects of sexual or physical abuse only in females because this model showed the strongest effect estimates in the SLCMA results and analyses of males would likely be uninformative, for reasons described later. Our hypothesis based on the SLCMA results was that the accumulation of sexual or physical abuse from 18 months to 6.75 years would predict social cognition at age 7.5, which in turn would predict social cognition at age 10.5, and subsequently social cognition at age 14. We also hypothesized based on the SLCMA results that the accumulation of sexual or physical abuse from 18 months to 8 years would also independently predict social cognition at 14 years (Figure 2). To evaluate goodness-of-fit, we used standard SEM fit statistics, including the normed comparative fit index (CFI) (Bentler, 1990), Tucker–Lewis index (TLI) (Tucker & Lewis, 1973), and root mean square error of approximation (RMSEA) (Steiger, 1990). We conducted these SEM analyses using the Lavaan package in R 3.3.2 (Yves, 2012).

Results

Sample characteristics

The analytic sample was gender balanced (49.71% female) and comprised predominantly white (97.1%) children from families whose parents were married and owned their home (Supplemental Table 1).

Distribution of exposure to maltreatment and social cognition skills

Over a quarter of the analytic sample (27%; $n = 1,182$) were exposed to at least one type of maltreatment, with 17.44% exposed to physical or emotional abuse, 13.29% exposed to sexual or physical abuse, and 5% exposed to both types.

Exposure to physical or sexual abuse was patterned by child sex (with males being more frequently exposed to either type) and exposure to physical or emotional abuse was patterned by several socio-demographic factors, including maternal material status, home ownership, and number of previous pregnancies ($p < .01$ for all chi-squared tests comparing children who were exposed to maltreatment to the entire analytic sample; Supplemental Table 1). Specifically, children who were exposed to maltreatment were less socioeconomically advantaged and less likely to be firstborn.

Age at exposure to maltreatment varied by type, with males having somewhat of an increase in sexual or physical abuse exposure as they aged. Alternatively, females experienced more physical or emotional abuse at preschool ages with more constant levels of exposure to sexual or physical abuse (Table 1).

Within each maltreatment type, exposures were highly correlated over time (Supplemental Table 5), with neighboring timepoints being generally more highly correlated than distant timepoints. Physical or emotional abuse by a caregiver had the highest correlation ($r = .73$) between ages 2.75 and 4 years, and sexual or physical by anyone had the highest correlation ($r = .59$) between ages 6.75 and 8 years.

Social cognition skills scores were moderately correlated across measurement timepoints ($r = .57-.67$). Notably, mean social cognition skill levels decreased at age 10.5 in both males and females, though significant sex differences were observed across all time points (Supplemental Table 6).

Association between maltreatment and social cognition

Results of the SLCMA analysis suggested a different patterning of associations between maltreatment and social cognition based on sex and the type of maltreatment (Table 2).

Among females, accumulation was the life course theoretical model consistently selected as the best-fitting one for both types of maltreatment. However, the effects of accumulation were only statistically significant at ages 7.5 and 14 for sexual or physical abuse ($\beta = 0.66$, $p < .01$ and $\beta = 0.65$, $p < .01$, respectively) and age 10.5 ($\beta = 0.27$, $p = .01$) for physical or emotional abuse by a caregiver, with less than 1% of the variance in social cognition explained by the accumulation of each type of maltreatment.

For males, no life course theoretical model achieved statistical significance between exposure to sexual or physical abuse and social cognition. However, an ever-exposed model was selected as the best fitting model for exposure to physical or emotional abuse on social cognition at age 7.5 ($\beta = 0.86$, $p < .01$), though this effect was not observed at the later measurements.

Of note, this general pattern of results was similar after winsorizing the social cognition score values to the 90th percentile,




Figure 2. Goodness of fit for structural equation model selected by the structured life course modeling approach (SLCMA). Abuse refers to sexual or physical abuse by anyone with the same timepoints of measurement as previous analyses. $**p < .001$ and $*p < .01$. Note, the Abuse measurements capture the same periods of abuse with the exception of the age-8 timepoint which is only included in “Abuse 18 m through 8 y summed”.

Table 2. Results of the structured life course modeling approach (SLCMA) for each measure of maltreatment on social cognition

		Sexual or physical abuse (by anyone)					
		Stage 1			Stage 2		
Model(s) selected		R^2	p value	β	SE	Lower CI	Upper CI
Female (N = 2,206)							
Age 7.5	Accumulation	.71%	<.01	0.66	0.13	0.41	0.91
Age 10.5	Accumulation	.22%	.08	0.60	0.12	0.30	0.90
Age 14	Accumulation	.81%	<.01	0.65	0.13	0.39	0.92
Male (N = 2,232)							
Age 7.5	None						
Age 10.5	None						
Age 14	None						
		Physical or emotional abuse by caregiver					
Female (N = 2,206)							
Age 7.5	None						
Age 10.5	Accumulation	.36%	.01	0.27	0.07	0.14	0.40
Age 14	None						
Male (N = 2,232)							
Age 7.5	Ever exposed	.43%	<.01	0.86	0.23	0.41	1.31
Age 10.5	Accumulation	.23%	.06	0.44	0.10	0.21	0.68
Age 14	None						

Stage 1 cell entries are r^2 values, p values. Stage 2 cell entries are betas, standard errors, and p values derived from multiple linear regression (one regression for each type of maltreatment and social cognition measurement). Models were only reported at Stage 2 when the covariance test p value was below the threshold of .1.

which reduced the effects of extreme scores (Supplemental Table 7).

Structural equation modeling (SEM)

Building from these results, we used SEM to examine the effect of sexual or physical abuse on females' social cognition across time using the theoretical models identified by the SLCMA. SEM modeling demonstrated the goodness of fit of our hypothesis: the main structural model (Figure 2) fit the data adequately (RMSEA = 0.09; 95% confidence interval (CI) [0.08; 0.10]; CFI = 0.98; TLI = 0.95), suggesting that sexual or physical abuse in early life explained lower social cognition scores later in life, via influences on earlier social cognition. Accumulation of sexual or physical abuse from 18 months to 6.75 years predicted social cognition at age 7.5 years and accumulation of sexual or physical abuse from 18 months to 8 years, as well as social cognition at age 10.5 predicted social cognition at 14 years. We also saw that social cognition scores at 7.5 years predicted social cognition scores at age 10.5. That is, abuse accumulated between 18 months and 6.75 years of age was associated with social cognition at 7.5 years ($\beta = 0.83, p < .01$), which in turn was associated with social cognition at 10.5 years ($\beta = 0.64, p < .01$), and subsequently linked to social cognition at 14 years ($\beta = 0.68, p < .01$). Social cognition at 14 years was predicted by abuse accumulated between 18 months and 8 years ($\beta = 0.13, p < .01$). Modification indices did not reveal any additional plausible paths between accumulation and the social cognition outcome that would improve model fit. The results of our initial SLCMA testing indicated that for males, the SEM test would unlikely be significant; unlike females, males did not have the same life course theoretical models selected for multiple time points of exposure.

Secondary analyses

A primary hypothesis tested in this paper is that childhood maltreatment predicts future social cognitive skills. However, children with poor social cognitive skills may also be more likely than their peers to be exposed to child maltreatment. To explore this possibility, we performed a secondary analysis to examine the association between social cognition and child maltreatment (see Supplemental Materials). Results suggested that poorer earlier social cognition skills were generally associated with lower levels of exposure to maltreatment (Supplemental Table 8).

Discussion

The current study used data from a large, population-based sample of children to examine associations between two types of child maltreatment and the development of subsequent social cognition skills from late childhood through early adolescence. Three main findings emerged from this work. First, childhood maltreatment differentially impacted males and female's social cognition development. Specifically, there were sex differences both in how the characteristics of maltreatment associated with social cognition and the strength of these associations. This sex-dependent pattern of findings was unsurprising in light of prior work showing differences between male and female social cognitive skills between the ages of 8 and 21 (Gur et al., 2012). Our findings are consistent with prior findings in social cognitive development, where females have been shown to outperform males in facial emotion recognition through age 16 (Lawrence, Campbell, & Skuse,

2015) and have increased social affect through early development (Messinger et al., 2015). Social cognitive skills in males typically develop later than females and thus we may see a smaller effect size of maltreatment when measuring early time points in males. Thus, we postulate that males and females may be similarly impacted by maltreatment, but that for males, such effects may not appear until later in development. Based on previous work, these effects of maltreatment on social cognition may not emerge until late in teenage years or emerging adulthood, as boys begin to catch-up in social cognitive skills with their female counterparts (Lawrence et al., 2015; Rose & Rudolph, 2006; Van Der Graaff et al., 2014).

Second, our results from the SLCMA suggest that more than any other feature of maltreatment, the accumulation of abuse plays the biggest role in explaining variations in social cognition development specifically for females. In females, the number of time periods exposed to physical or sexual abuse from 0–8 years was positively and linearly associated with social cognition deficits. While prior literature has primarily characterized the effects of the severity of abuse, with more severe abuse leading to greater social cognitive deficits, this is the first study to our knowledge to focus specifically on the number of time periods exposed in relation to social cognition abilities (Barahal et al., 1981; Luke & Banerjee, 2013). Such comparisons are likely to be informative for guiding the development of interventions to combat the negative consequences of exposure to abuse.

Third, our results from the SLCMA and SEM analyses pertaining to physical or sexual maltreatment not only emphasize the power of integrating different statistical models, but also expand upon previous work by suggesting that the duration of abuse and its effects on social cognition may persist from childhood to adolescence. Our findings are consistent with prior studies, which have examined the effect of abuse on social cognition during later life, and further suggest that the effects of adversity on social cognition can be observed shortly after abuse and may quickly lead to persistent differences in social cognition (Ford et al., 2011; Germine, Dunn, McLaughlin, & Smoller, 2015; Palmier-Claus et al., 2016). These findings emphasize the importance of early identification and intervention efforts to reach children, particularly girls, while these social cognitive deficits are developing, which would be expected to minimize any long-term repercussions of being exposed to adversity. Of note, our results are most likely driven by physical, rather than sexual abuse, due to the former being more common than the latter; this should be considered when interpreting results of this study. Though the effect size was small and there are likely other factors that have not been captured here that shape social cognitive development, our findings strengthen past research and help to inform future work on this topic.

Yet, two major questions also emerged from this analysis. The first is: why were there no significant effects of maltreatment on social cognition in males? Prior studies have shown that males lag substantially behind their female counterparts during the initial development of empathy (Rose & Rudolph, 2006; Van Der Graaff et al., 2014), which is one of the key social cognition skills developed during adolescence. Many items included in our social cognition measure capture experiences that involve empathy skills, including those asking about appropriate responses to other's emotions. Our finding that males had significantly poorer social cognition scores throughout time may suggest that our social cognition measure was capturing a domain that would both be immature in males and unlikely to be affected by

maltreatment. We further speculate that the development of social cognition in males may be less stable and more variable in late childhood and adolescence, given previous work noting dramatic changes in social behavior and interaction among males around puberty, such as increased antisocial and aggressive behaviors (Forbes & Dahl, 2010; Rowe, Maughan, Worthman, Costello, & Angold, 2004). Thus, it might be possible to see more expected adverse effects of maltreatment on social cognitive development reemerge once puberty ends.

A second question was: why do we see a shift in social cognition scores across time and further, why are there differences confined to age 10.5 years for the effects of sexual and physical abuse on social cognition? Univariate analyses revealed that for both males and females, social cognition scores worsened considerably at age 10.5. Moreover, among females, a model of exposure to physical or emotional abuse was selected as explaining a significant amount of variation in social cognition at 10.5, whereas no theoretical models were selected for the other outcome timepoints in the SLCMA modeling. These results are consistent with prior research suggesting that key social cognition skills, such as facial recognition, temporarily plateau or even decline in early adolescence (Carey & Diamond, 1977; Carey, Diamond, & Woods, 1980; Germine, Duchaine, & Nakayama, 2011). Thus, our measure of social cognition obtained at age 10.5 may fall within an expected aberrant time period, which could account for the unique pattern of results observed during this outcome time point.

There are several strengths of this study. Data came from a large, population-based prospective study. Previous work has tended to only include a single measurement of social cognition, involve retrospective maltreatment reports that do not include repeated measures, or focus on social cognition in clinical samples (e.g., children with autism or Turner's syndrome). To our knowledge, this is the first prospective, population-based study. The use of a prospective study was a particular strength, as it provided a stronger test of our research questions (relative to retrospective or cross-sectional work), and it enabled us to better contextualize these results and identify possibly causal relations. For example, as shown through our secondary analyses, we were able to learn that poorer social cognition skills could also predict lower levels of future exposure to maltreatment. Moreover, our use of a theory-driven analytic method (SLCMA) in combination with SEM was another major innovation. Sample code for implementing the SLCMA is publicly available through a GitHub page (<https://github.com/thedunnlab/SLCMA-pipeline>).

Some limitations must also be considered. First, the ALSPAC dataset comprised largely children of European-ancestry. Thus, our findings may not be generalizable to more racially and ethnically diverse populations. Second, the use of maternal self-report questionnaires to capture abuse may be problematic. Mothers may have underreported their child's exposure to abuse, particularly if mothers were implicated in the maltreatment or the maltreatment events occurred outside the home. Reliance on these subjective reports was also likely further complicated because ALSPAC did not provide reporters with clear guidelines or criteria to define these abuse experiences. Thus, some reporters could have downplayed their maltreating behavior and not identified it as abuse, whereas other reporters could have responded to these items considering behaviors that might not fit the traditional definition of maltreatment. Yet, even with these limitations, we found that the prevalence of maltreatment in this sample was comparable to nationally representative samples which use social service or other agency reporting (Gilbert *et al.*, 2009). These

similarities in prevalence estimates of maltreatment between our sample and nationally representative samples provide some confidence that our measures may be capturing true childhood maltreatment cases in this sample. Moreover, the use of maternal reporting could also lead to bias in social cognitive scoring, as mother's may expect boys to be less social than girls. Although there were sex differences in social cognition scores here, the magnitude of these differences were similar to those previously reported (Erwin *et al.*, 1992; Gur *et al.*, 2012; Williams *et al.*, 2009). However, previous work on the role of parental bias in social skills suggests that parents identify more social skill problems than children, but are less able to identify subtle symptoms such as withdrawal (Howells Wrobel & Lachar, 1998). Some research even suggests parental reports may be more accurate than child self-reports for social behaviors (Fisher, Mello, & Dykens, 2014). Third, the measures of maltreatment lacked details to characterize the specific nature of the abuse, including its severity. This has been a limitation of other large-scale epidemiological studies when examining abuse. However, by having the measurement of frequency of maltreatment (defined here as the number of occasions on which it was reported), we can distinguish between children who were exposed only a single time from those who were exposed multiple times, which can help clarify the effect of different maltreatment experiences (Dunn *et al.*, 2018). In future studies, more detailed assessments could prove valuable to document more specific effects of maltreatment features on social cognition and build upon these findings. Fourth, we modeled childhood adversity as an exposure that may impact social cognition; however, our results are only correlational and do not rule out the possibility that preliminary manifestations of poor social cognition in early childhood could also increase risk of maltreatment, which we explored briefly in our secondary analyses. Fifth, as with any longitudinal study, there was attrition over time. Although we attempted to address this attrition by using multiple imputation, the fluctuation in participants reporting outcomes across different outcome timepoints and source of reporting led to smaller sample sizes, since we only analyzed children with maternal reports at all three timepoints. However, we think the conservative exclusion criteria were necessary to minimize potential bias induced by heterogeneity in the mode of reporting. Removing participants who were missing the necessary longitudinal data for our analyses may exclude higher-risk children and/or limit broader generalizability. However, the included sample was comparable to the sample of participants who were excluded. Further work targeting the understanding of reporting patterns in social cognition research could guide the interpretation of our results as well as sample selection of future studies.

In summary, our results suggest that the accumulation of sexual or physical abuse in early childhood can have a lasting impact on female's social cognition skills. These findings underscore the need for further work to measure how maltreatment effects can persist across time and shape long-term developmental trajectories. These results also emphasize the importance of identifying females who have experienced maltreatment, who may be at risk for having social cognitive challenges.

Supplementary Material. The supplementary material for this article can be found at <https://doi.org/10.1017/S095457942000139X>.

Acknowledgments. The content is solely the responsibility of the authors and does not necessarily represent the official views of the National

Institutes of Health. The authors thank Alice Renaud for her assistance in preparing this manuscript for publication. We are extremely grateful to all the families who took part in this study, the midwives for their help in recruiting them, and the whole ALSPAC team, which includes interviewers, computer and laboratory technicians, clerical workers, research scientists, volunteers, managers, receptionists, and nurses. This publication is the work of the authors who will serve as guarantors for the contents of this paper.

Financial Statement. This research was specifically funded by the National Institute of Mental Health of the National Institutes of Health under Award Numbers K01MH102403 and 1R01MH113930 (Dunn). The UK Medical Research Council and the Wellcome Trust (Grant ref: 102215/2/13/2) and the University of Bristol provide core support for ALSPAC. A comprehensive list of grants funding is available on the ALSPAC website (<http://www.bristol.ac.uk/alspac/external/documents/grant-acknowledgements.pdf>).

Conflicts of Interest. None

References

- Addington, J., Girard, T. A., Christensen, B. K., & Addington, D. (2010). Social cognition mediates illness-related and cognitive influences on social function in patients with schizophrenia-spectrum disorders. *Journal of Psychiatry & Neuroscience : JPN*, *35*, 49. doi:10.1503/jpn.080039
- Barahal, R. M., Waterman, J., & Martin, H. P. (1981). The social cognitive development of abused children. *Journal of Consulting and Clinical Psychology*, *49*, 508–516. doi:10.1037/0022-006X.49.4.508
- Barresi, J., & Moore, C. (1996). Intentional relations and social understanding. *Behavioral and Brain Sciences*, *19*, 107–122. doi:10.1017/S0140525X00041790
- Ben-Shlomo, Y., & Kuh, D. (2002). A life course approach to chronic disease epidemiology: Conceptual models, empirical challenges, and interdisciplinary perspectives. *International Journal of Epidemiology*, *31*, 285–293. doi:10.1093/ije/31.2.285
- Bentler, P. M. (1990). Comparative fit indexes in structural models. *Psychological Bulletin*, *107*, 238–246. doi:10.1037/0033-2909.107.2.238
- Bosacki, S. L. (2000). *Theory of mind in preadolescence: connections among social understanding, self-concept, and social relations* (pp. 3602–3602). Toronto, Canada: University of Toronto.
- Bosch, N. M., Riese, H., Reijneveld, S. A., Bakker, M. P., Verhulst, F. C., Ormel, J., & Oldehinkel, A. J. (2012). Timing matters: Long term effects of adversities from prenatal period up to adolescence on adolescents' cortisol stress response. The TRAILS study. *Psychoneuroendocrinology*, *37*, 1439–1447. doi:10.1016/j.psyneuen.2012.01.013
- Boyd, A., Golding, J., Macleod, J., Lawlor, D. A., Fraser, A., Henderson, J., ... Davey Smith, G. (2013). Cohort profile: The 'children of the 90s' – the index offspring of the Avon Longitudinal Study of Parents and Children. *International Journal of Epidemiology*, *42*, 111–127. doi:10.1093/ije/dys064
- Briere, J., & Elliott, D. M. (2003). Prevalence and psychological sequelae of self-reported childhood physical and sexual abuse in a general population sample of men and women. *Child Abuse & Neglect*, *27*, 1205–1222. doi:10.1016/j.chiabu.2003.09.008
- Burnett, S., Sebastian, C., Cohen Kadosh, K., & Blakemore, S.-J. (2011). The social brain in adolescence: Evidence from functional magnetic resonance imaging and behavioural studies. *Neuroscience and Biobehavioral Reviews*, *35*, 1654–1664. doi:10.1016/j.neubiorev.2010.10.011
- Cabrera, N. J., Fagan, J., Wight, V., & Schadler, C. (2011). Influence of mother, father, and child risk on parenting and children's cognitive and social behaviors. *Child Development*, *82*, 1985–2005. doi:10.1111/j.1467-8624.2011.01667.x
- Carey, S., & Diamond, R. (1977). From piecemeal to configurational representation of faces. *Science*, *195*, 312–314. doi:10.1126/science.831281
- Carey, S., Diamond, R., & Woods, B. (1980). Development of face recognition: A maturational component? *Developmental Psychology*, *16*, 257–269. doi:10.1037/0012-1649.16.4.257
- Cicchetti, D., Cowell, R. A., Rogosch, F. A., & Toth, S. L. (2015). Childhood maltreatment and its effect on neurocognitive functioning: Timing and chronicity matter. *Development and Psychopathology*, *27*, 521–533.
- Cox, J. L., Holden, J. M., & Sagovsky, R. (1987). Detection of postnatal depression. Development of the 10-item Edinburgh postnatal depression scale. *The British Journal of Psychiatry*, *150*, 782–786. doi:10.1192/bjp.150.6.782
- Crick, N. R., & Dodge, K. A. (1994). A review and reformulation of social information-processing mechanisms in children's social adjustment. *Psychological Bulletin*, *115*, 74. doi:10.1037/0033-2909.115.1.74
- Dale, C. L., Findlay, A. M., Adcock, R. A., Vertinski, M., Fisher, M., Genevsky, A., ... Vinogradov, S. (2010). Timing is everything: Neural response dynamics during syllable processing and its relation to higher-order cognition in schizophrenia and healthy comparison subjects. *International Journal of Psychophysiology*, *75*, 183–193. doi:10.1016/j.ijpsycho.2009.10.009
- Dunn, E. C., McLaughlin, K. A., Slopen, N., Rosand, J., & Smoller, J. W. (2013). Developmental timing of child maltreatment and symptoms of depression and suicidal ideation in young adulthood: Results from the National Longitudinal Study of Adolescent Health. *Depression and Anxiety*, *30*, 955–964.
- Dunn, E. C., Nishimi, K., Powers, A., & Bradley, B. (2017). Is developmental timing of trauma exposure associated with depressive and post-traumatic stress disorder symptoms in adulthood? *Journal of Psychiatric Research*, *84*, 119–127. doi:10.1016/j.jpsychires.2016.09.004
- Dunn, E. C., Soare, T. W., Raffeld, M. R., Busso, D. S., Crawford, K. M., Davis, K. A., ... Susser, E. S. (2018). What life course theoretical models best explain the relationship between exposure to childhood adversity and psychopathology symptoms: Recency, accumulation, or sensitive periods? *Psychological Medicine*, *48*, 2562–2572. doi:10.1017/S0033291718000181
- Dvir, Y., Ford, J. D., Hill, M., & Frazier, J. A. (2014). Childhood maltreatment, emotional dysregulation, and psychiatric comorbidities. *Harvard Review of Psychiatry*, *22*, 149–161. doi:10.1097/HRP.0000000000000014
- Erwin, R. J., Gur, R. C., Gur, R. E., Skolnick, B., Mawhinney-Hee, M., & Smailis, J. (1992). Facial emotion discrimination: I. Task construction and behavioral findings in normal subjects. *Psychiatry research*, *42*(3), 231–240.
- Evans, G. W., Li, D., & Whipple, S. S. (2013). Cumulative risk and child development. *Psychological Bulletin*, *139*, 342–396. doi:10.1037/a0031808
- Finkelhor, D., Turner, H. A., Shattuck, A., & Hamby, S. L. (2013). Violence, crime, and abuse exposure in a national sample of children and youth: An update. *JAMA Pediatrics*, *167*, 614–621. doi:10.1001/jamapediatrics.2013.42
- Fisher, M. H., Mello, M. P., & Dykens, E. M. (2014). Who reports it best? A comparison between parent-report, self-report, and the real life social behaviors of adults with Williams syndrome. *Research in developmental disabilities*, *35*(12), 3276–3284.
- Forbes, E. E., & Dahl, R. E. (2010). Pubertal development and behavior: Hormonal activation of social and motivational tendencies. *Brain and Cognition*, *72*, 66–72. doi:10.1016/j.bandc.2009.10.007
- Ford, E., Clark, C., & Stansfeld, S. A. (2011). The influence of childhood adversity on social relations and mental health at mid-life. *Journal of Affective Disorders*, *133*, 320–327. doi:10.1016/j.jad.2011.03.017
- Fraser, A., Macdonald-Wallis, C., Tilling, K., Boyd, A., Golding, J., Davey Smith, G., ... Lawlor, D. A. (2013). Cohort profile: The Avon longitudinal study of parents and children: ALSPAC mothers cohort. *International Journal of Epidemiology*, *42*, 97–110. doi:10.1093/ije/dys066
- Frith, C. D., & Frith, U. (2007). Social cognition in humans. *Current Biology*, *17*, R724–R732. doi:10.1016/j.cub.2007.05.068
- Frith, C. D., & Frith, U. (2007). Social cognition in humans. *Current biology*, *17*(16), R724–R732.
- Germine, L. T., Duchaine, B., & Nakayama, K. (2011). Where cognitive development and aging meet: Face learning ability peaks after age 30. *Cognition*, *118*, 201–210. doi:10.1016/j.cognition.2010.11.002
- Germine, L., Dunn, E. C., McLaughlin, K. A., & Smoller, J. W. (2015). Childhood adversity is associated with adult theory of mind and social affiliation, but not face processing. *PLoS One*, *10*, e0129612. doi:10.1371/journal.pone.0129612
- Gilbert, R., Widom, C. S., Browne, K., Fergusson, D., Webb, E., & Janson, S. (2009). Burden and consequences of child maltreatment in high-income countries. *Lancet*, *373*, 68–81. doi:10.1016/S0140-6736(08)61706-7
- Green, M. F., Olivier, B., Crawley, J. N., Penn, D. L., & Silverstein, S. (2005). Social cognition in schizophrenia: Recommendations from the measurement and treatment research to improve cognition in schizophrenia work

- approaches conference. *Schizophrenia Bulletin*, 31, 882–887. doi:10.1093/schbul/sbi049
- Gur, R. C., Richard, J., Calkins, M. E., Chiavacci, R., Hansen, J. A., Bilker, W. B., ... Gur, R. E. (2012). Age group and sex differences in performance on a computerized neurocognitive battery in children age 8–21. *Neuropsychology*, 26, 251–265. doi:10.1037/a0026712
- Happé, F., & Frith, U. (2014). Annual Research Review: Towards a developmental neuroscience of atypical social cognition. *Journal of Child Psychology and Psychiatry*, 55(6), 553–577.
- Hibbeln, J. R., Davis, J. M., Steer, C., Emmett, P., Rogers, L., Williams, C., & Golding, J. (2007). Maternal seafood consumption in pregnancy and neurodevelopmental outcomes in childhood (ALSPAC study): An observational cohort study. *The Lancet*, 369, 578–585. doi:10.1016/S0140-6736(07)60277-3
- Kilford, E. J., Garrett, E., & Blakemore, S.-J. (2016). The development of social cognition in adolescence: An integrated perspective. *Neuroscience and Biobehavioral Reviews*, 70, 106–120. doi:10.1016/j.neubiorev.2016.08.016
- Kline, R. B. (2010). *Principles and practice of structural equation modeling*. New York, NY: Guilford Press.
- Kuh, D., Ben-Shlomo, Y., Lynch, J., Hallqvist, J., & Power, C. (2003). Life course epidemiology. *Journal of Epidemiology and Community Health*, 57, 778–783. doi:10.1136/jech.57.10.778
- Ladegaard, N., Larsen, E. R., Videbeck, P., & Lysaker, P. H. (2014). Higher-order social cognition in first-episode major depression. *Psychiatry Research*, 216, 37–43. doi:10.1016/j.psychres.2013.12.010
- Lang, C. A., Cox, M. J., & Flores, G. (2013). Maltreatment in multiple-birth children. *Child Abuse & Neglect*, 37, 1109–1113. doi:10.1016/j.chiabu.2013.03.002
- Lawrence, K., Campbell, R., & Skuse, D. (2015). Age, gender, and puberty influence the development of facial emotion recognition. *Frontiers in Psychology*, 6, 761. doi:10.3389/fpsyg.2015.00761
- Levin, A. R., Fox, N. A., Zeanah Jr, C. H., & Nelson, C. A. (2015). Social communication difficulties and autism in previously institutionalized children. *Journal of the American Academy of Child & Adolescent Psychiatry*, 54(2), 108–115.
- Lockhart, R., Taylor, J., Tibshirani, R. J., & Tibshirani, R. (2014). A significance test for the LASSO. *Annals of Statistics*, 42, 413–468. doi:10.1214/13-AOS1175
- Luke, N., & Banerjee, R. (2013). Differentiated associations between childhood maltreatment experiences and social understanding: A meta-analysis and systematic review. *Developmental Review*, 33, 1–28. doi:10.1016/j.dr.2012.10.001
- Magiati, I., Tay, X. W., & Howlin, P. (2014). Cognitive, language, social and behavioural outcomes in adults with autism spectrum disorders: A systematic review of longitudinal follow-up studies in adulthood. *Clinical Psychology Review*, 34, 73–86. doi:10.1016/j.cpr.2013.11.002
- Manly, J. T., Kim, J. E., Rogosch, F. A., & Cicchetti, D. (2001). Dimensions of child maltreatment and children's adjustment: Contributions of developmental timing and subtype. *Development and Psychopathology*, 13, 759–782. doi:10.1017/S0954579401004023
- McLaughlin, K. A., Green, J. G., Gruber, M. J., Sampson, N. A., Zaslavsky, A. M., & Kessler, R. C. (2010). Childhood adversities and adult psychiatric disorders in the National Comorbidity Survey Replication II: Associations with persistence of DSM-IV disorders. *Archives of General Psychiatry*, 67, 124–132. doi:10.1001/archgenpsychiatry.2009.187
- Messinger, D. S., Young, G. S., Webb, S. J., Ozonoff, S., Bryson, S. E., Carter, A., ... Zwaigenbaum, L. (2015). Early sex differences are not autism-specific: A Baby Siblings Research Consortium (BSRC) study. *Molecular Autism*, 6, 32. doi:10.1186/s13229-015-0027-y
- Mills, K. L., Lalonde, F., Clasen, L. S., Giedd, J. N., & Blakemore, S.-J. (2014). Developmental changes in the structure of the social brain in late childhood and adolescence. *Social Cognitive and Affective Neuroscience*, 9, 123–131. doi:10.1093/scan/nss113
- Munafò, M. R., & Davey Smith, G. (2018). Robust research needs many lines of evidence. *Nature*, 553, 399. doi:10.1038/d41586-018-01023-3
- Oliver, B. R., Barker, E. D., Mandy, W. P. L., Skuse, D. H., & Maughan, B. (2011). Social cognition and conduct problems: A developmental approach. *Journal of the American Academy of Child & Adolescent Psychiatry*, 50, 385–394. doi:10.1016/j.jaac.2011.01.006
- Palmier-Claus, J., Berry, K., Darrell-Berry, H., Emsley, R., Parker, S., Drake, R., & Bucci, S. (2016). Childhood adversity and social functioning in psychosis: Exploring clinical and cognitive mediators. *Psychiatry Research*, 238, 25–32. doi:10.1016/j.psychres.2016.02.004
- Piskulic, D., & Addington, J. (2011). Social cognition and negative symptoms in psychosis. *Psychiatry Research*, 188, 283–285. doi:10.1016/j.psychres.2011.04.028
- Prino, L. E., Rollé, L., Sechi, C., Patteri, L., Ambrosoli, A., Calderera, A. M., ... Brustia, P. (2016). Parental relationship with twins from pregnancy to 3 months: The relation among parenting stress, infant temperament, and well-being. *Frontiers in Psychology*, 7, 1628. doi:10.3389/fpsyg.2016.01628
- Rose, A. J., & Rudolph, K. D. (2006). A review of sex differences in peer relationship processes: Potential trade-offs for the emotional and behavioral development of girls and boys. *Psychological Bulletin*, 132, 98–131. doi:10.1037/0033-2909.132.1.98
- Rowe, R., Maughan, B., Worthman, C. M., Costello, E. J., & Angold, A. (2004). Testosterone, antisocial behavior, and social dominance in boys: Pubertal development and biosocial interaction. *Biological Psychiatry*, 55, 546–552. doi:10.1016/j.biopsych.2003.10.010
- Rubin, D. B. (1987). *Multiple imputation for nonresponse in surveys*. New York, NY: J. Wiley & Sons.
- Sasson, N. J., Nowlin, R. B., & Pinkham, A. E. (2013). Social cognition, social skill, and the broad autism phenotype. *Autism: The International Journal of Research and Practice*, 17, 655–667. doi:10.1177/1362361312455704
- Sergi, M. J., Rassovsky, Y., Nuechterlein, K. H., & Green, M. F. (2006). Social perception as a mediator of the influence of early visual processing on functional status in schizophrenia. *American Journal of Psychiatry*, 163, 448–454. doi:10.1176/appi.ajp.163.3.448
- Skuse, D. H., James, R. S., Bishop, D. V. M., Coppin, B., Dalton, P., Aamodt-Leeper, G., ... Jacobs, P. A. (1997). Evidence from Turner's syndrome of an imprinted x-linked locus affecting cognitive function. *Nature*, 387, 705–708. doi:10.1038/42706
- Smith, A. D., Hardy, R., Heron, J., Joinson, C. J., Lawlor, D. A., Macdonald-Wallis, C., & Tilling, K. (2016). A structured approach to hypotheses involving continuous exposures over the life course. *International Journal of Epidemiology*, 45, 1271–1279. doi:10.1093/ije/dyw238
- Smith, A. D., Heron, J., Mishra, G., Gilthorpe, M. S., Ben-Shlomo, Y., & Tilling, K. (2015). Model selection of the effect of binary exposures over the life course. *Epidemiology*, 26, 719–726. doi:10.1097/EDE.0000000000000348
- Steiger, J. H. (1990). Structural model evaluation and modification: An interval estimation approach. *Multivariate Behavioral Research*, 25, 173–180. doi:10.1207/s15327906mbr2502_4
- Stoltenborgh, M., Bakermans-Kranenburg, M. J., Alink, L. R. A., & Ijzendoorn, M. H. (2015). The prevalence of child maltreatment across the globe: Review of a series of meta-analyses. *Child Abuse Review* 24(1), 37–50. doi:10.1002/car.2353
- Teicher, M. H., Andersen, S. L., Polcari, A., Anderson, C. M., Navalta, C. P., & Kim, D. M. (2003). The neurobiological consequences of early stress and childhood maltreatment. *Neuroscience & Biobehavioral Reviews*, 27, 33–44. doi:10.1016/S0149-7634(03)00007-1
- Tucker, L. R., & Lewis, C. (1973). A reliability coefficient for maximum likelihood factor analysis. *Psychometrika*, 38, 1–10. doi:10.1007/BF02291170
- van Buuren, S., & Groothuis-Oudshoorn, K. (2011). Mice: Multivariate imputation by chained equations in R. *Journal of Statistical Software*, 45, 1–68.
- Van Der Graaff, J., Branje, S., De Wied, M., Hawk, S., Van Lier, P., & Meeus, W. (2014). Perspective taking and empathic concern in adolescence: Gender differences in developmental changes. *Developmental Psychology*, 50, 881–888. doi:10.1037/a0034325
- Vauth, R., Rüscher, N., Wirtz, M., & Corrigan, P. W. (2004). Does social cognition influence the relation between neurocognitive deficits and vocational functioning in schizophrenia? *Psychiatry Research*, 128, 155–165. doi:10.1016/j.psychres.2004.05.018
- Vetter, N. C., Leipold, K., Kliegel, M., Phillips, L. H., & Altgassen, M. (2013). Ongoing development of social cognition in adolescence. *Child Neuropsychology*, 19, 615–629. doi:10.1080/09297049.2012.718324

- Weijers, J., Fonagy, P., Eurelings-Bontekoe, E., Termorshuizen, F., Viechtbauer, W., & Selten, J. P. (2009). Mentalizing impairment as a mediator between reported childhood abuse and outcome in nonaffective psychotic disorder. *Psychiatry research, 259*, 463–469.
- Williams, L. M., Mathersul, D., Palmer, D. M., Gur, R. C., Gur, R. E., & Gordon, E. (2009). Explicit identification and implicit recognition of facial emotions: I. Age effects in males and females across 10 decades. *Journal of Clinical and Experimental Neuropsychology, 31*(3), 257–277.
- Wrobel Howells, N., & Lachar, D. (1998). Validity of self-and parent-report scales in screening students for behavioral and emotional problems in elementary school. *Psychology in the Schools, 35*(1), 17–27.
- Yves, R. (2012). Lavaan: An R package for structural equation modeling. *Journal of Statistical Software, 48*, 1–36.