Contributors

- Jan Fawcett, MD, Professor of Psychiatry, Rush-Presbyterian-St Luke's Medical Center, 1725 West Harrison, Suite 955, Chicago, Illinois 60612, USA
- Robert M. A. Hirschfeld, MD, Titus Harris Professor and Chairman of Psychiatry and Behavioral Sciences, University of Texas, Graves Building (Room 1200), 1014 Texas Avenue, Galveston, TX 77555, USA
- Martin B. Keller, MD, Psychiatrist-in-Chief, Butler Hospital, and Executive Psychiatrist-in-Chief, Brown Affiliated Hospitals, 345 Blackstone Boulevard, Providence, RI 02906, USA
- Ronald C. Kessler, PhD, Institute for Social Research, University of Michigan, 426 Thompson, Room 5062, Ann Arbor, Michigan 48105, USA
- Bärbel Knäuper, PhD, Max Planck Institute of Psychiatry, Clinical Institute, Kraepelinstrasse 10, 80804 München, Germany
- Joseph Mendels, MD, Medical Director, Philadelphia Medical Institute, 1015 Chestnut Street, Suite 1303, Philadelphia, PA 19107, USA
- Stuart A. Montgomery, MD, Academic Department of Psychiatry, St Mary's Hospital Medical School, Praed Street, London W2 1NY, UK
- Eugene S. Paykel, FRCPsych, Professor of Psychiatry, University of Cambridge, Addenbrooke's Hospital, Hills Road, Cambridge CB2 2QQ, UK
- Myrna M. Weissman, PhD, Professor of Psychiatry and Epidemiology, College of Physicians and Surgeons, Columbia University, and Chief, Division of Clinical and Genetic Epidemiology, New York State Psychiatric Institute, 722 West 168th Street, Box 14, New York, NY 10032, USA
- Hans-Ulrich Wittchen, PhD, Professor of Clinical Psychology, Max Planck Institute of Psychiatry, Clinical Institute, Kraepelinstrasse 10, 80804 München, Germany