

Pericarp structure of *Glebionis coronaria* (L.) Cass. ex Spach (Asteraceae) cypselae controls water uptake during germination

Giuseppe Puglia^{1,2*}, Simona Grimaldi¹, Angelino Carta³, Pietro Pavone¹ and Peter Toorop⁴

¹Department of Biological, Geological and Environmental Science, Plant Biology Section, University of Catania, via Empedocle 58, 95128, Catania, Italy; ²Consiglio Nazionale delle Ricerche, Istituto per i Sistemi Agricoli e Forestali del Mediterraneo (CNR-ISAFOM) U.O.S. Catania, via Empedocle, 58, 95128, Catania, Italy; ³Department of Biology, Unit of Botany, University of Pisa, via Luca Ghini 13, I-56126, Pisa, Italy; ⁴Seed Conservation Department, Royal Botanic Gardens, Kew, Wakehurst Place, Ardingly, West Sussex, RH17 6TN, UK

(Received 28 August 2014; accepted after revision 6 February 2015; first published online 14 April 2015)

Abstract

Glebionis coronaria (L.) Cass. ex Spach is a common Mediterranean weed producing distinctive central and peripheral dormant cypselae with a hard fruit coat, which was previously hypothesized to impose physical dormancy. Analysis of water uptake in cypselae and in naked seeds showed that it preferentially takes place at the basal end of the fruit; however, seeds within an intact pericarp do not fully imbibe when compared with naked seeds. Germination was not significantly different between the two heteromorphs, and after-ripening or cold stratification did not increase germination, while warm stratification at 35/20°C, as revealed by logistic regression, resulted in a significant improvement. However, loss of viability was also rapid at these high temperatures. Central and peripheral cypselae generally showed very low germination. In both heteromorphs, faster and higher germination (60–70%) was reached only after extensive scarification of pericarp tissue, and full germination was observed only after complete removal of pericarp tissue. Although the pericarp significantly reduced water uptake, no palisade layer(s) of macrosclereids could be observed. Xylem-vessel elements were found running through the basal end of the pericarp and forming the main point of water entry. We reject the hypothesis that *G. coronaria* cypselae have physical dormancy. Instead, water uptake and germination are impeded by: (1) directed water uptake, mainly through a pericarp-spanning channel-like structure; and (2) mechanical constraint on embryo growth exerted by

the hard pericarp. The channel-like structure forms the principal system for controlling seed germination.

Keywords: dormancy, garland chrysanthemum, germination, *Glebionis coronaria*, heterocarpy, imbibition, pericarp constraint

Introduction

Heterocarpy, or fruit heteromorphism on a single plant (McDonough, 1975; Imbert, 2002), is often associated with differential dispersal mechanisms (heterodispersy) (Lu *et al.*, 2010; Baskin *et al.*, 2013) and ecological behaviour (Picó and Koubek, 2003). This strategy increases the adaptability of the species to highly variable environments (Lloyd, 1984; Venable *et al.*, 1987), such as the Mediterranean seasonal climate with highly variable temperature and precipitation regimes (Skordilis and Thanos, 1995; Doussi and Thanos, 2002; Carta *et al.*, 2013). Fruit heteromorphism is particularly common in Asteraceae, occurring in several tribes and genera (Tanowitz *et al.*, 1987; Imbert, 2002). Moreover, a number of studies describe fruit heteromorphism in association with differences in germination behaviour, e.g. in *Bidens*, *Senecio*, *Picris* and *Crepis* spp. (McEvoy, 1984; Venable *et al.*, 1987; Rocha, 1996). Central cypselae have a thinner pericarp and usually exhibit higher dispersal ability, germinating over a wider range of temperatures immediately after favourable conditions occur, whereas peripheral ones are characterized by limited dispersal ability and more restricted germination conditions (Forsyth and Brown, 1982; Corkidi *et al.*, 1991; Brändel, 2007). As a consequence, the pericarp structure could be, in part, responsible for

*Correspondence
Email: puglia.giuseppe@gmail.com

the different germination responses between central and peripheral cypselae (McEvoy, 1984; Tanowitz *et al.*, 1987). In *Garhadiolus papposus* the thick pericarp and phyllary of the peripheral cypselae, as well as the thick pericarp of the intermediate cypselae, were theorized to be the reason why these two cypselae types are more dormant than the central cypselae, by means of imposing a mechanical restraint on the embryo (Sun *et al.*, 2009). Similarly, the morphological and anatomical variability of the thickness and structure of the pericarp in dark cypselae of *Anthemis chrysantha*, belonging to the same tribe as *Glebionis coronaria*, was hypothesized to effect differences in dormancy and germination behaviour, leading to variations in imbibition time, oxygen exchange and mechanical restriction (Aguado *et al.*, 2011). In *Raphanus raphanistrum*, a Brassicaceae species growing in disturbed environments and bearing persistent fruits, the fruit wall was shown to restrict water uptake (Cousens *et al.*, 2010). Seed dormancy, which spreads seed germination over time and enables synchronization of germination and seedling growth with seasons of the year when conditions are favourable (Baskin and Baskin, 1998; Schütz *et al.*, 2002), is another effective adaptive mechanism (Venable *et al.*, 1987; Mandák, 2003), especially in highly disturbed environments (Hawes *et al.*, 2005; Toorop *et al.*, 2012). *G. coronaria* Cass. ex L. Spach (Asteraceae) is a heterocarpic annual weed with dormant cypselae. Although native to the Mediterranean region (Meusel *et al.*, 1965; Pignatti, 1982), nowadays it is naturalized far beyond its 'native' range and, due to its abundance and persistence, it is reported as an invasive outside Europe (Wang *et al.*, 2009; Cook and Talley, 2014). Knowledge of its germination mechanisms is important in limiting the distribution of this species in habitats where its presence could alter the ecosystem equilibrium (Cook and Talley, 2014).

G. coronaria grows in cultivated fields, along roadsides and in disturbed sites on a variety of soil types. Its heteromorphic cypselae differ in dispersal behaviour, pericarp size and level of dormancy (Bastida and Menéndez, 2004; Bastida *et al.*, 2010). In previous studies, it was hypothesized that *G. coronaria* cypselae display physical dormancy imposed by the pericarp tissue (Bastida and Menéndez, 2004; Bañón *et al.*, 2009). However, according to the classification system proposed by Baskin and Baskin (2004) physical dormancy is defined as the impermeability of the seed (or fruit) covers to water, and it is frequently associated with the presence of one or multiple layers of, usually, Malpighian palisade cells isolating the embryo (or seed) from water uptake. However, even though fresh matured cypselae of *G. coronaria* display an evident inhibition of germination, physical dormancy is not known to occur in the Asteraceae family (Baskin *et al.*, 2000). Therefore, the aims of this study were to

investigate: (1) whether physical dormancy plays a role in the germination of *G. coronaria* seeds; and (2) whether physiological dormancy plays a role in the germination of *G. coronaria* seeds. The two heteromorphs were both evaluated separately in this investigation.

Materials and methods

Plant material

Freshly matured cypselae were collected from *G. coronaria* plants growing in natural populations in an abandoned field at road margins in the province of Catania (37°32'49.65"N, 14°56'45.60"E), 252 m above sea level in June 2013 and used within 1 month. After cleaning, cypselae were separated into two collections stored at different conditions: one seed lot, used only for afterripening experiments, was kept in paper bags in the laboratory at ambient conditions [19–24°C, 25–50% relative humidity (RH)], while the other lot was maintained at 15% RH and a temperature of 15°C and it was used for all the other tests.

Cypselae germination tests

Germination experiments consisted, where not otherwise specified, of three samples of 50 cypselae each sown on two layers of moist filter paper (Whatman No. 1) in 90-mm-diameter Petri dishes. Central and peripheral heteromorphs were always tested separately. To reduce mould proliferation during germination tests, cypselae were washed with 1% commercial chlorine and then rinsed three times in distilled water before sowing. Filter papers were soaked in double-distilled water, and extra water was added when needed during the experiments to maintain the moisture of the paper and cypselae. Experiments were conducted in temperature ($\pm 1^\circ\text{C}$) and light ($40 \mu\text{mol m}^{-2} \text{s}^{-1}$) controlled conditions using a 8/16 h daily thermo- and photo-period (=light hereafter).

For analysing the germination response of fresh matured cypselae, the two heteromorphs were immediately cleaned after collection and imbibed at constant temperatures of 5, 10, 15, 20 or 25°C and alternating temperatures of 15/5 or 20/10°C in light for 100 d. Germination in constant darkness was examined by wrapping Petri dishes in two aluminium-foil layers, kept inside a thick-textured black bag for 100 d.

The effect of afterripening on germination response was tested by storing cypselae at ambient conditions (see above) for 0, 4, 12 or 18 weeks and then imbibing them in light at constant temperature regimes of 5, 10, 15 or 20°C, or alternating temperatures of 15/5 or 20/10°C. Cold stratification treatment was carried out

by sowing fresh matured central and peripheral cypselae at 5°C for 0, 4, 8 or 12 weeks in constant darkness. After each cold stratification period, cypselae were incubated in light at the constant temperature regimes of 5, 10, 15, 20 or 25°C or at an alternating temperature regime of 15/5 or 20/10°C. Warm stratification treatment was applied by imbibing fresh matured central and peripheral cypselae at 35/20°C for 0, 3, 5 or 8 weeks in constant darkness with 8/16 h of alternating temperatures. After each warm stratification period, cypselae were incubated in light at the constant temperature of 5, 10, 15 or 20°C or alternating temperatures of 15/5 or 20/10°C.

In all experiments, dishes were examined for germinated seeds daily during the first 2 weeks and with an interval of at least 2 d subsequently. For dark experiments, germination was checked once at the end of the test. Seeds of Asteraceae species are characterized by a thin seed coat and a reduced or absent endosperm within a thick pericarp (Boesewinkel and Boumann, 1995; Debeaujon *et al.*, 2007). Thus, protrusion of the radicle >1 mm through the seed coat was the criterion for seed germination. All emerged seedlings were counted and removed from the dish. At the end of the test, seeds from ungerminated cypselae were cut and checked for viability on the basis of embryo appearance, paying special attention to colour and turgor. Those cypselae that looked viable were tested further for viability using the tetrazolium test with a 1.0% (w/v) solution; defective seeds (i.e. empty, damaged and infected) were excluded from all calculations. Seed coat rupture was always observed after the rupture of the fruit coat (FCR), which disclosed the seed inside (see supplementary Fig. S1). This event was also recorded during germination tests and used for subsequent data analysis. To examine the influence of high temperatures on any physiological dormancy, intact cypselae were sown in continuous darkness at constant (35, 40, 45 or 50°C) and 8/16 h alternating (35/20, 40/20, 45/20 or 50/20°C) regimes for 5 d and the FCR and seed death rate were recorded.

Water uptake and pericarp scarification

Water imbibition of *G. coronaria* cypselae was monitored under six different conditions: (1) intact cypselae, in order to test whether the fruit coats were permeable to water; (2) excised seeds (from dry cypselae) to examine seed coat permeability; (3) only pericarp tissue (from dry cypselae), to analyse the influence of fruit coat during imbibition; and (4–6) different blocking conditions, where either (4) basal, (5) apical or (6) the entire pericarp surface except for the basal region was blocked with Super Glue® (de Souza *et al.*, 2012), in order to localize the fruit coat portion that contributes most to water uptake. In all

the conditions, the weight increase of four samples of 25 tissues per group [cypselae in (1), (4), (5) and (6); fruit coats in (3); and seeds in (2)] was monitored with a Sartorius A200S four-place electronic precision balance (Sartorius, Göttingen, Germany). Weight measurements were taken of dry seeds and, subsequently, after various imbibition times at 15°C (8/16 h photoperiod) at regular time intervals up to 72 h. Water uptake (%) was calculated as [(final mass – initial mass)/initial mass] × 100.

To further investigate the permeability of pericarp structure, intact cypselae from both heteromorphs were immersed for 1, 2, 4, 8, 16, 24 or 48 h in a 1% methylene blue solution (Sigma–Aldrich, Milan, Italy) at 15°C (8/16 h photoperiod), as previously described by Orozco-Segovia *et al.* (2007). Afterwards, at each time interval, central and peripheral cypselae were dissected and the staining pattern of intact seeds (still enveloped by intact testa) was compared with that of longitudinally sectioned seeds under a dissecting microscope (Olympus szx12) equipped with a digital camera (Olympus u-tv0.5xc-3; Olympus, Tokyo, Japan), to monitor any internal staining of seeds.

To determine whether the pericarp inhibits seed germination mechanically, for both heteromorphs three samples of 50 seeds were excised from the pericarp and imbibed in light at constant temperatures (5, 10, 15, 20, 25 or 30°C). A control experiment was carried out by sowing intact cypselae under the same conditions. Due to the higher speed of germination in this test, dishes were examined twice a day until radicle protrusion. At the end of the test period, the number of days to reach 50% of the total germination (t_{50}) was calculated. Furthermore, to analyse the role of pericarp integrity, the germination response was tested after different degrees of pericarp scarification. Four samples of 25 cypselae of both heteromorphs were scarified at the apical end (where cotyledons are located), at the basal end (where the radicle is located) or by removing a lateral half of the pericarp tissue. Scarified cypselae were subsequently imbibed and incubated in light at 10°C for 3 weeks.

Pericarp characterization

To estimate the relevant contribution of the pericarp tissue and the seed mass to the total cypselae mass, 100 randomly selected dry cypselae of each type were weighed, with a Sartorius A200S four-place electronic precision balance, before and after pericarp removal. To investigate the anatomical structure of the pericarp tissue, 50 randomly selected cypselae from each cypselae type were cut longitudinally and transversely. Afterwards the samples were dehydrated through a graded series of ethanol and then placed on an aluminium stub, sputter-coated with gold plate (100 Å) and observed

Table 1. Mean final germination (%) of fresh matured peripheral and central cypselae incubated in light (8/16 h photo- and thermo-period) and in darkness for 100 d under different temperature regimes. Upper and lower 95% binomial confidence intervals are reported in parentheses

	Central cypselae in light	Central cypselae in dark	Peripheral cypselae in light	Peripheral cypselae in dark
5°C	3.3 (2.4–4.9)	0	0.7 (0.7–3.4)	0
10°C	1.7 (1.5–4.4)	1.3 (1.1–3.3)	0	0
15°C	1.6 (1.4–4.2)	0	0	0
20°C	0	0	0	0
25°C	1.9 (1.6–4.8)	0	1.9 (1.5–3.7)	0
30°C	0	0.6 (0.6–2.8)	0	1.3 (1.1–3.4)
15/5°C	12.6 (4.8–6.4)	9.8 (4.2–5.8)	0	7.3 (3.6–5.4)
20/10°C	3.3 (2.2–4.2)	4.7 (2.7–4.7)	1.3 (1.1–3.4)	4.0 (2.5–4.5)

by a Zeiss EVO10-LS environmental scanning electron microscope (Zeiss, Göttingen, Germany). To study the pericarp tissue composition, longitudinal and transverse sections were made from 50 randomly selected cypselae from each cypselae type. Pericarp sections were stained in a solution of 1% phloroglucinol, and 1N HCl was added until tissue sections acquired a pink/red-purplish colouring. Images were taken with a camera mounted on a Primo Star – Carl Zeiss microscope at 40× magnification.

Data analysis

We analysed the effects of treatments on final germination, fruit coat rupture (FCR) and seed mortality by fitting factorial generalized linear models (GLMs, logit link function, binomial distribution) to the germination data (or total FCR events or number of dead seeds) with incubation conditions (mean temperature and temperature regimes) and pre-treatments (dry afterripening, cold stratification, warm stratification), where present, as predictors. We fitted full models with all variables pooled together and then we excluded some of them based on the Akaike Information Criterion (AIC). Alternating and constant temperatures were contrasted. Mean temperature was included as a continuous variable. Significant differences in germination of scarified cypselae were tested with the χ^2 -test. Analysis of variance (ANOVA) was used to analyse the cypselae, pericarp and seed mass. All measurements were obtained from viable seeds and analysed using the R environment for statistical computing (R Development Core Team, 2013).

Results

Effect of temperature and light on the germination of fresh cypselae

Central and peripheral cypselae did not display a significant difference in the germination response; they

occasionally reached 12% and 7% germination respectively, at 15/5°C, but generally germinated to less than 5% (Table 1). The GLM model showed a significant improvement ($P < 0.001$) of germination at alternating regimes compared with constant temperatures, especially with cooler mean temperatures (15/5°C; $P < 0.01$). Overall, the presence of light produced a small but significant ($P < 0.05$) increase in germination of central cypselae. In contrast, peripheral cypselae germinated better in the dark ($P < 0.001$). In all germination tests conducted, FCR at the basal end of the cypselae was always observed to precede seed coat rupture. Seed viability was always above 90% for fresh matured cypselae in all of the tested conditions.

Effect of afterripening and stratification treatments on dormancy release

GLM analysis revealed that, both for central and peripheral cypselae, afterripening did not produce any significant improvement of germination, reaching 15.3, 15.6 and 17.3% for central cypselae and 1.2, 2.0 and 1.9% for peripheral cypselae respectively, after 4, 12 and 18 week-long treatments across the temperature treatments (Fig. 1A, Table 2). Cold stratification had a negative coefficient in the GLM model since it significantly inhibited ($P < 0.001$) germination of cypselae (Fig. 1B, Table 2). Cypselae germination reached 7.1, 3.1 and 3.2% for central cypselae and 2.0, 2.0 and 0.67% for peripheral cypselae after 4, 8 and 12 week-long treatments, respectively. In contrast, warm stratification treatment had a highly significant positive effect ($P < 0.0001$). Indeed, after warm stratification, cypselae germination reached 14.3, 46.6 and 57.6% respectively, after 3, 5 and 8 week-long treatments for central cypselae, whereas for peripheral morphs 29.0, 37.2 and 44.2% germination was obtained respectively, after the same time intervals (Fig. 1C, Table 2). Warm stratification led to a highly significant ($P < 0.0001$) promotion of fruit-coat rupture for both heteromorphs, with the FCR percentage significantly




Figure 1. Mean final germination (%) of central and peripheral cypselae (white bars and dark bars respectively) and mean final FCR (fruit coat rupture) (%) of central and peripheral cypselae (circles and triangles respectively). (A) Afterripened cypselae for short-, intermediate- and long-term treatments (4, 12, 18 weeks, respectively) and imbibed in light (8/16 h photoperiod) at constant and alternating temperatures. (B) Cold-stratified cypselae for short-, intermediate- and long-term treatments (4, 8, 12 weeks, respectively) and imbibed in light (8/16 h photoperiod) at constant and alternating temperatures. (C) Warm-stratified cypselae for short-, intermediate- and long-term treatments (3, 5, 8 weeks, respectively) and imbibed in light (8/16 h photoperiod) at constant and alternating temperatures. Error bars indicate upper and lower 95% binomial confidence intervals. Germination control data (no treatment) are provided in Table 1.

increasing ($P < 0.0001$) with the applied stratification time period. Incubation at higher temperatures for 5 d, regardless of the analysed heteromorph, produced a highly significant increase of FCR events and seed mortality with higher temperatures ($P < 0.0001$; Fig. 2).

Water uptake

Imbibition of intact cypselae showed a significant ($P < 0.001$) mass increase after 1 h for both heteromorphs. After 48 h of imbibition the weight increase attained a stable plateau, while after 72 h it reached

43.0 and 38.0% for central and peripheral cypselae respectively (Fig. 3). Excised seeds showed a significantly ($P < 0.0001$) higher capability to take up water compared with intact cypselae, with increased water content of 54.1 and 52.1% respectively, in central and peripheral seeds (Fig. 3). Pericarp tissues retained a significantly ($P < 0.0001$) lower amount of water compared with intact cypselae, increasing their weight by 17.2 and 12.7% respectively, for pericarp derived from central and peripheral cypselae (Fig. 3). Blocking the apical region of pericarp surface resulted only in a slightly significant ($P = 0.036$ for central and $P = 0.029$ for peripheral cypselae) weight decrease after 72 h

Table 2. Generalized linear model fitted to germination data comparing mean temperature, alternating temperature contrasted with mean temperature, afterripening, cold stratification and warm stratification treatments. AIC, Akaike Information Criterion; SE, standard error; Z, Wald statistic

Model	Effect	Coefficient	SE	Z	P value
AIC = 815.87	Intercept	-2.003	0.197	-10.118	<0.001
	Mean temperature	-0.055	0.014	-3.928	<0.001
	Mean temperature × alternating temperatures	-0.080	0.024	-3.306	<0.001
	Afterripening	-0.018	0.030	-0.613	0.539
	Cold stratification	-0.356	0.057	-6.156	<0.001
	Warm stratification	0.246	0.034	7.226	<0.0001

(reaching 41.7% for central and 36.0% for peripheral cypselae; Fig. 3). However, cypselae of both heteromorphs that were blocked in the basal region of the pericarp showed a highly significant ($P < 0.0001$) limited water uptake, achieving only 24.0 and 17.0% respectively, in central and peripheral cypselae (Fig. 3). On the other hand, blocking the entire pericarp surface except for basal region produced an increase in weight of 34.7 and 27.0% for central and peripheral cypselae respectively, which was significantly higher ($P < 0.0001$) than the cypselae with the blocked basal region, but at the same time significantly lower ($P < 0.0001$) than the intact cypselae (Fig. 3).

Excised seeds from both cypselae heteromorphs stained with methylene blue showed seed coat coloration after 1 h at the basal tip. This spread progressively towards the apical end (Fig. 4), showing no evident difference in the colouring between the two heteromorphs. Embryo coloration, as shown in longitudinally sectioned seeds, showed a considerable delay compared with seed coat coloration, as demonstrated in intact seeds. Methylene blue penetration into the seed coat was observed at the radicle end of seeds 8 h after onset of imbibition, while in the embryo after 16 h of imbibition, and it was complete after 48 h (Fig. 4).




Figure 2. Mean final fruit-coat rupture (white bars) and seed mortality (dark bars) of central (A, B) and peripheral (C and D) cypselae incubated in continuous darkness, at constant (A, C) and alternating regimes (B, D), for 5 d (no germination was observed). Error bars indicate upper and lower 95% binomial confidence intervals.




Figure 3. Mean water uptake during 72 h at 15°C (8/16 h photoperiod) after sowing central cypselae/seed (A) and peripheral cypselae/seed (B) with different parts of the pericarp blocked. Data are means of four samples of 25 seeds; error bars indicate standard deviation.

Effect of pericarp on germination response

Regardless of the morphotype, excising seeds from fruit tissue resulted in a highly significant ($P < 0.0001$) improvement of germination percentage (Table 3). In particular at 5, 10, 15, 20 and 25°C cypselae germination always exceeded 90%. At 30°C seeds excised from central and peripheral morphs germinated to 84.9% and 89.3% respectively. Values of germination speed (t_{50}) indicated a faster response at the temperature range between 20 and 30°C (Table 3).

Scarification of the pericarp at the apical end for both cypselae heteromorphs did not produce any significant increase of germination with respect to intact cypselae. In contrast, after removal of a lateral portion of the pericarp, a highly significant improvement ($P < 0.0001$) in cypselae germination was observed, up to 59% for central and 67% for peripheral

cypselae. Scarification at the basal end induced only a moderately significant ($P < 0.01$) germination increase in both central and peripheral cypselae to 9 and 4% respectively.

Analysis of the masses of seed, pericarp tissue and whole cypselae in both heteromorphs revealed that the pericarp tissue is the main component of the total cypselae mass. Furthermore, ANOVA analyses showed that, even though the cypselae weight of the two heteromorphs was highly significantly different ($P < 0.001$), the weight of seeds excised from central and peripheral morphs (1.10 ± 0.26 mg and 1.90 ± 0.46 mg respectively) was not significantly different. On the other hand, the pericarp mass of the two heteromorphs showed a highly significant ($P < 0.001$) difference, 3.49 ± 0.41 mg and 8.53 ± 0.62 mg for central and peripheral cypselae respectively.




Figure 4. Intact seeds (enveloped by the seed coat) and two seed halves, after manual seed extraction from central (left figure) and peripheral (right figure) cypselae that were stained in methylene blue solution (1%) for different time intervals, as indicated in the figure. For each sampling time point, one intact seed and two longitudinal seed halves are shown, the latter two demonstrate the stain penetration into the embryo.

Table 3. Mean final germination (%) and days to reach 50% of total germination (t_{50}) of excised seeds from central and peripheral cypselae incubated in light under different temperature regimes. For germination results upper and lower 95% binomial confidence intervals, and standard error for time to achieve 50% of the final germination level, are reported in parentheses

Test condition	Central seeds germination (%)	Central seeds t_{50}	Peripheral seeds germination (%)	Peripheral seeds t_{50}
5°C	100	7.2 (± 0.1)	94.0 (5.0–3.2)	7.8 (± 0.5)
10°C	100	5.3 (± 0.2)	100	5.3 (± 0.2)
15°C	100	2.4 (± 0.4)	99.3 (2.9–0.6)	2.0 (± 0.4)
20°C	100	1.6 (± 0.2)	100	1.5 (± 0.2)
25°C	100	0.9 (± 0.2)	100	1.6 (± 0.7)
30°C	84.9 (6.6–5.2)	1.5 (± 0.2)	89.3 (6.0–4.4)	0.9 (± 0.1)

Anatomical characterization of cypselae pericarp

In both morphotypes, the exocarp consisted of flat epidermal cells with an indistinctly reticulate ornamentation, which presents glands irregularly spread in the intercostal gaps (exocarpic region between two longitudinal prominences). The inner endocarp surface of the two heteromorphs generally showed a similar structure (Figs 5A and 6A). In contrast, remarkable differences were observed in the mesocarp tissue thickness between the two morphotypes. The central cypselae mesocarp is formed, in the lateral portion, by 8–10 layers of flattened sclereid cells with a total thickness from 200 μm up to 550 μm (in the winged area), whereas in the basal part the mesocarp is 200–250 μm thick and it is mainly made up of isodiametric cells. Peripheral cypselae presented a broader mesocarp that consisted of 12–18 layers of flattened sclereid cells with a lateral thickness from 300 μm up to 1.8 mm (in the wing area), and 400–500 μm in the basal area, which is formed of isodiametric cells (Figs 5B and 6B). High-magnification images of the pericarp tissue showed the presence of xylem vessels with a lateral pseudoscalariform wall pitting (Carlquist, 2001) (Figs 5C and 6C), which form a channel-like structure at the basal end, running from the endocarp through the external mesocarp (Figs 5B and 6B). At the apical end, bundles of vessel elements were found but they remained in the internal part of the mesocarp (Figs 5D and 6D). In the median portion of the cypselae pericarp, only tangential vessel elements were observed (Figs 5E and 6E). Transverse sections of the above-mentioned pericarp region confirmed that vessels remain in the mesocarpic tissue without entering into the seed cavity. Further anatomical characterization by light microscopy highlighted the presence of brachysclereid-lignified cells in the mesocarp area, more flattened at the internal portion (Figs 5F and 6F), while transverse sections showed the presence of lumen and pits in these brachysclereid-lignified cells.

Discussion

Freshly matured cypselae of *G. coronaria* with high viability demonstrated very limited germination when sown at a range of temperature conditions. Obviously, some form of dormancy is present. Even though heterocarpy in Asteraceae has often been associated with differences in germination behaviour (McEvoy, 1984; Brändel, 2007), in this study the germination response of the two morphotypes was not significantly different under the tested conditions. Afterripening and cold stratification treatments did not considerably improve germination, whereas warm-stratified cypselae displayed a marked improvement, with a higher response after longer treatments. A similar trend was observed for fruit-coat rupture frequency, which had an even higher value at supra-optimal stratification temperatures of 35°C or higher, where, conversely, extensive cypselae mortality was observed. Regardless of the cypselae morphotypes, progressive scarification of the pericarp tissue resulted in germination promotion that is likely to be a consequence of the loss of mechanical constraint against radicle protrusion exerted by the pericarp. In this perspective, warm stratification seems to act as a scarification agent, loosening the pericarp constriction and allowing radicle emergence after incubation at cooler regimes.

Anatomical analyses of central and peripheral cypselae revealed the absence in seed covers of a palisade layer(s) of macrosclereid cells, which is a characteristic trait of physical dormancy (*sensu* Baskin and Baskin, 2004), whereas a point of water entry was found, located at the basal end of the cypselae. Both morphotypes exhibited a considerable and rapid mass increase after sowing in water, with the latter entering mainly through the basal tip, as shown in Fig. 3. The staining experiments with methylene blue solution confirmed the direction of water penetration, and showed that the seed coat delays embryo imbibition, which is complete 48 h after sowing seeds in water.




Figure 5. Anatomical characterization of central cypselae of *G. coronaria* observed by scanning electron microscopy (A, B, C, D, E) and by light microscopy (F). (A) Micrograph of a longitudinal section of a central cypselum. (B) Magnification of the basal end of the pericarp. Arrowheads indicate xylem-vessel bundles running through the pericarp tissue into the seed cavity. (C) Detail of a longitudinal section of a vessel bundle; insert: transverse section of vessels. (D) Inside view of apical cypselum pericarp; insert: transverse section of apical portion of pericarp showing only tangential vessels in mesocarpy. (E) Transversal section of pericarp median portion showing only tangential vessels in mesocarpy. (F) Longitudinal section of pericarp stained with phloroglucinol, showing lignified brachysclereid cells in the mesocarpy; insert: detail of transverse section stained with phloroglucinol, showing lignified brachysclereid cells in the mesocarpy. Abbreviations: Ex, exocarpy; M, mesocarpy; Bs, brachysclereid cells; En, endocarpy; V, vessels; Ve, vessel element.

Further examination revealed that the seed mass is similar in the two heteromorphs and the difference in fruit weight is mainly due to the pericarp tissue. In fact, anatomical analyses revealed a similar composition of mesocarpy, which is thicker in peripheral cypselae than

in the central ones. Moreover both morphotypes share the presence of a channel-like structure composed of a xylem-vessel system, which connects the seed within the cypselum with the external environment. The channel-like structure runs from the basal end, where




Figure 6. Anatomical characterization of peripheral cypsels of *G. coronaria* observed by scanning electron microscopy (A, B, C, D, E) and by light microscopy (F). (A) Micrograph of a longitudinal section of a peripheral cypsel. (B) Magnification of the basal end of the pericarp. Arrowheads indicate xylem-vessel bundles running through the pericarp tissue into the seed cavity. (C) Detail of a longitudinal section of a vessel bundle (arrowheads). (D) Inside view of apical cypsel pericarp from a longitudinal section; insert: transverse section of apical portion of pericarp showing only tangential vessels in mesocarpy. (E) Transversal section of pericarp median portion showing only tangential vessels in mesocarpy. (F) Longitudinal section of pericarp stained with phloroglucinol, showing lignified brachysclereid cells in the mesocarpy; insert: detail of transverse section stained with phloroglucinol, showing lignified brachysclereid cells in the mesocarpy. Abbreviations: Ex, exocarpy; M, mesocarpy; Bs, brachysclereid cells; En, endocarpy; V, vessels; Ve, vessel element.

vessel bundles allow water entrance throughout the pericarp towards the seed part where the radicle is located. In addition, vessel elements located in the apical and lateral portions and brachysclereid cells, spread across the mesocarpy, allow a further water

supply through lateral parts of the pericarp. This system forms a gateway through the pericarp and allows a similar degree of seed imbibition for both cypsels types. Consequently, a similar germination response is observed under laboratory conditions.

Nevertheless, unlimited water availability and full imbibition of cypselae are very unusual in the natural environment of this species (Doussi and Thanos, 2002; Aguado *et al.*, 2011; Carta *et al.*, 2013). Under natural soil conditions, with limited and unstable water soil saturation, central morphs bearing a thinner pericarp may germinate earlier, while viable peripheral ungerminated cypselae are more prone to be seasonally incorporated in the soil seed bank (Bastida *et al.*, 2010). Rapid loss of viability at 35°C or higher under non-limiting water availability suggests that exclusion of water in the summer is essential for survival of cypselae in the soil. Therefore, the hard pericarp that restricts the water passage represents an essential tool in controlling seed germination and survival upon occasional rains, thus providing an ecological advantage.

We conclude that in *G. coronaria* the pericarp plays a decisive role in controlling seed germination by exerting a mechanical constraint against radicle protrusion. Warm stratification, mimicking the environmental conditions experienced by cypselae at the end of summer, overcomes mechanical (physiological) dormancy by weakening the hard pericarp scaffold, resulting in a higher number of fruit-coat rupture events and, as a consequence, in higher germinability. Warm-stratified cypselae respond to a broad range of germination temperatures, although the optimum seems to be 10°C. Even though physical dormancy was not observed, water uptake through the lateral part, as well as the apical end, of the pericarp was very limited. A point of water entry was found located only at the basal end of the pericarp, resulting in longer times for water to reach the apical end of the seed within the intact cypselae and thus contributing to temporal spread in germination.

Supplementary material

To view supplementary material for this article, please visit <http://dx.doi.org/10.1017/S0960258515000148>

Acknowledgements

We are grateful to Dr Girolamo Fichera and Professor Antonia Cristaudo (Department of Biological, Geological and Environmental Science, University of Catania) respectively, for scanning electron microscopy analyses and for support in light microscopy. Moreover, we would like to thank Professor Cristina Salmeri (University of Palermo) for her advice on anatomical features.

Financial support

This work was financed with founding provided by Department of Biological, Geological and

Environmental Science, University of Catania and supported by the Millennium Seed Bank Project, Kew.

Conflicts of interest

None.

References

- Aguado, M., Martínez-Sánchez, J.J., Reig-Arminana, J., García-Breiño, F.J., Franco, J.A. and Vicente, M.J. (2011) Morphology, anatomy and germination response of heteromorphic achenes of *Anthemis chrysantha* J. Gay (Asteraceae), a critically endangered species. *Seed Science Research* **21**, 283–294.
- Bañón, S., Martínez-Sánchez, J.J., Vicente, M.J., Conesa, E., Franco, J.A. and Fernández, J.A. (2009) Effect of applying commercial gibberellins on seed germination of *Chrysanthemum coronarium* L. *Acta Horticulturae* **813**, 435–439.
- Baskin, C.C. and Baskin, J.M. (1998) *Seeds: Ecology, biogeography and evolution of dormancy and germination*. San Diego, Academic Press.
- Baskin, J.M. and Baskin, C.C. (2004) A classification system for seed dormancy. *Seed Science Research* **14**, 1–16.
- Baskin, J.M., Baskin, C.C. and Li, X. (2000) Taxonomy, anatomy and evolution of physical dormancy in seeds. *Plant Species Biology* **15**, 139–152.
- Baskin, J.M., Lu, J.J., Baskin, C.C. and Tan, D.Y. (2013) The necessity for testing germination of fresh seeds in studies on diaspore heteromorphism as a life-history strategy. *Seed Science Research* **23**, 83–88.
- Bastida, F. and Menéndez, J. (2004) Germination requirements and dispersal timing in two heterocarpic weedy Asteraceae. *Communications in Agricultural and Applied Biological Sciences* **69**, 67–76.
- Bastida, F., Gonzáles-Andújar, J.L., Monteagudo, F.J. and Menéndez, J. (2010) Aerial seed bank dynamics and seedling emergence patterns in two annual Mediterranean Asteraceae. *Journal of Vegetation Science* **21**, 541–550.
- Boesewinkel, F.D. and Bouman, F. (1995) The seed: structure and function. pp. 1–24 in Kigel, J.; Galili, G. (Eds) *Seed development and germination*. New York, Marcel Dekker.
- Brändel, M. (2007) Ecology of achene dimorphism in *Leontodon saxatilis*. *Annals of Botany* **100**, 1189–1197.
- Carlquist, S. (2001) *Comparative wood anatomy: systematic, ecological, and evolutionary aspects of dicotyledon wood*. New York, Springer.
- Carta, A., Bedini, G., Müller, J.V. and Probert, R.J. (2013) Comparative seed dormancy and germination of eight annual species of ephemeral wetland vegetation in a Mediterranean climate. *Plant Ecology* **214**, 339–349.
- Cook, R. and Talley, T.S. (2014) The invertebrate communities associated with a *Chrysanthemum coronarium*-invaded coastal sage scrub area in Southern California. *Biological Invasions* **16**, 365–380.
- Corkidi, L., Rincon, E. and Vazquez-Yanes, C. (1991) Effects of light and temperature on germination of heteromorphic achenes of *Bidens odorata* (Asteraceae). *Canadian Journal of Botany* **69**, 574–579.
- Cousens, R.D., Young, K.R. and Tadayyon, A. (2010) The role of the persistent fruit wall in seed water regulation

- in *Raphanus raphanistrum* (Brassicaceae). *Annals of Botany* **105**, 101–108.
- de Souza, T.V., Voltolini, C.H., Santos, M. and Silveira Paulilo, M.T.** (2012) Water absorption and dormancy-breaking requirements of physically dormant seeds of *Schizolobium parahyba* (Fabaceae – Caesalpinioideae). *Seed Science Research* **22**, 169–176.
- Debeaujon, I., Lepiniec, L., Pourcel, L. and Routaboul, J.M.** (2007) Seed coat development and dormancy. pp. 25–49 in Bradford, K.J.; Nonogaki, H. (Eds) *Seed development, dormancy and germination*. Oxford, Blackwell Publishing.
- Doussi, M.A. and Thanos, C.A.** (2002) Ecophysiology of seed germination in Mediterranean geophytes. 1. *Muscari* spp. *Seed Science Research* **12**, 193–201.
- Forsyth, C. and Brown, N.A.C.** (1982) Germination of the dimorphic fruits of *Bidens pilosa* L. *New Phytologist* **90**, 151–164.
- Hawes, C., Begg, G.S., Squire, G.R. and Iannetta, P.P.M.** (2005) Individuals as the basic accounting unit in studies of ecosystem function: functional diversity in shepherd's purse, *Capsella*. *Oikos* **109**, 521–534.
- Imbert, E.** (2002) Ecological consequences and ontogeny of seed heteromorphism. *Perspectives in Plant Ecology, Evolution and Systematics* **5**, 13–36.
- Lloyd, D.G.** (1984) Variation strategies of plants in heterogeneous environments. *Biological Journal of the Linnean Society* **21**, 357–385.
- Lu, J., Tan, D., Baskin, J.M. and Baskin, C.C.** (2010) Fruit and seed heteromorphism in the cold desert annual ephemeral *Diptychocarpus strictus* (Brassicaceae) and possible adaptive significance. *Annals of Botany* **105**, 999–1014.
- Mandák, B.** (2003) Germination requirements of invasive and non-invasive *Atriplex* species: a comparative study. *Flora* **198**, 45–54.
- McDonough, W.T.** (1975) Germination polymorphism in *Grindelia squarrosa*. *Northwest Science* **49**, 190–200.
- McEvoy, P.B.** (1984) Dormancy and dispersal in dimorphic achenes of tansy ragwort *Senecio jacobaea*. *Oecologia* **61**, 160–168.
- Meusel, H., Jäger, E. and Weinert, E.** (1965) *Comparative chorology of the Central European flora*. Jena, Gustav Fischer.
- Orozco-Segovia, A., Márquez-Guzmán, J., Sánchez-Coronado, M.E., De Buen, A.G., Baskin, J.M. and Baskin, C.C.** (2007) Seed anatomy and water uptake in relation to seed dormancy in *Opuntia tomentosa* (Cactaceae, Opuntioideae). *Annals of Botany* **99**, 581–592.
- Picó, F. and Koubek, T.** (2003) Inbreeding effects on fitness traits in the heterocarpic herb *Leontodon autumnalis* L. (Asteraceae). *Acta Oecologica* **24**, 289–294.
- Pignatti, S.** (1982) *Flora d'Italia*. Bologna, Edagricole.
- R Development Core Team** (2013) *R: A language and environment for statistical computing*. Vienna, Austria, R Foundation for Statistical Computing.
- Rocha, O.J.** (1996) The effects of achene heteromorphism on the dispersal capacity of *Bidens pilosa* L. *International Journal of Plant Sciences* **157**, 316–322.
- Schütz, W., Milberg, P. and Lamont, B.B.** (2002) Seed dormancy, after-ripening and light requirements of four annual Asteraceae in south-western Australia. *Annals of Botany* **90**, 707–714.
- Skordilis, A. and Thanos, C.A.** (1995) Seed stratification and germination strategy in the Mediterranean pines *Pinus brutia* and *P. halepensis*. *Seed Science Research* **5**, 151–160.
- Sun, H.Z., Lu, J.J., Tan, D.Y., Baskin, J.M. and Baskin, C.C.** (2009) Dormancy and germination characteristics of the trimorphic achenes of *Garhadiolus papposus* (Asteraceae), an annual ephemeral from the Junggar Desert, China. *South African Journal of Botany* **75**, 537–545.
- Tanowitz, B.D., Salopek, P.F. and Mahall, B.E.** (1987) Differential germination of ray and disc achenes in *Hemizonia increscens* (Asteraceae). *American Journal of Botany* **74**, 303–312.
- Toorop, P.E., Cuerva, R.C., Begg, G.S., Locardi, B., Squire, G.R. and Iannetta, P.P.** (2012) Co-adaptation of seed dormancy and flowering time in the arable weed *Capsella bursa-pastoris* (shepherd's purse). *Annals of Botany* **109**, 481–489.
- Venable, D.L., Burquez, A., Corral, G., Morales, E. and Espinosa, F.** (1987) The ecology of seed heteromorphism in *Heterosperma pinnatum* in central Mexico. *Ecology* **68**, 65–76.
- Wang, Z.B., Chen, Y.F. and Chen, Y.H.** (2009) Functional grouping and establishment of distribution patterns of invasive plants in China using self-organizing maps and indicator species analysis. *Archives of Biological Sciences* **61**, 71–78.