

## CROPS AND SOILS RESEARCH PAPER

# Seed production of barnyardgrass (*Echinochloa crus-galli*) in response to time of emergence in cotton and rice

M. V. BAGAVATHIANNAN\*<sup>1</sup>, J. K. NORSWORTHY<sup>1</sup>, K. L. SMITH<sup>2</sup> AND P. NEVE<sup>3</sup>

<sup>1</sup> Department of Crop, Soil and Environmental Sciences, University of Arkansas, 1366 W Altheimer Drive, Fayetteville, AR 72704, USA

<sup>2</sup> University of Arkansas – Cooperative Extension Service, P.O. Box 3508, Monticello, AR 71656, USA

<sup>3</sup> Crop and Environmental Sciences, Warwick HRI, University of Warwick, Wellesbourne CV35 9EF, UK

(Received 5 July 2011; revised 20 October 2011; accepted 2 November 2011;  
first published online 29 November 2011)

## SUMMARY

The spread of herbicide resistance in barnyardgrass (*Echinochloa crus-galli* (L.) Beauv.) poses a serious threat to crop production in the southern United States. A thorough knowledge of the biology of barnyardgrass is fundamental for designing effective resistance-management programmes. In the present study, seed production of barnyardgrass in response to time of emergence was investigated in cotton and rice, respectively, in Fayetteville and Rohwer, Arkansas, over a 2-year period (2008–09). Barnyardgrass seed production was greater when seedlings emerged with the crop, but some seed production was observed even if seedlings emerged several weeks after crop emergence. Moreover, barnyardgrass seed production was highly variable across environments. When emerging with the crop (0 weeks after crop emergence (WAE)), barnyardgrass produced c. 35 500 and 16 500 seeds/plant in cotton, and c. 39 000 and 2900 seeds/plant in rice, in 2008 and 2009, respectively. Seed production was observed when seedlings emerged up to 5 WAE (2008) or 7 WAE (2009) in cotton and up to 5 WAE (2008, 2009) in rice; corresponding seed production was c. 2500 and 1500 seeds/plant in cotton, and c. 14 700 and 110 seeds/plant in rice, in 2008 and 2009, respectively. The results suggest that cultural approaches that delay the emergence of barnyardgrass or approaches that make the associated crop more competitive will be useful in integrated management programmes. In the context of herbicide resistance management, it may be valuable to prevent seed return to the seedbank, irrespective of cohorts. The findings are vital for parameterizing herbicide resistance simulation models for barnyardgrass.

## INTRODUCTION

The time of weed emergence is an important consideration for successful weed management. Because competition for above and below ground resources can affect the growth and development of weeds, individuals that emerge during the early crop growth stages have the ability to compete well with crops (Gibson *et al.* 2002). However, the degree of such interaction might vary considerably across species and environments (Clay *et al.* 2005). A greater understanding of specific weed–crop interaction as a function of the time of weed emergence will aid the formulation of effective weed management strategies. Several attempts have been made to document the time of emergence for many

agriculturally important weed species (Blackshaw *et al.* 1981; Ciuberkis *et al.* 2007). Traditionally, these studies have aimed at quantifying the effect of weed emergence timing on the growth and yield of crops, and consequently, management attempts were typically focused on reducing crop yield loss, while little emphasis was given to limiting weed seed production.

Seed production is an important determinant of long-term weed population dynamics (Davis *et al.* 2003), and weed management programmes that do not aim beyond a single growing season will probably be ineffective (Gallandt 2006). In addition, the increased spread of herbicide resistance in weed populations warrants a strict focus on limiting weed seed production and seedbank size. Resistance-management programmes are likely to fail if seedbank renewal of resistant individuals is not completely arrested.

\* To whom all correspondence should be addressed. Email: muthu@uark.edu

Managing herbicide-resistant weeds has become one of the top challenges to weed management programmes in the USA, and this is forcing weed scientists to revisit current weed management approaches. In this respect, an understanding of weed biology is critical for devising effective weed management strategies (Van Acker 2009; Gressel 2011). Most farmers will benefit from a greater understanding of the seed production ecology of specific weed species.

Barnyardgrass (*Echinochloa crus-galli* (L.) Beauv.) is a serious weed in major cropping systems in the southern USA, including rice and cotton. Because barnyardgrass is a C<sub>4</sub> weed, it is capable of competing well with C<sub>3</sub> crop plants; it will compete even better under conditions of higher temperatures and limited water availability, scenarios projected to be caused by climate change (Rodenburg *et al.* 2011). It is noteworthy that barnyardgrass infestation can cause yield reductions ranging from 30 to 100% in rice (Johnson *et al.* 1998) and from 21 to 97% in cotton (Keeley & Thullen 1991). Moreover, in the USA, barnyardgrass has evolved resistance to several herbicide families (Heap 2011). The effectiveness of barnyardgrass management programmes rely on the knowledge of its dynamics, including its reproductive potential as affected by the time of emergence relative to crops. Such knowledge will be useful for manipulating crop competition as a valuable tool in integrated weed management programmes (Gibson *et al.* 2002).

According to Mitich (1990), barnyardgrass can produce up to 1 million seeds/plant under ideal growing conditions, but seed production can be highly plastic depending on local growing conditions, nutrient availability and day length (Maun & Barrett 1986). Moreover, the fecundity of barnyardgrass can be severely affected by competition from the associated crop (Lindquist & Kropff 1996; Gibson *et al.* 2003); the extent of competition depends on the time of weed emergence relative to the crop (Cowan *et al.* 1998) and on the crop type (Clay *et al.* 2005). In Ontario, Canada, Bosnic & Swanton (1997) investigated the seed production of barnyardgrass in maize and reported that at a density of 10 plants/m<sup>2</sup>, barnyardgrass produced up to 34 600 seeds/m<sup>2</sup> when it emerged by the 3-leaf stage of maize, whereas only 2800 seeds/m<sup>2</sup> were produced when it emerged after the 4-leaf stage. In Greece, at a similar density (5–10 barnyardgrass plants/m of maize row), barnyardgrass produced up to 1300 seeds/plant when it emerged with the crop and c. 170 seeds/plant when emergence was delayed till the 6-leaf stage of maize (Travlos *et al.* 2011). Likewise, in South Dakota,

USA, barnyardgrass (density: 1.3 plants/m<sup>2</sup>) seed production in maize ranged from 3385 seeds/plant (planted prior to maize emergence) to 158 seeds/plant (planted at the 2-leaf stage of maize) (Clay *et al.* 2005). In rice, barnyardgrass seed production ranged from 2800 seeds/plant when it emerged with the crop to c. 100 seeds/plant when it emerged 45 days after rice emergence (Chauhan & Johnson 2010). In soybean, however, Clay *et al.* (2005) reported that barnyardgrass failed to produce any mature seeds.

These evidences suggests that barnyardgrass seed production is highly variable across crops and environments, yet no such investigations have been carried out in the Midsouth of the USA, where barnyardgrass is an important weed. The objective of the present study was to establish the relationships between barnyardgrass time of emergence and reproductive potential in rice and cotton, two of the most economically important crops in Arkansas, USA.

## MATERIALS AND METHODS

### Experimental set-up and data collection

Field experiments were conducted in the summer of 2008 and 2009 at Rohwer (33°49'N, 91°16'W) and Fayetteville (36°06'N, 94°10'W), Arkansas, USA, to determine the impact of relative time of emergence on the reproduction of barnyardgrass in rice (Rohwer) and cotton (Fayetteville). The soil types of Rohwer and Fayetteville were a Sharkey clay and Taloka silt loam, respectively (NRCS 2010). Rainfall data (Fig. 1) were obtained from the permanent weather units installed in respective experimental stations.

The timing of important agronomic and experimental activities for rice and cotton are given in Tables 1 and 2. Cotton (variety: Stoneville 4554 B2RF) was seeded in rows 1 m wide at a seeding rate of 150 seeds/m of row. Rice (variety: Wells) was also seeded in rows with row spacing of 190 mm and c. 80 seeds/m of row. In Arkansas rice production, emergence of barnyardgrass ceases when fields are flooded at about 6 weeks after crop emergence (WAE), while emergence of barnyardgrass could be observed up to 7 WAE in cotton (J. K. Norsworthy, unpublished results). To represent this, barnyardgrass cohorts were established at weekly intervals from 0 to 5 WAE in rice and from 0 to 7 WAE in cotton. At 0 WAE, barnyardgrass was planted such that it would emerge in synchrony with the crop as much as possible. After planting, barnyardgrass seeds were watered

Table 1. *Timing of important agronomic and experimental activities for rice*

No.	Activity	Year	
		2008	2009
1	Seeding rice	5 May	11 May
2	Barnyardgrass planting (0 WAE)	5 May	11 May
3	First irrigation (subsequent irrigations as required)	5 May	11 May
4	Barnyardgrass planting (1 WAE)	12 May	18 May
5	Barnyardgrass planting (2 WAE)	19 May	26 May
6	Barnyardgrass planting (3 WAE)	26 May	1 Jun
7	Barnyardgrass planting (4 WAE)	2 Jun	8 Jun
8	Barnyardgrass planting (5 WAE)	9 Jun	15 Jun
9	Flooding	16 Jun	23 Jun
10	Bagging barnyardgrass panicles	1 Aug	5 Aug
11	Termination of flooding	13 Aug	20 Aug
12	Harvesting barnyardgrass samples	28 Aug	1 Sep

WAE: weeks after crop emergence.

Table 2. *Timing of important agronomic and experimental activities for cotton*

No.	Activity	Year	
		2008	2009
1	Cotton planting	8 May	11 May
2	First irrigation (subsequent irrigations as required)	8 May	11 May
3	Barnyardgrass planting (0 WAE)	10 May	12 May
4	Barnyardgrass planting (1 WAE)	16 May	19 May
5	Barnyardgrass planting (2 WAE)	23 May	26 May
6	Barnyardgrass planting (3 WAE)	30 May	2 Jun
7	Barnyardgrass planting (4 WAE)	6 Jun	9 Jun
8	Barnyardgrass planting (5 WAE)	13 Jun	16 Jun
9	Barnyardgrass planting (6 WAE)	20 Jun	23 Jun
10	Barnyardgrass planting (7 WAE)	27 Jun	30 Jun
11	Bagging barnyardgrass panicles	19 Aug	14 Aug
12	Harvesting barnyardgrass samples	3 Oct	9 Oct

WAE: weeks after crop emergence.




Fig. 1. Daily precipitation in Rohwer (rice) and Fayetteville (cotton) in 2008 and 2009.

immediately to ensure timely emergence. Standard production practices for the southern USA were used for rice (Slaton 2001) and cotton (UGA 2008). Cotton was irrigated as required using an overhead-sprinkler irrigation system. Rice was initially grown as an upland crop with irrigation as necessary and the fields were permanently flooded from about 6 WAE until maturity.

The current study was conducted in a randomized complete block design (RCBD) with four replications.

The treatments (i.e. time of emergence of barnyardgrass) were randomly assigned to each plot (1 m<sup>2</sup> area) and each replication consisted of seven plots in cotton and five plots in rice (one treatment per plot). The barnyardgrass seeds used in the present study came from a naturally occurring barnyardgrass population in Stuttgart, AR, USA. The seeds were collected in autumn 2007 and stored in a cold room at 4 °C until used in the experiment. In each plot, two barnyardgrass plants were established in such a way that

intra-specific competition was negligible. Ten barnyardgrass seeds per hill (two hills in total) were manually planted at c. 50 mm from the crop row in cotton and c. 100 mm from the row in rice. Over half the seeds germinated and the emerged seedlings were thinned to two plants per hill at the two-leaf stage and then to one plant at the three-leaf stage, allowing for a single barnyardgrass to grow further at each hill. In 2009, it was not possible to establish a barnyardgrass stand at 3 WAE in rice due to waterlogging.

The plots were kept free from other weeds throughout the season by the application of glyphosate in cotton (glyphosate-resistant) and propanil/fenoxaprop in rice. Glyphosate (Roundup™) was applied at 1525 g a.i./ha at 1-leaf, 4-leaf and 8-leaf stages. In rice, propanil (Riceshot™) was applied at 4480 g a.i./ha at the 2- to 3-leaf stage and at the 4- to 5-leaf stage. Propanil (as above) was tank mixed with fenoxaprop (Ricestar™) at 90 g a.i./ha. Prior to spraying, all emerged barnyardgrass plants were covered with plastic containers. Two representative panicles per plant were randomly selected prior to seed maturation and covered with mesh bags to capture the shattering seeds. At maturity, the seed heads were clipped and the seeds from each panicle were removed and counted. The total number of panicles and seeds/plant were calculated for each sampled plant from each plot. The experimental set-up in 2009 was the same as in 2008.

#### Data analysis

Data were analysed using the Statistical Analysis Software (SAS) version 9.1 (SAS Institute, 2003). Regression models are appropriate for establishing the relationship between predicted and predictor variable(s) obtained from time series experiments; therefore, data on the number of panicles/plant and seed production/plant were regressed against the time of emergence of barnyardgrass in cotton and rice. The NLIN procedure of SAS was used in this regard. The regression model that provided the best fit to the data was chosen for each crop (i.e. cotton and rice), because the aim was to understand the response of barnyardgrass when emerging under each of these crops and environments, not to establish an overall relationship across the crops/environments.

In non-linear models, a measure closely corresponding to the coefficient of determination (known as pseudo- $R^2$ ) is used for expressing the quality of fit (Chism *et al.* 1992). Therefore, the best model fit was

identified based on the pseudo- $R^2$  value that was calculated using Eqn (1) (Schabenberger *et al.* 1999). Means were plotted for each emergence time for each year and crop.

$$\text{Pseudo-}R^2 = 1 - \left( \frac{\text{Residual sum of square}}{\text{Total sum of square [corrected]}} \right) \quad (1)$$

The reproductive variables of barnyardgrass (number of panicles and seeds/plant) relative to the time of emergence in rice were non-linearly regressed using a logarithmic function (Eqn 2).

$$y = y^0 + a \ln(x) \quad (2)$$

where  $y$  is the predicted variable (number of panicles or seeds/plant),  $y^0$  is the initial value that is equal to the value of  $y$  when the weed emerges with the crop,  $a$  is the regression coefficient and  $x$  is the time of weed emergence, presented as WAE.

In cotton, the reproductive attributes of barnyardgrass (number of panicles and seeds/plant) relative to the time of emergence was regressed using an exponential function (Eqn 3).

$$y = ae^{-bx} \quad (3)$$

where  $y$  is the predicted variable (number of panicles or seeds/plant),  $a$  represents the reproductive potential of barnyardgrass when it emerges with the crop,  $e$  is the exponent,  $b$  is a fitted constant and  $x$  is the time of weed emergence (WAE).

In each crop, the differences in panicles/plant and seed production/plant between the 2 years of study were examined using a sum of square reduction test (two-curve comparison) (Schabenberger *et al.* 1999). To do this, full and reduced models were fitted wherein the reduced model was a constrained version of the full model. The test statistic  $F_{\text{obs}}$  was calculated as given in Eqn (4).

$$F_{\text{obs}} = \frac{\text{SS(Residual)}_{\text{Reduced}} - \text{SS(Residual)}_{\text{Full}}}{\text{DF(Residual)}_{\text{Reduced}} - \text{DF(Residual)}_{\text{Full}}} / \frac{\text{MS(Residual)}_{\text{Full}}}{\text{DF(Residual)}_{\text{Full}}} \quad (4)$$

where SS is the sum of squares, DF is degrees of freedom and MS is the mean square. The calculated  $F_{\text{obs}}$  was compared with the cut-offs from an  $F$  distribution considering  $\text{DF(Residual)}_{\text{Reduced}} - \text{DF(Residual)}_{\text{Full}}$  as numerator and  $\text{DF(Residual)}_{\text{Full}}$  as denominator DF. The analysis showed significant year effects ( $P \leq 0.05$ ) on the reproduction of barnyardgrass; therefore, data were analysed and presented separately for each year. However, statistical comparisons of barnyardgrass

reproduction were not made between the two crops due to location differences (cotton was grown in Fayetteville and rice in Rohwer).

## RESULTS

The reproductive attributes (number of panicles/plant and number of seeds/plant) of barnyardgrass generally declined for each delay in emergence relative to the crop, but some seed production was still observed when barnyardgrass emerged several weeks after cotton or rice. For each year and crop, barnyardgrass panicle and seed production followed a similar trend except for cotton in 2009, where greater panicle production was not reflected in seed production (Fig. 2). Panicle and seed production, however, differed between the two study years (Figs 2 and 3). In particular, the seed production of barnyardgrass was lower in 2009 compared with 2008, with 54% reduction in cotton and 92% reduction in rice at 0 WAE.

Overall, seed production was observed when barnyardgrass emerged up to 5 WAE (2008) or 7 WAE (2009) in cotton and up to 5 WAE (2008 and 2009) in rice. The coefficients for reproductive attributes of barnyardgrass conformed to an exponential relationship for cotton and logarithmic relationship for rice (Figs 2 and 3). In general, seed production in barnyardgrass was greater in the seedlings that emerged with the crop (i.e. 0 WAE) compared with the later-emerging cohorts, and the decline in seed production was very prominent when the seedlings emerged after 3 WAE (Figs 2 and 3).

Barnyardgrass seed production at the last cohort was reduced by 93 and 91% (2008 and 2009, respectively) in cotton, and by 62 and 92% (2008 and 2009, respectively) in rice, compared with the seedlings that emerged at 0 WAE. In cotton, barnyardgrass produced 35 500 (in 2008) or 16 500 (in 2009) seeds/plant at 0 WAE and 2500 (in 2008) or 1500 (in 2009) seeds/plant at 5 WAE; in rice, barnyardgrass produced 39 000 (in 2008) or 2900 (in 2009) seeds/plant at 0 WAE and 14 750 or 110 seeds/plant at 5 WAE, respectively in 2008 and 2009.

## DISCUSSION


The present results support previous findings that barnyardgrass seed production reduces with delayed emergence relative to the crop (Bosnic & Swanton 1997; Clay *et al.* 2005; Chauhan & Johnson 2010; Travlos *et al.* 2011). Generally, the barnyardgrass seed

production/per plant observed in the present study were greater than that of other similar studies, possibly because the present study had only two barnyardgrass plants in a 1 m<sup>2</sup> area (whereas the densities were greater and variable in other studies), leading to greater seed output on a per plant basis.


When emerging simultaneously with the crop, barnyardgrass would be expected to experience less competition for resources compared with cohorts that emerged later, allowing for more successful growth and reproduction. Conversely, when emerging after the crop was already established, barnyardgrass seedlings would be expected to experience greater competition from the crop. Previous studies have indicated that competition for light, nutrients and moisture, as well as day length, are the major factors that typically affect the seed production of later-emerging weed seedlings (Maun & Barrett 1986; Lindquist & Kropff 1996; Gibson *et al.* 1999, 2002; Gibson & Fischer 2001; Clay *et al.* 2005).

Shading by the crop canopy is an important mechanism of interference between crops and weeds. Jha & Norsworthy (2009) have shown that crop canopy formation typically reduces the amount and quality of light filtering through the canopy, and the growth and reproduction of shaded plants are severely affected through phytochrome-mediated processes (Ballare *et al.* 1990). Moreover, it appears that the effect of shading is more severe on seedlings that emerge later than it is on cohorts that emerge early. For instance, Gibson *et al.* (1999) demonstrated that when it emerged with rice, *Echinochloa* compensated for shading via plasticity in leaf morphology and biomass partitioning. As a result, competition for nutrients was also required to affect growth and reproduction in early emerging cohorts.

For later cohorts, however, competition both above and below ground would have been severe. Assemat *et al.* (1981) and Perera *et al.* (1992) showed that root competition plays an important role in interference between rice and barnyardgrass. As a result, early establishment confers a competitive advantage to the crop over the weeds. The effect of day length would have been minimal in the present study because barnyardgrass cohorts were established before July, allowing them to produce sufficient vegetative growth prior to the onset of short-day conditions in early autumn, which triggers flowering in this species (Maun & Barrett 1986). However, a non-competition control, which was not included in the present experiment, would have been helpful to explain the effects of crop



**Fig. 2.** Regression curve for barnyardgrass panicle and seed production at different times of emergence in cotton in 2008 and 2009. The data conformed to an exponential relationship ( $y = a e^{-bx}$ ), where  $a$  is the initial value that starts the exponential function and  $b$  is the fitted constant. The quality of the model fit was expressed using the pseudo- $R^2$  value.



**Fig. 3.** Regression curve for barnyardgrass panicle and seed production at different times of emergence in rice in 2008 and 2009. The data conformed to a logarithmic relationship ( $y = y^o + a \ln(x)$ ), where  $y^o$  is the initial value that corresponds to the value of  $y$  when the weed emerges with the crop and  $a$  is the regression coefficient. The quality of the model fit was expressed using the pseudo- $R^2$  value.

competition and day length on barnyardgrass seed production.

The relationships between time of emergence and seed production of barnyardgrass have been described using separate models for cotton and rice based on the quality of model fit. Cotton and rice were grown in two

different locations with different environmental conditions; therefore, it was difficult to separate the effect of crop type alone on barnyardgrass seed production. The differences observed between these two systems could have been due to a combined effect of crop type, soil type, management (particularly planting density

and row spacing) and local growing conditions. Studies have indicated that competitive interactions are greater within dicots or monocots rather than between them (Toler *et al.* 1996; Cowan *et al.* 1998). Nevertheless, crop types that can exhibit a rapid canopy formation may confer a competitive advantage over the weeds.

The significant reduction in barnyardgrass seed production in 2009 compared with 2008 could have been due to the changes in the extent of inter-specific competition as influenced by environmental conditions. However, it is not clear what combination of environmental factors/crop  $\times$  environment interactions caused the variability between the years, and the study was not intended to capture this. In 2009, high levels of early-season precipitation (in May; Fig. 1) slowed the establishment of cotton, the growth of which is known to be severely affected by prolonged wet conditions (Bange *et al.* 2004). Barnyardgrass, on the other hand, grows well under wet conditions (Rahn *et al.* 1968) compared with cotton, which led to more tillers and thereby more panicles/plant, but seed production/panicle was reduced as cotton growth recovered soon after.

In rice, a different situation occurred in 2009. Intensive early-season precipitation (Fig. 1) and a near-submerged condition (intermittently) favoured the early growth of rice. Although barnyardgrass continued to grow under submerged conditions, the plants grew upright with low seed production, and the results concur with Rahn *et al.* (1968). In 2008, the rice fields were submerged only following the establishment of flood at 6 WAE, which may explain the greater seed production in the early emerging cohorts in 2008 compared with 2009. A non-competition control would have been helpful to explain the causes for these differences, yet the results suggest that creating a submerged condition during the early growth stages of rice may be a useful strategy for combating barnyardgrass.

The present study provides important data on the reproductive ecology of barnyardgrass. However, the effects were highly variable over the years and environments, leading to a range of parameter values. This is not surprising because barnyardgrass seed production has been shown to be highly variable across environments (Holm *et al.* 1977; Maun & Barrett 1986; Norris 1992; Clay *et al.* 2005), and the present study was not designed for or aimed at exploring the causes for such variation. Nevertheless, it provided an opportunity to understand the extent of environmental stochasticity associated with

barnyardgrass seed production, which is highly valuable in parameterizing the population dynamic models for barnyardgrass. System- or environment-specific data may be vital if such models are sensitive to this parameter. Given the variability observed in barnyardgrass seed production within a confined geographical region, future studies aiming at predicting seed production as a function of crop  $\times$  environmental interactions will be useful.

The present study also provides valuable information that can be used to devise effective weed management practices. Barnyardgrass seedlings that emerge within a month of crop emergence contribute most to the seed rain and enriching the seedbank. Therefore, greater emphasis is warranted on early season weed control. The results are in accordance with those of Cowan *et al.* (1998), Clay *et al.* (2005) and Travlos *et al.* (2011). Nevertheless, in the context of herbicide resistance management, preventing seed production and seedbank renewal of the resistant individuals is the key, irrespective of their time of emergence. The results also suggest that substantial weed control can be attained simply by delaying the timing of weed emergence relative to the crop and/or by making the crop more competitive. There are opportunities for breeding crop cultivars with desirable agronomic traits, perhaps by exploiting the existing intra-cultivar variation (Tokatlidis *et al.* 2011). Additionally, production practices including but not limited to deep tillage, flooding (for rice), and altering planting date, row spacing and planting density can be utilized to improve crop competitiveness over weeds.

The authors are grateful to Bayer CropScience, Dow AgroSciences, Syngenta, Monsanto, Valent U.S.A. and BASF for the financial support to collect biological data and model the evolution of herbicide resistance in barnyardgrass.

## REFERENCES

- ASSEMAT, L., MORISHIMA, H. & OKA, H. (1981). Neighbor effects between rice (*Oryza sativa* L.) and barnyardgrass (*Echinochloa crus-galli* Beauv.) strains. II. Some experiments of interaction between plants. *Acta Oecologica Plantarum* **2**, 63–78.
- BANGE, M. P., MILROY, S. P. & THONGBAI, P. (2004). Growth and yield of cotton in response to waterlogging. *Field Crops Research* **88**, 129–142.
- BLACKSHAW, R. E., STOBLE, E. H. & STURKO, A. R. W. (1981). Effects of seeding dates and densities of green foxtail (*Setaria viridis*) on the growth and productivity of spring wheat (*Triticum aestivum*). *Weed Science* **29**, 212–217.

- BOSNIC, A. & SWANTON, C. J. (1997). Influence of barnyardgrass (*Echinochloa crus-galli*) time of emergence and density on corn (*Zea mays*). *Weed Science* **45**, 276–282.
- BALLARE, C. L., SCOPEL, A. L. & SANCHEZ, R. A. (1990). Far-red radiation reflected from adjacent leaves: an early signal of competition in plant canopies. *Science* **247**, 329–332.
- CHAUHAN, B. S. & JOHNSON, D. E. (2010). Implications of narrow crop row spacing and delayed *Echinochloa colona* and *Echinochloa crus-galli* emergence for weed growth and crop yield loss in aerobic rice. *Field Crops Research* **117**, 177–182.
- CHISM, W. J., BIRCH, J. B. & BINGHAM, S. W. (1992). Nonlinear regressions for analyzing growth stage and quinclorac interactions. *Weed Technology* **6**, 898–903.
- CIUBERKIS, S., BERNOTAS, S., RAUDONIUS, S. & FELIX, J. (2007). Effect of weed emergence time and intervals of weed and crop competition on potato yield. *Weed Technology* **21**, 213–218.
- CLAY, S. A., KLEINJAN, J., CLAY, D. E., FORCELLA, F. & BATCHELOR, W. (2005). Growth and fecundity of several weed species in corn and soybean. *Agronomy Journal* **97**, 294–302.
- COWAN, P., WEAVER, S. E. & SWANTON, C. J. (1998). Interference between pigweed (*Amaranthus spp.*), barnyardgrass (*Echinochloa crus-galli*), and soybean (*Glycine max*). *Weed Science* **46**, 533–539.
- DAVIS, A. S., DIXON, P. M. & LIEBMAN, M. (2003). Cropping system effects on giant foxtail (*Setaria faberi*) demography: II. Retrospective perturbation analysis. *Weed Science* **51**, 930–939.
- GALLANDT, E. R. (2006). How can we target the weed seedbank? *Weed Science* **54**, 588–596.
- GIBSON, K. D. & FISCHER, A. J. (2001). Relative growth and photosynthetic response of water-seeded rice and *Echinochloa oryzoides* (Ard.) Fritsch to shade. *International Journal of Pest Management* **47**, 305–309.
- GIBSON, K. D., FOIN, T. C. & HILL, J. E. (1999). The relative importance of root and shoot competition between water-seeded rice and *Echinochloa phyllopogon*. *Weed Research* **39**, 181–190.
- GIBSON, K. D., FISCHER, A. J., FOIN, T. C. & HILL, J. E. (2002). Implications of delayed *Echinochloa* spp. Germination and duration of competition for integrated weed management in water-seeded rice. *Weed Research* **42**, 351–358.
- GIBSON, K. D., FISCHER, A. J., FOIN, T. C. & HILL, J. E. (2003). Crop traits related to weed suppression in water-seeded rice (*Oryza sativa* L.). *Weed Science* **51**, 87–93.
- GRESSEL, J. (2011). Global advances in weed management. *Journal of Agricultural Science, Cambridge* **149**, 47–53.
- HEAP, I. (2011). *The International Survey of Herbicide Resistant Weeds*. Online resource available online at <http://www.weedscience.org> (verified 24 October 2011).
- HOLM, L. G., PLUCKNETT, D. L., PANCHO, J. V. & HERBERGER, J. P. (1977). *The World's Worst Weeds: Distribution and Biology*. Honolulu, USA: University Press of Hawaii.
- JHA, P. & NORSWORTHY, J. K. (2009). Soybean canopy and tillage effects on emergence of palmer amaranth (*Amaranthus palmeri*) from a natural seed bank. *Weed Science* **57**, 644–651.
- JOHNSON, D. E., DINGKUHN, M., JONES, M. P. & MAHAMANE, M. C. (1998). The influence of rice plant type on the effect of weed competition on *O. sativa* and *O. glaberrima*. *Weed Research* **38**, 207–216.
- KEELEY, P. E. & THULLEN, R. J. (1991). Growth and interaction of barnyardgrass (*Echinochloa crus-galli*) with cotton (*Gossypium hirsutum*). *Weed Science* **39**, 369–375.
- LINDQUIST, J. L. & KROPFF, M. J. (1996). Applications of an ecophysiological model for irrigated rice (*Oryza sativa*) *Echinochloa* competition. *Weed Science* **44**, 52–56.
- MAUN, M. A. & BARRETT, S. C. H. (1986). The biology of Canadian weeds. 77. *Echinochloa crus-galli* (L.) Beauv. *Canadian Journal of Plant Science* **66**, 739–759.
- MITICH, L. W. (1990). Intriguing world of weeds: barnyardgrass. *Weed Technology* **4**, 918–920.
- NORRIS, R. F. (1992). Case history for weed competition/population ecology: barnyardgrass (*Echinochloa crus-galli*) in sugarbeets (*Beta vulgaris*). *Weed Technology* **6**, 220–227.
- NRCS. (2010). *Keys to Soil Taxonomy*, 11th edition. Natural Resources Conservation Service of the United States Department of Agriculture, Online resource available online at <http://soils.usda.gov/technical/classification/taxonomy/> (verified 31 October 2011).
- PERERA, K. K., AYRES, P. G. & GUNASENA, H. P. M. (1992). Root growth and the relative importance of root and shoot competition in interactions between rice (*Oryza sativa*) and *Echinochloa crus-galli*. *Weed Research* **32**, 67–76.
- RAHN, E. M., SWEET, R. D., VENGRIS, J. & DUNN, S. (1968). *Life History Studies as Related to Weed Control in the Northeast. 5 – Barnyardgrass*. Bulletin 368. Newark, USA: Agricultural Experiment Station University of Delaware.
- RODENBURG, J., MEINKE, H. & JOHNSON, D. E. (2011). Challenges for weed management in African rice systems in a changing climate. *Journal of Agricultural Science, Cambridge* **149**, 427–435.
- SAS INSTITUTE. (2003). *SAS 9.1 for Windows*. Cary, NC: SAS.
- SCHABENBERGER, O., THARP, B. E., KELLIS, J. J. & PENNER, D. (1999). Statistical test for hormesis and effective dosage in herbicide dose-response. *Agronomy Journal* **91**, 713–721.
- SLATON, N. A. (2001). *Rice Production Handbook*. Arkansas Cooperative Extension Service Miscellaneous Publication 192. Little Rock, AR: University of Arkansas.
- TRAVLOS, I. S., ECONOMOU, G. & KANATAS, P. J. (2011). Corn and barnyardgrass competition as influenced by relative time of weed emergence and corn hybrid. *Agronomy Journal* **103**, 1–6.
- TOKATLIDIS, I. S., TSIKRIKONI, C., LITHOURGIDIS, A. S., TSIALTAS, J. T. & TZANTRAMAS, C. (2011). Intra-cultivar variation in cotton: response to single-plant yield selection at low density. *Journal of Agricultural Science, Cambridge* **149**, 197–204.
- TOLER, J. E., GUICE, J. B. & MURDOCK, E. C. (1996). Interference between johnsongrass (*Sorghum halepense*), smooth pigweed (*Amaranthus hybridus*), and soybean (*Glycine max*). *Weed Science* **44**, 331–338.
- UGA. (2008). *Georgia Cotton Production Guide*. Online resource available online at <http://commodities.caes.uga.edu/fieldcrops/cotton/2008cottonguide/2008CottonGuide.htm> (verified 31 October 2011).
- VAN ACKER, R. C. (2009). Weed biology serves practical weed management. *Weed Research* **49**, 1–5.