

## Research Article

\*Contributed equally.

**Cite this article:** Wang J-F, Lin R-H, Zhang X, Hide G, Lun Z-R, Lai D-H (2022). Novel insertions in the mitochondrial maxicircle of *Trypanosoma musculi*, a mouse trypanosome. *Parasitology* **149**, 1546–1555. <https://doi.org/10.1017/S0031182022001019>

Received: 6 March 2022  
Revised: 14 July 2022  
Accepted: 17 July 2022  
First published online: 4 August 2022




**Key words:**

Divergent regions; indels; maxicircle; minicircle; palindrome; *Trypanosoma lewisi*; *Trypanosoma musculi*

**Author for correspondence:**

Zhao-Rong Lun,  
E-mail: [lsslzr@mail.sysu.edu.cn](mailto:lsslzr@mail.sysu.edu.cn);  
De-Hua Lai,  
E-mail: [laidehua@mail.sysu.edu.cn](mailto:laidehua@mail.sysu.edu.cn)

# Novel insertions in the mitochondrial maxicircle of *Trypanosoma musculi*, a mouse trypanosome

Ju-Feng Wang<sup>1,\*</sup> , Ruo-Hong Lin<sup>1,\*</sup>, Xuan Zhang<sup>1</sup>, Geoff Hide<sup>2</sup> ,  
Zhao-Rong Lun<sup>1,2</sup> and De-Hua Lai<sup>1</sup> 

<sup>1</sup>Guangdong Provincial Key Laboratory of Aquatic Economic Animals, State Key Laboratory of Biocontrol, School of Life Sciences, Sun Yat-Sen University, Guangzhou 510275, The People's Republic of China and <sup>2</sup>Ecosystems and Environment Research Centre and Biomedical Research Centre, School of Science, Engineering and Environment, University of Salford, Salford M5 4WT, UK

**Abstract**

*Trypanosoma musculi* is a, globally distributed, mouse-specific haemoflagellate, of the family Trypanosomatidae, which shares similar characteristics in morphology with *Trypanosoma lewisi*. The kinetoplast (mitochondrial) DNA of Trypanosomatidae flagellates is comprised of catenated maxicircles and minicircles. However, genetic information on the *T. musculi* kinetoplast remains largely unknown. In this study, the *T. musculi* maxicircle genome was completely assembled, with PacBio and Illumina sequencing, and the size was confirmed at 34 606 bp. It consisted of 2 distinct parts: the coding region and the divergent regions (DRs, DRI and II). In comparison with other trypanosome maxicircles (*Trypanosoma brucei*, *Trypanosoma cruzi* and *T. lewisi*), the *T. musculi* maxicircle has a syntenic distribution of genes and shares 73.9, 78.0 and 92.7% sequence identity, respectively, over the whole coding region. Moreover, novel insertions in *MURF2* (630 bp) and in *ND5* (1278 bp) were found, respectively, which are homologous to minicircles. These findings support an evolutionary scenario similar to the one proposed for insertions in *Trypanosoma cruzi*, the pathogen of American trypanosomiasis. These novel insertions, together with a deletion (281 bp) in *ND4*, question the role of Complex I in *T. musculi*. A detailed analysis of DRII indicated that it contains numerous repeat motifs and palindromes, the latter of which are highly conservative and contain A<sub>5</sub>C elements. The comprehensively annotated kinetoplast maxicircle of *T. musculi* reveals a high degree of similarity between this parasite and the maxicircle of *T. lewisi* and suggests that the DRII could be a valuable marker for distinguishing these evolutionarily related species.

**Introduction**

Trypanosomes are protozoan parasites that are distributed globally which infect humans, vertebrate animals and intermediate invertebrate hosts. Among them, members of the subgenus *Herpetosoma*, such as *Trypanosoma lewisi* and *Trypanosoma musculi*, are commonly found in rodents (Hoare, 1972; Kostygov *et al.*, 2021). These 2 trypanosomes cannot be easily distinguished due to their high degree of similarity in morphological characteristics and genetic markers such as the SSU rDNA sequences (Hong *et al.*, 2017). However, they do significantly differ in many aspects. In particular, *T. lewisi* infects only rats and sometimes humans (Sarataphan *et al.*, 2007; Verma *et al.*, 2011), while *T. musculi* infects only mice and is unlikely to be pathogenic to humans (Zhang *et al.*, 2018). A few research studies have indicated that both *T. musculi* and *T. lewisi* can modulate the host immune response in coinfections with various other infectious agents, to potentially cause more harm to the hosts, by altering the infection kinetics and increasing the duration of colonization in the host (Lowry *et al.*, 2014; Vaux *et al.*, 2016; Nzoumbou-Boko *et al.*, 2017; Gao *et al.*, 2021). To gain a better understanding of the biological characteristics, the kinetoplast DNA (kDNA) of *T. lewisi* has been comprehensively analysed (Lin *et al.*, 2015; Li *et al.*, 2020). However, little is known of the details of the kDNA in *T. musculi*.

Trypanosomes are members of the kinetoplastea group of protozoa, named due to the presence of the kDNA. Trypanosome kDNA is a specific network structure of interlocking mitochondrial DNA circles, which consists of thousands of minicircles with dozens of maxicircles (Lukes *et al.*, 2002). Earlier research on *Trypanosoma brucei* has shown that kDNA comprises at least 5% of the total cellular DNA, while most other eukaryotic mitochondrial DNA accounts for no more than 1% (Lukeš *et al.*, 2018). In general, kDNA maxicircles encode functional homologues of mitochondrial genes which are flanked by non-coding regions that diverge significantly in sequence and size amongst trypanosome species (Simpson *et al.*, 1987; Sloof *et al.*, 1992; Westenberger *et al.*, 2006). One of the unusual features of the kinetoplast is that most of the maxicircle gene transcripts are not mature and do not encode functional open reading frames. These encrypted transcripts become translatable only after post-transcriptional processing, namely RNA editing, that inserts and deletes uridine residues

(Stuart *et al.*, 1997, 2005). RNA editing was first discovered in cytochrome oxidase subunit 2 (*COII*) gene of *Trypanosoma brucei* and *Crithidia fasciculata*, whose mRNA transcripts have 4 uridine insertions (Benne *et al.*, 1986). The minicircles, recognized by a conserved motif of 12 nucleotides (GGGGTTGGTGTA) (Ray, 1989), encode guide RNA (gRNA) molecules that accurately position the editing machinery to ensure correct maxicircle transcripts are produced (Blum and Simpson, 1990).

Here, using PacBio and Illumina sequencing reads, the complete maxicircle sequence of *T. musculi* was assembled and annotated, including the repetitive non-coding variable region. Comparative analyses indicate that the gene organization and distribution in *T. musculi* maxicircles are highly conserved with *T. brucei*, *T. cruzi* and *T. lewisi*. The maxicircle kDNA gene organization of *T. musculi* and comparison with its species relatives was also presented. In addition, the genetic information on the divergent region (DR) II reveals that it may provide a good marker for molecular diagnosis and molecular epidemiological investigation of trypanosomes.

## Materials and methods

### Parasites, ultrastructure, kDNA extraction and restriction endonuclease digestion

*Trypanosoma musculi* Partinico II strain was gifted by Professor Philippe Vincendeau of Université de Bordeaux, France, which was originally obtained from the London School of Hygiene and Tropical Medicine (Krampitz, 1969). *T. musculi* Partinico 2, Lincicome and CDC strains were purchased from American Type Culture Collection (ATCC). Trypanosomes were harvested from the blood of infected mice and cultured in RPMI-1640 medium at 37°C supplemented with 10% fetal bovine serum (FBS) and a feeder layer of mouse macrophages as modified from Behr (Behr *et al.*, 1990). Protocols for the use of mice were approved by the Institutional Review Board for Animal Care at Sun Yat-Sen University under license 31672276. For transmission electron microscopy, trypanosome specimens were prepared according to the method of Bozzola (Bozzola, 2014), and observed by using the JEM-100CX-II microscope system. For *T. musculi* DNA preparations, total DNA was purified using a phenol–chloroform method and kDNA was extracted by sucrose gradient ultracentrifugation according to previously published methods (Pérez-Morga and Englund, 1993). The isolated kDNA network was visualized on a 1% agarose gel and analysed with restriction enzymes *HindIII*, *EcoRI*, *BamHI*, *RsaI*, *HaeIII* and *TaqI* (New England Biolabs, USA). A computer-simulated restriction enzyme digestion map of *T. musculi* maxicircle was generated using the Dnaman 9.0 software (Lynnon Corporation, Quebec, Canada) based on the sequence assembled in this study.

### Immunofluorescence assay

Trypanosome cells ( $1 \times 10^7$  cells mL<sup>-1</sup>) were centrifuged for 5 min at 3000 × *g* and washed twice in phosphate-buffered saline (PBS). The cells were then transferred onto clean slides, which were left to air-dry in a fume hood, following fixation by methanol for 10 min. Dried slides were rehydrated and washed twice in PBS for 5 min at room temperature. The slides were then incubated with primary mAb-anti-L8C4 (1:800) followed by incubation with Cy3-Conjugated goat anti-mouse IgG (A10521, Thermo Fisher) (1:400) followed by counterstaining consisting of 1 × PBS with 3 μg mL<sup>-1</sup> 4,6-diamidino-2-phenylindole (DAPI) (Kohl *et al.*, 1999). They were then photographed using a Leica fluorescence microscope.

### Deep sequencing, sequence assembly and PCR verification

To generate a high-quality maxicircle assembly, a kDNA Illumina library was constructed and sequenced using Illumina HiSeq2000 technology commercially (Novogene, China). Also, a PacBio Sequel library was constructed using total DNA and sequenced commercially (Annoroad, China). The Illumina reads were checked for quality and trimmed using fastqc (<http://www.bioinformatics.babraham.ac.uk/projects/fastqc/>) and Trimmomatic (Bolger *et al.*, 2014). Canu 2.0 software was used in *de novo* assembly of the *T. musculi* genome with parameter ‘genomeSize = 30 m minReadLength = 600 minOverlapLength = 300 corOutCoverage = 100 corMinCoverage = 2 correctedErrorRate = 0.035’ using the PacBio reads (Koren *et al.*, 2017). Then, the genome contigs, assembled from PacBio reads, were polished using the Illumina reads by Pilon software to improve genome assembly (Walker *et al.*, 2014). Finally, the assembly sequences were aligned, with BLAST, to a previously obtained *T. lewisi* maxicircle assembly (KR072974) and redundant overlap deleted by MEGA 7.0 to yield the complete maxicircle sequence (Camacho *et al.*, 2009; Kumar *et al.*, 2016). To obtain *T. b. rhodesiense*, *Trypanosoma grayi* and *T. lewisi* complete maxicircle genomes, processed reads were assembled from WGS data (SRX3199071, SRX620256 and SRR11918574, respectively) freely available on NCBI using SPAdes 3.12.0 with parameter ‘--plasmid --careful -t 16 -m 200’ (Antipov *et al.*, 2016). Then, alignment and trimming were also carried out as completed for *T. musculi*. The sequences of the *T. musculi* maxicircle coding region were also corrected by PCR verification using 12 pairs of primers. Meanwhile, *MURF2*, *ND4* and *ND5* genes were also amplified in 3 additional *T. musculi* strains (*T. musculi* Partinico 2, Lincicome and CDC). Primers used are summarized in Table S1.

### Gene annotation

Annotation of *T. musculi* maxicircle coding regions was performed by comparison with *T. brucei* (EATRO 427, M94286.1), *T. cruzi* (CL, DQ343645.1) and *T. lewisi* (CPO02, KR072974.1) manually using BLAST. Patterns of RNA editing of *T. musculi* maxicircle genes were predicted according to GC% and RNA editing pattern of *T. lewisi* (Li *et al.*, 2020).

### Data analysis

Dotplot graphs of the *T. musculi* maxicircle sequence plotted against itself and 3 other Trypanosomatidae species were generated by YASS software with default parameter (allow 10% indels, 25% mutations and e-value  $< 1 \times 10^{-5}$  in alignment) (Noé and Kucherov, 2005). GC percentage, assembly coverage and homology search algorithms were drawn using Circos v0.69 (Krzywinski *et al.*, 2009). Regions (>300 bp) with sufficient sequence identity were plotted as coloured ribbons, denoting the percentage of sequence identity. BioEdit software was used to create alignments and calculate nucleotide percentage identity matrices among different Trypanosomatidae species (Hall, 1999). Curation of the palindromes and inserted sequence homology analysis was performed using BLAST. MEME software was used to identify motifs and generate LOGO diagrams in the DR I region (Bailey *et al.*, 2015).

### Phylogenetic analysis

The entire coding region of the kinetoplast maxicircles were aligned using ClustalO 1.2.4 (Sievers *et al.*, 2011) and the alignment was trimmed using Gblocks 0.91b with option ‘-t = d -b4 = 5 -b5 = h’ (Talavera and Castresana, 2007). Maximum likelihood trees were generated by RAxML 8.2.12 with 1000 bootstrap

replicates (Stamatakis, 2014). Neighbour joining and Minimum evolution trees were performed using MEGA 7.0 including 1000 bootstrap pseudo-replicates. Maxicircle genome sequences used in this work are summarized in Table S2.

## Results

### Morphology, ultrastructure, kDNA isolation and restriction enzyme digestion

In culture, *T. muscui* cells tend to attach to each other and form a rosette-like pattern (Fig. 1A), via their flagella, as determined using specific antibodies against the paraflagellar rod (Fig. 1C). At this stage, *T. muscui* is at the epimastigote stage in which the kinetoplast lies closely beside the nucleus. Ultrastructural analysis showed that the kinetoplast DNA disc was  $660 \pm 99$  nm in length and  $152 \pm 26$  nm in width ( $n = 50$ ) (Fig. 1B), which is similar to closely related species such as *T. lewisi* (Lin *et al.*, 2015).


A total of  $10^{10}$  *T. muscui* were harvested and high-quality kDNA was obtained with a 260/280 absorbance ratio of 1.80. Kinetoplast DNA was found to be intact and free from contamination with nuclear or host DNAs as judged by agarose electrophoresis (Fig. S1A). Meanwhile, kDNA was incubated with endonucleases of *HindIII*, *EcoRI*, *BamHI*, *RsaI*, *HaeIII* and *TaqI* and a computer-simulated restriction enzyme digestion map was generated based on the maxicircle assembly which is described later (Figs S1B–C). Some bands smaller than 4.0 kb did not correspond to the computed simulated patterns of *T. muscui* maxicircle, which implies the presence of a high number of possible heterogeneous minicircles in the kDNA of *T. muscui*. The bands consistently observed in Fig. S1B at  $\sim 1.3$  kb suggest the presence of minicircles of a similar size as reported in *T. lewisi* (Li *et al.*, 2020). Patterns that are free of kDNA in the wells were

achieved using *HaeIII* and *TaqI* and indicated a high frequency of cleavage of kDNA minicircles. Most likely, the bands with molecular sizes of  $>4$  kb correlated with the computer-simulated patterns (Fig. S1C), are derived from kDNA maxicircles, except a band ( $\sim 5$  kb) in *RsaI* lane, a potential result of incomplete digestion. Moreover, the presence of 4 large molecular weight bands, in the *EcoRI* digestion, with sizes of  $>10$ ,  $\sim 7$ ,  $\sim 6$  and  $\sim 4$  kb, indicated that the full-size kDNA maxicircle is larger than the sum of 27 kb.

### Assembly and annotation of the kDNA maxicircle


Genomic DNA from the *T. muscui* Partinico II strain was sequenced on PacBio Sequel and Illumina platforms and contigs were assembled with the long-read assembler Canu 2.0 and corrected with the Illumina reads in Pilon. Then, a contig in length of 38 603 bp was identified in a BLAST search against the *T. lewisi* maxicircle (KR072974). This had 2 overlapping regions of 4002 bp (covering positions from  $-3341$  or  $31\ 266$  to  $661$  bp) at each end (Fig. S2) confirming completion of the circle. Meanwhile, the maxicircle sequence has also been confirmed using 5 overlapping raw reads from the PacBio library (Fig. S2) and the maxicircle coding region sequences were also further refined using 12 pairs of primers and Sanger sequencing validation (Fig. S3). Finally, a 34 606 bp-long complete *T. muscui* maxicircle sequence was obtained, with an average coverage of 13.2X from Illumina reads and 268X from PacBio reads, including the coding regions (16 975 bp) and the DRs (17 631 bp). The overall GC content of the maxicircle was 23.7%, with 27.5% in coding regions and 20.1% in DRs (Fig. 2).

Twenty genes were annotated in the *T. muscui* maxicircle by comparison with known Trypanosomatidae species (*T. brucei*,



**Fig. 1.** Morphology and ultrastructure of *T. muscui*. (A) Giemsa staining of the epimastigote form of *T. muscui* from *in vitro* culture (scale bar  $5\ \mu\text{m}$ ). Flagellum (F), nucleus (N) and kinetoplast (K) are indicated. (B) Electron micrograph of the *T. muscui* trypomastigote form (scale bar  $200\ \text{nm}$ ). Basal body (BB), flagellum (F), flagellar pocket (FP) and kinetoplast (K) are indicated. (C) Immunofluorescence analysis (IFA) shows epimastigote-like forms of *T. muscui* from *in vitro* cultivation with antibody L8C4 detecting paraflagellar rods (green) and DAPI detecting kinetoplast DNA and nuclear DNA (blue) (scale bars  $5\ \mu\text{m}$ ).





**Fig. 2.** Circos plot of the *T. muscui* maxicircle. The outer track indicates the gene arrangements and gene distribution; the middle track represents the GC content (orange for above-average and green for below-average) and the inner track is a histogram of assembly coverage.

*T. cruzi* and *T. lewisi*), as listed in Table 1. All genes were found to be syntenic with the maxicircles of the comparator Trypanosomatidae species *T. brucei*, *T. cruzi* and *T. lewisi* (Fig. 2, Fig. S4). A sequence homology analysis (Fig. S5), showed that the *T. muscui* maxicircle has 92.7% identity to *T. lewisi* (blue ribbons). The ribbons change to yellow when compared with *T. cruzi* (78.0% identity) and *T. brucei* (73.9% identity), largely due to the low similarity in extensively edited genes (Table 2). Moreover, 3 breaks shown as discontinuations of lines or ribbons (Figs S4 and S5) appear in *T. muscui* genes *MURF2*, *ND5* and *ND4*, with 2 sections (630 and 1278 bp) inserted and 1 section deleted (281 bp), respectively (Fig. 3A and Table 1).

The confirmation of mutations in *MURF2*, *ND5* and *ND4* was performed with 3 other strains (*T. muscui* Particino 2, Lincicome and CDC). PCR results showed that the insertion in *MURF2* is specific to *T. muscui* Particino II strain and not present in other 3 strains, while the insertion in *ND5* and the deletion in *ND4* exist in all tested strains (Fig. 3B). Furthermore, these insertions and the deletion have also been confirmed by inspecting alignments of the raw reads mapped back to the maxicircle assembly. Alignment analysis of insertions showed that a fragment (150 bp) at the 5' end region of *MURF2* insertion sequence is homologous to both the 5' end and middle regions of *ND5* insertion sequence (Fig. 3C). Moreover, those sequences in *MURF2* and *ND5*, respectively, share 95.3, 96 and 95.3% identity with conserved regions of *T. lewisi* minicircles (MN447336.1, MN447339.1 and MN447386.1), and these 150 bp homologous regions cover 3 conserved sequence blocks (CSBs) of minicircles, indicating a minicircle origin of both insertions in *MURF2* and *ND5*. Together with the data shown in Fig. S1 and Fig. 3C, it seems that, unsurprisingly, *T. muscui* has a similar size and structure to minicircles reported for *T. lewisi* (Li *et al.*, 2020), i.e. ~1.3-kb category I minicircles that have 2 conserved regions

with CSB1-3 motifs (and perhaps also ~1.5 kb category II minicircles with only 1 conserved region; such a band is also apparent in Fig. S1B). It therefore appears that the ~1.3 kb *ND5* insertion corresponds to a (degenerated) category I minicircle and the 630 bp *MURF2* insertion corresponds to half a category I minicircle.

RNA editing patterns of the maxicircle have been well studied in *T. brucei* and *T. lewisi* (Gerasimov *et al.*, 2018; Li *et al.*, 2020) and they are well correlated with GC%. The GC% pattern in *T. muscui* is fairly similar to *T. brucei* and *T. lewisi*. However, unexpectedly high GC contents were noticed in *MURF2* and *ND5* (Fig. S6), and they are precisely attributed to the insertions in both genes. In another region, *COII* and its *cis*-acting gRNA were identified (Table 1).

The whole coding region of the maxicircle is considered as a valuable marker for phylogenetic relationships of Trypanosomatidae species (Kaufer *et al.*, 2019). To further confirm the evolutionary relationship of *T. muscui* and other Trypanosomatidae species, sequences corresponding to whole coding region of *T. muscui* were aligned with the sequences from other Trypanosomatidae species to infer phylogenetic relationships. In the tree, *T. muscui* and *T. lewisi* are identified as belonging to the same subgenus *Herpetosoma*, clustered with the sister groups of *Schizotrypanum* and *Aneza* (Fig. 4).

### Sequence analysis of the maxicircle DRs

A common theme of the maxicircle DR is the presence of various repeat arrays, which is also the case for *T. muscui*. The full map of DR of *T. muscui* was built by the YASS and Circos packages to identify homologous regions and to show global patterns of DR organization (Fig. 5A and B). Dot-plot analyses of the DR showed 2 typical sections (I, II), flanking either the 12S rRNA or *ND5*. DR I is in a length of about 1.6 kb, which is composed of short and

**Table 1.** Gene positions and lengths in the maxicircle of *T. muscili*

Gene	RNA editing	<i>T. muscili</i> position	<i>T. muscili</i> length	<i>T. lewisi</i> length	<i>T. cruzi</i> length	<i>T. brucei</i> length
<b>12S rRNA</b>	–	1–1164	1164	1168	1161	1149
<b>9S rRNA</b>	–	1210–1818	609	608	608	611
<b>ND8</b>	Extensive	1859–2130	272	285	279	266
<b>ND9<sup>a</sup></b>	Extensive	2195–2536	342	350	338	321
<b>uS3m<sup>a</sup></b>	None	2575–2823	249	241	264	234
<b>ND7</b>	Extensive	2874–3647	774	771	755	702
<b>COIII</b>	Extensive	3721–4135	415	414	424	439
<b>Cyb</b>	Minor	4222–5301	1080	1080	1080	1080
<b>ATPase6</b>	Extensive	5342–5644	303	304	336	369
<b>ND2</b>	None	5687–7038	1352	1341	1341	1237
<b>CR3<sup>b</sup></b>	Extensive	6979–7111	~133	~123	~119	/
<b>ND1<sup>a</sup></b>	None	7129–8112	984	942	942	957
<b>COII</b>	Minor	8111–8739	629	629	629	626
<b>COII gRNA</b>	–	8746–8758	–	–	–	–
<b>MURF2</b>	Minor	8767–10 439	1673	1053	1056	1041
<b>(Insertion)</b>	–	8897–9526	630	–	–	–
<b>COI<sup>a</sup></b>	None	10 429–12 081	1653	1650	1650	1734
<b>CR4<sup>a</sup></b>	Extensive	12 102–12 319	218	212	207	185
<b>ND4<sup>c</sup></b>	None	12 428–13 464	1037	1314	1314	1311
<b>ND3<sup>a</sup></b>	Extensive	13 453–13 636	184	187	193	256
<b>RPS12</b>	Extensive	13 711–13 908	198	190	191	172
<b>ND5</b>	None	13 929–16 975	3047	1773	1770	1770
<b>(Insertion)</b>	–	14 923–16 200	1278	–	–	–

Gene positions are shown relative to the start of the 12S rRNA.

<sup>a</sup>These genes are encoded on the reverse strand.

<sup>b</sup>CR3 2 end positions from *T. muscili*, *T. lewisi*, *T. cruzi* and *T. brucei* are uncertain.

<sup>c</sup>A fragment deletion is found in the *T. muscili* ND4 gene.

highly repetitive units of about 107 bp, with 2 motifs being found (Fig. 5C). While DRII is in a length of about 14 kb, it consists of a series of tandem elements, namely  $\alpha$ ,  $\beta$ ,  $\gamma$ ,  $\sigma$ , and short version  $\alpha'$ ,  $\beta'$ ,  $\gamma'$  (Fig. S7).

Palindromes are a typical structure already found in *T. cruzi*, *T. lewisi* and *T. rangeli*. Based on identifying homologues using BLAST, 4 AT-rich conserved palindromes showed up in the DRII (Fig. 5B and D). Palindromes I and IV have the same perfect palindrome structure, 34 bp long, and are located at 19 898 and 28 055 bp. While palindromes II and III have 1 T-to-A substitution, they are located at 23 648 and 26 061 bp. A further BLAST analysis with the maxicircles of *T. b. brucei* (Lister 427, MN904526.1), *T. b. equiperdum* (STIB 818, EU185799.1), *Trypanosoma congolense* (IL3000, GCA\_003013265.1) and

*Trypanosoma vivax* (Y486, MT090068.1) enabled the identification of similar palindromes in these species (Fig. 5D), only 1 of each species is shown for illustrative purposes. These palindromes are highly conserved and contain an A<sub>5</sub>C element.

Unlike highly conserved coding regions, DRs show species specificity among trypanosomes (Fig. S5). It displays about 70% sequence identity in DRI between the *T. muscili* and *T. lewisi* maxicircle, while there are only some similar sequences (~400 bp) in DRII. Moreover, there are no other homologous sequences between *T. muscili* and the other 2 species (*T. cruzi* and *T. brucei*) in the DRs (Fig. S5). Therefore, these results suggest that DRII is highly divergent among trypanosomes, which may have the potential to be a good molecular marker for distinguishing *T. muscili* from related species.

**Table 2.** Comparison of the average percentage identities of *T. muscili* kDNA maxicircle with those from the other 3 Trypanosomatidae species

Comparison of <i>T. muscili</i>	Whole coding region	5'-edited genes	Extensively edited genes	rRNAs	Non-edited genes
vs <i>T. lewisi</i>	92.7%	95.8%	92.4%	95.4%	92.2%
vs <i>T. cruzi</i>	78.0%	84.5%	73.7%	84.0%	78.4%
vs <i>T. brucei</i>	73.9%	84.4%	60.2%	80.5%	74.8%


Entire coding region: starting from 5' end of 12S rRNA to 3' end of ND5.

5'-edited genes: *Cyb*, *COII*.

Extensively edited genes: *ND8*, *ND9*, *ND7*, *COIII*, *ATPase6*, *CR3*, *CR4*, *ND3*, *RPS12*.

Non-edited genes: *uS3m*, *ND2*, *ND1*, *COI*.

*MURF2*, *ND4* and *ND5* genes are not calculated in *T. muscili* (5'-edited genes or non-edited genes) due to insertions/deletion.




**Fig. 3.** Analysis of insertions or deletions in the *T. muscui* maxicircle genes *MURF2*, *ND5* and *ND4*. (A) A diagram of insertions or deletions in the *T. muscui* maxicircle genes *MURF2* (630 bp insertion), *ND5* (1278 bp insertion) and *ND4* (281 bp deletion). Insertion regions are labelled as grey boxes. (B) PCR amplification of the *T. muscui* *MURF2*, *ND5* and *ND4* genes from *T. muscui* Partinico II (1), *T. muscui* Partinico 2 (2), *T. muscui* Lincicome (3), *T. muscui* CDC (4), and fragments are analysed on a 1.0% agarose gel. M, DL2000 marker (Takara, China). N, negative control. The positions of primers are shown in (A) and Table S1. (C) Alignment of the conserved regions from the *T. lewisi* minicircle (MN447335.1) and insertion sequences of *MURF2* and *ND5*. Conserved sequence blocks (CSB-I, II, III) are highlighted in grey.

**Discussion**

In this study, a sequence of the 34 606 bp kDNA maxicircle genome from *T. muscui* was reported and an in-depth investigation of *T. muscui* maxicircle sequences and comparative analysis with

other Trypanosomatidae species were also undertaken. The size of the total coding region of the *T. muscui* kDNA maxicircle is 16 975 bp with 2 pronounced insertions in *T. muscui* *MURF2* (630 bp), *ND5* (1278 bp) and 1 deleted fragment of *ND4*



**Fig. 4.** Phylogenetic analysis of the Trypanosomatidae species using the maxicircle coding regions. Phylogenetic tree is performed based on Maximum likelihood/Neighbour joining/Minimum evolution methods with 1000 bootstrap replicates with the respective bootstrap confidences indicated at branch points. Branch lengths are indicated by the black line and the scale bar represents the number of nucleotide substitutions per site.





as non-edited genes in other known Trypanosomatidae species and it is inconceivable that these large insertions/deletions could be corrected by U-insertion/deletion editing of the mRNAs. Nevertheless, all of the above insertions/deletions are found in *ND4*, *ND5* as well as *ND7* genes, and these genes all encode subunits of the mitochondrial respiratory chain NADH-dehydrogenase (Complex I). Since the presence of a functional Complex I in Trypanosomatidae species has long been debated (Oppendoes and Michels, 2008; Duarte and Tomás, 2014). Deletions in kDNA encoding Complex I subunits were identified in some strains of *T. cruzi* that seem no impact in mitochondrial bioenergetics, ROS production or redox state in this parasite (César Carranza *et al.*, 2009). Although the presence of Complex I and its involvement in respiration has been clearly demonstrated in *T. brucei*, it appears to be non-essential for procyclic forms (Beattie and Howton, 1996; Verner *et al.*, 2011; Surve *et al.*, 2012). The lack of editing in several Complex I subunits in *L. tarentolae* UC strain also suggests that it may not be essential (Simpson *et al.*, 2015). Therefore, the possibility that the role of Complex I subunits is less important in *T. muscui* was favoured with the presence of insertions/deletions in *ND5* and *ND4*. In addition, another insertion occurs in *MURF2*, whose function remains uncertain but hypothesis could be risen. *MURF2* might be a new component in Complex I. The insertion in *MURF2* may be a recent event as it is only found in Partinico II strain, but not other 3 strains of *T. muscui*. The loss of conservation in *MURF2* could probably be attributed to the loss of function of Complex I components and consequently on selection pressures on the gene. To verify this hypothesis, a highly sensitive and accurate identification of the functioning of Complex I in Trypanosomatidae species would be interesting to investigate.

The DR of the kinetoplast maxicircle was initially described as a variable and non-coding region and the DR structure seemed to be drastically different in various species (Borst *et al.*, 1980, 1982; Stuart and Gelvin, 1982; Muhich *et al.*, 1983; Maslov *et al.*, 1984). Therefore, the function of the DRs remains as an enigma. Studies on *T. brucei*, *Crithidia oncopelti*, *Leptomonas collosoma* and *Leishmania seymouri* revealed some CSBs-like sequences in their maxicircle DRs. As CSBs are essential for minicircle replication (Ryan *et al.*, 1988), CSBs-like sequences may play a similar role in maxicircle replication (Gorbat *et al.*, 1990; Sloof *et al.*, 1992; Myler *et al.*, 1993; Flegontov *et al.*, 2006). However, CSB-I or III-like regions were not identified in *T. muscui* DRs, instead, only a CSB-II-like region (CCCGTGT) is located at 19 817 bp. CSB-I or III-like regions were also not found in DRs of the closely related *T. lewisi*, suggesting a CSB-independent maxicircle replication mechanism exists in these species. Therefore, although CSB-like sequences were present in the insertions of *T. muscui* *MURF2* and *ND5*, it is not clear whether they are also involved in maxicircle replication.

It has been demonstrated that hairpins or cruciform structures (palindromes) are frequently associated with promoters and may also act as protein-binding sites (Wadkins, 2000). Palindromes with an A<sub>5</sub>C-element in DRs are suggested as recognition sites for binding of transcription factors or transcription initiation (Vasil'eva *et al.*, 2004; Flegontov *et al.*, 2006). Some palindromes also have been identified in *T. muscui* as well as in a variety of other Trypanosomatidae species, where each consists of 1 A<sub>5</sub>C-element. It may be speculated that these palindromes play a significant role in Trypanosomatidae species maxicircles, judged by their high degree of sequence conservation in the evolution of Kinetoplastida species.

The trypanosome maxicircle presents itself as a complex evolutionary system, and it may be an excellent taxonomic marker in phylogenetic analysis. The coding region of the maxicircle in phylogenetic analyses provides a robust evolutionary insight

into the relationships within Trypanosomatidae species (Lin *et al.*, 2015; Kaufer *et al.*, 2019; Kay *et al.*, 2020). A close affinity between *T. muscui* and *T. lewisi* in *Herpetosoma* was also supported, which clustered with the sister groups of *Schizotrypanum* and *Aneza*. Unlike the highly conserved coding region, the DRs of maxicircle, especially DRII sequence, was found to be significantly divergent and species-specific (Kay *et al.*, 2020). The homologies in DRII between closely related species, e.g., *T. muscui* and *T. lewisi*, phylogenetic clades of *T. cruzi*, are limited (Figs S5 and S8). Such a characteristic of DRII provides an opportunity for developing a valuable molecular marker for distinguishing closely related species and subspecies. Actually, a preliminary test on 3 *T. muscui* strains and 6 *T. lewisi* strains revealed a consistent amplification of DRII fragments, which could enable them to be distinguished from each other and 13 strains of other trypanosomes (Hong *et al.*, 2017).

In general, this study reports the first detailed description and analysis of the kDNA maxicircle genome of *T. muscui* and reveals a relatively high overall conservation of gene content and synteny with other trypanosome species. Furthermore, the divergence of DRII suggests its potential as a valuable marker for distinguishing these evolutionarily related species.

**Supplementary material.** The supplementary material for this article can be found at <https://doi.org/10.1017/S0031182022001019>

**Acknowledgements.** The authors would like to thank Dr Ling-Ling Zheng for help with bioinformatics analysis. Thanks to all the members in the laboratories who provided useful help during the field and laboratory work. We would also like to appreciate the critical comments from the anonymous reviewers which have greatly helped to improve our paper.

**Author's contributions.** J-F Wang, R-H Lin, D-H Lai and Z-R Lun designed the study. J-F Wang, R-H Lin and X Zhang conducted data gathering and performed statistical analyses. J-F Wang, R-H Lin and D-H Lai drafted the manuscript and undertook data extraction and screening. G Hide, ZR Lun and DH Lai critically reviewed the paper. All authors approved the final version and agree to be accountable for all aspects of the work.

**Financial support.** The project was supported by grants from the National Natural Science Foundation of China (31672276, 31720103918) and the Natural Sciences Foundation of Guangdong Province (2022A1515011874).

**Conflict of interest.** The authors declare there are no conflicts of interest.

**Ethical standards.** The animal ethical approval of sample collection was obtained from the Institutional Review Board of Animal Care at Sun Yat-Sen University (License no. 31672276).

**Data availability.** Nucleotide sequence data reported in this paper are available in GenBank databases under accession numbers: *Trypanosoma muscui* maxicircle sequence (OM000218), *Trypanosoma lewisi* maxicircle sequence (OM000219), *Trypanosoma grayi* maxicircle sequence (OM049542), and *Trypanosoma brucei rhodesiense* maxicircle sequence (OM049543). PacBio and Illumina sequencing data have been deposited in NCBI's Sequence Read Archive (SRA) with BioProject ID PRJNA792722.

## References

- Antipov D, Hartwick N, Shen M, Raiko M, Lapidus A and Pevzner PA (2016) plasmidSPAdes: assembling plasmids from whole genome sequencing data. *Bioinformatics (Oxford, England)* **32**, 3380–3387.
- Bailey TL, Johnson J, Grant CE and Noble WS (2015) The MEME suite. *Nucleic Acids Research* **43**, W39–W49.
- Baptista CS, Vêncio RZN, Abdala S, Carranza JC, Westenberger SJ, Silva MN, Pereira CAdb, Galvão LMC, Gontijo ED, Chiari E, Sturm NR and Zingales B (2006) Differential transcription profiles in *Trypanosoma cruzi* associated with clinical forms of Chagas disease: maxicircle NADH dehydrogenase subunit 7 gene truncation in asymptomatic patient isolates. *Molecular and Biochemical Parasitology* **150**, 236–248.
- Beattie DS and Howton MM (1996) The presence of rotenone-sensitive NADH dehydrogenase in the long slender bloodstream and the procyclic



- forms of *Trypanosoma brucei brucei*. *European Journal of Biochemistry* **241**, 888–894.
- Behr MA, Mathews SA and D'Alessandro PA** (1990) A medium for the continuous cultivation of bloodstream forms of *Trypanosoma lewisi* at 37°C. *Journal of Parasitology* **76**, 711–716.
- Benne R, Van den Burg J, Brakenhoff JP, Sloof P, Van Boom JH and Tromp MC** (1986) Major transcript of the frameshifted coxII gene from trypanosome mitochondria contains four nucleotides that are not encoded in the DNA. *Cell* **46**, 819–826.
- Berná L, Greif G, Pita S, Faral-Tello P, Díaz-Viraqué F, Souza RDCMD, Vallejo GA, Alvarez-Valin F and Robello C** (2021) Maxicircle architecture and evolutionary insights into *Trypanosoma cruzi* complex. *PLoS Neglected Tropical Diseases* **15**, e0009719.
- Blum B and Simpson L** (1990) Guide RNAs in kinetoplastid mitochondria have a nonencoded 3' oligo(U) tail involved in recognition of the preedited region. *Cell* **62**, 391–397.
- Bolger AM, Lohse M and Usadel B** (2014) Trimmomatic: a flexible trimmer for Illumina sequence data. *Bioinformatics (Oxford, England)* **30**, 2114–2120.
- Borst P, Fase-Fowler F, Hoeijmakers JH and Frasch AC** (1980) Variations in maxi-circle and mini-circle sequences in kinetoplast DNAs from different *Trypanosoma brucei* strains. *Biochimica et Biophysica Acta (BBA)* **610**, 197–210.
- Borst P, Weijers PJ and Brakenhoff GJ** (1982) Analysis by electron microscopy of the variable segment in the maxi-circle of kinetoplast DNA from *Trypanosoma brucei*. *Biochimica et Biophysica Acta (BBA)* **699**, 272–280.
- Bozzola JJ** (2014) Conventional specimen preparation techniques for transmission electron microscopy of cultured cells. In Kuo J (ed.), *Electron Microscopy: Methods and Protocols*. Totowa, NJ: Humana Press, pp. 1–19.
- Camacho C, Coulouris G, Avagyan V, Ma N, Papadopoulos J, Bealer K and Madden TL** (2009) BLAST+: architecture and applications. *BMC Bioinformatics* **10**, 421.
- César Carranza J, Kowaltowski AJ, Mendonça MAG, de Oliveira TC, Gadelha FR and Zingales B** (2009) Mitochondrial bioenergetics and redox state are unaltered in *Trypanosoma cruzi* isolates with compromised mitochondrial complex I subunit genes. *Journal of Bioenergetics and Biomembranes* **41**, 299–308.
- Duarte M and Tomás AM** (2014) The mitochondrial complex I of trypanosomatids – an overview of current knowledge. *Journal of Bioenergetics and Biomembranes* **46**, 299–311.
- Flegontov PN, Guo Q, Ren L, Strelkova MV and Kolesnikov AA** (2006) Conserved repeats in the kinetoplast maxicircle divergent region of *Leishmania* sp. and *Leptomonas seymouri*. *Molecular Genetics and Genomics* **276**, 322–333.
- Gao JM, Yi SQ, Geng GQ, Xu ZS, Hide G, Lun ZR and Lai DH** (2021) Infection with *Trypanosoma lewisi* or *Trypanosoma muscili* may promote the spread of *Toxoplasma gondii*. *Parasitology* **148**, 703–711.
- Gerasimov ES, Gasparyan AA, Kaurov I, Tichý B, Logacheva MD, Kolesnikov AA, Lukeš J, Yurchenko V, Zimmer SL and Flegontov P** (2018) Trypanosomatid mitochondrial RNA editing: dramatically complex transcript repertoires revealed with a dedicated mapping tool. *Nucleic Acids Research* **46**, 765–781.
- Gorbat A, Maslov DA, Peters LS, Gaviernik P, Viustenkhagen T and Kolesnikov AA** (1990) Analysis of the sequence of repeats in divergent regions of maxi-circular DNA from kinetoplasts of *Crithidia oncopelti*. *Molekuliarnaia Biologiia* **24**, 1539–1548.
- Hall TA** (1999) BioEdit: a user-friendly biological sequence alignment editor and analysis program for Windows 95/98/NT. *Nucleic Acids Symposium Series* **41**, 95–98.
- Hoare CA** (1972) *The Trypanosomes of Mammals. A Zoological Monograph*. Oxford, UK: Blackwell Scientific Publications.
- Hong XK, Zhang X, Fusco OA, Lan YG, Lun ZR and Lai DH** (2017) PCR-based identification of *Trypanosoma lewisi* and *Trypanosoma muscili* using maxicircle kinetoplast DNA. *Acta Tropica* **171**, 207–212.
- Kaufner A, Stark D and Ellis J** (2019) Evolutionary insight into the Trypanosomatidae using alignment-free phylogenomics of the kinetoplast. *Pathogens (Basel, Switzerland)* **8**, 157.
- Kay C, Williams TA and Gibson W** (2020) Mitochondrial DNAs provide insight into trypanosome phylogeny and molecular evolution. *BMC Evolutionary Biology* **20**, 161.
- Kohl L, Sherwin T and Gull K** (1999) Assembly of the paraflagellar rod and the flagellum attachment zone complex during the *Trypanosoma brucei* cell cycle. *Journal of Eukaryotic Microbiology* **46**, 105–109.
- Koren S, Walenz BP, Berlin K, Miller JR, Bergman NH and Phillippy AM** (2017) Canu: scalable and accurate long-read assembly via adaptive -mer weighting and repeat separation. *Genome Research* **27**, 722–736.
- Koslowsky D, Sun Y, Hindenach J, Theisen T and Lucas J** (2014) The insect-phase gRNA transcriptome in *Trypanosoma brucei*. *Nucleic Acids Research* **42**, 1873–1886.
- Kostygov AY, Karnkowska A, Votýpka J, Tashyreva D, Maciszewski K, Yurchenko V and Lukeš J** (2021) Euglenozoa: taxonomy, diversity and ecology, symbioses and viruses. *Open Biology* **11**, 200407.
- Krampitz HE** (1969) Geographical distribution, host-parasite relationship and multiplication of Sicilian strains of *Trypanosoma (Herpetosoma) duttoni* Thiroux 1950 (Protozoa, Trypanosomatidae). *Zeitschrift für Parasitenkunde (Berlin, Germany)* **32**, 297–315.
- Krzywinski M, Schein J, Birol I, Connors J, Gascoyne R, Horsman D, Jones SJ and Marra MA** (2009) Circos: an information aesthetic for comparative genomics. *Genome Research* **19**, 1639–1645.
- Kumar S, Stecher G and Tamura K** (2016) MEGA7: molecular evolutionary genetics analysis version 7.0 for bigger datasets. *Molecular Biology and Evolution* **33**, 1870–1874.
- Li SJ, Zhang X, Lukeš J, Li BQ, Wang JF, Qu LH, Hide G, Lai DH and Lun ZR** (2020) Novel organization of mitochondrial minicircles and guide RNAs in the zoonotic pathogen *Trypanosoma lewisi*. *Nucleic Acids Research* **48**, 9747–9761.
- Lin RH, Lai DH, Zheng LL, Wu J, Lukeš J, Hide G and Lun ZR** (2015) Analysis of the mitochondrial maxicircle of *Trypanosoma lewisi*, a neglected human pathogen. *Parasites & Vectors* **8**, 665.
- Lowry JE, Leonhardt JA, Yao C, Belden EL and Andrews GP** (2014) Infection of C57BL/6 mice by *Trypanosoma muscili* modulates host immune responses during *Brucella abortus* cocolonization. *Journal of Wildlife Diseases* **50**, 11–20.
- Lukes J, Guilbride DL, Votýpka J, Ziková A, Benne R and Englund PT** (2002) Kinetoplast DNA network: evolution of an improbable structure. *Eukaryotic Cell* **1**, 495–502.
- Lukeš J, Wheeler R, Jirsová D, David V and Archibald JM** (2018) Massive mitochondrial DNA content in diplomemid and kinetoplastid protists. *IUBMB Life* **70**, 1267–1274.
- Maslov DA, Kolesnikov AA and Zaitseva GN** (1984) Conservative and divergent base sequence regions in the maxicircle kinetoplast DNA of several trypanosomatid flagellates. *Molecular and Biochemical Parasitology* **12**, 351–364.
- Muhich ML, Simpson L and Simpson AM** (1983) Comparison of maxicircle DNAs of *Leishmania tarentolae* and *Trypanosoma brucei*. *Proceedings of the National Academy of Sciences of the United States of America* **80**, 4060–4064.
- Myler PJ, Glick D, Feagin JE, Morales TH and Stuart KD** (1993) Structural organization of the maxicircle variable region of *Trypanosoma brucei*: identification of potential replication origins and topoisomerase II binding sites. *Nucleic Acids Research* **21**, 687–694.
- Neboháčová M, Kim CE, Simpson L and Maslov DA** (2009) RNA editing and mitochondrial activity in promastigotes and amastigotes of *Leishmania Donovanii*. *International Journal for Parasitology* **39**, 635–644.
- Noé L and Kucherov G** (2005) YASS: enhancing the sensitivity of DNA similarity search. *Nucleic Acids Research* **33**, W540–W543.
- Nzoumbou-Boko R, De Muylder G, Semballa S, Lecordier L, Dauchy FA, Gobert AP, Holzmüller P, Lemesre JL, Bras-Gonçalves R, Barnabé C, Courtois P, Daulouède S, Beschin A, Pays E and Vincendeau P** (2017) *Trypanosoma muscili* infection in mice critically relies on mannose receptor-mediated arginase induction by a TbKHC1 kinesin H chain homolog. *Journal of Immunology* **199**, 1762–1771.
- Oppendoes FR and Michels PAM** (2008) Complex I of Trypanosomatidae: does it exist? *Trends in Parasitology* **24**, 310–317.
- Pérez-Morga D and Englund PT** (1993) The structure of replicating kinetoplast DNA networks. *Journal of Cell Biology* **123**, 1069–1079.
- Ray DS** (1989) Conserved sequence blocks in kinetoplast minicircles from diverse species of trypanosomes. *Molecular and Cellular Biology* **9**, 1365–1367.
- Ryan KA, Shapiro TA, Rauch CA and Englund PT** (1988) Replication of kinetoplast DNA in trypanosomes. *Annual Review of Microbiology* **42**, 339–358.
- Sarataphan N, Vongpakorn M, Nuansrichay B, Autarkool N, Keowkarnkah T, Rodtian P, Stich RW and Jittapalpong S** (2007) Diagnosis of a *Trypanosoma lewisi*-like (*Herpetosoma*) infection in a sick infant from Thailand. *Journal of Medical Microbiology* **56**, 1118–1121.

- Sievers F, Wilm A, Dineen D, Gibson TJ, Karplus K, Li W, Lopez R, McWilliam H, Remmert M, Söding J, Thompson JD and Higgins DG (2011) Fast, scalable generation of high-quality protein multiple sequence alignments using Clustal Omega. *Molecular Systems Biology* **7**, 539.
- Simpson L, Neckelmann N, de la Cruz VF, Simpson AM, Feagin JE, Jasmer DP and Stuart K (1987) Comparison of the maxicircle (mitochondrial) genomes of *Leishmania tarentolae* and *Trypanosoma brucei* at the level of nucleotide sequence. *Journal of Biological Chemistry* **262**, 6182–6196.
- Simpson L, Douglass SM, Lake JA, Pellegrini M and Li F (2015) Comparison of the mitochondrial genomes and steady state transcriptomes of two strains of the trypanosomatid parasite, *Leishmania tarentolae*. *PLoS Neglected Tropical Diseases* **9**, e0003841.
- Sloof P, de Haan A, Eier W, van Iersel M, Boel E, van Steeg H and Benne R (1992) The nucleotide sequence of the variable region in *Trypanosoma brucei* completes the sequence analysis of the maxicircle component of mitochondrial kinetoplast DNA. *Molecular and Biochemical Parasitology* **56**, 289–299.
- Stamatakis A (2014) RAxML version 8: a tool for phylogenetic analysis and post-analysis of large phylogenies. *Bioinformatics (Oxford, England)* **30**, 1312–1313.
- Stuart KD and Gelvin SB (1982) Localization of kinetoplast DNA maxicircle transcripts in bloodstream and procyclic form *Trypanosoma brucei*. *Molecular and Cellular Biology* **2**, 845–852.
- Stuart K, Allen TE, Heidmann S and Seiwert SD (1997) RNA editing in kinetoplastid protozoa. *Microbiology and Molecular Biology Reviews* **61**, 105–120.
- Stuart KD, Schnauffer A, Ernst NL and Panigrahi AK (2005) Complex management: RNA editing in trypanosomes. *Trends in Biochemical Sciences* **30**, 97–105.
- Surve S, Heestand M, Panicucci B, Schnauffer A and Parsons M (2012) Enigmatic presence of mitochondrial complex I in *Trypanosoma brucei* bloodstream forms. *Eukaryotic Cell* **11**, 183–193.
- Talavera G and Castresana J (2007) Improvement of phylogenies after removing divergent and ambiguously aligned blocks from protein sequence alignments. *Systematic Biology* **56**, 564–577.
- Vasil'eva MA, Bessolitsina EA, Merzlyak EM and Kolesnikov AA (2004) Identification of the 12S rRNA gene promoter in *Leptomonas seymouri* mitochondrial DNA. *Molecular Biology* **38**, 839–843.
- Vaux R, Schnoeller C, Berkachy R, Roberts LB, Hagen J, Gounaris K and Selkirk ME (2016) Modulation of the immune response by nematode secreted acetylcholinesterase revealed by heterologous expression in *Trypanosoma muscili*. *PLoS Pathogens* **12**, e1005998.
- Verma A, Manchanda S, Kumar N, Sharma A, Goel M, Banerjee PS, Garg R, Singh BP, Balharbi F, Lejon V, Deborggraeve S, Singh Rana UV and Puliyl J (2011) *Trypanosoma lewisi* or *T. lewisi*-like infection in a 37-day-old Indian infant. *American Journal of Tropical Medicine and Hygiene* **85**, 221–224.
- Verner Z, Čermáková P, Škodová I, Kriegová E, Horváth A and Lukeš J (2011) Complex I (NADH:ubiquinone oxidoreductase) is active in but non-essential for procyclic *Trypanosoma brucei*. *Molecular and Biochemical Parasitology* **175**, 196–200.
- Wadkins RM (2000) Targeting DNA secondary structures. *Current Medicinal Chemistry* **7**, 1–15.
- Walker BJ, Abeel T, Shea T, Priest M, Abouelliel A, Sakthikumar S, Cuomo CA, Zeng Q, Wortman J, Young SK and Earl AM (2014) Pilon: an integrated tool for comprehensive microbial variant detection and genome assembly improvement. *PLoS One* **9**, e112963.
- Westenberger SJ, Cerqueira GC, El-Sayed NM, Zingales B, Campbell DA and Sturm NR (2006) *Trypanosoma cruzi* mitochondrial maxicircles display species- and strain-specific variation and a conserved element in the non-coding region. *BMC Genomics* **7**, 60.
- Zhang X, Hong XK, Li SJ, Lai DH, Hide G, Lun ZR and Wen YZ (2018) The effect of normal human serum on the mouse trypanosome *Trypanosoma muscili* *in vitro* and *in vivo*. *Experimental Parasitology* **184**, 115–120.