

Review Article

Cite this article: Hielscher E, DeVylder JE, Saha S, Connell M, Scott JG (2018). Why are psychotic experiences associated with self-injurious thoughts and behaviours? A systematic review and critical appraisal of potential confounding and mediating factors. *Psychological Medicine* **48**, 1410–1426. <https://doi.org/10.1017/S0033291717002677>

Received: 12 June 2017

Revised: 8 August 2017

Accepted: 15 August 2017

First published online: 20 September 2017

Key words:

Confounders; psychotic experiences; self-harm; self-injurious behaviour; suicide

Author for correspondence:

E. Hielscher, E-mail: e.hielscher@uq.edu.au

Why are psychotic experiences associated with self-injurious thoughts and behaviours? A systematic review and critical appraisal of potential confounding and mediating factors

E. Hielscher^{1,2,3}, J. E. DeVylder⁴, S. Saha^{2,5}, M. Connell^{1,6} and J. G. Scott^{1,2,6}

¹Centre for Clinical Research, Faculty of Medicine, The University of Queensland, Brisbane, QLD, Australia; ²Queensland Centre for Mental Health Research (QCMHR), The Park Centre for Mental Health, Brisbane, QLD, Australia; ³School of Public Health, The University of Queensland, Brisbane, QLD, Australia; ⁴Graduate School of Social Service, Fordham University, New York, NY, USA; ⁵Department of Psychiatry, and Queensland Brain Institute, University of Queensland, St Lucia, QLD, Australia and ⁶Metro North Mental Health, Royal Brisbane and Women's Hospital, Herston, QLD, Australia

Abstract

Psychotic experiences (PEs), including hallucination- and delusion-like experiences, are robustly associated with self-injurious thoughts and behaviours (SITB) in the general population. However, it remains unclear as to why there is an association. The purpose of this systematic review was to elucidate the role of other factors that influence the association between PEs and SITB and, in doing so, highlight potential mechanisms underlying the relationship. A search of electronic international databases was undertaken, including PubMed, PsycINFO and EMBASE, and eligible studies were grouped according to seven confounder categories: sociodemographics, mental disorders, alcohol and substance use, environmental, psychological, intervention and family history/genetic factors. The systematic search strategy identified 41 publications reporting on 1 39 427 participants from 16 different countries. In the majority of studies, where adjustment for other variables occurred, the association between PEs and SITB persisted, suggesting PEs have an independent role. Common mental disorders, psychological distress and negative environmental exposures explained a substantial amount of the variance and therefore need to be considered as potential underlying mechanisms. There was high variability in the variables adjusted for in these studies, and so the question still remains as to whether the association between PEs and self-harm/suicidality can be attributed (fully or in part) to confounding and mediating factors or directly causal mechanisms. Regardless of causality, the now extensive literature reporting an association between these two clinical phenomena supports the broad usefulness of PEs as an indicator of risk for SITB.

Introduction

The high rates of self-harm, suicide attempts and completed suicides among individuals with psychotic disorders (Palmer *et al.* 2005) also extend to those in the general population with psychotic experiences (PEs), i.e. hallucinatory and delusional experiences (Honings *et al.* 2016a). PEs are more common than psychotic disorders, reported by 5–8% of adolescents and adults (Kelleher *et al.* 2012a; McGrath *et al.* 2015), and are strongly associated with self-injurious thoughts and behaviours (SITB), including non-suicidal self-injury, suicidal ideation, plans and attempts (Koyanagi *et al.* 2015a; DeVylder *et al.* 2015b). Understanding the mechanisms underlying the association between PEs and SITB may assist identification of people at high risk of suicide and also inform interventions for those with PEs.

A meta-analysis reported increased likelihood of suicidal ideation and suicidal behaviour in those with any PE [odds ratios (ORs) and 95% confidence interval (CI) 2.47; 1.71–3.59 and 3.03; 2.08–4.41, respectively] (Honings *et al.* 2016a). Studies have also reported dose–response relationships with regard to both number of PEs (Nishida *et al.* 2010; Saha *et al.* 2011a; Cederlöf *et al.* 2016) and severity of suicidality (Kelleher *et al.* 2012b; DeVylder & Hilimire, 2015). Less is known about the association between PEs and non-suicidal self-injury, typically defined as deliberate self-injury without suicidal intent (Peterson *et al.* 2008), although two studies report increased risk of non-suicidal self-injury in adolescents (Martin *et al.* 2015) and adults (Koyanagi *et al.* 2015a) with PEs.

The association between PEs and SITB has been well replicated, although underlying mechanisms remain unclear. It may be directly causal where PEs induce self-injurious thoughts or acts via such symptoms as command hallucinations, i.e. voices directing a person to harm themselves (Kelleher *et al.* 2012b). Alternatively, shared risk factors such as depression




Fig. 1. Three potential pathways to explain the association between psychotic experiences and self-injurious thoughts and behaviours (SITB).

may result in indirect pathways (see Fig. 1) (Jang *et al.* 2014). A longitudinal study found the PE-suicidality association was only present in those reporting common mental disorders at baseline (Honings *et al.* 2016b), and adding a measure of depressive symptoms to PEs significantly improved prediction of later suicide attempts (Sullivan *et al.* 2015). Conversely, a study of the Adult Psychiatric Morbidity Survey ($n = 7403$) reported the strong association between hallucinations and suicide attempts was independent of common mental disorders such as depression (Kelleher *et al.* 2017). This was somewhat supported by Honings *et al.*'s (2016a) meta-analysis, which found adjusting for depression attenuated but did not negate the PE–SITB association. These studies, however, did not adjust for other co-occurring diagnoses or other confounding factors.

Other potential confounders and mediators (e.g. traumatic experiences, emotional dysregulation) likely play a role in the PE–SITB association (DeVylder *et al.* 2015a), but to date, have been inconsistently investigated: studies have either not adjusted for the same factors (e.g. self-esteem *v.* emotional dysregulation) or not adjusted beyond the more typical variables of sociodemographics, depression, and substance use. In light of these gaps in the literature, we conducted a systematic review and critically analysed third variables (confounders, mediators) that have been examined to date which influence the PE–SITB relationship.

This review aimed to elucidate the role of confounding and mediating factors in the association between PEs and SITB. Specifically, we aimed to extend previous work (Honings *et al.* 2016a) by: (1) including additional studies that have been subsequently published or initially omitted; (2) improving definitions and categorisation of SITB types; (3) presenting associations between SITB and different types of PEs, as well as different populations (adolescent *v.* adult); and (4) systematically examining different types of confounders and mediators not considered in the previous review. In conducting this review, we also identified gaps in the literature where potential mechanisms for the PE–SITB association have yet to be investigated, so as to guide future research.

Methods

Data sources

This systematic review adhered to the PRISMA Guidelines (Moher *et al.* 2009). An electronic literature search was initially conducted in January 2017, which covered relevant international databases, including PubMed, PsycINFO and EMBASE (final search conducted in May 2017). Additional records were identified through citation databases, such as the Web of Science and Scopus, as well as through Google Scholar. Reference lists of included articles were also hand-searched. Authors of included articles were contacted to obtain further details about their analyses.

Definitions

Self-injurious thoughts and behaviours

SITB were defined as actions where an individual deliberately initiates a non-socially or culturally sanctioned behaviour that, without intervention from others, will harm or injure their own body tissue (St. Germain & Hooley, 2012; Larkin *et al.* 2014), as well as thoughts about harming oneself or ending one's life (Nock *et al.* 2007). This definition is consistent with the WHO/EURO multicentre study on self-harm (Platt *et al.* 1992; Hawton *et al.* 2003), and was incorporated to investigate how PEs relate to a wide spectrum of distinct but overlapping self-injury-related constructs.[†] The separation of self-injurious thoughts from behaviours is recommended as they have distinct prevalence rates, functions, correlates, and outcomes (Klonsky *et al.* 2016). For this reason, SITB was divided into two categories. The first was self-harming and suicidal ideation, which included all cognitions (thoughts, feelings, plans) about self-harm and suicide, without engagement in self-injurious behaviour. No study

[†]The notes appear after the main text.

specifically captured self-harming thoughts² and therefore this category was titled 'suicidal ideation'.

The second category was labelled non-accidental self-injury (NASI) and included non-suicidal self-injury, deliberate self-harm (with/without suicidal intent), and suicide attempt. There is expanding evidence for the Gateway Theory (Kapur *et al.* 2013; Whitlock *et al.* 2013), which proposes that non-suicidal self-injury and fatal suicide sit at two ends of the same spectrum, with non-suicidal self-injury being an antecedent to suicide (Linehan, 1986; Stanley *et al.* 1992). Some of the evidence includes considerable overlap in risk factors of non-suicidal and suicidal behaviour (Grandclerc *et al.* 2016), comparable prevalence rates of non-suicidal self-injury and deliberate self-harm (Muehlenkamp *et al.* 2012), and in prospective studies, non-suicidal self-injury is one of the more robust predictors of suicide attempts (Guan *et al.* 2012; Scott *et al.* 2015). Following the evidence, in this review we presented NASI as a construct that does not assume or preclude suicidal intent. It includes behaviours on a continuum of suicidal intent, ranging from non-suicidal self-injury to suicide attempt, as opposed to discrete intent categories.

Psychotic experiences

PEs were defined as hallucinations and delusions (e.g. persecution, referential and grandiose) endorsed by individuals in the general population that do not occur in the context of sleep or substance use (Johns *et al.* 2004; Linscott & van Os, 2013). They may be appraised as clinically relevant symptoms or as subclinical/sub-threshold experiences, not prompting help-seeking behaviour (Linscott & van Os, 2010). Although phenotypically PEs lie on a continuum with psychotic disorders, most with PEs do not transition to psychosis (Yung & Lin, 2016). More commonly, PEs are associated with a broader range of adverse health and social outcomes (McGrath *et al.* 2016; Saha *et al.* 2011b).

Eligibility criteria

Inclusion criteria were studies that: (a) were peer-reviewed and published in English language, from inception until May 2017; (b) measured both SITB and PEs; (c) reported on the PE–SITB association; and (d) were conducted in a general population or non-clinical community sample.

Exclusion criteria were studies limited to: (a) body modification (i.e. cutting, piercing, tattooing) as a cultural practice or expression of creativity; (b) assisted suicide by a physician; (c) indirect forms of self-injurious behaviour (e.g. binge-drinking, excessive exercise); (d) skin picking or hair pulling³; (e) self-injurious behaviour arising in those with development disorder; (f) studies conducted in clinical, mental health or help-seeking populations; or (g) assessment of schizotypy and related personality assessments.⁴

Search strategy

A literature search was conducted in PubMed, PsycINFO and EMBASE⁵ using the following terms: (psychotic exp* OR subclinical psych* OR psychotic-like OR delusion* OR hallucinat*) AND (self-injur* OR self-harm* OR suic*). Search terms were sourced from Honings *et al.*'s (2016a) review, as well as other systematic reviews on PEs (Kaymaz *et al.* 2012; Linscott & van Os, 2010, 2013) and SITB (Larkin *et al.* 2014; Taylor *et al.* 2015; Zetterqvist, 2015). See online Supplementary Material I for full PubMed search.

Data extraction

One author (EH) located eligible studies using the search strategy and downloaded these into EndNote. Titles and abstracts were screened, with at least 10% done independently by two authors (EH and SS) against the predefined inclusion criteria. A similar process was undertaken for the full-text screen. Data on study design, SITB and PEs measurement, third variables, and key findings were extracted and entered into an excel template by EH.

Data synthesis and confounder analysis

The literature was critically reviewed by narrative synthesis of aggregate-level data. Due to the wide range of third variables measured, and the fact that most studies simultaneously adjusted for multiple variables, meta-analysis was precluded. Potential confounders and mediators were examined using a protocol developed for analysis of epidemiologic studies of parental smoking (Witorsch & Witorsch, 1993). The key aims of our analysis were to determine: (1) which potential confounders/mediators have been considered; (2) frequency at which confounders/mediators have been considered; and (3) whether these variables influenced the PE–SITB relationship.

Considering the variability of third variables, they were grouped into seven higher-level categories: sociodemographics, mental disorders, alcohol and substance use, environmental (e.g. trauma), psychological, intervention, and family history/genetic factors. These categories were derived from extracted data of included studies, as well as known risk factors in the broader self-harm and suicide literature (Klonsky, 2007; Franklin *et al.* 2017). We examined the influence of these seven categories on: (1) suicidal ideation, and (2) NASI, which was further divided into non-suicidal self-injury, deliberate self-harm and suicide attempt. Despite substantial overlap between risk factors for non-suicidal self-injury and suicide attempts (Grandclerc *et al.* 2016), it was important to investigate these constructs separately, as some variables may be important for the association between PEs and one self-injurious outcome but not the other.

We labelled all seven categories as confounders, as this was how most studies described these variables. However, some of these variables are mediators, which explain the association between PEs and SITB. Online Supplementary Material II presents a more detailed discussion of each category, and we further address this in the 'Discussion'.

Results

Characteristics of included studies

The systematic search strategy identified 38 individual studies (from 41 publications) reporting on 1 39 427 participants ($M = 3873$, range = 66–16 131) from 16 countries (see Fig. 2). Most studies were cross-sectional (73.7%), with ten reporting longitudinal data (follow-up range = 3 months–27 years). There was a mixture of data sources: 26.3% of studies reporting national household surveys, 15.8% reporting birth cohorts, and the remainder (57.9%) reporting community samples, e.g. high school and university students. Seven of the 41 publications performed analyses on the same three datasets (see Fig. 2). Because the authors presented different variables of interest (e.g. ideation *v.* attempts) and adjusted for different confounders, the remaining results section presents the literature in terms of number of publications, as opposed to individual studies/datasets.




Fig. 2. PRISMA flow chart of literature search.

Most studies ($n = 24$; 58.5%) reported aggregated PEs, as opposed to the association with individual symptoms (e.g. visual hallucinations). All but one study (Kelleher *et al.* 2012b)⁶ reported on different types of SITB separately. The level of adjustment varied across studies (see Tables 1 and 2), with 11 (26.8%) not controlling for any confounding variables. Only one study (Jang *et al.* 2014) treated their third variable of interest (depressive symptoms) as a mediating variable.

PEs and suicidal ideation

When broken down by SITB outcomes, 23 studies reported the PE–suicidal ideation association after adjustment for confounders; with just over half (52.2%) confined to an adolescent and/or young adult sample (see Table 3). Three of 23 studies reported the association between suicidal ideation and different types of PEs, yielding mixed findings (Nishida *et al.* 2010; Capra *et al.* 2015; Koyanagi *et al.* 2015b). Most ($n = 14$; 60.9%) controlled for three confounder categories or less, with demographics, mental disorders and/or substance use most common. Only two cross-sectional studies controlled for six out of the seven confounder categories (Nishida *et al.* 2008; DeVlyder *et al.* 2015a). Adjusting for these confounders attenuated the strength of the PE–suicidal ideation association (OR range = 1.03–2.10, $p < 0.05$) compared with unadjusted associations and associations found in most other studies.

PEs and NASI

Table 4 outlines the studies ($n = 20$; non-suicidal self-injury = 2, deliberate self-harm = 4, suicide attempt = 15) reporting the PE–

NASI association, adjusted for confounders. All deliberate self-harm studies focused on adolescents, non-suicidal self-injury studies equally covered adolescents and adults, whereas most suicide attempt studies ($n = 10$; 66.7%) reported adult populations. Four of 20 studies compared the association of NASI with different PEs, and all found individual PEs were differentially associated with all three NASI outcomes (Nishida *et al.* 2010; Capra *et al.* 2015; Koyanagi *et al.* 2015a, b). Similar to suicidal ideation studies, most NASI studies ($n = 15$; 75.0%) controlled for three or less confounder categories, with demographics, mental disorders and substance use again the most common (see Table 4). Only two studies controlled for six of seven confounder categories, reporting barely significant associations between both deliberate self-harm (OR 1.37, 95% CI 1.00–1.86, Nishida *et al.* 2008) and suicide attempts (OR 1.01, 95% CI 0.98–1.04, DeVlyder *et al.* 2015a) with PEs. Each confounder category is further discussed below with reference to all SITB outcomes.

Sociodemographics

Most studies ($n = 28$; 68.3%) adjusted for sociodemographics, incorporating diverse variables ranging from age and sex to sexual orientation and employment status. Four studies (9.8%) presented point estimates of the PE–SITB relationship after adjusting for demographics (see Tables 3 and 4) and associations remained significant (OR range: 2.07–7.79). Further adjustment for such variables as mental disorders and substance use substantially attenuated the strength of the association (OR range: 2.49–3.37; Fisher *et al.* 2013; DeVlyder *et al.* 2015b), with some estimates becoming non-significant (OR range: 1.32–1.60; Nishida *et al.*

Table 1. Frequency of consideration of potential confounders in studies examining the association between psychotic experiences and self-injurious thoughts and behaviours

Confounder no.	Potential confounder ^a	Studies (41 total)
1.	Sociodemographics <ul style="list-style-type: none"> Sex, age, race/ethnicity, years of education (participant and parents), marital status, income (participant and household or family), migrant status (participant and parents), socioeconomic status of family and neighbourhood, urbanicity, sexual orientation, employment status, country of birth (participant and parents), family structure (i.e. living apart from one or both parents), household type, primary language, English language proficiency, school grade, religious/spiritual affiliation 	Twenty-eight (68.3% of studies)
2.	Mental disorders <ul style="list-style-type: none"> Depression (symptoms, episode and diagnosed disorder; both 12 months and lifetime), any anxiety disorder and mixed anxiety/depression (symptoms and diagnosed disorder; past month, 12 months, and lifetime), generalised anxiety disorder, phobic disorder or phobia (specific and social), panic disorder, obsessive-compulsive disorder, lifetime mood (depression, mania), borderline personality (symptoms and diagnosed disorder), agoraphobia, behavioural disorders (ADHD, oppositional defiant, conduct), internalising/externalising psychopathology, eating disorders (anorexia and bulimia nervosa), somatisation, separation anxiety, PTSD, schizophrenia spectrum disorders, bipolar disorder, antisocial personality disorder, self-induced vomiting for dieting purposes, ataque de nervios, count of lifetime psychiatric disorders, self-reported mental health 	Twenty (48.8% of studies) ^b
3.	Alcohol and substance use <ul style="list-style-type: none"> Alcohol and substance use (cannabis, ecstasy, methamphetamines, amphetamines, cocaine, opiates, hallucinogens, sedatives, tranquilisers, amyl nitrate, anabolic steroids, glues, cigarettes, polysubstance; past month, 12 months, and lifetime), alcohol use disorder (past 6 months, 12 months and lifetime), other substance use disorder (both 12 months and lifetime), any substance (alcohol/drug) use disorder (lifetime), hazardous drinking 	Fifteen (36.6% of studies)
4.	Environmental factors <ul style="list-style-type: none"> Stressful and traumatic life events (e.g. death of a family member, raped or assaulted), history of bullying and victimisation (past 12 months and lifetime), violence from adults in the home (past month), total support group (i.e. size of support group), history of childhood trauma (sexual, physical), school mobility (no. of schools attended between kindergarten and high school graduation), school problems, bullying others/physically assaulting others (past 12 months), family relationship/support, time spent watching television or using personal computers 	Ten (24.4% of studies)
5.	Psychological factors <ul style="list-style-type: none"> Psychological distress (GHQ-12), self-esteem (Rosenberg Self-Esteem Scale), stress (Psychological Stress Index), emotional regulation (suppression and reappraisal), irritability when exchanging emails (past week), trait anger (Anger Expression Inventory), self-awareness of mental distress, stereotype awareness (i.e. mental illness stigma), difficulty falling asleep due to hypersensitivity to noises (past month), difficulty concentrating due to hypersensitivity to noises (past month), interpersonal sensitivity, hostility, WRAT-4 reading score, preference for being alone or with others 	Nine (21.9% of studies)
6.	Intervention factors <ul style="list-style-type: none"> Hospitalisation admissions (lifetime psychiatric and general), current contact with medical services, ever talked with professional (school counsellor, psychologist, social work, psychiatrist or other) for feelings or problems with mood or behaviour, current help-seeking behaviour (both with professionals and non-professionals, e.g. friends), lifetime psychiatric medication 	Five (12.2% of studies)
7.	Family history/genetic factors <ul style="list-style-type: none"> Family history of mental illness 	Two (4.9% of studies)

Note: All confounders in eligible studies were captured in this table, except for Children's Global Assessment Scale (OR 0.96, 95% CI 0.95–0.97, Calkins *et al.* 2014, 2017), self-reported physical health (OR 0.88, 95% CI 0.60–1.29), and count of lifetime chronic medical conditions (OR 1.20, 95% CI 1.06–1.33; Lewis-Fernández *et al.* 2009). None of these variables fitted our confounder categorisation, and most did not significantly contribute to the fully adjusted models.

^aConfounders are ordered according to how frequently they appear in the literature, both within and across categories.

^bFour of the studies included depressive symptoms only.

2014; Honings *et al.* 2016b), suggesting considerable residual confounding when only sociodemographics were included in the model.

Mental disorders

Less than half of the studies ($n = 20$; 48.8%) adjusted for mental disorders, most commonly depression (12-month or lifetime diagnosis, episode, or symptoms), anxiety disorders and mixed

anxiety/depression diagnosis. Five studies controlled for mental disorders individually, or in separate steps in their models (Polanczyk *et al.* 2010; Fisher *et al.* 2013; Koyanagi *et al.* 2015a; Honings *et al.* 2016b; Kelleher *et al.* 2017). Most studies adjusting for comorbid depressive symptoms and common mental disorders (depression, anxiety, personality disorders) reported reduced ORs, but the PE–SITB association remained significant (OR range: 2.00–3.20; Polanczyk *et al.* 2010; Fisher *et al.* 2013; Kelleher *et al.* 2017). This was somewhat supported by Honings

Table 2. Consideration of potential confounding variables in studies examining the relationship between psychotic experiences and self-injurious thoughts and behaviours

Study	Confounder no. ^a						
	1	2	3	4	5	6	7
Calkins <i>et al.</i> (2014)	+	+	–	–	+	+	–
Calkins <i>et al.</i> (2017)	+	+	–	+	+	+	–
Capra <i>et al.</i> (2015)	+	–	+	–	–	–	+
Cederlöf <i>et al.</i> (2016)	–	–	–	–	–	–	–
Connell <i>et al.</i> (2016)	+	–	+	–	–	–	–
DeVylder <i>et al.</i> (2015a)	+	+	+	+	+	–	+
DeVylder <i>et al.</i> (2015b)	+	+	+	–	–	–	–
DeVylder <i>et al.</i> (2015c)	–	–	–	–	–	–	–
DeVylder & Hilimire (2015)	+	–	–	–	+	–	–
DeVylder & Kelleher (2016)	+	–	+	–	–	–	–
Engin <i>et al.</i> (2009)	+	+	–	+	+	–	–
Fisher <i>et al.</i> (2013)	+	+	–	–	–	–	–
Freeman <i>et al.</i> (2011)	–	–	–	–	–	–	–
Gatta <i>et al.</i> (2016)	–	–	–	–	–	–	–
Honings <i>et al.</i> (2016b)	+	+	–	–	–	–	–
Jang <i>et al.</i> (2014)	+	+	–	–	–	–	–
Jenkins <i>et al.</i> 2015)	+	+	–	+	–	–	–
Kelleher <i>et al.</i> (2012b) ^{b,d}	+	–	–	–	–	–	–
Kelleher <i>et al.</i> (2012b) ^{c,d}	+	–	–	–	–	–	–
Kelleher <i>et al.</i> (2013a) ^d	+	–	+	–	–	–	–
Kelleher <i>et al.</i> (2014)	+	+	+	–	–	–	–
Kelleher <i>et al.</i> (2017)	–	+	–	–	–	–	–
Kerr & Muehlenkamp (2010)	–	–	–	–	–	–	–
Koyanagi <i>et al.</i> (2015a)	+	+	+	+	–	–	–
Koyanagi <i>et al.</i> (2015b)	+	+	+	+	–	–	–
Krabbendam <i>et al.</i> (2004)	–	–	–	–	–	–	–
Lewis-Fernández <i>et al.</i> (2009)	+	+	+	+	–	–	–
Martin <i>et al.</i> (2015)	+	–	–	–	+	–	–
Nishida <i>et al.</i> (2008)	+	+	+	+	+	+	–
Nishida <i>et al.</i> (2010)	+	–	+	+	+	–	–
Nishida <i>et al.</i> (2014)	+	–	–	–	+	+	–
Núñez <i>et al.</i> (2016)	–	–	–	–	–	–	–
Olfson <i>et al.</i> (1996)	–	–	–	–	–	–	–
Olfson <i>et al.</i> (2002)	+	+	+	–	–	–	–
Östling & Skoog (2002)	–	–	–	–	–	–	–
Östling <i>et al.</i> (2007)	–	–	–	–	–	–	–
Polanczyk <i>et al.</i> (2010)	–	+	–	–	–	–	–
Saha <i>et al.</i> (2011a)	+	+	+	+	–	–	–
Sharifi <i>et al.</i> (2015)	+	+	+	–	–	+	–
Sullivan <i>et al.</i> (2015)	+	+	–	–	–	–	–

(Continued)

Table 2. (Continued.)

Study	Confounder no. ^a						
	1	2	3	4	5	6	7
Temmingh <i>et al.</i> (2011)	+	–	–	–	–	–	–
Zhang <i>et al.</i> (2016)	–	–	–	–	–	–	–

+, Indicates the potential confounding variables were considered in a particular study.

^aCorresponds to numbers in Table 1.

^bAdolescent Brain Development (ABD) study reported in Kelleher *et al.* (2012b).

^cChallenging Times (CT) study reported in Kelleher *et al.* (2012b).

^dConfounders adjusted for using a stratified analysis were only considered if stratification was performed as part of the main analysis (i.e. overall estimate for psychotic experiences), and if the authors presented the odds ratios in each strata i.e. with and without the third variable of interest. These two publications did not present such information in their psychiatric disorder/psychopathology stratified subanalysis, and therefore 'mental disorders' were not counted in their confounder categorisation.

et al.'s (2016a) meta-analysis, where they included only studies that controlled for depression and found an attenuated (but significant) pooled OR for all SITB outcomes except deliberate self-harm (OR for SITB = 2.02, 95% CI 2.33–4.40). However, most studies controlled for a number of confounders simultaneously, making it difficult to interpret the proportion of the PE–SITB relationship attributable to depression.

Analyses stratified by presence of mental disorder provided further insight. A longitudinal study reported the association was only present in the subsample of participants with PEs who reported mental disorders at baseline (Honings *et al.* 2016b). By contrast, other cross-sectional and longitudinal studies found the PE–SITB relationship persisted even when the sample was restricted to those without mental disorders (Saha *et al.* 2011a; Sullivan *et al.* 2015; Kelleher *et al.* 2017). Mental disorders consistently explain, in part, the PE–SITB association, although the association is frequently evident even in those without mental illness endorsing PEs.

Alcohol and substance use

Almost 40% of studies adjusted for substance use, most controlling for past 12 month or lifetime substance use as opposed to abuse or dependence. Substance use was typically adjusted for in the same step as mental disorders preventing any conclusions about their role in the PE–SITB relationship. Alcohol and cannabis use were not significant contributors in DeVlyder *et al.*'s (2015a) fully adjusted model, whereas Capra *et al.* (2015) found lifetime cannabis use was a significant predictor in both suicidal ideation and attempts models (β -coefficient range: 0.10–0.18). The strong association between substance use and both suicidality (Vijayakumar *et al.* 2011) and PEs (Rognli *et al.* 2017) suggests it may be important in the PE–SITB relationship.

Environmental factors

Although traumatic life events are associated with both suicidality and PEs (Scott *et al.* 2007; Kelleher *et al.* 2013b), fewer than 25% of studies adjusted for these and no study independently controlled for such factors. In DeVlyder *et al.*'s (2015a) fully adjusted models, bullying, school mobility and childhood sexual trauma were key variables, resulting in large attenuation of all suicidal outcomes with only broadly defined suicidal ideation remaining significant (OR 1.03, 95% CI 1.01–1.05). Although PEs and suicidal behaviour appear to co-occur in the context of severe social

stressors, these environmental exposures did not entirely account for the PE–suicidality association (β -coefficient range: 0.19–2.23).

Psychological factors

Psychological factors (e.g. psychological distress) were adjusted for in 21.9% of studies. Martin *et al.* (2015) found PEs, in the absence of psychological distress, were not associated with increased risk of future non-suicidal self-injury and suicide attempts (OR range: 1.63–1.65). The only other study independently controlling for psychological distress (Nishida *et al.* 2014) reported attenuated ORs of deliberate self-harm and suicidal ideation in those with PEs, with only suicidal ideation remaining significant (OR 2.1, 95% CI 1.5–2.9). Some suggest that PEs are manifestations of distress (Saha *et al.* 2011a; Koyanagi *et al.* 2015b). DeVlyder *et al.* (2015a) investigated other psychological factors such as self-esteem and stereotype awareness (i.e. mental illness stigma), which explained a significant amount of variance between PEs and intensity of suicidal ideation (β -coefficient range: 0.10–0.29), but did not entirely account for the relationship.

Intervention factors

Only five studies adjusted for intervention factors, including hospitalisation history and help-seeking behaviour. Only Nishida *et al.* (2014) individually adjusted for such factors, by stratifying their sample by help-seeking. Risk of both suicidal ideation and deliberate self-harm were highest among non-help-seeking adolescents (who had a need for care) with PEs; however, the PE–SITB relationship persisted regardless of help-seeking status.

Family history/genetic factors

Only two studies adjusted for family history of mental illness (Capra *et al.* 2015; DeVlyder *et al.* 2015a), with both simultaneously adjusting for multiple confounders; however, family history explained little of the variance.

Discussion

There is consistent evidence that PEs are an important clinical indicator of suicide risk. However, no study has explained if the relationship is causal or if their co-occurrence arises from shared risk factors. This distinction is critical to inform the clinical assessment and interventions for those with PEs. In this review,

Table 3. Strength of evidence in studies (n = 23) examining the psychotic experiences (PEs) – suicidal ideation relationship, after adjustment for confounders

Study	N, Age (years)	Design	Temporality	Confounder (s) ^a	Psychotic experience	Outcome measure	Association between PEs and outcome measure
<i>Studies with adolescents and young adults (≤25 years)</i>							
Calkins <i>et al.</i> (2014) USA	4848 11–21	CS	N/A	1, 2, 5, 6	Psychosis spectrum (LT and past year)	LT suicidal ideation	OR 1.45, 95% CI 1.07–1.96
Calkins <i>et al.</i> (2017) USA	249 14–18	L	N	1, 2, 4, 5, 6	Persistence of psychosis spectrum over 2 years	LT suicidal ideation	OR 0.85, 95% CI 0.37–1.98
Capra <i>et al.</i> (2015) Australia	1610 M = 22.1 ± 5.1	CS	N/A	1, 3, 7	LT perceptual abnormalities	LT suicidal ideation	OR 1.30, 95% CI 1.12–1.50
					LT persecutory ideation		OR 1.26, 95% CI 1.18–1.35
					LT bizarre experiences		OR 1.00, 95% CI 0.94–1.06
DeVylder & Hilimire (2015) USA	622 M = 18.8	CS	N/A	1, 5	LT auditory hallucinations	LT suicidal ideation	OR 1.90, 95% CI 1.19–3.0
						LT suicidal plan	OR 2.18, 95% CI 1.00–4.78
Engin <i>et al.</i> (2009) Turkey	1992 M = 20.02 ± 1.54	CS	N/A	1, 2, 4, 5	Past week psychotic symptoms/psychoticism	LT suicidal thoughts	OR 1.27, 95% CI 1.11–1.45
Jang <i>et al.</i> (2014) South Korea	6977 14–19	CS	N/A	1, 2	Current psychotic-like experiences (last 4 weeks)	Current suicidal ideation	Beta coefficient (β) = 0.13. Depressive symptoms partially mediated association (β = 0.56)
Kelleher <i>et al.</i> (2013a) Ireland	1112 13–16	L	Y	1, 3	LT auditory hallucinations at baseline	Suicidal ideation (often/always during past 2 weeks) at 3-month follow-up	OR 8.79, 95% CI 2.61–29.60
						Suicidal ideation (often/always during past 2 weeks) at 12-month follow-up	OR 2.55, 95% CI 0.24–27.36
Kelleher <i>et al.</i> (2014) Sweden	109 16–17; 179 19–20	L	Y	1, 2, 3	6-month auditory hallucinations at age 13–14	6-month suicidal ideation at age 16–17	OR 0.94, 95% CI 0.19–4.78
					6-month auditory hallucinations at age 16–17	6-month suicidal ideation at age 19–20	OR 5.53, 95% CI 1.33–23.0
Nishida <i>et al.</i> (2008) ^b Japan	4894 12–15	CS	N/A	1, 2, 3, 4, 5, 6	LT psychotic-like experiences	LT suicidal ideation	OR 2.10, 95% CI 1.83–2.41
Nishida <i>et al.</i> (2010) ^b Japan	4894 12–15	CS	N/A	1, 3, 4, 5	LT any psychotic experiences	LT suicidal feelings	OR 3.1, 95% CI 2.2–4.5
							OR 2.47, 95% CI 1.40–4.34
					LT thoughts read		OR 1.93, 95% CI 0.83–4.47
					LT special messages		OR 2.44, 95% CI 1.87–3.18
					LT spied-upon		OR 2.26, 95% CI 1.79–2.87
				LT hearing voices			

(Continued)

Table 3. (Continued.)

Study	N, Age (years)	Design	Temporality	Confounder (s) ^a	Psychotic experience	Outcome measure	Association between PEs and outcome measure
Nishida et al. (2014) Japan	16 131 12–18	CS	N/A	1, 6 ^c	LT psychotic-like experiences	Current suicidal ideation	OR 3.1, 95% CI 2.2–4.2
				1, 5, 6 ^c	LT psychotic-like experiences	Current suicidal ideation	OR 2.1, 95% CI 1.5–2.9
Sullivan et al. (2015) England	3171 12 and 16	L	N	1, 2	6-month psychotic experiences at age 12	LT suicidal ideation at age 16	OR 1.31, 95% CI 0.99–1.73
<i>Studies with adults (≥16 years)^d</i>							
Connell et al. (2016) Australia	654 30–33	L	N	1, 3	Auditory or visual hallucinations at age 14	LT suicidal ideation at age 30	OR 1.44, 95% CI 0.93–2.23
						LT suicidal plans at age 30	OR 1.30, 95% CI 0.67–2.52
						LT suicidal ideation at age 30	OR 2.89, 95% CI 1.66–5.02
						LT suicidal plans at age 30	OR 2.89, 95% CI 1.37–6.09
DeVylder et al. (2015b) ^e USA	11 716 18+	CS	N/A	1	12-month psychotic experiences	12-month suicidal ideation	OR 4.25, 95% CI 2.28–7.89
				1, 2, 3	12-month psychotic experiences	12-month suicidal ideation	OR 2.49, 95% CI 1.01–6.13
DeVylder et al. (2015a) USA	590 17–37	CS	N/A	1, 2, 3, 4, 5, 7	Past month psychotic experiences	LT suicidal ideation	OR 1.03, 95% CI 1.01–1.05
						LT ideation with intent/plan	OR 1.01, 95% CI 0.98–1.03
						Intensity of worst-point ideation	$t = 1.84, p > 0.05$
DeVylder & Kelleher (2016) ^e USA	11 776 18+	CS	N/A	1, 3	LT psychotic experiences	LT suicidal ideation	OR 2.53, 95% CI 1.85–3.47
Honings et al. (2016b) Germany and Netherlands	15 837 $M = 39.4 \pm 13.8$	L	Y	1	LT psychotic experiences at baseline	Suicidal ideation at 3-year or 5-year follow-up	OR 2.07, 95% CI 1.50–2.87
				1, 2	LT psychotic experiences at baseline	Suicidal ideation at 3-year or 5-year follow-up	OR 1.32, 95% CI 0.93–1.87
Jenkins et al. (2015) Kenya	1133 $M = 42.0 \pm 18.1$	CS	N/A	1, 2, 4	Past year psychotic symptoms	LT suicidal thoughts	OR 1.4, 95% CI 0.79–2.43
Koyanagi et al. (2015b) ^f England	7363 16+	CS	N/A	1, 4	12-month any psychotic symptom	12-month suicidal ideation	OR 5.47, 95% CI 3.78–7.91
					12-month mania/hypomania	OR 5.21, 95% CI 2.21–12.32	
					12-month thought control	OR 5.74, 95% CI 2.29–14.37	
					12-month paranoia	OR 6.72, 95% CI 3.81–11.84	

(Continued)

Table 3. (Continued.)

Study	N, Age (years)	Design	Temporality	Confounder (s) ^a	Psychotic experience	Outcome measure	Association between PEs and outcome measure
					12-month strange experience		OR 6.09, 95% CI 3.92–9.46
					12-month auditory hallucination		OR 6.54, 95% CI 2.65–16.13
				1, 2, 3, 4	12-month any psychotic symptom	12-month suicidal ideation	OR 3.13, 95% CI 2.09–4.68
					12-month mania/hypomania		OR 4.00, 95% CI 1.41–11.37
					12-month thought control		OR 4.20, 95% CI 1.65–10.72
					12-month paranoia		OR 4.03, 95% CI 2.31–7.03
					12-month strange experience		OR 3.22, 95% CI 1.95–5.32
					12-month auditory hallucination		OR 4.05, 95% CI 1.76–9.32
Lewis-Fernández <i>et al.</i> (2009) USA	2554 18+	CS	N/A	1, 2, 3, 4	LT psychotic symptoms	LT suicidal ideation	OR 2.30, 95% CI 1.47–3.60
					12-month psychotic symptoms	12-month suicidal ideation	OR 2.04, 95% CI 1.12–3.74
Olfson <i>et al.</i> (2002) USA	1005 18–70	CS	N/A	1, 2, 3	LT psychotic symptoms	Current suicidal ideation	OR 3.4, 95% CI 1.9–6.1
Saha <i>et al.</i> (2011a) Australia	8773 18+	CS	N/A	1, 2, 3, 4	LT delusional-like experiences	LT suicidal ideation	OR 2.73, 95% CI 1.89–3.94
						LT suicide plan	OR 1.96, 95% CI 1.15–3.35
Temmingh <i>et al.</i> (2011) South Africa	4250 18+	CS	N/A	1	LT auditory or visual hallucinations	LT suicidal ideation	OR 2.44, 95% CI 1.7–3.5

CS, cross-sectional; L, longitudinal; Y, yes, temporality considered; N, no, temporality not considered; N/A, not applicable; LT, lifetime; OR, odds ratio; 95% CI, 95% confidence interval.

Bold represents a significant finding.

^aCorresponds to numbers in Table 1.

^bReports on data from the 2006 Mie Prefecture high school survey.

^cVariable used as stratification variable in the main analysis, e.g. if mental disorder was the stratification variable, we reported on the influence of psychotic experiences on SITB within the 'no mental disorders' strata.

^dThe adult category (16+) overlaps with the adolescent and young adult category (≤25) because most general population samples range from 16 (or 18) to 95 years.

^eReports on data from the Collaborative Psychiatric Epidemiology Surveys (CPES).

^fReports on data from the 2007 Adult Psychiatric Morbidity Survey (APMS).

Table 4. Strength of evidence in studies (n=20) examining the psychotic experiences (PEs) – non-accidental self-injury (NASI) relationship, after adjustment for confounders

Study	N, Age (years)	Design	Temporality	Confounder (s) ^a	Psychotic experience	Outcome measure	Association between PEs and outcome measure
<i>Non-suicidal self-injury [NSSI; studies with adolescents and young adults (<=25 years)]</i>							
Martin et al. (2015) Australia	1896 12–17	L	Y	1, 5 ^b	LT psychotic experiences at baseline	LT NSSI at baseline	OR 2.88, 95% CI 1.51–5.49
						NSSI at 12-month follow-up	OR 1.63, 95% CI 0.73–3.64
<i>Non-suicidal self-injury [NSSI; studies with adults (>=16 years)]^f</i>							
Koyanagi et al. (2015a) ^d England	7363 16+	CS	N/A	1, 4	12-month any psychotic-like experience	LT NSSI	OR 2.78, 95% CI 1.88–4.11
					12-month mania/hypomania		OR 2.85, 95% CI 0.98–8.23
					12-month thought control		OR 2.45, 95% CI 1.05–5.74
					12-month paranoia		OR 3.57, 95% CI 1.96–6.52
					12-month strange experience		OR 3.13, 95% CI 1.99–4.93
					12-month auditory hallucination		OR 4.03, 95% CI 1.56–10.42
				1, 2, 3, 4	12-month any psychotic-like experience	LT NSSI	OR 1.30, 95% CI 0.83–2.04
					12-month mania/hypomania		OR 1.13, 95% CI 0.34–3.74
					12-month thought control		OR 1.19, 95% CI 0.39–3.63
					12-month paranoia		OR 2.02, 95% CI 1.03–3.96
					12-month strange experience		OR 1.27, 95% CI 0.73–2.20
					12-month auditory hallucination		OR 2.23, 95% CI 0.75–6.65
<i>Deliberate self-harm [DSH; with/without suicidal intent; studies with adolescents and young adults (<=25 years)]</i>							
Nishida et al. (2008) ^e Japan	4894 12–15	CS	N/A	1, 2, 3, 4, 5, 6	LT psychotic-like experiences	Past year self-harming behaviours	OR 1.37, 95% CI 1.00–1.86
Nishida et al. (2010) ^e Japan	4894 12–15	CS	N/A	1, 3, 4, 5	LT any psychotic experience	Past year DSH	OR 3.1, 95% CI 2.0–4.8
					LT thoughts read		OR 1.56, 95% CI 0.71–3.43
					LT special messages		OR 1.62, 95% CI 0.48–5.50
					LT spied-upon		OR 1.93, 95% CI 1.34–2.77
					LT hearing voices		OR 2.32, 95% CI 1.67–3.22
Nishida et al. (2014) Japan	16 131 12–18	CS	N/A	1, 6 ^b	LT psychotic-like experiences	12-month DSH	OR 2.1, 95% CI 1.1–3.9
				1, 5, 6 ^b	LT psychotic-like experiences	12-month DSH	OR 1.6, 95% CI 0.8–3.1
Polanczyk et al. (2010) England and Wales	2232 12	CS	N/A	2	LT psychotic symptoms at age 12	LT DSH at age 12 (parent response)	OR 2.0, 95% CI 1.0–3.9

(Continued)

Table 4. (Continued.)

Study	N, Age (years)	Design	Temporality	Confounder (s) ^a	Psychotic experience	Outcome measure	Association between PEs and outcome measure
<i>Suicide attempt [studies with adolescents and young adults (<25 years)]</i>							
Capra <i>et al.</i> (2015) Australia	1610 M = 22.1 ± 5.1	CS	N/A	1, 3, 7	LT perceptual abnormalities	LT suicidal attempt	OR 1.61, 95% CI 1.32–1.97
					LT persecutory Ideation		OR 1.34, 95% CI 1.26–1.54
					LT bizarre Experiences		OR 0.92, 95% CI 0.84–1.02
DeVylder & Hilimire (2015) USA	622 M = 18.8	CS	N/A	1, 5	LT auditory hallucinations	LT suicide attempt	OR 3.99, 95% CI 1.37–11.61
Kelleher <i>et al.</i> (2013a) Ireland	1112 13–16	L	N	1, 3	LT auditory hallucinations at baseline	LT suicide attempt at 3-month follow-up	OR 10.01, 95% CI 2.24–45.49
						LT suicide attempt at 12-month follow-up	OR 11.27, 95% CI 4.44–28.62
Martin <i>et al.</i> (2015) Australia	1896 12–17	L	Y	1, 5 ^b	LT psychotic experiences at baseline	LT attempted suicide at baseline	OR 3.31, 95% CI 0.55–19.94
						Attempted suicide at 12-month follow-up	OR 1.65, 95% CI 0.33–8.26
Sullivan <i>et al.</i> (2015) England	3171 12 and 16	L	N	1, 2	6-month psychotic experiences at age 12	LT suicide attempt at age 16	OR 1.75, 95% CI 1.20–2.54
<i>Suicide attempt [studies with adults (≥16 years)]^c</i>							
Connell <i>et al.</i> (2016) Australia	654 30–33	L	N	1, 3	Auditory or visual hallucinations at age 14	LT suicide attempt at age 30	OR 2.63, 95% CI 1.04–6.63
					Auditory or visual hallucinations at ages 14 and 21	LT suicide attempt at age 30	OR 7.11, 95% CI 2.68–18.83
DeVylder <i>et al.</i> (2015b) ^f USA	11 716 18+	CS	N/A	1	12-month psychotic experiences	12-month suicide attempt	OR 7.79, 95% CI 2.97–20.41
				1, 2, 3	12-month psychotic experiences	12-month suicide attempt	OR 3.37, 95% CI 1.17–9.70
DeVylder <i>et al.</i> (2015a) USA	590 17–37	CS	N/A	1, 2, 3, 4, 5, 7	Past month psychotic experiences	LT suicidal behaviours (attempts, aborted or interrupted)	OR 1.01, 95% CI 0.98–1.04
DeVylder & Kelleher (2016) ^f USA	11 776 18+	CS	N/A	1, 3	LT psychotic experiences	LT suicide attempt	OR 3.47, 95% CI 1.56–7.75
Fisher <i>et al.</i> (2013) New Zealand	1037 11 and 38	L	Y	1	LT psychotic symptoms at age 11	Past year suicide attempts/completes at age 38	RR 3.82, 95% CI 1.67–8.75
				1, 2	LT psychotic symptoms at age 11	Past year suicide attempts/completes at age 38	RR 2.58, 95% CI 1.10–6.08
		L	Y	1			OR 3.95, 95% CI 1.85–8.43

(Continued)

Table 4. (Continued.)

Study	N, Age (years)	Design	Temporality	Confounder (s) ^a	Psychotic experience	Outcome measure	Association between PEs and outcome measure
Honings <i>et al.</i> (2016b) Germany and Netherlands	15 837 M= 39.4 ± 13.8				LT psychotic experiences at baseline	Suicide attempt at 3-year or 5-year follow-up	
				1, 2	LT psychotic experiences at baseline	Suicide attempt at 3-year or 5-year follow-up	OR 3.44, 95% CI 1.54–7.69
Jenkins <i>et al.</i> (2015) Kenya	1133 M= 42.0 ± 18.1	CS	N/A	1, 4	Past year psychotic symptoms	LT suicide attempt	OR 5.1, 95% CI 2.00–13.03
Kelleher <i>et al.</i> (2017) ^d England	7403 16–95	CS	N/A	2 ^b	Past year auditory or visual hallucinations	LT suicide attempt	OR 3.20, 95% CI 1.71–5.98
Koyanagi <i>et al.</i> (2015b) ^d England	7363 16+	CS	N/A	1, 4	12-month any psychotic symptom	12-month suicide attempt	OR 6.52, 95% CI 2.84–14.95
					12-month mania/hypomania		OR 15.67, 95% CI 3.40–72.14
					12-month thought control		OR 2.88, 95% CI 0.71–11.69
					12-month paranoia		OR 19.06, 95% CI 7.48–48.58
					12-month strange experience		OR 6.58, 95% CI 2.57–16.85
					12-month auditory hallucination		OR 6.14, 95% CI 1.36–27.76
				1, 2, 3, 4	12-month any psychotic symptom	12-month suicide attempt	OR 3.84, 95% CI 1.67–8.83
					12-month mania/hypomania		OR 10.23, 95% CI 2.23–46.93
					12-month thought control		OR 1.71, 95% CI 0.45–6.49
					12-month paranoia		OR 9.89, 95% CI 3.48–28.11
					12-month strange experience		OR 3.95, 95% CI 1.44–10.85
					12-month auditory hallucination		OR 5.84, 95% CI 1.25–27.21
Saha <i>et al.</i> (2011a) Australia	8773 18+	CS	N/A	1, 2, 3, 4	LT delusional-like experiences	LT suicide attempt	OR 2.77, 95% CI 1.58–4.85

CS, cross-sectional; L, longitudinal; Y, yes, temporality considered; N, no, temporality not considered; N/A, not applicable; LT, lifetime; OR, odds ratio; RR, relative risk; 95% CI, 95% confidence interval.

Bold represents a significant finding.

^aCorresponds to numbers in Table 1.

^bVariable used as stratification variable in the main analysis, e.g. if mental disorder was the stratification variable, we reported on the influence of psychotic experiences on SITB within the 'no mental disorders' strata.

^cThe adult category (16+) overlaps with the adolescent and young adult category (≤25) because most general population samples range from 16 (or 18) to 95 years.

^dReports on data from the 2007 Adult Psychiatric Morbidity Survey (APMS).

^eReports on data from the 2006 Mie Prefecture high school survey.

^fReports on data from the Collaborative Psychiatric Epidemiology Surveys (CPES).

almost 30% of all general population and community sample studies did not adjust for any confounding variables, possibly resulting in an assumption that SITB is directly attributable to PEs, without taking into consideration factors which confound or mediate this association (see confounder *v.* mediator discussion below). In the majority of studies, where adjustment for confounders occurred, the association persisted albeit with attenuated effect sizes, suggesting PEs are independently associated with self-harm and suicidality, in both adolescent and adult populations. There was high variability in the confounders adjusted for in these studies, with less than half controlling for mental disorders, and less than one-quarter controlling for environmental and psychological factors. Because most studies simultaneously adjusted for multiple variables, no conclusions could be reached as to the influence of an individual factor. However, mental disorders, environmental and psychological factors explained a substantial amount of the variance, whereas most sociodemographic (except for age of PEs in adolescence and sexual orientation) and family history variables were of less importance.

Understanding the role of confounders is important because such variables may mask an actual association or, more commonly, falsely demonstrate an apparent association between the exposure and outcome measure (Skelly *et al.* 2012). When studies appropriately control for such variables, they present a more accurate estimate of the true association due to the exposure (Skelly *et al.* 2012). Results of the current review supported pathway one (see Fig. 1) where PEs are directly causal of SITB. However, the inconsistent and incomplete adjustment for confounders prevents rejection of pathway three (fully explained by shared risk factors) as an explanatory model. Also of note, the associations between PEs and SITB were essentially eliminated in two adolescent/young adult studies with exhaustive adjustments (Nishida *et al.* 2008; DeVylder *et al.* 2015a), suggesting the relationship may indeed be spurious, although these findings would benefit from replication in more broadly representative samples. Alternatively, reverse causality may also be possible, in which the mind's way of adapting to severe distress of SITB results in PEs (Forman *et al.* 2004) (pathway two in Fig. 1). No study to date has tested the reverse causality pathway in the general population, although this explanation is less consistent with broad clinical assumptions.

It is also important to recognise that even if the PE–SITB association becomes non-significant after adjustment, this does not necessarily mean that PEs are not causally important to SITB. Rather it could be the case that the confounders of interest are in fact mediators which explain the association between PEs and SITB on the causal pathway (Christenfeld *et al.* 2004). Jang *et al.* (2014), for example, found support for depressive symptoms as a partial mediator of the PE–suicidal ideation relationship; however, the authors incorporated a cross-sectional mediation analysis, which has faced considerable criticism (Cole & Maxwell, 2003). Confounders and mediators cannot be differentiated statistically, but rather this differentiation is derived from theory and an understanding of the temporal precedence among variables (MacKinnon *et al.* 2000; Babyak, 2009). As explained in online Supplementary Material II, sociodemographics and family history/genetic factors should be treated as confounders, as PEs cannot precede these variables. All other categories should likely be treated as mediating variables, except for substance use and certain environmental factors (e.g. childhood abuse), which have both been consistently shown to precede PEs (Kelleher *et al.* 2013b; McGrath *et al.* 2016). It is also

important to consider that some variables may in fact be moderators (e.g. intervention factors), which influence the strength or direction of the association. All these factors need to be considered in order to advance our understanding of the relationships at play.

Studies with consistent and comprehensive adjustment of third variables are required to explain the PE–SITB relationship. Furthermore, future studies should examine associations of SITB with different PE characteristics, as these have been differentially associated with each NASI outcome. Such PE characteristics include the frequency and emotional valence of content (DeVylder *et al.* 2015a). Theoretical models of self-harm and suicidality may also provide frameworks to better understand the PE–SITB relationship. This includes theories which focus on the transition from suicidal thoughts to actions (Three-Step Theory, Klonsky & May, 2015; Interpersonal Theory, Van Orden *et al.* 2010), as well those which integrate both non-suicidal and suicidal behaviours (Hamza *et al.* 2012), which ideally should be tested using longitudinal data. Schimanski *et al.* (2017) focused on the schizotypy–suicidal ideation relationship and found stronger support for the Interpersonal Theory than for the Three-Step Theory of suicide, suggesting constructs such as thwarted belongingness and perceived burdensomeness should be considered in future investigations. In addition to these theoretical components, there are other potentially confounding variables (e.g. genetic factors), which have yet to be investigated (see the section 'Future research').

Strengths and weaknesses of the review

This systematic review considered and classified all confounding factors, enabling interpretation of individual variables in the PE–SITB relationship. However, meta-analysis was precluded, preventing quantitative measurement of the impact of individual variables on the association. Statistical approaches (e.g. structural equation modelling) are required in existing datasets, as well as in future studies to further test the specific effect of individual confounders and mediators.

Studies should also give consideration to the temporal association between the variables in order to inform possible causality using a discrete-time survival modelling approach. Approximately 30% of all longitudinal studies did not account for SITB at baseline measurement of PEs, and then administered lifetime SITB measures at follow-up, which prevented conclusions as to which occurred first. Also, many studies did not adjust for borderline personality disorder (BPD) or psychotic disorders (only seven studies excluded participants with psychotic disorders). Both BPD and psychosis have high rates of suicide (Hor & Taylor, 2010) and for low prevalence outcomes such as suicide attempts, inclusion of participants with these diagnoses may inflate the PE–SITB association. Koyanagi *et al.* (2015a) found BPD symptoms had a substantial influence on the PE–non-suicidal self-injury association, whereas Kelleher *et al.* (2017) found the PE–suicide attempt association persisted in those without BPD diagnosis or traits. Future studies should consistently account for these diagnoses, where BPD is of particular interest in the PE–non-suicidal self-injury association.

Future research

The PE–SITB association is in part attributable to common mental disorders such as depression; however, more research is

required to determine their specific role as either a confounder or mediator (Jang et al. 2014), particularly given evidence that PEs are indicative of greater severity of underlying depressive and anxiety disorders. To date, environmental factors have not been controlled for in longitudinal studies which consider temporality. In addition, the role of social isolation in the PE–SITB association has not been examined. Isolation and loneliness are associated with both PEs (Lim & Gleeson, 2014) and SITB (Trout, 1980), and are relevant to the Interpersonal Theory construct of thwarted belongingness. Only two studies controlled for emotional dysregulation (Nishida et al. 2008; DeVlyder et al. 2015a), which is surprising considering its key role in the development of non-suicidal and suicidal self-injury (Klonsky, 2007).

There are no studies examining shared genetic risk factors underlying the PE–SITB relationship. Mental disorders, environmental and psychological factors explained much of the variance; however, it is possible that these risk factors are themselves partly explained by genetic factors (DeVlyder et al. 2015a). In addition, cognition, including IQ and other neurocognitive function (e.g. working memory), has not been properly investigated. Calkins et al. (2014, 2017) included an academic achievement variable (WRAT-4 reading subtest) into their PE-suicidal ideation model, and found it had negligible effects (OR range: 0.99–1.00, $p \geq 0.05$). Considering that this variable is modestly related to IQ (Reilly et al. 2014; Olsen et al. 2015), cognitive intelligence may be of less importance. However, other aspects of cognitive function such as working memory and executive function may be of more interest for future research. Finally, poor sleep is a strong risk factor for both PEs (Reeve et al. 2015) and SITB (Hysing et al. 2015) and may be important in the PE–SITB relationship.

Conclusion

This systematic review critically examined confounders and mediators adjusted for in the PE–SITB association. In most studies, the association between PEs and both suicidal ideation and NASI remained significant after adjustments, suggesting an independent association. A substantial amount of the variance was explained by mental disorders, psychological distress and environmental exposures. Interventions to target these symptoms of mental ill health in those with PEs may assist in reducing risk of self-harm and suicide. In order to fully understand the mechanisms underpinning the PE–SITB association, hypothesis driven studies with more consistent and comprehensive adjustment for third variables are required, as well as studies which clarify the phenomenological aspects of different types of PEs. Such efforts will inform intervention, which in turn may reduce the risk of morbidity and mortality by suicide in those with PEs.

Notes

¹ This broader term takes into account the fact that motivation for self-injurious behaviour is often complex (Hjelmeland et al. 2002).

² For example, have you thought about hurting yourself?

³ Skin picking (or dermatillomania/excoriation disorder) and hair pulling (or trichotillomania) have been established as separate disorders in the clinical literature (Zucker et al. 2011; Grant & Chamberlain 2015).

⁴ Schizotypy is typically defined as traits stable over time with anomalies across multiple dimensions (perceptual, interpersonal), whereas PEs are more transitional anomalies (Pedrero & Debbane 2017).

⁵ Separate search strategies were devised for each database. Relevant index terms were incorporated, including MeSH Terms for PubMed, Emtree terms for EMBASE, and the Thesaurus of Psychological Index Terms for PsycINFO.

⁶ Because this study (Kelleher et al. 2012b) collapsed across SITB outcomes (suicidal ideation, suicide attempt, non-suicidal self-injury), it was excluded from results in Tables 3 and 4.

Supplementary material. The supplementary material for this article can be found at <https://doi.org/10.1017/S0033291717002677>

Acknowledgements. We thank Nicky Foxlee, a medical librarian, who assisted with devising the search strategy for this systematic review. We also thank Professor John McGrath for providing comments on earlier versions of the manuscript. EH is supported by the Dr F and Mrs ME Zaccari Scholarship, Australia. JGS is supported by a National Health and Medical Research Council Practitioner Fellowship Grant (grant number 1105807).

Declaration of Interest. None.

References

- Babjak MA (2009) Understanding confounding and mediation. *Evidence-Based Mental Health* 12, 68–71.
- Calkins ME et al. (2014) The psychosis spectrum in a young U.S. community sample: findings from the Philadelphia Neurodevelopmental Cohort. *World Psychiatry* 13, 296–305.
- Calkins ME et al. (2017) Persistence of psychosis spectrum symptoms in the Philadelphia Neurodevelopmental Cohort: a prospective two-year follow-up. *World Psychiatry* 16, 62–76.
- Capra C et al. (2015) Subtypes of psychotic-like experiences are differentially associated with suicidal ideation, plans and attempts in young adults. *Psychiatry Research* 228, 894–898.
- Cederlöf M et al. (2016) A longitudinal study of adolescent psychotic experiences and later development of substance use disorder and suicidal behavior. *Schizophrenia Research* 181, 13–16.
- Christenfeld NJ et al. (2004) Risk factors, confounding, and the illusion of statistical control. *Psychosomatic Medicine* 66, 868–875.
- Cole DA and Maxwell SE (2003) Testing mediational models with longitudinal data: questions and tips in the use of structural equation modeling. *Journal of Abnormal Psychology* 112, 558–577.
- Connell M et al. (2016) Hallucinations in adolescents and risk for mental disorders and suicidal behaviour in adulthood: prospective evidence from the MUSP birth cohort study. *Schizophrenia Research* 176, 546–551.
- DeVlyder JE and Hilimire MR (2015) Suicide risk, stress sensitivity, and self-esteem among young adults reporting auditory hallucinations. *Health & Social Work* 40, 175–181.
- DeVlyder JE et al. (2015a) Social and psychological contributions to the co-occurrence of sub-threshold psychotic experiences and suicidal behavior. *Social Psychiatry and Psychiatric Epidemiology* 50, 1819–1830.
- DeVlyder JE and Kelleher I (2016) Clinical significance of psychotic experiences in the context of sleep disturbance or substance use. *Psychological Medicine* 46, 1761–1767.
- DeVlyder JE et al. (2015b) Suicidal ideation and suicide attempts among adults with psychotic experiences: data from the collaborative psychiatric epidemiology surveys. *JAMA Psychiatry* 72, 219–225.
- DeVlyder JE et al. (2015c) Psychotic experiences as indicators of suicidal ideation in a non-clinical college sample. *Psychiatry Research* 226, 489–493.
- Engin E et al. (2009) University students' suicidal thoughts and influencing factors. *Journal of Psychiatric and Mental Health Nursing* 16, 343–354.
- Fisher HL et al. (2013) Specificity of childhood psychotic symptoms for predicting schizophrenia by 38 years of age: a birth cohort study. *Psychological Medicine* 43, 2077–2086.
- Forman EM et al. (2004) History of multiple suicide attempts as a behavioral marker of severe psychopathology. *American Journal of Psychiatry* 161, 437–443.
- Franklin JC et al. (2017) Risk factors for suicidal thoughts and behaviors: a meta-analysis of 50 years of research. *Psychological Bulletin* 143, 187–232.
- Freeman D et al. (2011) Concomitants of paranoia in the general population. *Psychological Medicine* 41, 923–936.
- Gatta M et al. (2016) Non-suicidal self-injury among Northern Italian high school students: emotional, interpersonal and psychopathological correlates. *Journal of Psychopathology* 22, 185–190.

- St. Germain SA and Hooley JM** (2012) Direct and indirect forms of non-suicidal self-injury: evidence for a distinction. *Psychiatry Research* **197**, 78–84.
- Grandclerc S et al.** (2016) Relations between nonsuicidal self-injury and suicidal behavior in adolescence: a systematic review. *PLoS ONE* **11**, e0153760.
- Grant JE and Chamberlain SR** (2015) Trichotillomania and skin picking disorder: different kinds of OCD. *The Journal of Lifelong Learning in Psychiatry* **13**, 184–189.
- Guan K, Fox KR and Prinstein MJ** (2012) Nonsuicidal self-injury as a time-invariant predictor of adolescent suicide ideation and attempts in a diverse community sample. *Journal of Consulting and Clinical Psychology* **80**, 842–849.
- Hamza CA, Stewart SL and Willoughby T** (2012) Examining the link between nonsuicidal self-injury and suicidal behavior: a review of the literature and an integrated model. *Clinical Psychology Review* **32**, 482–495.
- Hawton K et al.** (2003) Deliberate self-harm in Oxford, 1990–2000: a time of change in patient characteristics. *Psychological Medicine* **33**, 987–995.
- Hjelmeland H et al.** (2002) Why people engage in parasuicide: a cross-cultural study of intentions. *Suicide and Life-Threatening Behavior* **32**, 380–393.
- Honings S et al.** (2016a) Psychotic experiences and risk of self-injurious behaviour in the general population: a systematic review and meta-analysis. *Psychological Medicine* **46**, 237–251.
- Honings S et al.** (2016b) Psychotic experiences and incident suicidal ideation and behaviour: disentangling the longitudinal associations from connected psychopathology. *Psychiatry Research* **245**, 267–275.
- Hor K and Taylor M** (2010) Suicide and schizophrenia: a systematic review of rates and risk factors. *Journal of Psychopharmacology* **24**, 81–90.
- Hysing M et al.** (2015) Sleep problems and self-harm in adolescence. *British Journal of Psychiatry* **207**, 306–312.
- Jang JH et al.** (2014) Psychotic-like experiences and their relationship to suicidal ideation in adolescents. *Psychiatry Research* **215**, 641–645.
- Jenkins R et al.** (2015) Tedium vitae, death wishes, suicidal ideation and attempts in Kenya-prevalence and risk factors. *BMC Public Health* **15**, 759.
- Johns LC et al.** (2004) Prevalence and correlates of self-reported psychotic symptoms in the British population. *British Journal of Psychiatry* **185**, 298–305.
- Kapur N et al.** (2013) Non-suicidal self-injury v. attempted suicide: new diagnosis or false dichotomy? *British Journal of Psychiatry* **202**, 326–328.
- Kaymaz N et al.** (2012) Do subthreshold psychotic experiences predict clinical outcomes in unselected non-help-seeking population-based samples? A systematic review and meta-analysis, enriched with new results. *Psychological Medicine* **42**, 2239–2253.
- Kelleher I, Cederlöf M and Lichtenstein P** (2014) Psychotic experiences as a predictor of the natural course of suicidal ideation: a Swedish cohort study. *World Psychiatry* **13**, 184–188.
- Kelleher I et al.** (2012a) Prevalence of psychotic symptoms in childhood and adolescence: a systematic review and meta-analysis of population-based studies. *Psychological Medicine* **9**, 1–7.
- Kelleher I et al.** (2013a) Psychotic symptoms and population risk for suicide attempt: a prospective cohort study. *JAMA Psychiatry* **70**, 940–948.
- Kelleher I et al.** (2013b) Childhood trauma and psychosis in a prospective cohort study: cause, effect, and directionality. *American Journal of Psychiatry* **170**, 734–741.
- Kelleher I et al.** (2012b) Psychotic symptoms in adolescence index risk for suicidal behavior: findings from 2 population-based case-control clinical interview studies. *Archives of General Psychiatry* **69**, 1277–1283.
- Kelleher I, Ramsay H and DeVlyder J** (2017) Psychotic experiences and suicide attempt risk in common mental disorders and borderline personality disorder. *Acta Psychiatrica Scandinavica* **135**, 212–218.
- Kerr PL and Muehlenkamp JJ** (2010) Features of psychopathology in self-injuring female college students. *Journal of Mental Health Counselling* **32**, 290–308.
- Klonsky ED** (2007) The functions of deliberate self-injury: a review of the evidence. *Clinical Psychology Review* **27**, 226–239.
- Klonsky ED and May AM** (2015) The three-step theory (3ST): a new theory of suicide rooted in the 'Ideation-to-Action' framework. *International Journal of Cognitive Therapy* **8**, 114–129.
- Klonsky ED, May AM and Saffer BY** (2016) Suicide, suicide attempts, and suicidal ideation. *Annual Review of Clinical Psychology* **12**, 307–330.
- Koyanagi A, Stickle A and Haro JM** (2015a) Psychotic-like experiences and nonsuicidal self-injury in England: results from a national survey. *PLoS ONE* **10**, e0145533.
- Koyanagi A, Stickle A and Haro JM** (2015b) Subclinical psychosis and suicidal behavior in England: findings from the 2007 adult psychiatric morbidity survey. *Schizophrenia Research* **168**, 62–67.
- Krabbendam L et al.** (2004) Dimensions of depression, mania and psychosis in the general population. *Psychological Medicine* **34**, 1177–1186.
- Larkin C, Di Blasi Z and Arensman E** (2014) Risk factors for repetition of self-harm: a systematic review of prospective hospital-based studies. *PLoS ONE* **9**, e84282.
- Lewis-Fernández R et al.** (2009) Significance of endorsement of psychotic symptoms by US Latinos. *Journal of Nervous and Mental Disease* **197**, 337–347.
- Lim MH and Gleeson JF** (2014) Social connectedness across the psychosis spectrum: current issues and future directions for interventions in loneliness. *Frontiers in Psychiatry* **5**, 154.
- Linehan MM** (1986) Suicidal people: one population or two? *Annals of the New York Academy of Sciences* **487**, 16–27.
- Linscott RJ and van Os J** (2010) Systematic reviews of categorical versus continuum models in psychosis: evidence for discontinuous subpopulations underlying a psychometric continuum. Implications for DSM-V, DSM-VI, and DSM-VII. *Annual Review of Clinical Psychology* **6**, 391–419.
- Linscott RJ and van Os J** (2013) An updated and conservative systematic review and meta-analysis of epidemiological evidence on psychotic experiences in children and adults: on the pathway from proneness to persistence to dimensional expression across mental disorders. *Psychological Medicine* **43**, 1133–1149.
- MacKinnon DP, Krull JL and Lockwood CM** (2000) Equivalence of the mediation, confounding and suppression effect. *Prevention Science* **1**, 173–181.
- Martin G et al.** (2015) Psychotic experiences and psychological distress predict contemporaneous and future non-suicidal self-injury and suicide attempts in a sample of Australian school-based adolescents. *Psychological Medicine* **45**, 429–437.
- McGrath JJ et al.** (2015) Psychotic experiences in the general population: a cross-national analysis based on 31,261 respondents from 18 countries. *JAMA Psychiatry* **72**, 697–705.
- McGrath JJ et al.** (2016) The bidirectional associations between psychotic experiences and DSM-IV mental disorders. *American Journal of Psychiatry* **173**, 997–1006.
- Moher D et al.** (2009) Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *PLoS Medicine* **6**, e1000097.
- Muehlenkamp JJ et al.** (2012) International prevalence of adolescent non-suicidal self-injury and deliberate self-harm. *Child and Adolescent Psychiatry and Mental Health* **6**, 10.
- Nishida A et al.** (2010) Psychotic-like experiences are associated with suicidal feelings and deliberate self-harm behaviors in adolescents aged 12–15 years. *Acta Psychiatrica Scandinavica* **121**, 301–307.
- Nishida A et al.** (2014) Risk for suicidal problems in poor-help-seeking adolescents with psychotic-like experiences: findings from a cross-sectional survey of 16,131 adolescents. *Schizophrenia Research* **159**, 257–262.
- Nishida A et al.** (2008) Associations between psychotic-like experiences and mental health status and other psychopathologies among Japanese early teens. *Schizophrenia Research* **99**, 125–133.
- Nock MK et al.** (2007) Self-Injurious thoughts and behaviors interview: development, reliability, and validity in an adolescent sample. *Psychological Assessment* **19**, 309–317.
- Núñez D, Arias VB and Campos S** (2016) The reliability and validity of Liu's self-report questionnaire for screening putative pre-psychotic states (BQSPS) in adolescents. *PLoS ONE* **11**, e0167982.
- Olson M et al.** (2002) Psychotic symptoms in an urban general medicine practice. *American Journal of Psychiatry* **159**, 1412–1419.
- Olson M et al.** (1996) Psychotic symptoms in primary care. *The Journal of Family Practice* **43**, 481–488.
- Olsen JP et al.** (2015) Reading ability as an estimator of premorbid intelligence: does it remain stable among ethnically diverse HIV+ adults? *The Clinical Neuropsychologist* **29**, 1034–1052.

- Östling S, Börjesson-Hanson A and Skoog I** (2007) Psychotic symptoms and paranoid ideation in a population-based sample of 95-year-olds. *The American Journal of Geriatric Psychiatry* **15**, 999–1004.
- Östling S and Skoog I** (2002) Psychotic symptoms and paranoid ideation in a nondemented population-based sample of the very old. *Archives of General Psychiatry* **59**, 53–59.
- Palmer BA, Pamkratz S and Botswick JM** (2005) The lifetime risk of suicide in schizophrenia: a reexamination. *Archives of General Psychiatry* **62**, 247–253.
- Pedrero EF and Debbané M** (2017) Schizotypal traits and psychotic-like experiences during adolescence: an update. *Psicothema* **29**, 5–17.
- Peterson J *et al.*** (2008) Nonsuicidal self-injury in adolescents. *Psychiatry* **5**, 20–26.
- Platt S *et al.*** (1992) Parasuicide in Europe: the WHO/EURO multicentre study on parasuicide. I. Introduction and preliminary analysis for 1989. *Acta Psychiatrica Scandinavica* **85**, 97–104.
- Polanczyk G *et al.*** (2010) Etiological and clinical features of childhood psychotic symptoms. *Archives of General Psychiatry* **67**, 328–338.
- Reeve S, Sheaves B and Freeman D** (2015) The role of sleep dysfunction in the occurrence of delusions and hallucinations: a systematic review. *Clinical Psychology Review* **42**, 96–115.
- Reilly C *et al.*** (2014) Academic achievement in school-aged children with active epilepsy: a population-based study. *Epilepsia* **55**, 1910–1917.
- Rognli EB *et al.*** (2017) Substance use and sociodemographic background as risk factors for lifetime psychotic experiences in a non-clinical sample. *Journal of Substance Abuse Treatment* **74**, 42–47.
- Saha S *et al.*** (2011a) The association between delusional-like experiences and suicidal thoughts and behaviour. *Schizophrenia Research* **132**, 197–202.
- Saha S *et al.*** (2011b) The association between trauma and delusional-like experiences. *Psychiatry Research* **189**, 259–264.
- Schimanski ID *et al.*** (2017) Preliminary evidence that schizophrenia liability at age 15 predicts suicidal ideation two years later. *Schizophrenia Research* **181**, 60–62.
- Scott JG *et al.*** (2007) Association between trauma exposure and delusional experiences in a large community-based sample. *British Journal of Psychiatry* **190**, 339–343.
- Scott LN *et al.*** (2015) Non-suicidal self-injury and suicidal ideation as predictors of suicide attempts in adolescent girls: a multi-wave prospective study. *Comprehensive Psychiatry* **58**, 1–10.
- Sharifi V *et al.*** (2015) Psychotic experiences and risk of death in the general population: 24–27 year follow-up of the Epidemiologic Catchment Area study. *British Journal of Psychiatry* **207**, 30–36.
- Skelly AC, Dettori JR and Brodt ED** (2012) Assessing bias: the importance of considering confounding. *Evidence-Based Spine-Care Journal* **3**, 9–12.
- Stanley B *et al.*** (1992) Suicide and the self-harm continuum: phenomenological and biochemical evidence. *International Review of Psychiatry* **4**, 149–155.
- Sullivan SA *et al.*** (2015) The longitudinal association between psychotic experiences, depression and suicidal behaviour in a population sample of adolescents. *Social Psychiatry and Psychiatric Epidemiology* **50**, 1809–1817.
- Taylor PJ, Hutton P and Wood L** (2015) Are people at risk of psychosis also at risk of suicide and self-harm? A systematic review and meta-analysis. *Psychological Medicine* **45**, 911–926.
- Temmingh H *et al.*** (2011) The prevalence and correlates of hallucinations in a general population sample: findings from the South African Stress and Health Study. *African Journal of Psychiatry* **14**, 211–217.
- Trout DL** (1980) The role of social isolation in suicide. *Suicide and Life-Threatening Behavior* **10**, 10–23.
- Van Orden KA *et al.*** (2010) The interpersonal theory of suicide. *Psychological Review* **117**, 575–600.
- Vijayakumar L, Kumar MS and Vijayakumar V** (2011) Substance use and suicide. *Current Opinion in Psychiatry* **24**, 197–202.
- Whitlock J *et al.*** (2013) Nonsuicidal self-injury as a gateway to suicide in young adults. *Journal of Adolescent Health* **52**, 486–492.
- Witorsch P and Witorsch RJ** (1993) Analysis of potential confounding variables in epidemiologic studies of parental/ household smoking and respiratory health in preschool children. *Indoor and Built Environment* **2**, 71–91.
- Yung AR and Lin A** (2016) Psychotic experiences and their significance. *World Psychiatry* **15**, 130–131.
- Zetterqvist M** (2015) The DSM-5 diagnosis of nonsuicidal self-injury disorder: a review of the empirical literature. *Child and Adolescent Psychiatry and Mental Health* **9**, 31.
- Zhang J, Song J and Wang J** (2016) Adolescent self-harm and risk factors. *Asia-Pacific Psychiatry* **8**, 287–295.
- Zucker N *et al.*** (2011) The significance of repetitive hair-pulling behaviors in eating disorders. *Journal of Clinical Psychology* **67**, 391–403.