

# Elimination of urogenital schistosomiasis in Iran: past history and the current situation

SHAHRAM KHADEMVATAN<sup>1,2</sup>, SHOKROLLAH SALMANZADEH<sup>1\*</sup>,  
MASOUD FOROUTAN-RAD<sup>3,4</sup> and MOHAMMAD GHOMESHI<sup>5</sup>

<sup>1</sup> Health Research Institute, Infectious and Tropical Diseases Research Center, Ahvaz Jundishapur University of Medical Sciences, Ahvaz, Iran

<sup>2</sup> Department of Medical Parasitology and Mycology & Cellular and Molecular Research Center, Urmia University of Medical Sciences, Urmia, Iran

<sup>3</sup> Student Research Committee, Ahvaz Jundishapur University of Medical Sciences, Ahvaz, Iran

<sup>4</sup> Department of Medical Parasitology, Faculty of Medicine, Ahvaz Jundishapur University of Medical Sciences, Ahvaz, Iran

<sup>5</sup> CDC Department, Deputy of Health, Ahvaz Jundishapur University of Medical Sciences, Ahvaz, Iran

(Received 10 February 2016; revised 5 April 2016; accepted 19 April 2016; first published online 9 June 2016)

## SUMMARY

In recent years, through a national programme for schistosomiasis control, this infection has been eliminated from Iran. The aim of this study was to report the process of significant decrease of urogenital schistosomiasis in southwestern Iran. During national programme surveillance for urogenital schistosomiasis control which was implemented by Centres for Disease Control and Prevention (CDC) of Khuzestan province from 1975 to 2013, more than 1·3 million urine samples were taken from inhabitants of high risk foci. All urine samples were gathered between 10:00 a.m and 02:00 p.m and, after centrifuging, specimens were tested under optical microscope in order to detect *Schistosoma haematobium* eggs. Data analysis was performed using SPSS 18 software. In this retrospective study significant reduction was seen in number of infections between 1975 and 2013. During the years 1975–1980, 1981–1990 and 1991–2000 there were 1582, 761 and 79 cases of *S. haematobium*, respectively. In 2001 only one case was reported from Ahvaz and indeed this was the last case of urogenital schistosomiasis in Khuzestan and of course, in Iran. Prevalence from 1·064% between 1975 and 1980 slumped to 0% in 2012–2013. During several projects for surveillance of urogenital schistosomiasis, selective population chemotherapy, snail control, population education, environmental improvement, etc were carried out throughout the surveillance period. According to elimination of *S. haematobium* in Khuzestan province, the only endemic region of Iran, control of disease, especially the campaign with intermediate host snails should be continued. Iran can be a successful model for countries suffering from this disease.

Key words: *Schistosoma haematobium*, *Bulinus truncatus*, elimination, control, Iran, Khuzestan.

## INTRODUCTION

Schistosomiasis or bilharziasis, is a water borne parasitic infection and, as one of the most common tropical diseases after malaria is placed in the second rank. World Health Organization considers it a Neglected Tropical Disease (NTD) (Fenwick *et al.* 2009). Schistosomiasis is endemic in 78 developing countries throughout the world and over 240 million persons are infected in rural and semi-urban regions. From this number, about 20 million people have severe complications and approximately 120 million have signs of the infection. It is estimated that 800 million individuals live in endemic and high risk regions throughout the world, and therefore may acquire the disease and among 200 000–280 000 deaths as a result of disease occur annually. Also, more than 85% of cases happened in sub-Saharan

Africa (Fenwick *et al.* 2009; Mo *et al.* 2014). Schistosomiasis is considered as one of the main infections of public health in Eastern Mediterranean Region (EMR) countries and has socio-economic importance. There are two main human *Schistosoma* species in EMR region, *Schistosoma haematobium* and *Schistosoma mansoni*. *Schistosoma mansoni* exists in 9 out of 23 EMR countries (Egypt, Libya, Saudi Arabia, Oman, Djibouti, Yemen, Sudan, Southern Sudan and Somalia) and *S. haematobium* exists in 15 EMR countries (Libya, Egypt, Saudi Arabia, Oman, Sudan, Southern Sudan, Yemen, Somalia, Jordan, Morocco, Tunisia, Lebanon, Syria, Iraq and Iran) (WHO, 2013). In recent years a significant decline has been observed in the prevalence of urogenital schistosomiasis in many EMR endemic regions and some of these countries have reached elimination of infection (Al Ghahtani and Amin, 2005; EMRO/WHO, 2007; Niaz *et al.* 2010; Amarir *et al.* 2011; WHO, 2013; Othman and Soliman, 2015). Based on recent policy by WHO, Iran is categorized among countries that require evaluation in order to verify if interruption

\* Corresponding author: Health Research Institute, Infectious and Tropical Diseases Research Center, Ahvaz Jundishapur University of Medical Sciences, Ahvaz, Iran. Tel: (+98 613)3367543-50. Fax: (+98 613) 3332036. E-mail: [salmanidmd@yahoo.com](mailto:salmanidmd@yahoo.com)

of transmission has been achieved and updating planning and implementation purposes (WHO, 2013).

### *Urogenital schistosomiasis in Iran*

In Iran, Khuzestan is the only province in which urogenital schistosomiasis has been reported (Arfaa *et al.* 1967; Massoud *et al.* 1982; Mombeni and Kheradmand, 2005; Gholamreza *et al.* 2008). Of the different species of Schistosomes, only *S. haematobium* has been reported in Iran so only urinary schistosomiasis has been diagnosed in this area (Arfaa *et al.* 1967; Mombeni and Kheradmand, 2005; Hamidinia *et al.* 2014). The presence of raging rivers, cane sugar fields, hydroelectric and agricultural dams and the optimum temperature and humidity have made Khuzestan a favourable habitat for the snail *Bulinus truncatus*, which is the intermediate host for *S. haematobium* (Hamidinia *et al.* 2014). According to previous studies, the urogenital schistosomiasis was limited to a few foci of Khuzestan province and it is estimated that 25 000–30 000 individuals were infected (Massoud *et al.* 1982). Iraq, a country located to the west of Iran, is considered as an endemic region for urogenital schistosomiasis (WHO, 2013). In recent years, by the implementation of a national programme for schistosomiasis control, this infection has been eliminated from Iran. The aim of this study was to report some aspects of the control interventions and the process of dramatic reduction of urogenital schistosomiasis in southwestern Iran, Khuzestan province.

## MATERIALS AND METHODS

### *Study region*

Khuzestan province is located in the southwest of Iran and can be basically divided into two regions, the rolling hills and mountainous regions north of the Ahvaz Ridge and the plains and marshlands to its south. The province covers an area of about 64 055 Km<sup>2</sup> and has a population of approximately 4 531 720 inhabitants (2 286 209 male and 2 245 511 female). It is a continuation of the Mesopotamian plain, and is bordered on the south by the Persian Gulf, on the west by Iraq, and on the north and east by the Zagros mountains (Salmanzadeh *et al.* 2015). The area is irrigated by the Karoun, Karkheh, Jarahi, Kheirabad, Dez and Maroun rivers (Fig. 1). Khuzestan has great potential for agricultural expansion, which is almost unrivalled by the country's other provinces. Large and permanent rivers flow over the entire territory contributing to the fertility of the land. The Karun, Iran's most effluent river, 850 kms long, flows into the Persian Gulf through this province. The climate of Khuzestan is generally hot and occasionally humid, particularly in the south, while winters are much more pleasant and dry. Summertime temperatures

routinely exceed 50°C and in the winter it can drop below freezing, with occasional snowfall all the way south to Ahvaz. Khuzestan province is known to master the hottest temperatures on record for a populated city anywhere in the world. Many sandstorms and dust storms are frequent with the arid and desert-style terrains ([http://en.wikipedia.org/wiki/Khuzestan province](http://en.wikipedia.org/wiki/Khuzestan_province)).

### *Data collection*

The current data about epidemiological status of schistosomiasis were gathered from Centres for Disease Control and Prevention (CDC) of Khuzestan province, in the southwest of Iran during 1975–2013. This centre is responsible for annual surveillance of urogenital schistosomiasis in high risk regions of the province. Also, active and passive case finding, follow-up of the patients, detection of natural habitats for *B. truncatus* snails and mollusciciding are the other activities of CDC. Briefly, in the period of surveillance more than 1.3 million urine samples were taken from inhabitants, mainly suspected persons with hematuria and/or bladder irritability symptoms of high risk zones, including: Ahvaz, Andimeshk, Shush, Shushtar and Dezful. All urine samples were gathered between 10:00 a.m and 02:00 p.m for ova detection based on sedimentation method and under light microscope. Also, the aforementioned regions were searched in order to detect *B. truncatus* habitats.

## RESULTS

In this retrospective cross-sectional study a significant decrease was seen in number of *S. haematobium* infections between 1975 and 2013. During these years more than 1.3 million urine samples were examined in five high risk foci of *S. haematobium* to detect ova. Between the years 1975–1980, 1981–1990 and 1991–2000 there were 1582, 761 and 79 cases of *S. haematobium*, respectively (Table 1). In 2001, only one case was reported from the west of Ahvaz and indeed this was the last case of urogenital schistosomiasis in Khuzestan and of course, in Iran (Table 1). In the past decade from 2002 to 2013, from more than 0.4 million urine samples taken, no eggs were detected and no patient was diagnosed (Table 1). Also, prevalence of *S. haematobium* from 1.064% between 1975 and 1980 slumped to 0% in 2012–2013 and total prevalence from 1975 to 2013 was 0.176% (Fig. 2). Despite the absence of urogenital schistosomiasis in Khuzestan province, urine sampling from suspected cases and investigation of previously infected villages and the natural habitats for *B. truncatus* snail are currently being continued (Table 2). From multiple foci in the past, at present (during 2012–2013) only 83 foci are diagnosed as natural habitats for *B. truncatus* snail in



Fig. 1. Location of Khuzestan province in Iran.

Table 1. Number of urine samples for *S. haematobium* case finding/number of confirmed *S. haematobium* patients in Khuzestan province, southwestern Iran (1975–2013)

Year	City						Total (n)
	Ahvaz	Andimeshk	Shush	Shushtar	Dezful		
1975–1980	116603/704	5964/0	–/174	–/28	26076/676		148643/1582
1981–1990	136382/134	11250/2	–/318	–/43	73740/264		221372/761
1991–2000	175703/35	17956/1	96987/19	–/5	218012/19		508658/79
2001–2011	72317/1	33842/0	196399/0	34056/0	125793/0		462407/1
2012–2013	611/0	6128/0	8338/0	2360/0	12649/0		30086/0
Total	501616/874	75140/3	301724/511	36416/76	456270/959		1371166/2423




Fig. 2. Prevalence of *S. haematobium* in Khuzestan province, southwestern Iran (1975–2013).

Khuzestan, 32 foci in Andimeshk and 51 foci in Dezful districts (Table 2).

DISCUSSION

The purpose of current study was to report the process of dramatic reduction of urogenital schistosomiasis in Iran. According to our finding, prevalence of *S. haematobium* has significantly declined during the four recent decades in the only endemic region in the southwest of Iran, Khuzestan province and showed elimination had been achieved. Previous investigations demonstrated the elimination of urinary schistosomiasis in this region of Iran (Gholamreza *et al.* 2008; Mombeni and Kheradmand, 2005; Hamidinia *et al.* 2014); although, transitional bladder cell carcinoma and schist-ogranuloma of the fallopian tubes due to

Table 2. Number of villages investigated for *B. truncatus*/number of natural habitats for *B. truncatus* snails detected in Khuzestan province, southwestern Iran (1975–2013)

Year	City					Total (n)
	Ahvaz	Andimeshk	Shush	Shushtar	Dezful	
1975–1980	15/81	17/17	–	–	12/109	44/207
1981–1990	–	13/14	–	5/6	19/245	37/265
1991–2000	9/67	12/12	3249/449	65/9	21/332	3356/869
2001–2011	–	8/8	4009/62	65/1	14/201	4096/272
2012–2013	18/0	108/32	253/0	49/0	242/51	670/83
Total (n)	42/148	158/83	7511/511	184/16	308/938	8203/1696

*S. haematobium* in Iran were reported (Maraghi *et al.* 2004; Ketabchi and Moshtaghi-Kashanian, 2012). Based on Masooud *et al.*'s study in selected high risk regions of Khuzestan province, in the southwest of Iran during 1970–1979, prevalence of urinary schistosomiasis was 8.3% in 1970 and fell to 0.74% in 1979. Moreover, overall incidence of schistosomiasis infection dramatically declined from 3.5% in 1970 to 0% in 1977 (Massoud *et al.* 1982). In a follow-up survey conducted by Gholamreza *et al.* in some areas of Khuzestan province, out of 3400 urine samples collected during 2005–2007 from villagers residing in the province, no *S. haematobium* positive cases were seen (Gholamreza *et al.* 2008). *S. haematobium* is still endemic in Iraq, a country located to the west of Iran (WHO, 2013), thus; in order to eradicate disease greater control of traffic between the two countries is required and this process till elimination in Iraq should be continued. According to recommended strategies of WHO, significant advances against schistosomiasis in EMR countries have been achieved during the past four decades (WHO, 2013). Schistosomiasis in some EMR countries has been eliminated like Lebanon, Tunisia, Morocco and Iran and no positive cases have been diagnosed during the past few years (Gholamreza *et al.* 2008; Amarir *et al.* 2011; Rollinson *et al.* 2013; WHO, 2013; Hamidinia *et al.* 2014). Also, some EMR countries such as Jordan, Egypt, Iraq, Oman, Libya, Syria and Saudi Arabia have reached low endemicity. For example in Egypt, prevalence of infection before starting national schistosomiasis control programme (NSCP) in 1967 was calculated approximately 40% and after implementation of this programme, overall prevalence fell down to less than 3%. Also, in Egypt prevalence of *S. haematobium* between 1988 and 2001 declined from 11.9 to 1.3% (EMRO/WHO, 2007; WHO, 2013; Othman & Soliman, 2015). Niclosamide molluscicide is widely used for snail control and affects all stages of the snail life cycle (McCullough *et al.* 1980). Despite the negative effects of niclosamide on fishes and environment (Woolhouse *et al.* 1998), it has a key role in control and elimination of disease in some countries like Zanzibar (Knopp *et al.* 2013), Mauritius (Dhunpath, 1994), Morocco (Amarir *et al.* 2011), Saudi Arabia (Al Ghahtani and Amin, 2005), Tunisia (Rollinson

*et al.* 2013) and our country, Iran (Massoud *et al.* 1982; Mombeni & Kheradmand, 2005; Gholamreza *et al.* 2008).

In a study conducted by Amarir *et al.* (2011) in Morocco, from 2382 individuals participated, no positive cases were reported, which proved elimination. In another survey performed by Niaz *et al.* in Pakistan, no positive cases (0%) were seen and authors declared Pakistan free from human schistosomiasis (Niaz *et al.* 2010). These findings are in agreement with our study. In contrast to our results, *S. haematobium* has a high rate in some countries and is endemic, like Senegal (Meurs *et al.* 2012), Cameroon (Cunin *et al.* 2003), Nigeria (Chidozie and Daniyan, 2008; Singh and Muddasiru, 2014), Sudan (Ismail *et al.* 2014), Zambia (Chomba and Mutale, 2014), Zanzibar (Rudge *et al.* 2008) and Yemen (Sady *et al.* 2013).

Virtually the same strategies have been employed in countries, which reached to schistosomiasis elimination. For instance, during the 1950s an interdisciplinary, intersectoral and nationwide collaboration was implemented by the Japanese government, which diminished the rate of *Schistosoma japonicum* significantly (Kasai *et al.* 2007; Rollinson *et al.* 2013). Also, *S. haematobium* control strategies in Tunisia were launched from 1970 and were focused on interrupting transmission by treatment of afflicted persons, usage from molluscicides in endemic foci, improvement of water resources and agricultural infrastructure (EMRO/WHO, 2007; Rollinson *et al.* 2013). Japan and Morocco could be considered as successful models for multi sectoral engagement and cooperation in elimination of *S. japonicum* and *S. haematobium*, respectively (Rollinson *et al.* 2013).

#### *The urogenital schistosomiasis control programme of Iran*

First efforts to control urogenital schistosomiasis in Iran were started in 1959 by Massoud *et al.* (Massoud *et al.* 1982) and considerable amount of baseline data was collected before experimental and large scale control procedures were presented. Control measures were begun in the high risk

regions using a combination of chemotherapy with different antischistosomal drugs and snail control, environmental improvement including sanitary measures, and focal mollusciciding of snail habitats. During several projects for surveillance of urogenital schistosomiasis, selective population chemotherapy and mollusciciding were carried out throughout the surveillance period (1975–2013), but the main activities were concentrated in spring and autumn seasons. During this programme more than 1.3 million urine samples were taken from residents in 5 high risk foci (Ahvaz, Andimeshk, Shush, Shush-tar and Dezful) (Fig. 1) under the supervision of CDC of the region and many of the cases were treated. All urine samples were gathered during the years between 10:00 a.m and 02:00 p.m to detect *S. haematobium* infection. Positive cases were treated with niridazole (30 mg kg BW<sup>-1</sup>) daily for 4 consecutive days. Because of the side effects of this drug and its poor tolerability it was replaced by praziquantel in the early years of 1980s, (Massoud *et al.* 1982). Follow-up of urine samples was also done for all positive patients to confirm their cure. Each patient had his own folder and all epidemiological data, laboratory findings, prescribed drug and the outcome of the therapy were recorded in it. Also, active case finding was conducted by testing of urine samples from individuals less than 15 years old from numerous villages in different parts of endemic regions with known cases of *S. haematobium* infection or in areas in which the intermediate snail was detected. All children who were negative in initial exam were reexamined in order to determine the disease incidence at yearly intervals. In addition, reports from physician's offices and hospitals were recorded and treated.

Our success in elimination of *S. haematobium* originated from constant implementation of disease control, briefly: traffic control between Iran and Iraq (endemic region), campaign with snails as intermediate hosts (*B. truncatus*), health education, appropriate disposal of feces and urine, pipe borne water for all parts of rural and urban regions. Besides these efforts education programmes were conducted for the people living in the endemic areas to break the transmission cycle through reduced human–water contact and diminished environmental contamination with excreta. Projects for improvements in water supply and sanitation and water-related infrastructures and water resources development were also conducted. *Bulinus truncatus* are intermediate host snails of *S. haematobium* in Iran. *Bulinus* snails live in several habitats like canals, swamps and borrow-pits (Chu *et al.* 1968; Massoud *et al.* 1982). The highest snail populations were observed among May–July and October–December months. All habitats of *B. truncatus* in Khuzestan were investigated two times a year and were treated with molluscicide agents such as niclosamide. At the

same time, improvement of environmental hygiene such as safe water supply, draining of the canals and building of standard latrines was considered. The percentage of people who use from optimized water resources is 93.1 and 99.6% in rural and urban regions, respectively in Khuzestan province, south-west of Iran. Also 93.8 and 100% of the rural and urban inhabitants, possess optimized sewage disposal system, respectively. Literacy rate in Khuzestan province was estimated 90.9 (92.1 and 89.8% for males and females, respectively) among people of 15–24 years of age (Rashidian *et al.* 2012).

National guideline for urogenital schistosomiasis control was revised in 2006 and schistosomiasis control programme was changed from its previous vertical form to an integrated form in the present health service delivery system. The health service delivery system in the Islamic Republic of Iran is characterized by its tiered design with parallel structure in urban and rural areas. The first tier of delivery encompasses primary health care (PHC) and comprises two levels of care. The first level includes rural health houses with a defined served population of 1400 individuals. Health houses are staffed by Behvarzes (allied health workers). Behvarzes are selected then trained in Behvarz training centres for 2 years, which consists of training and service in health houses. This level also includes rural health centres, which are staffed by a physician and a health worker. These centres supervise a number of health houses with a defined service population of 6000–10 000 individuals. The second level of the PHC tier is the district health centre, which undertakes planning and supervising functions. In urban areas there is a similar structure that includes health posts serving a population of 12 000 individuals, then urban health centres (with a defined service population of 40 000–60 000 individuals) and district health centres (Fig. 3) (Shadpour, 2000; Mehrdad, 2009; Rahbar, 2009). In Khuzestan there are 26 district health centres, 79 urban health centres, 121 rural health centres, 62 urban-rural health centres, 209 health posts and 859 health houses. By using the capacity of such a health network, PHC is distributed all over the country from the most peripheral villages (health house) to the big cities. Surveillance programmes for many diseases are done through this network and of course it helps the schistosomiasis control programmes in elimination phase for more accurate case finding and early warning if a suspected case is reported. New surveillance system for urogenital schistosomiasis in elimination phase that was recently published by the CDC of Ministry of Health is based on active and passive case findings in previously infected areas, monitoring snails population, distribution and activity using Geographic Information System and Global Positioning System and implementation of appropriate control measures, education including community education and training courses for




Fig. 3. Diagram of health system network in Iran.

health care workers, performing practical researches by use of classic, serologic and molecular methods, and also multicentre researches and increasing national, regional and international cooperation.

#### Challenges in elimination phase

The most important challenges for maintaining elimination status includes development of irrigation and agricultural projects such as sugarcane in Khuzestan province, which may lead to increasing snail population (Mowlavi *et al.* 2009). As urogenital schistosomiasis is eliminated and new medical students are not familiar with the disease, the knowledge of physicians and health workers about this infection is decreasing. Presence of active foci of urogenital schistosomiasis in southern regions of Iraq, increased trans-border traffic and people travelling to visit shrine sites between the two countries are other challenges in elimination phase.

In conclusion, by using a national programme for urogenital schistosomiasis control based on active and passive case findings, patient treatment, population education, conducting training courses for health workers, snail control programmes including sanitary measures, environmental improvement, focal mollusciciding of snail habitats and improvement of water supply for people living in endemic areas, the infection has been eliminated from Iran and this can be a successful model for countries suffering from this disease. It should be mentioned that maintaining long-term monitoring of the situation in light of immigration of infected persons from other areas would be useful. In regard to results of previous surveys in different regions, both microscopic and molecular technique should be applied in future years (Meurs *et al.* 2015; Utzinger *et al.* 2015). In future, to prove eradication, studies based on serology methods in people in order to more accurately detect the disease should be

designed. Also, molecular evaluation like polymerase chain reaction-based techniques in *B. truncatus* snails to determine the presence of intermediate stages of parasite is required.

#### AUTHORS' CONTRIBUTION

S. S. and S. K. conceived the study; S. S., M. F. R. and S. K. designed the study protocol; M. G. collected the data; M. F. R., S. K. and M. G. analysed and interpreted the data; S. S., M. F. R. and S. K. wrote the manuscript; S. S., M. F. R. and S. K. critically revised the manuscript. All authors read and approved the final manuscript.

#### ACKNOWLEDGEMENTS

The authors would like to thank all staff at the health care centres in Khuzestan Province. Also, the authors sincerely acknowledge their gratitude to the efforts of Centres for Disease Control in Ahvaz Jundishapur University of Medical Sciences.

#### FINANCIAL SUPPORT

This study was financially supported by Student Research Committee of Ahvaz Jundishapur University of Medical Science, Iran (NO: 93S87) to M. F. R. and approved by Ethical committee NO: IR.AJUMS.REC.1395.151.

#### CONFLICT OF INTERESTS

The authors declare no conflict of interests.

#### REFERENCES

- Al Ghahtani, A. G. and Amin, M. A. (2005). Progress achieved in the elimination of schistosomiasis from the Jazan region of Saudi Arabia. *Annals of Tropical Medicine and Parasitology* **99**, 483–490.
- Amarir, F., El Mansouri, B., Fella, H., Sebt, F., Mohammed, L., Handali, S., Wilkins, P., El Idrissi, A.L., Sadak, A. and Rhajaoui, M. (2011). National serologic survey of Haematobium schistosomiasis in Morocco: evidence for elimination. *The American Journal of Tropical Medicine and Hygiene* **84**, 15–19.

- Arfaa, F., Bijan, H. and Farahmandian, I. (1967). Present status of urinary bilharziasis in Iran. *Transactions of the Royal Society of Tropical Medicine and Hygiene* **61**, 358–367.
- Chidozie, E. U. and Daniyan, S. Y. (2008). Urinary schistosomiasis epidemiological survey of urinary schistosomiasis among children in selected schools: a preliminary study in Minna, Nigeria. *African Journal of Biotechnology* **7**, 2773–2776.
- Chomba, C. and Mutale, S. (2014). Factors characterising high prevalence rates of urinary schistosomiasis in Mufumbwe district, north western province of Zambia. *Global Institute for Research & Education* **3**, 86–95.
- Chu, K. Y., Massoud, J. and Arfaa, F. (1968). Distribution and ecology of *Bulinus truncatus* in Khuzestan, Iran. *Bulletin of the World Health Organization* **39**, 607–637.
- Cunin, P., Tchuem Tchuente, L. A., Poste, B., Djibrilla, K. and Martin, P. M. (2003). Interactions between *Schistosoma haematobium* and *Schistosoma mansoni* in humans in north Cameroon. *Tropical Medicine & International Health* **8**, 1110–1117.
- Dhunpath, J. (1994). Progress in the control of schistosomiasis in Mauritius. *Transactions of the Royal Society of Tropical Medicine and Hygiene* **88**, 507–509.
- EMRO/WHO (2007). *Inter-Country Meeting on Strategies to Eliminate Schistosomiasis from the Eastern Mediterranean Region*. Muscat, Oman.
- Fenwick, A., Webster, J. P., Bosque-Oliva, E., Blair, L., Fleming, F. M., Zhang, Y., Garba, A., Stothard, J. R., Gabrielli, A. F., Clements, A. C., Kabatereine, N. B., Toure, S., Dembele, R., Nyandindi, U., Mwansa, J. and Koukounari, A. (2009). The schistosomiasis control initiative (SCI): rationale, development and implementation from 2002–2008. *Parasitology* **136**, 1719–1730.
- Gholamreza, M., Jafar, M., Allahbedasht, M., Nikkhooy, A., Mohammad, K. A. and Rokni, M. B. (2008). Survey of urinary schistosomiasis in Iran. *Pakistan Journal of Medical Sciences*, **24**, 857–860.
- Hamidinia, D., Maraghi, S., Azimi, F., Ai, A. and Shirian, S. (2014). The role of climate on prevalence or eradication of vesical schistosomiasis in Khuzestan Province of Iran. *Journal of Parasitic Diseases* 1–3. doi: 10.1007/s12639-014-0515-6.
- Ismail, H. A., Hong, S. T., Babiker, A. T., Hassan, R. M., Sulaiman, M. A., Jeong, H. G., Kong, W. H., Lee, S. H., Cho, H. I., Nam, H. S., Oh, C. H. and Lee, Y. H. (2014). Prevalence, risk factors, and clinical manifestations of schistosomiasis among school children in the White Nile River basin, Sudan. *Parasites & Vectors*, **7**, 478.
- Kasai, T., Nakatani, H., Takeuchi, T. and Crump, A. (2007). Research and control of parasitic diseases in Japan: current position and future perspectives. *Trends in Parasitology* **23**, 230–235.
- Ketabchi, A. and Moshtaghi-Kashanian, G. (2012). Urinary schistosomiasis with simultaneous bladder squamous cell carcinoma and transitional cell carcinoma. *Iranian Journal of Parasitology* **7**, 96–98.
- Knopp, S., Person, B., Ame, S. M., Mohammed, K. A., Ali, S. M., Khamis, I. S., Rabone, M., Allan, F., Gouvras, A., Blair, L., Fenwick, A., Utzinger, J. and Rollinson, D. (2013). Elimination of schistosomiasis transmission in Zanzibar: baseline findings before the onset of a randomized intervention trial. *PLoS Neglected Tropical Diseases* **7**, e2474.
- Maraghi, S., Ranjbar, N. and Haddad, M. H. (2004). Schisto-granuloma of the fallopian tube from a low-endemic area in Iran. *Acta Tropica* **92**, 89–90.
- Massoud, J., Arfaa, F., Farahmandian, I., Ardalan, A. and Mansoorian, A. (1982). Progress in the national schistosomiasis control programme of Iran. *Bulletin of the World Health Organization* **60**, 577–582.
- McCullough, F. S., Gayral, P., Duncan, J. and Christie, J. D. (1980). Molluscicides in schistosomiasis control. *Bulletin of the World Health Organization* **58**, 681–689.
- Mehrdad, R. (2009). Health system in Iran. *Japan Medical Association Journal* **52**, 69–73.
- Meurs, L., Mbow, M., Vereecken, K., Menten, J., Mboup, S. and Polman, K. (2012). Epidemiology of mixed *Schistosoma mansoni* and *Schistosoma haematobium* infections in northern Senegal. *International Journal for Parasitology* **42**, 305–311.
- Meurs, L., Brienen, E., Mbow, M., Ochola, E. A., Mboup, S., Karanja, D. M., Secor, W. E., Polman, K. and van Lieshout, L. (2015). Is PCR the next reference standard for the diagnosis of schistosoma in stool? A comparison with microscopy in Senegal and Kenya. *PLoS Neglected Tropical Diseases* **9**, e0003959.
- Mo, A. X., Agosti, J. M., Walson, J. L., Hall, B. F. and Gordon, L. (2014). Schistosomiasis elimination strategies and potential role of a vaccine in achieving global health goals. *American Journal of Tropical Medicine and Hygiene* **90**, 54–60.
- Mombeni, H. and Kheradmand, A. (2005). *Schistosoma haematobium* control in Khoozestan province in Iran: prosperities and failures. *Medical Journal of the Islamic Republic of Iran* **19**, 19–22.
- Mowlavi, G., Mansoorian, A., Mahmoodi, M., Pourshojaei, R. and Salehi, M. (2009). Identification of freshwater snails in cane-sugar fields in the northern part of Khuzestan Province from a public health perspective. *Journal of School of Public Health & Institute of Public Health Research* **7**, 69–76.
- Niaz, S., Tanveer, A. and Qureshi, A. (2010). Prevalence of humans schistosomiasis in different areas of Punjab Pakistan. *Pakistan Journal of Science* **62**, 160–162.
- Othman, A. A. and Soliman, R. H. (2015). Schistosomiasis in Egypt: a never-ending story?. *Acta Tropica* **148**, 179–190.
- Rahbar, M. (2009). *Primary health care in Islamic Republic of Iran. Series of health report in Islamic Republic of Iran*, 2nd Edn. Ministry of Health and Medical Education, Deputy for Health, Tehran, Iran.
- Rashidian, A., Khosravi, A., Khabiri, R., Khodayari-Moez, E., Elahi, E., Arab, M. and Radaie, Z. (2012). *Islamic Republic of Iran's multiple indicator demographic and health survey (IrMIDHS) 2010*, Ministry of Health and Medical Education. Tehran.
- Rollinson, D., Knopp, S., Levitz, S., Stothard, J. R., Tchuente, L. A. T., Garba, A., Mohammed, K. A., Schur, N., Person, B., Colley, D. G. and Utzinger, J. (2013). Time to set the agenda for schistosomiasis elimination. *Acta Tropica* **128**, 423–440.
- Rudge, J. W., Stothard, J. R., Basanez, M. G., Mgeni, A. F., Khamis, I. S., Khamis, A. N. and Rollinson, D. (2008). Micro-epidemiology of urinary schistosomiasis in Zanzibar: local risk factors associated with distribution of infections among schoolchildren and relevance for control. *Acta Tropica* **105**, 45–54.
- Sady, H., Al-Mekhlafi, H. M., Mahdy, M. A., Lim, Y. A., Mahmud, R. and Surin, J. (2013). Prevalence and associated factors of Schistosomiasis among children in Yemen: implications for an effective control programme. *PLoS Neglected Tropical Diseases* **7**, e2377.
- Salmanzadeh, S., Foroutan-Rad, M., Khademvatan, S., Moogahi, S. and Bigdeli, S. (2015). Significant decline of malaria incidence in south-west of Iran (2001–2014). *Journal of Tropical Medicine* **2015**, 523767.
- Shadpour, K. (2000). Primary health care networks in the Islamic Republic of Iran. *Eastern Mediterranean Health Journal* **6**, 822–825.
- Singh, K. and Muddasiru, D. (2014). Epidemiology of schistosomiasis in school aged children in some riverine areas of Sokoto, Nigeria. *Journal of Public Health and Epidemiology* **8**, 197–201.
- Utzinger, J., Becker, S. L., van Lieshout, L., Van Dam, G. J. and Knopp, S. (2015). New diagnostic tools in schistosomiasis. *Clinical Microbiology and Infection* **21**, 529–542.
- WHO (2013). Schistosomiasis: progress report 2001–2011, strategic plan 2012–2020.
- Woolhouse, M. E., Etard, J. F., Dietz, K., Ndhlovu, P. D. and Chandiwana, S. K. (1998). Heterogeneities in schistosome transmission dynamics and control. *Parasitology* **117** (Pt 5), 475–482.