

Effort valuation and psychopathology in children and adults

Nicholas H. Nguyen¹, Avery B. Albert^{1,2}, Sarah Van Orman¹, Patricia Forken¹, Steven D. Blatt¹, Wanda P. Fremont¹, Stephen V. Faraone^{1,*} and Stephen J. Glatt^{1,*}

Original Article

*These authors contributed equally to this work.

Cite this article: Nguyen NH, Albert AB, Van Orman S, Forken P, Blatt SD, Fremont WP, Faraone SV, Glatt SJ (2019). Effort valuation and psychopathology in children and adults. *Psychological Medicine* **49**, 2801–2807. <https://doi.org/10.1017/S0033291718003884>

Received: 1 May 2018

Revised: 8 August 2018

Accepted: 27 November 2018

First published online: 14 January 2019

Key words:

EEfRT; effort valuation; psychopathology; RDoC; reward

Author for correspondence:

Stephen J. Glatt, E-mail: stephen.glatt@psychgenelab.com

¹Department of Psychiatry and Behavioral Sciences, State University of New York – Upstate Medical University, Syracuse, New York, USA and ²Department of Psychology, Syracuse University, Syracuse, New York, USA

Abstract

Background. The Research Domain Criteria initiative was launched by the US National Institute of Mental Health to establish a multi-level framework for understanding psychological constructs relevant to human psychiatric disorders, and identified ‘effort valuation/willingness to work’ as a clinically useful construct worthy of further study. This construct encompasses the processes by which the cost(s) of obtaining an outcome are calculated, and the tendency to overcome response costs to obtain a reinforcer. The current study aims to examine effort valuation as a correlate of psychopathology in children and adults, and the moderating effects of sex on this relationship.

Methods. Participants were 1215 children aged 6–12 and their parents ($n = 1044$). All participants completed the Effort Expenditure for Rewards Task as a measure of effort expenditure. Child psychopathology was measured *via* the Child Behavior Checklist, while adult psychopathology was measured *via* the Adult Self Report. Additionally, the Social Adjustment Inventory for Children and Adolescents and Injury Behavior Checklist were used to examine child social impairments/problem behaviors.

Results. In children, significant interactions between reward sensitivity and sex were observed in association with anxiety and thought problems, specifically at low reward sensitivity levels. In adults, main effects of effort expenditure were seen in drug and alcohol abuse, where higher effort was associated with higher degrees of abuse.

Conclusions. These results establish effort valuation as a relevant psychological construct for understanding psychopathology, but with different profiles of associated psychopathology across sex in children and adults.

Introduction

The Research Domain Criteria (RDoC) initiative was launched by the US National Institute of Mental Health (NIMH) as a framework for approaching both research on and diagnosis of mental disorders. Much research challenges the categorical representation of disorders of the Diagnostic and Statistical Manual (DSM) of the American Psychiatric Association and the International Classification of Diseases (ICD). The goal of the RDoC program is to facilitate the research-based validation of constructs, defined by neurobiological and behavioral measures to support revisions of current diagnostic systems (Cuthbert, 2014). This initiative presently outlines five major domains of human functioning and behavior intended to reflect the contemporary understanding of major neurobiological systems. A key premise of RDoC is that these ‘cross-disorder’ constructs are relevant to multiple diagnoses. This premise has been supported by research indicating common risk factors for multiple disorders (Lee *et al.*, 2013).

The RDoC Positive Valence Systems domain primarily concerns responses to positive motivational situations, consisting of constructs such as Approach Motivation (AM), Initial Responsiveness to Reward Attainment, Sustained Responsiveness to Reward Attainment, Reward Learning, and Habit. AM is a multi-faceted construct that includes component processes such as reward valuation, effort valuation, reward prediction error, and preference-based decision-making. The current study focuses on the effort valuation component of the AM construct, which is defined as the processes by which the cost(s) of attaining an outcome is computed, and the tendency to overcome those response costs to obtain a positive reinforcer (NIMH, 2016).

Previous research regarding the role of effort valuation in psychopathology has consistently found differences in effort-based decision-making between patients with mental disorders and healthy comparison subjects. A study examining reward motivation in major depressive disorder (MDD) found that MDD patients are less willing to expend effort for rewards, and are less effective in their use of information about the magnitude and probability of rewards

to guide choice behavior (Treadway *et al.*, 2012). For autism spectrum disorders (ASDs), research has found a relationship between effort-based decision-making and repetitive behavior symptoms in individuals with ASD. Affected individuals were more willing to expend effort to obtain a reward regardless of differences in reward value and probability (Damiano *et al.*, 2012). Several studies have examined effort valuation in schizophrenia, noting that impaired effort allocation is a consistent finding in individuals with the disorder relative to healthy comparison participants (Treadway *et al.*, 2015; McCarthy *et al.*, 2016). These research findings suggest that measures of effort valuation could be useful predictors of psychiatric disorder; yet, it remains unclear if effort valuation relates to specific diagnoses, to dimensions of psychopathology, or if any associations with psychopathology are consistent across, age, sex, and race.

The impact of effort evaluation on psychopathology has been observed in schizophrenia, where an overall reduction in effort allocation was associated with worse community and work function (Barch *et al.*, 2014). Evidence of a reward-processing deficit in adults with ASD has been demonstrated in research regarding decision-making. Researchers observed a positive association between inefficient effort-based decision-making and decreased sensitivity to reward parameters. These findings provide a possible explanation for the characteristic reduction in motivation to seek social interaction and stimuli (Damiano *et al.*, 2012). Therefore, understanding the relationship between psychopathology and aberrations in effort valuation may provide valuable insight into behavioral impairments spanning several diagnostic categories.

Sex differences have been observed across multiple forms of psychopathology, especially in disorders that have previously also been linked to aberrant reward behavior. These differences include rates of occurrence, type and degree of symptom severity, and treatment response. Examples of differing prevalence would be the 3:1 male to female ratio observed in ADHD or the 2:1 female to male ratio for anxiety/depression (Altemus *et al.*, 2014; Ji *et al.*, 2018). With respect to symptomatology, previous research in ASD has found that externalizing problems, such as aggressive behavior and hyperactivity, were more prominent in males, whereas internalizing problems, such as emotional issues in anxiety and depression, were greater in females (Werling and Geschwind, 2013). Our current understanding of the mechanisms behind these differences is limited, and it is still unclear how and to what degree they are influenced by genetics, physiological factors, and social conditions. In light of that, we believe that it is critical to include investigation of sex differences in relating reward behavior to psychopathology. To our knowledge, no research has examined the effects of sex on the relationship between effort valuation and psychopathology. Building upon that, by specifically focusing on how sex impacts effort valuation and reward processing, we hope to gain further understanding of the underlying mechanisms of abnormal behavior associated with psychopathology.

As an RDoC-based study, our intent was to take a broad approach to relating reward behavior to psychopathology. To this end, we included the Social Adjustment Inventory for Children and Adolescents (SAICA) and Injury Behavior Checklist (IBC) to also examine the relationship between social behavior, aggression, and injurious behavior in order to gain a more complete understanding of human reward behavior. Aberrant social behavior and aggression have been linked with atypical reward behavior in previous research. In depression, a reduced neural response to social reward has been found among offspring of depressed parents as compared with those whose

parents had no history of depression (Olinio *et al.*, 2015). Similar dysregulated reward mechanisms have been observed between substance use disorders and impulsive-aggressive behavior, and to date, reward behavior has been understudied in aggression research (Venables, 2017). These findings suggest that reward behavior could be a key component of many areas of human brain function and investigating atypical reward responses may be important to multiple research domains, including psychopathology.

Against this backdrop, the current study aims to examine effort valuation as a correlate of psychopathology in children and adults, and the moderating effects of sex on the relationship between effort valuation and psychopathology. We hypothesized that effort valuation would be a significant correlate of multiple dimensions of psychopathology, and aim to investigate how sex modulated that relationship. If aberrant reward behavior was indeed linked to psychopathology, we anticipated that we might also observe sex-related differences, based on findings from previous research regarding gender differences in mental disorders. By understanding these connections, we hoped to further establish the validity of effort valuation as a relevant construct in psychiatric research. To that end, we also aim to investigate the cross-generational and familial continuity of the construct by comparing effort valuation between parents and their children, as well as between siblings. We hypothesized that effort valuation would be predictive of similar domains of psychopathology between adults and children and that effort evaluation would be significantly correlated among siblings and between parents and children.

Methods

Procedure

Participants were recruited from a variety of sources within the greater Syracuse area, including the Child and Adolescent Psychiatry Clinic at SUNY Upstate Medical University and child psychiatrists and mental health clinicians working in private practice in the community, as well as from community events (local fairs, festivals, etc.). Children with the following characteristics and conditions were excluded from the study: adopted, sensorimotor disabilities, a diagnosed neurological condition, a history of head injury with documented loss of consciousness lasting more than 10 min, an uncontrolled medical condition, use of psychotropic medications, or an inability to understand the English language. Exclusionary criteria were the same for parents, with the exception of the adoption criterion and the addition of two other criteria: parents who did not have the ability to independently complete study tasks and women who were pregnant or gave birth within 6 months prior to the study visit were excluded from participation. An estimate of intelligence quotient (IQ) was obtained from scores on the vocabulary and abstraction subtests of the Shipley-2 (a validated, age-appropriate instrument for subjects between the ages of 7 and 89 years). This was applied to both adults and children within the recommended age range. As the mean of these two tests correlates 0.90 with full scale IQ, subjects with an estimated IQ below 80 were excluded. Informed consent was obtained from all parents and assent was given by all children upon arrival for their study visit.

Participants

A total of 1215 children and 1044 parents, with and without a history of psychiatric problems, participated in this study. Our study

population is drawn from a convenience sample that was purposely enriched for psychopathology via recruitment in local clinics. All children were between the ages of 6 and 12 years (mean age = 9 years, *s.d.* = 2.2), and their parents were between the ages of 23 and 59 years (mean age = 37 years, *s.d.* = 7.2). Parental age was capped at 59 years of age to avoid the possibility of cognitive decline. While there were approximately equal numbers of female and male children (49% *v.* 51%), significantly more of the participating parents were female than male (69% *v.* 31%). Participants were diverse in their ancestral backgrounds, with 65% of parents identifying as White, 24% Black, and 11% other or multiple races, and 55% of children identifying as White, 25% Black, and 20% other or multiple races. Additionally, 7% of parents and 11% of children were Hispanic. Parents reported both for themselves and for their children whether they had ever sought mental health care for emotional or behavioral problems, with 41% of children and 45% of parents reporting such psychiatric history. The dataset includes a total of 770 different families, with an average family size of 2.93.

Measures

Study visits were approximately 3 h in length, and involved the completion of a variety of computerized inventories and behavioral paradigms.

Effort Expenditure for Rewards Task

The Effort Expenditure for Rewards Task (EEfRT) was developed by Treadway and colleagues to measure effort-based decision-making (Treadway *et al.*, 2009). The task is a multi-trial game in which participants are asked to choose between a 'hard task' and an 'easy task' in order to obtain monetary rewards. Each trial consists of a required number of repeated manual presses of a keyboard button within a time limit in order to be eligible to win a small monetary reward. Easy task trials required 30 button presses with the dominant index finger within 7 s with an assigned reward value of \$1.00, while hard-task trials required 100 button presses with the non-dominant little finger within 21 s with a varying assigned reward value between \$1.24 and \$4.30. Subjects were not guaranteed to win the reward upon task completion; some trials were 'no-win' trials in which the subject receives no reward, and others were 'win' trials in which the subject receives the assigned reward amount. At the beginning of each trial in the adult version of the EEfRT task only, subjects are provided with one of three levels of probability for actually obtaining a reward upon trial completion: 'high' (88%) probability of it being a win trial, 'medium' (50%), and 'low' (12%). Thus, not every trial may result in a reward, even if the participant successfully completed the task. Probability levels apply to both the hard and easy task, with equal proportions of each probability level across the experiment. Probability of choosing the hard task was calculated for each subject overall, across all trials. This probability component is not present in the child version of the EEfRT task. All subjects received trials presented in the same randomized order. Our analysis variables for the EEfRT task are effort expenditure and reward sensitivity. Effort expenditure for each individual was calculated as the percentage of trials for which that participant chose the hard task. Individuals who chose the hard task more than 50% of the time were considered to have high effort expenditure, and those who chose the hard task less than 50% of the time were considered to have low effort expenditure. To derive reward sensitivity, a logistic betaweight was calculated via

a generalized linear model with hard task choice as the outcome (dependent) variable and reward magnitude (monetary amount; dollars) as the predictor (independent) variable.

Adult Self Report

The Adult Self Report (ASR) was used to measure psychopathology in adult participants. This 126-item self-report measure is well validated and widely used in clinical practice to assess symptoms of psychopathology and adaptive functioning in individuals aged 18–59 years (Rescorla and Achenbach, 2004). Participants were asked to respond to each item on a three-point scale from *not true* to *very true/often true*. The ASR provides T-scores for seven syndrome scales (anxious/depressed, withdrawn, somatic complaints, thought problems, attention problems, aggressive behavior, rule-breaking behavior), six scales specific to symptoms of DSM diagnoses (affective disorders, anxiety disorders, somatic problems, avoidant personality features, attention/deficit hyperactivity problems, antisocial personality features), three composite scores (internalizing composite, externalizing composite, total problems composite), and four scales assessing substance use (tobacco, alcohol, recreational drugs, substance use composite). An adaptability composite is also provided along with subscale scores for friendships, spouse, family, and employment. Due to the nature of the ASR assessment, the minimum possible T-score generated is 50, which would represent a participant whose level of self-reported psychopathology is that of a typically developing, psychiatrically unaffected individual (no psychopathology). A T-score of 50 then represented a 'zero value' in our study. These outcome variables are simply T-scores; they are not count data.

Child Behavior Checklist

The Child Behavior Checklist (CBCL) was used to measure psychopathology in children. This 113-item parent-report measure is well validated and widely used in clinical practice to assess emotional and behavioral functioning in children ages 6–18 years (Achenbach, 1991). Parents were asked to respond to each item on a three-point scale from *not true* to *very true/often true*, indicating how true the item is for their child. The CBCL provides T-scores for eight syndrome scales (anxious/depressed, withdrawn/depressed, somatic complaints, social problems, thought problems, attention problems, rule-breaking behavior, aggressive behavior), nine DSM-oriented scales (affective problems, anxiety problems, somatic problems, attention deficit/hyperactivity problems, oppositional defiant problems, conduct problems, sluggish cognitive tempo, obsessive compulsive problems, post-traumatic stress problems), and three composite scores (internalizing composite, externalizing composite, total problems composite). The CBCL shares the same minimum T-score as the ASR (50), so our outcome variables follow the same logic described in the section above.

Social Adjustment Inventory for Children and Adolescents

The SAICA was used to measure children's functioning across several domains. This 78-item semi-structured interview was designed to assess social functioning in children ages 6–18 years (John *et al.*, 1987). Direct responses to the interview questions from the children were recorded on a four-point scale, with higher scores indicating greater impairment. Mean scores were created for four domains of functioning: current school functioning, spare time activities, peer functioning, and current home behavior. An overall functioning score was calculated by taking the mean of these subscale scores.

Injury Behavior Checklist

The IBC was used to measure dangerous and problem behavior in children. This 24-item parent-report measure asks respondents to rate the frequency of injurious behaviors displayed by their child (Potts et al., 1997). For each item, parents were asked to rate their children on a five-point scale from *not at all* to *more than once a week*. Items were sum-scored, with possible scores ranging from 24 to 120, and with higher scores indicating more frequent injurious and problem behaviors.

Statistical analyses

Statistical analyses used R version 3.4.1. A series of negative binomial regression models examined effort valuation as a predictor of psychopathology in children. Negative binomial regression was used to account for non-normal distribution of the outcome variables, which were T-scores. These variables also had an over-representation of zeros because both the ASR and CBCL T-scores less than the mean (zero) are fixed at zero by the developers of the ASR and CBCL scoring algorithms. These extra zeros were modeled as zero inflation within the negative binomial framework. Main effects of effort expenditure, reward sensitivity, and sex were tested, as well as interaction effects of effort expenditure, reward sensitivity, and sex. All child models covaried for parent income and education, but not age, as there was no association between child's age and EEfRT scores.

Similar to the children, negative binomial regression was used to account for non-normal distribution of outcome variables in parents. Main effects of effort expenditure, reward sensitivity, and sex were tested, as well as interaction effects of effort expenditure, reward sensitivity, and sex. All adult models covaried for demographic variables including age, IQ, education, employment, income, race, and marital status.

Robust standard errors were used across all models to account for non-independence in the data such as those caused by familial relationships. Paired-sample correlation tests using intraclass correlations as well as Kendall's τ were also conducted to determine the degree of sibling-sibling and parent-child correlations.

Multiple imputation was utilized to handle missing adult income data, as 25% of participants chose not to report their household income. Ten imputations were conducted. Demographic variables including education, employment, marital status, age, IQ, ancestry, and sex were used to predict income in the imputation procedure. When modeled together, these variables explained a large proportion of the variance in income ($R^2 = 0.698$). Multiple imputation was not used for any other variable, as missing data was not an issue for any other variables included in these analyses.

The Benjamini Hochberg False Discovery Rate method (Benjamini and Hochberg, 1995) corrected for multiple testing. This procedure is a recommended alternative to Bonferroni-type corrections, which are often criticized for increasing the likelihood of type-II errors, particularly when a large number of tests are conducted (Perneger, 1998). A 5% false discovery rate was utilized for determining the significance of findings across both groups. The total number of tests conducted was 18 for the adults and 21 for the children. p values reported in the results are false discovery rate-adjusted.

Results

See online Supplementary Table S1 for more demographic information about the participants of this study. Online




Fig. 1. Reward sensitivity and sex in CBCL DSM anxiety problems.

Supplementary Tables S2 and S3 include bivariate correlations between measures of effort expenditure, reward sensitivity, and each of the outcome variables tested.

EEfRT task statistics

For the adult participant population, average proportion of hard task choices was 36.76%, and the average reward sensitivity betaweighted was 0.765. Mean percent completion rate among adults was 93.47%. On average, adult participants timed out in their choice of the hard *v.* easy task in 5.90% of trials. For the child participant population, average proportion of hard task choices was 58.34%, and the average reward sensitivity betaweighted was 0.118. Mean percent completion rate among children was 87.27%. On average, child participants timed out in their choice of the hard *v.* easy task in 1.38% of trials. Additionally, we calculated the number of participants in each population who chose only easy or only hard tasks (possibly indicating lack of task comprehension), to confirm that these subjects' choices did not drive the reported associations with psychopathology. For adults, 0.7% of participants chose only easy tasks, and 0.0% of participants chose only hard tasks. For children, 0.6% of participants chose only easy tasks, and 3.5% of participants chose only hard tasks.

Models predicting child psychopathology

Additional interactions between reward sensitivity and sex were observed in the models predicting DSM-5 anxiety problems. At low reward sensitivity, increased anxiety problems were observed for male subjects relative to female subjects ($\beta = 0.522$, $p = 0.041$) (Fig. 1). Similarly, we found increased thought problems at low reward sensitivity for male subjects in comparison to female subjects ($\beta = 0.438$, $p = 0.045$) (Fig. 2).

Models predicting adult psychopathology

A main effect of effort expenditure was observed in the models predicting ASR alcohol use. At high hard task choice percentage, increased alcohol usage was observed for our subjects ($\beta = 0.531$, $p = 0.0083$) as compared with those with low hard task percentage (Fig. 3). Additionally, a main effect of effort expenditure was observed in the models predicting ASR drug use. At high hard task choice percentage, increased drug usage was observed for our subjects ($\beta = 3.398$, $p = 0.014$) as compared with those with low hard task choice percentage (Fig. 4).




Fig. 2. Reward sensitivity and sex in CBCL thought problems.




Fig. 3. Effort in ASR alcohol use.

Familial transmission

In examining the transmission of effort valuation within families, we found significant sibling–sibling correlations but no significant parent–child correlations. When examining associations of effort valuation between siblings using EEfRT hard task choice percentage, we found a significant positive correlation between both full siblings ($r = 0.13$, $p = 0.039$) and half siblings ($r = 0.19$, $p = 0.008$). However, in examining associations of effort valuation between these children and their parents using EEfRT hard task choice percentage, we found no significant correlations between children and their respective mothers and fathers ($p > 0.05$).

Discussion

The current study sought to test hypotheses regarding the relationship between effort valuation and psychopathology in children and adults, and to determine the impact of sex on these relationships. As we hypothesized, effort valuation was a significant predictor of multiple dimensions of psychopathology, some of which were moderated by sex. However, our hypothesis that effort valuation would be linked to the same psychopathologies between adults and children was not supported. Additionally, while we found significant relationships between effort valuation and several areas of psychopathology and functioning that varied by sex, many of our tests of association were negative and correlations between effort valuation and psychopathological outcomes




Fig. 4. Effort in ASR drug use.

were generally small. Given the large size of our sample, these negative findings cannot be attributed to low statistical power. These results suggest that dysfunction in effort valuation may be a contributing factor rather than a driving force for a range of psychopathology and impairment in children and adults.

Contrary to our hypothesis, effort valuation was associated with different psychopathologies for adults and children. Additionally, while a positive sibling–sibling correlation was found for effort valuation, no significant correlation was found between parents and their children. Given that many psychiatric conditions are transmitted in families, the lack of parent–child correlations for effort valuation presents important questions to consider for future research, especially since there have been no prior familial studies of effort valuation in psychopathology to date. This result may mean that the child and adult assessments for effort valuation do not gauge the same constructs. For example, the parent version of the EEfRT included an additional reward evaluation component (probability of trial actually resulting in reward) in each trial which was not present in the child version. That difference may have impacted adult subjects' decision-making process. Alternatively, the lack of parent–child correlation could indicate that effort valuation, and its correlates, change as a child moves through the stages of development. At the time of assessment, children were between the ages of 6 and 12 years. Changes in effort valuation have been previously observed in a comparison study between 4- and 6-year old children where the older children displayed greater ability to evaluate effort and reward quality (Benozio and Diesendruck, 2015).

Previous studies in children have found sex and reward sensitivity to be linked to anxiety disorder, such that clinically anxious male youth displayed both decreased risk taking and sensitivity to reward in comparison to clinically anxious females, as well as typically developing male and female youths (Dorfman *et al.*, 2016). Our results provide further support for the relationship between sex, reward sensitivity, and anxiety disorder, where at low reward sensitivity we found that male children reported greater anxiety problems than female children. It has become increasingly apparent that the study of gender differences in adolescent anxiety is important, given that anxiety disorders often begin to present symptomatically in adolescence. Additionally, a longitudinal study of adolescents demonstrated a greater level of functional impairments among anxious males than anxious females with regards to academic performance, self-esteem, sense of well-being, and socialization with friends (Derdikman-Eiron *et al.*, 2012). As a result, the gender-specific alterations in the reward system

measures observed both here and in previous research suggest that treatments for anxiety that target the reward system may need to be tailored to gender.

While past studies have also found relationships between ASDs and both aberrant effort valuation and diminished reward processing in children (Scott-Van Zeeland *et al.*, 2010; Mosner *et al.*, 2017), we found that this relationship may differ based on sex. Our results revealed that lesser reward sensitivity was associated with a higher degree of thought problems in male children *v.* females. Thought problems according to the CBCL scale include seeing or hearing things, repeating acts, and strange ideas/behavior. In children, thought problems have been linked with relatively high sensitivity and specificity to ASD (Mazefsky *et al.*, 2011). To our knowledge, no previous studies have specifically examined the moderating effects of sex in relating effort expenditure or reward sensitivity with ASD in children. However, with ASD being a predominantly male disorder (Loomes *et al.*, 2017), understanding gender-specific alterations in reward behavior may be important for treatment development.

Limitations/future directions

The current work has several limitations. The use of self-reported race, rather than genetic data, to understand ancestry may have placed greater weight on the social over the biological component of an individual's ancestral background. Participants' self-reported racial identity may not fully align with their genetic ancestry, instead representing their personal understanding of their ancestral background. For example, Hispanic participants may only identify with their Hispanic ethnicity rather than any racial category. Future research should compare the implications of self-identified ancestry *v.* genetic ancestry to determine if these differing ways of defining one's heritage diverge in relation to effort valuation and associated psychopathological outcomes.

In the current study, we were unable to examine the impact of racial identities other than White/Caucasian and Black/African-American due to the low number of individuals who identified with other races. Also, our adult sample was disproportionately female, so future studies should seek to recruit a more balanced sample in addition to correcting the under-representation of other racial minorities besides Black/African-American. Few participants displayed elevated symptoms for some of the psychological problems examined in this study, so future research collecting data from more individuals who experience particular psychological problems may reveal new or clearer patterns between effort valuation and specific forms of psychopathology.

With regards to our assessments of psychopathology, we acknowledge that our usage of parent-reports as measures for both child and parent psychopathology is an important limitation as it represents an indirect rather than direct measure of child psychopathology. While the CBCL and ASRs cover a wide range of psychopathological outcomes, further research on effort valuation should be conducted in the context of additional psychopathologies not specifically included in the CBCL or ASR, such as schizophrenia or ASD. Relationships with those two disorders in particular have been established in adults using the EEfRT paradigm (Damiano *et al.*, 2012; Barch *et al.*, 2014; Treadway *et al.*, 2015; McCarthy *et al.*, 2016), but studies of children have not been conducted to date.

In addition, future studies should explore the relationship between other Positive Valence System constructs and psychopathology and impairment using a similar approach, to determine if these constructs differentially associate with various forms of

psychopathology and functioning. It will be important to consider longitudinal work to investigate the stability of effort valuation over time, especially with regards to children passing through stages of their development. Such work will be key to understanding how effort valuation at various stages of life might be used to predict future psychopathology and functioning in individuals.

Conclusions

The current study establishes effort valuation as a relevant psychological construct for understanding psychopathology and functioning in adults and, especially, in children. Findings from this study both confirm and expand upon the current state of knowledge in the field with regards to examining sex when considering how effort valuation relates to anxiety disorder and ASDs in children, and alcohol and drug use in adults. The associations observed with regards low effort expenditure, low reward sensitivity, and these types of psychopathology indicate that deficits in reward processing behavior may be an important therapeutic target. Furthermore, effort expenditure and reward sensitivity may be potentially modifiable risk factors, and our data demonstrates that the degree to which these risk factors impact pathology might be a key factor in developing targeted treatment strategies, as well as an important consideration in further research to understand the biological mechanisms underlying these behavioral deviations. These results highlight the importance of examining effort valuation as it relates to areas of psychopathology and functioning not previously examined, and provides evidence in favor of longitudinal study of effort valuation to further determine the stability of the construct over time and through developmental stages.

Supplementary material. The supplementary material for this article can be found at <https://doi.org/10.1017/S0033291718003884>.

Author ORCIDs.  Nicholas H. Nguyen 0000-0002-0744-1920
Stephen J. Glatt 0000-0002-0360-7567

Acknowledgements. The authors would like to thank our subjects for their participation in this research study, which was funded by NIH grants and supplements (R01MH101519-01A1 and R01MH101519-01A1S1). The authors also wish to thank Tom Weickert for his helpful and critical discussion.

References

- Achenbach TM (1991) Integrative guide to the 1991 CBCL/4-18 YSR, and TRF profiles. *University of Vermont, Department of Psychology Pediatrics* 108, e14.
- Altemus M, Sarvaiya N and Neill Epperson C (2014) Sex differences in anxiety and depression clinical perspectives. *Frontiers in Neuroendocrinology* 35, 320–330.
- Barch DM, Treadway MT and Schoen N (2014) Effort, anhedonia, and function in schizophrenia: reduced effort allocation predicts amotivation and functional impairment. *Journal of Abnormal Psychology* 123, 387–397.
- Benjamini Y and Hochberg Y (1995) Controlling the false discovery rate: a practical and powerful approach to multiple testing. *Journal of the Royal Statistical Society B* 57, 289–300.
- Benozio A and Diesendruck G (2015) From effort to value. *Psychological Science* 26, 1423–1429.
- Cross-Disorder Group of the Psychiatric Genomics Consortium, Lee SH, Ripke S, Neale BM, Faraone SV, Purcell SM, Perlis RH, Mowry BJ, Thapar A, Goddard ME, Witte JS, Absher D, Agartz I, Akil H, Amin F, Andreassen OA, Anjorin A, Anney R, Anttila V, Arking DE, Asherson P, Azevedo MH, Backlund L, Badner JA, Bailey AJ, Banaschewski T, Barchas JD, Barnes MR, Barrett TB, Bass N, Battaglia A, Bauer M, Bayés M, Bellivier F, Bergen SE, Berrettini W, Betancur C, Bettecken T, Biederman J, Binder EB, Black DW, Blackwood DH, Bloss CS, Boehnke M, Boomsma DI, Breen G,

- Breuer R, Bruggeman R, Cormican P, Buccola NG, Buitelaar JK, Bunney WE, Buxbaum JD, Byerley WF, Byrne EM, Caesar S, Cahn W, Cantor RM, Casas M, Chakravarti A, Chambert K, Choudhury K, Cichon S, Cloninger CR, Collier DA, Cook EH, Coon H, Cormand B, Corvin A, Coryell WH, Craig DW, Craig IW, Crosbie J, Cuccaro ML, Curtis D, Czamara D, Datta S, Dawson G, Day R, De Geus EJ, Degenhardt F, Djurovic S, Donohoe GJ, Doyle AE, Duan J, Dudbridge F, Duketis E, Ebbstein RP, Edenberg HJ, Elia J, Ennis S, Etain B, Fanous A, Farmer AE, Ferrier IN, Flickinger M, Fombonne E, Foroud T, Frank J, Franke B, Fraser C, Freedman R, Freimer NB, Freitag CM, Friedl M, Frisén L, Gallagher L, Gejman PV, Georgieva L, Gershon ES, Geschwind DH, Giegling I, Gill M, Gordon SD, Gordon-Smith K, Green EK, Greenwood TA, Grice DE, Gross M, Grozeva D, Guan W, Gurling H, De Haan L, Haines JL, Hakonarson H, Hallmayer J, Hamilton SP, Hamshire ML, Hansen TF, Hartmann AM, Hautzinger M, Heath AC, Henders AK, Herms S, Hickie IB, Hipolito M, Hoefels S, Holmans PA, Holsboer F, Hoogendijk WJ, Hottenga JJ, Hultman CM, Hus V, Inganos A, Ising M, Jamain S, Jones EG, Jones I, Jones L, Tzeng JY, Kähler AK, Kahn RS, Kandaswamy R, Keller MC, Kennedy JL, Kenny E, Kent L, Kim Y, Kirov GK, Klauck SM, Klei L, Knowles JA, Kohli MA, Koller DL, Konte B, Korszun A, Krabbendam L, Krasucki R, Kuntsi J, Kwan P, Landén M, Långström N, Lathrop M, Lawrence J, Lawson WB, Leboyer M, Ledbetter DH, Lee PH, Lencz T, Lesch KP, Levinson DF, Lewis CM, Li J, Lichtenstein P, Lieberman JA, Lin DY, Linszen DH, Liu C, Lohoff FW, Loo SK, Lord C, Lowe JK, Lucae S, MacIntyre DJ, Madden PA, Maestrini E, Magnusson PK, Mahon PB, Maier W, Malhotra AK, Mane SM, Martin CL, Martin NG, Mattheisen M, Matthews K, Mattingsdal M, McCarroll SA, McGhee KA, McGough JJ, McGrath PJ, McGuffin P, McInnis MG, McIntosh A, McKinney R, McLean AW, McMahan FJ, McMahan WM, McQuillin A, Medeiros H, Medland SE, Meier S, Melle I, Meng F, Meyer J, Middeldorp CM, Middleton L, Milanova V, Miranda A, Monaco AP, Montgomery GW, Moran JL, Moreno-De-Luca D, Morken G, Morris DW, Morrow EM, Moskvina V, Muglia P, Mühlisen TW, Muir JW, Müller-Myhsok B, Murtha M, Myers RM, Myin-Germeys I, Neale MC, Nelson SF, Nievergelt CM, Nikolov I, Nimgaonkar V, Nolen WA, Nöthen MM, Nurnberger JI, Nwulia EA, Nyholt DR, O'Dushlaine C, Oades RD, Olincy A, Oliveira G, Olsen L, Ophoff RA, Osby U, Owen MJ, Palotie A, Parr JR, Paterson AD, Pato CN, Pato MT, Penninx BW, Pergadia ML, Pericak-Vance MA, Pickard BS, Pimm J, Piven J, Posthuma D, Potash JB, Poustka F, Propping P, Puri V, Quedest DJ, Quinn EM, Ramos-Quiroga JA, Rasmussen HB, Raychaudhuri S, Rehnström K, Reif A, Ribasés M, Rice JP, Rietschel M, Roeder K, Roeyers H, Rossin L, Rothenberger A, Rouleau G, Ruderfer D, Rujescu D, Sanders AR, Sanders SJ, Santangelo SL, Sergeant JA, Schacher R, Schalling M, Schatzberg AF, Scheftner WA, Schellenberg GD, Scherer SW, Schork NJ, Schulze TG, Schumacher J, Schwarz M, Scolnick E, Scott LJ, Shi J, Shilling PD, Shyn SI, Silverman JM, Slager SL, Smalley SL, Smit JH, Smith EN, Sonuga-Barke EJ, St Clair D, State M, Steffens M, Steinhausen HC, Strauss JS, Strohmaier J, Stroup TS, Sutcliffe JS, Szatmari P, Szelinger S, Thirumalai S, Thompson RC, Todorov AA, Tozzi F, Treutlein J, Uhr M, van den Oord EJ, Van Grootheest G, Van Os J, Vicente AM, Vieland VJ, Vincent JB, Visscher PM, Walsh CA, Wassink TH, Watson SJ, Weissman MM, Werge T, Wienker TF, Wijsman EM, Willemsen G, Williams N, Willsey AJ, Witt SH, Xu W, Young AH, Yu TW, Zammit S, Zandi PP, Zhang P, Zitzman FG, Zöllner S, Devlin B, Kelsoe JR, Sklar P, Daly MJ, O'Donovan MC, Craddock N, Sullivan PF, Smoller JW, Kendler KS, Wray NR, International Inflammatory Bowel Disease Genetics Consortium (IBDGC) (2013) Genetic relationship between five psychiatric disorders estimated from genome-wide SNPs. *Nature Genetics* 45, 984–994.
- Cuthbert BN (2014) The RDoC framework: facilitating transition from ICD/DSM to dimensional approaches that integrate neuroscience and psychopathology. *World Psychiatry* 13, 28–35.
- Damiano CR, Aloï J, Treadway M, Bodfish JW and Dichter GS (2012) Adults with autism spectrum disorders exhibit decreased sensitivity to reward parameters when making effort-based decisions. *Journal of Neurodevelopmental Disorders* 4, 1–10. doi: 10.1186/1866-1955-4-13.
- Derdikman-Eiron R, Indredavik MS, Bakken IJ, Bratberg GH, Hjemdal O and Colton M (2012) Gender differences in psychosocial functioning of adolescents with symptoms of anxiety and depression: longitudinal findings from the Nord-Trøndelag Health Study. *Social Psychiatry and Psychiatric Epidemiology* 47, 1855–1863.
- Dorfman J, Rosen D, Pine D and Ernst M (2016) Anxiety and gender influence reward-related processes in children and adolescents. *Journal of Child and Adolescent Psychopharmacology* 26, 380–390.
- Ji Y, Hong X, Wang G, Chatterjee N, Riley AW, Lee L, Surkan PJ, Bartell TR, Zuckerman B and Wang X (2018) A prospective birth cohort study on early childhood lead levels and attention deficit hyperactivity disorder: new insight on sex differences. *The Journal of Pediatrics* 199, 124–131.
- John K, Gammon GD, Prusoff BA and Warner V (1987) The Social Adjustment Inventory for Children and Adolescents (SAICA): testing of a new semistructured interview. *Journal of the American Academy of Child and Adolescent Psychiatry* 26, 898–911. Available at <http://www.ncbi.nlm.nih.gov/pubmed/3429410> (Accessed 16 April 2018).
- Loomes R, Hull L and Mandy WPL (2017) What is the male-to-female ratio in autism spectrum disorder? A systematic review and meta-analysis. *Journal of the American Academy of Child & Adolescent Psychiatry* 56, 466–474.
- Mazefsky CA, Anderson R, Conner CM and Minshew N (2011) Child behavior checklist scores for school-aged children with autism: preliminary evidence of patterns suggesting the need for referral. *Journal of Psychopathology and Behavioral Assessment* 33, 31–37.
- McCarthy JM, Treadway MT, Bennett ME and Blanchard JJ (2016) Inefficient effort allocation and negative symptoms in individuals with schizophrenia. *Schizophrenia Research* 170, 278–284.
- Mosner MG, Kinard JL, McWeeny SS, Jasmine S, Markiewitz ND, Damiano-Goodwin CR, Burchinal MR, Rutherford HJV, Greene RK, Treadway MT and Dichter GS (2017) Vicarious effort-based decision-making in autism spectrum disorders. *Journal of Autism and Developmental Disorders* 47, 2992–3006.
- NIMH (2016) *NIMH Subconstruct: Effort*. [online] Available at: <https://www.nimh.nih.gov/research-priorities/rdoc/constructs/effort.shtml> [Accessed 1 Mar. 2018].
- Olino TM, Silk J, Osterritter C and Forbes EE (2015) Social reward in youth at risk for depression: a preliminary investigation of subjective and neural differences. *Journal of Child and Adolescent Psychopharmacology* 25, 711–721.
- Perneger TV (1998) What's wrong with Bonferroni adjustments. *BMJ* 316, 1236–1238.
- Potts R, Martinez IG, Dedmon A, Schwarz L, DiLillo D and Swisher L (1997) Brief report: cross-validation of the injury behavior checklist in a school-age sample. *Journal of Pediatric Psychology* 22, 533–540. Available at <http://jpepsy.oxfordjournals.org/content/22/4/533.abstract%5Cnpapers3://publication/uuid/56A47B28-1953-4CAE-9167-4A62688147E9>.
- Rescorla LA and Achenbach TM (2004) The Achenbach System of Empirically Based Assessment (ASEBA) for ages 18 to 90 years. In *The Use of Psychological Testing for Treatment Planning and Outcomes Assessment, Volume 3: Instruments for adults*, pp. 115–152, Mahwah, NJ, US: Lawrence Erlbaum Associates Publishers.
- Scott-Van Zeeland AA, Dapretto M, Ghahremani DG, Poldrack RA and Bookheimer SY (2010) Reward processing in autism. *Autism Research: Official Journal of the International Society for Autism Research* 3, 53–67.
- Treadway MT, Buckholz JW, Schwartzman AW, Lambert WE and Zald DH (2009) Worth the “EEfRT”? The effort expenditure for rewards task as an objective measure of motivation and anhedonia. *PLoS ONE* 4, 1–9. doi: 10.1371/journal.pone.0006598.
- Treadway MT, Bossaller NA, Shelton RG and Zald DH (2012) Effort-based decision-making in major depressive disorder: a translational model of motivational anhedonia. *Journal of Abnormal Psychology* 121, 553–558.
- Treadway MT, Peterman JS, Zald DH and Park S (2015) Impaired effort allocation in patients with schizophrenia. *Schizophrenia Research* 161, 382–385.
- Venables NC (2017) Reward mechanisms across aggressive and addictive forms of externalizing psychopathology. *Biological Psychiatry* 82, e25–e27.
- Werling DM and Geschwind DH (2013) Sex differences in autism spectrum disorders. *Current Opinion in Neurology* 26, 146–153.