

UNIVERSAL COUNTABLE BOREL QUASI-ORDERS

JAY WILLIAMS

Abstract. In recent years, much work in descriptive set theory has been focused on the Borel complexity of naturally occurring classification problems, in particular, the study of countable Borel equivalence relations and their structure under the quasi-order of Borel reducibility. Following the approach of Louveau and Rosendal for the study of analytic equivalence relations, we study countable Borel quasi-orders.

In this paper we are concerned with universal countable Borel quasi-orders, i.e., countable Borel quasi-orders above all other countable Borel quasi-orders with regard to Borel reducibility. We first establish that there is a universal countable Borel quasi-order, and then establish that several countable Borel quasi-orders are universal. An important example is an embeddability relation on descriptive set theoretic trees.

Our main result states that embeddability of finitely generated groups is a universal countable Borel quasi-order, answering a question of Louveau and Rosendal. This immediately implies that biembeddability of finitely generated groups is a universal countable Borel equivalence relation. The same techniques are also used to show that embeddability of countable groups is a universal analytic quasi-order.

Finally, we show that, up to Borel bireducibility, there are 2^{\aleph_0} distinct countable Borel quasi-orders, which symmetrize to a universal countable Borel equivalence relation.

§1. Introduction. A countable Borel quasi-order Q is a quasi-order defined on a Polish space X (or more generally, a standard Borel space) such that Q is Borel when viewed as a subset of X^2 and for every $x \in X$, the set of predecessors of x , $\{y \mid y Q x\}$, is countable. There are several natural examples, such as \leq_T and \leq_1 on $2^{\mathbb{N}}$, as well as the embeddability relation on the space \mathcal{G} of finitely generated groups.

As in the case of Borel equivalence relations, we are largely interested in how countable Borel quasi-orders are related to each other under Borel reducibility.

DEFINITION 1.1.

a) Suppose that Q is a Borel quasi-order on a Polish space X and Q' is a Borel quasi-order on a Polish space Y . We say that Q is Borel reducible to Q' , written $Q \leq_B Q'$, if there is a Borel function $f: X \rightarrow Y$ such that

$$x Q y \Leftrightarrow f(x) Q' f(y).$$

b) A countable Borel quasi-order Q' is universal if for every countable Borel quasi-order Q , $Q \leq_B Q'$.

Borel reducibility is intended to capture the notion of the relative complexity of quasi-orders, so that if $Q \leq_B Q'$, then we consider Q' to be more complicated than Q . A universal countable Borel quasi-order can then be thought of as

Received February 26, 2013.

Key words and phrases. Borel equivalence relations, quasi-orders, monoids, group theory, small cancellation.

© 2014, Association for Symbolic Logic
0022-4812/14/7903-0015/\$3.40
DOI:10.1017/jsl.2013.35

a countable Borel quasi-order which is as complicated as possible. It was shown by Dougherty, Jackson, and Kechris in [3] that there is a universal countable Borel equivalence relation, and in [9] Louveau and Rosendal showed there is a universal analytic quasi-order. In section 2, using an analog of the Feldman-Moore Theorem, we prove the following result.

THEOREM 1.2. *There is a universal countable Borel quasi-order.*

Given a quasi-order Q on X , the corresponding equivalence relation E_Q on X is defined by

$$x E_Q y \Leftrightarrow x Q y \wedge y Q x.$$

For example, Turing equivalence \equiv_T is E_{\leq_T} . It is easily checked that if Q is a universal countable Borel quasi-order, then E_Q is a universal countable Borel equivalence relation, and this provides another source of examples for such equivalence relations.

Much of this paper is dedicated to proving that various countable Borel quasi-orders are universal. One universal countable Borel quasi-order, in particular, is used in several of these proofs. Recall that given a discrete space X , a subset $T \subseteq X^{<\mathbb{N}}$ is said to be a tree on X if it is closed under initial segments.

DEFINITION 1.3. Given a discrete space X , the quasi-order \preceq_X^{tree} on the space of trees on X is defined by

$$T \preceq_X^{tree} T' \Leftrightarrow (\exists u \in X^{<\mathbb{N}}) T = T'_u,$$

where $T'_u = \{v \in X^{<\mathbb{N}} \mid u \frown v \in T'\}$. (Here, $u \frown v$ indicates the string u followed by v .)

THEOREM 1.4. \preceq_2^{tree} is a universal countable Borel quasi-order.

This quasi-order is combinatorially simple and thus easy to work with. This makes it useful for establishing other quasi-orders are universal.

Our ultimate goal is to show the biembeddability relation for finitely generated groups is a universal countable Borel equivalence relation. Although it may be possible to prove this only using results on countable Borel equivalence relations, the use of quasi-orders seems to be the most direct route to this result. We first use small cancellation methods from group theory to establish the following theorem.

THEOREM 1.5. *Embeddability of countable groups is a universal analytic quasi-order.*

It follows that biembeddability of countable groups is a universal analytic equivalence relation. This is strictly more complex than isomorphism of countable groups; for a more thorough exploration of the relation between biembeddability and isomorphism in the Borel reducibility context see [6]. The ideas developed to prove Theorem 1.5 are then used to reduce \preceq_2^{tree} to the embeddability relation for finitely generated groups. Thus we have the following theorem.

THEOREM 1.6. *Embeddability of finitely generated groups is a universal countable Borel quasi-order.*

Thus the embeddability structure of finitely generated groups is as complicated as possible. We find as a corollary that the biembeddability relation for finitely generated groups is a universal countable Borel equivalence relation. This answers

a question of Louveau and Rosendal in [9]. It was previously shown in [12] that the isomorphism relation for finitely generated groups is a universal countable Borel equivalence relation, and so this result can be seen as saying the two relations have precisely the same complexity, in contrast with the case of countable groups.

The rest of the paper is organized as follows. In section 2, we establish that there is a universal countable Borel quasi-order by proving a Feldman-Moore-type result relating countable Borel quasi-orders to Borel actions of countable monoids. In section 3, we show that \preceq_2^{tree} is universal. In section 4, we show that various group-theoretic countable Borel quasi-orders, which arise as simple generalizations of well-known universal countable Borel equivalence relations, are universal. In sections 5 and 6, we prove the theorems on the embeddability relations for countable groups and finitely generated groups. Finally, in section 7 we show that, up to Borel bireducibility, there are 2^{\aleph_0} countable Borel quasi-orders, which give rise to a universal countable Borel equivalence relation.

§2. A universal countable Borel quasi-order. We start by proving an analog of the Feldman-Moore Theorem [5] for countable Borel quasi-orders. Although to the author’s knowledge, this result is not in the literature, it is a straightforward application of the well-known Lusin-Novikov theorem (see Theorem 18.10 in Kechris [8]), and should perhaps be considered as folklore.

THEOREM 2.1. *If \preceq is a countable Borel quasi-order on the Polish space X , there is a monoid M which acts on X in a Borel way such that*

$$x \preceq y \iff (\exists m \in M) x = m \cdot y.$$

PROOF. First, note that by definition for all $y \in X$, $\preceq_y = \{x \mid x \preceq y\}$ is countable, which implies the set $\preceq \subseteq X \times X$ has countable sections with respect to its second coordinate. By the Lusin-Novikov theorem, $\succcurlyeq = \cup_n f_n$, where each $f_n: E_n \rightarrow X$ is a Borel function, with $E_n \subseteq X$ Borel.

We can extend these to functions defined on all of X by letting $f_n(y) = y$ for $y \in X \setminus E_n$. These functions are still Borel, and their union is still equal to \succcurlyeq by reflexivity. We may also add the identity function to our collection without changing the union, again by reflexivity. With all this in place, the collection f_n generates a monoid M under composition, and M acts on X by $m \cdot x = m(x)$. If $x \preceq y$, then there exists $m \in M$ such that $x = m \cdot y = m(y)$, and the transitivity of \preceq ensures that for all $m \in M$ and $x \in X$, $m \cdot x \preceq x$. \dashv

We wish to use this result to show that there is a universal countable Borel quasi-order. Our approach closely follows the proof of Dougherty, Jackson, and Kechris in [3] that there is a universal countable Borel equivalence relation.

DEFINITION 2.2. For every standard Borel space X and countable monoid M , the corresponding canonical Borel action of M on X^M is defined by $(m \cdot f)(s) = f(sm)$ for $m, s \in M$ and $f \in X^M$. We denote the corresponding quasi-order by \preceq_M^X , i.e., for $f, g \in X^M$,

$$f \preceq_M^X g \iff (\exists m \in M) f = m \cdot g.$$

To see that this is an action, let $m, n \in M$ and $f \in X^M$. Then

$$\begin{aligned} (m \cdot (n \cdot f))(s) &= (n \cdot f)(sm) \\ &= f(sm) \\ &= (mn \cdot f)(s), \end{aligned}$$

as desired.

Note that in the above definition, if M is a group, then \preceq_M^X is in fact an equivalence relation. In particular, $\preceq_{\mathbb{F}_2}^2$ is the universal countable Borel equivalence relation E_∞ .

DEFINITION 2.3 (The quasi-order \preceq_ω). Let M_ω be the free monoid on countably many generators. Then define \preceq_ω to be $\preceq_{M_\omega}^{2^{\mathbb{N}}}$.

THEOREM 2.4. \preceq_ω is a universal countable Borel quasi-order.

PROOF. Let \preceq be a countable Borel quasi-order on a Polish space X . By Theorem 2.1, there is a countable monoid M such that \preceq is the quasi-order induced by a Borel action of M on X . Let $f : M_\omega \rightarrow M$ be a surjective homomorphism. Then we can define an action of M_ω on X by

$$m \cdot x = f(m) \cdot x.$$

This action is Borel and also induces \preceq , and so without loss of generality we may assume that $M = M_\omega$.

Let $\{U_i\}_{i \in \mathbb{N}}$ be a sequence of Borel sets in X , which separates points. Then we define $\phi : X \rightarrow (2^{\mathbb{N}})^{M_\omega}$ by $x \mapsto \phi_x$, with

$$\phi_x(s)(i) = 1 \iff s \cdot x \in U_i.$$

This map is Borel, and since the collection U_i separates points, we see it is injective. Furthermore, if $t \in M_\omega$, then $t \cdot \phi_x = \phi_{t \cdot x}$. To see this, let $s \in M_\omega$, and $i \in \mathbb{N}$. Then

$$\begin{aligned} \phi_{t \cdot x}(s)(i) = 1 &\iff s \cdot t \cdot x \in U_i \\ &\iff \phi_x(st)(i) = 1 \\ &\iff t \cdot \phi_x(s)(i) = 1. \end{aligned}$$

Now suppose that $x \preceq y$. Then there exists $m \in M_\omega$ such that $x = m \cdot y$. It follows that $\phi_x = \phi_{m \cdot y} = m \cdot \phi_y$, and so $\phi_x \preceq_\omega \phi_y$. The same reasoning works in reverse, and hence $\phi_x \preceq_\omega \phi_y$ implies that $x \preceq y$. Thus ϕ is a Borel reduction. \dashv

Thus there exists a universal countable Borel quasi-order. Next we wish to find universal countable Borel quasi-orders which are easier to work with. We proceed by a series of easily proven lemmas, which are the analogs of propositions 1.4–1.8 in Dougherty, Jackson, and KeCHRIS [3]. The proofs of most of them are virtually the same, and so we omit them here.

LEMMA 2.5. If M, N are monoids and M is a homomorphic image of N , then $\preceq_M^X \leq_B \preceq_N^X$.

LEMMA 2.6. For any countable monoid M , $\preceq_M^{2^{\mathbb{Z}-\{0\}}} \leq_B \preceq_{M \times \mathbb{Z}}^3$.

LEMMA 2.7. For any countable monoid M , $\preceq_M^3 \leq_B \preceq_{M \times \mathbb{Z}_2}^2$.

LEMMA 2.8. *Let M_2 denote the free monoid on 2 generators. Then*

$$\preceq_{M_\omega}^2 \leq_B \preceq_{M_2}^2.$$

PROOF. We start by embedding M_ω into M_2 in order to view it as a submonoid of M_2 . Let $M_\omega = \langle x_1, x_2, \dots \rangle$ and $M_2 = \langle a, b \rangle$. We define our embedding by $e \mapsto e$ and $x_n \mapsto ab^n$ for all $n \in \mathbb{N}^+$.

Next we note that if $h \in M_2$, then we can canonically write h as a product $h = h'g$, with $g \in M_\omega$ and $h' \in M_2 \setminus M_\omega$, possibly with $g = e$ or $h' = e$, by finding the longest word in M_ω at the end of h . Define $L: M_2 \rightarrow \mathbb{N}$ by

$$L(h) = \text{the length of } h',$$

where $h = h'g$ is the canonical form of h . This function has the desirable property that multiplying an element $h \in M_2$ on the right by an element $g \in M_\omega$ does not change the given length, i.e., $L(h) = L(hg)$.

Define $f: 2^{M_\omega} \rightarrow 2^{M_2}$ by $p \mapsto p^*$, where

$$p^*(h) = \begin{cases} p(h) & \text{if } L(h) = 0 \\ 1 & \text{if } L(h) = 1, \\ 0 & \text{if } L(h) > 1. \end{cases}$$

Suppose that $p \preceq_{M_\omega}^2 q$. Then $\exists g \in M_\omega$ such that $p = g \cdot q$. So if $h \in M_\omega$,

$$\begin{aligned} (g \cdot q^*)(h) &= q^*(hg) \\ &= q(hg) \\ &= (g \cdot q)(h) \\ &= p(h) \\ &= p^*(h). \end{aligned}$$

If $h \in M_2 \setminus M_\omega$, then since $L(h) = L(hg)$, we find

$$\begin{aligned} (g \cdot q^*)(h) &= q^*(hg) \\ &= p^*(h). \end{aligned}$$

So $p^* \preceq_{M_2}^X q^*$.

Now suppose that $p^* \preceq_{M_2}^2 q^*$. Then there exists $g \in M_2$ such that $p^* = g \cdot q^*$. Clearly if $g \in M_\omega$, then $p = g \cdot q$. If instead $g \in M_2 \setminus M_\omega$, then $L(g) \geq 1$, and we should have that $p^*(b) = (g \cdot q^*)(b) = q^*(bg)$. But $p^*(b) = 1$, while $L(bg) > 1$, and so $q^*(bg) = 0$. Thus this case cannot happen, and hence f is a Borel reduction. ⊥

THEOREM 2.9. $\preceq_\omega \leq_B \preceq_{M_2}^2$. *It follows that $\preceq_{M_2}^2$ is universal.*

PROOF. Using the preceding lemmas, we find that

$$\begin{aligned} \preceq_{M_\omega}^{2^{\mathbb{N}}} &\leq_B \preceq_{M_\omega}^{2^{\mathbb{Z}-\{0\}}} \\ &\leq_B \preceq_{M_\omega \times \mathbb{Z}}^3 \text{ by Prop. 2.6} \\ &\leq_B \preceq_{M_\omega \times \mathbb{Z} \times \mathbb{Z}_2}^2 \text{ by Prop. 2.7} \\ &\leq_B \preceq_{M_\omega}^2 \text{ by Prop. 2.5} \\ &\leq_B \preceq_{M_2}^2 \text{ by Prop. 2.8.} \end{aligned}$$
⊥

The quasi-order $\preceq_{M_2}^2$ is easier to work with than \preceq_ω , as both the monoid and the space being acted on are simpler. Using $\preceq_{M_2}^2$, we will find another universal countable Borel quasi-order, this one of a more combinatorial nature.

§3. A quasi-order on trees. In this section, we will reduce $\preceq_{M_2}^2$ to \preceq_2^{tree} , the quasi-order on descriptive-set-theoretic trees from definition 1.3. This has the advantage of moving us away from working with monoids and towards more classical areas of mathematics. We must first make a few intermediate reductions.

DEFINITION 3.1. The quasi-order \preceq_2^s (the s is for “suffix”) on $\mathcal{P}(M_2)$ is defined by

$$A \preceq_2^s B \iff (\exists m \in M_2) Am = B^m,$$

where

$$B^m = B \cap M_2m.$$

REMARK 3.2. Note that if we made a similar definition for a group M , then we would always have that $B^m = B$, since in this case $Mm = M$. So this definition is only interesting when dealing with a monoid.

If we identify $\mathcal{P}(M_2)$ with 2^{M_2} , then this quasi-order is the same as $\preceq_{M_2}^2$. Writing it in this way brings out the fact that knowing a set $A \in \mathcal{P}(M_2)$ and that $A \preceq_{M_2}^2 B$ only gives partial information about B . This differs from E_∞ , the analogous equivalence relation, since knowing $A \in \mathcal{P}(\mathbb{F}_2)$ and that $A E_\infty B$ gives information about all of B .

Next, we modify this quasi-order slightly, in order to make it somewhat easier to work with.

DEFINITION 3.3. The quasi-order \preceq_2^p (the p is for “prefix”) on $\mathcal{P}(M_2)$ is defined by

$$A \preceq_2^p B \iff (\exists m \in M_2) mA = B_m,$$

where

$$B_m = B \cap mM_2.$$

As before, this definition is only interesting when working with a monoid.

THEOREM 3.4. $\preceq_2^s \sim_B \preceq_2^p$

PROOF. Every nontrivial element $w \in M_2$ may be written as $w = a^{n_0}b^{m_0} \dots a^{n_k}b^{m_k}$, where $n_i, m_j \in \mathbb{N}$, and only n_0 or m_k may be 0. Define $\bar{w} = b^{m_k}a^{n_k} \dots b^{m_0}a^{n_0}$, and $\bar{e} = e$. Then the bijection $f : M_2 \rightarrow M_2$ defined by $f(w) = \bar{w}$ induces a Borel bijection $f^* : \mathcal{P}(M_2) \rightarrow \mathcal{P}(M_2)$ such that if $Am = B^m$, then $\bar{m}f^*(A) = f^*(B)_{\bar{m}}$. Similarly, if $wf^*(A) = f^*(B)_w$, then $A\bar{w} = B^{\bar{w}}$. Thus f^* is a Borel reduction from \preceq_2^s to \preceq_2^p . Since f^* is its own inverse, we see that it is also a Borel reduction from \preceq_2^p to \preceq_2^s . \dashv

One can view M_2 as the complete binary tree $2^{<\mathbb{N}}$, with each word in M_2 corresponding to a node in the tree. From this point of view, when looking at $A \subseteq M_2$, we see that A_m is simply the set of words in A , which are above the node corresponding to m . (See figure 1.) This natural interpretation of one of the sets involved in \preceq_2^p in terms of trees leads us to consider the quasi-order \preceq_X^{tree} . Recall that for a countable discrete space X , a tree on X is a (nonempty) collection of finite sequences of




FIGURE 1. The set $A = \{a, b, aa, abb, bab, \dots\}$ in the binary tree corresponding to M_2 . Note that, for example, $A_{ba} = \{bab, \dots\}$ is the set of words in A above ba .

elements of X , which is closed under initial segments. Let $\Lambda(X)$ be the Borel set of infinite trees in $Tr(X)$, the Polish space of trees on X .

Note that if we have $A, B \in \mathcal{P}(M_2)$ and $m \in M_2$ such that $mA = B_m$, and furthermore A, B are both trees on $\{a, b\}$, then m witnesses that $A \preceq_{\{a,b\}}^{tree} B$. If A or B is not a tree, then it does not make sense to compare them using $\preceq_{\{a,b\}}^{tree}$, but this is only a minor difficulty, as we will see in the proof of Theorem 3.5.

THEOREM 3.5. $\preceq_2^p \leq_B \preceq_2^{tree} \upharpoonright \Lambda(2)$. It follows that \preceq_2^{tree} is universal.

PROOF. We will create our Borel reduction in two parts. First we will define a Borel reduction from \preceq_2^p to $\preceq_3^{tree} \upharpoonright \Lambda(3)$. Given $A \in \mathcal{P}(M_2)$, we define the tree $T_A \in Tr(3)$ as follows. We start with the complete binary tree $2^{<\mathbb{N}}$, and add to it the sequence $\hat{w} \frown 2$ iff $w \in A$, where \hat{w} is the sequence in $2^{<\mathbb{N}}$ corresponding to the word w in M_2 . This set is closed under initial segments and so is a tree. Clearly it is infinite. Let T_A be this collection of sequences.

Suppose that $A \preceq_2^p B$. Then there exists $m \in M_2$ such that $mA = B_m$. First note that $2^{<\mathbb{N}}$ is contained in both T_A and $(T_B)_{\hat{m}}$. Next suppose that $w \in M_2$. Then

$$\begin{aligned} \hat{w} \frown 2 \in T_A &\iff w \in A \\ &\iff m \frown w \in B \\ &\iff \widehat{m \frown w} \frown 2 = \hat{m} \frown \hat{w} \frown 2 \in T_B \\ &\iff \hat{w} \frown 2 \in (T_B)_{\hat{m}}. \end{aligned}$$

So $T_A = (T_B)_{\hat{m}}$.

Conversely, suppose that $T_A = (T_B)_\alpha$ for some $\alpha \in 3^{<\omega}$. If α contains a 2, then $(T_B)_\alpha$ is $\{\emptyset\}$ or \emptyset , since the only sequences in T_B containing 2 are leaves of the tree. However, T_A is infinite. So $\alpha \in 2^{<\omega}$, which means that there is a word $w \in M_2$ such that $\hat{w} = \alpha$. Now

$$\begin{aligned} wx \in B &\iff \widehat{wx} \frown 2 \in T_B \\ &\iff \hat{x} \frown 2 \in (T_B)_{\hat{w}} \\ &\iff \hat{x} \frown 2 \in T_A \\ &\iff x \in A, \end{aligned}$$

so $wA = B_w$. Thus the map $t : \mathcal{P}(M_2) \rightarrow Tr(3)$ sending A to T_A is a Borel reduction.

Next, we define a map $C : Tr(3) \rightarrow Tr(2)$, which when composed with t will be a Borel reduction from \preceq_2^P to $\preceq_2^{tree} \upharpoonright \Lambda(2)$. First we inductively define a map $c : 3^{<\mathbb{N}} \rightarrow 2^{<\mathbb{N}}$. Let $c(e) = e, c(0) = 00, c(1) = 01$, and $c(2) = 10$. Now assume that c has been defined for all words of length $\leq n$, and let $w = x \frown u$, where $x \in \{a, b, c\}$ and $u \in 3^{<\mathbb{N}}$ has length n . Define $c(w) = c(x) \frown c(u)$. Given $T \in Tr(3)$, apply c to the elements of T and close the resulting set under initial segments to get a tree $C(T) \in Tr(2)$.

Suppose that $t(A) \preceq_3^{tree} t(B)$, so there exists $u \in 3^{<\mathbb{N}}$ (in fact, $u \in 2^{<\mathbb{N}}$) such that $t(A) = t(B)_u$. Then for all $w \in 3^{<\mathbb{N}}$

$$\begin{aligned} c(w) \in c(t(A)) &\iff w \in t(A) \\ &\iff u \frown w \in t(B) \\ &\iff c(u \frown w) = c(u) \frown c(w) \in c(t(B)). \end{aligned}$$

Hence $C(t(A)) = C(t(B))_{c(u)}$ and thus $C(t(A)) \preceq_2^{tree} C(t(B))$.

Now suppose that $C(t(A)) \preceq_2^{tree} C(t(B))$, and so there exists $w \in 2^{<\mathbb{N}}$ such that $C(t(A)) = C(t(B))_w$. Suppose that w is not in the image of c . Then we either have $C(t(B))_w = \emptyset$, which is impossible, or w is an initial segment of odd length of something in the image of c . If w ends in a 0, then $100 \in C(t(B))_w$, but this is not in $C(t(A))$. If w ends in a 1, then $00 \notin C(t(B))_w$, but $00 \in C(t(A))$. Thus w is in the image of c , say $w = c(u)$. Then

$$\begin{aligned} u \frown v \in t(B) &\iff c(u) \frown c(v) \in C(t(B)) \\ &\iff c(v) \in C(t(A)) \\ &\iff v \in t(A). \end{aligned}$$

Thus $t(A) = t(B)_u$, and so $t(A) \preceq_3^{tree} t(B)$. ◻

§4. Universal quasi-orders from group theory. We have seen that E_∞ is the same as the quasi-order $\preceq_{\mathbb{F}_2}^2$, and so our universal quasi-order $\preceq_{M_2}^2$ is a natural generalization of E_∞ . At this point, we will turn our attention to other quasi-orders, which can be seen as generalizations of E_∞ . The most obvious generalization is the quasi-order $\subseteq_{\mathcal{P}(\mathbb{F}_2)}^{\mathbb{F}_2, t}$ on $\mathcal{P}(\mathbb{F}_2)$ defined by

$$A \subseteq_{\mathcal{P}(\mathbb{F}_2)}^{\mathbb{F}_2, t} B \iff (\exists g \in \mathbb{F}_2) gA \subseteq B.$$

Replacing the \subseteq symbol on the right-hand side of the definition with the $=$ symbol gives E_∞ . Unfortunately for our purposes, the above quasi-order is clearly not countable, and in fact has been shown to be a universal K_σ quasi-order (see Louveau-Rosendal [9]). Consequently, $\subseteq_{\mathcal{P}(\mathbb{F}_2)}^{\mathbb{F}_2, t}$ is much more complex than any countable Borel quasi-order. So we instead consider the following (countable Borel) quasi-order.

DEFINITION 4.1. If G is a countable group, then \preceq_t^G is the countable Borel quasi-order on $\mathcal{P}(G)$ defined by

$$A \preceq_t^G B \iff (\exists g_1, \dots, g_n \in G) A = g_1 B \cap \dots \cap g_n B.$$

For any group G , let $\Omega(G)$ be the set of infinite subsets of G . In order to show that $\preceq_t^{\mathbb{F}_2}$ is a universal countable Borel quasi-order, we will reduce $\preceq_2^{tree} \upharpoonright \Lambda(2)$ to $\preceq_t^{\mathbb{F}_2} \upharpoonright \Omega(\mathbb{F}_2)$.

Every tree on 2 is isomorphic to a tree T on $\{a, b\}$, and these can easily be identified with subsets of $\mathbb{F}_2 = \langle a, b \rangle$. If we take a subset $T \subset \mathbb{F}_2$ corresponding to a tree and multiply it on the left by w^{-1} , then the positive words in $w^{-1}T$ are precisely T_w . Unfortunately, there is no natural way to pick out the positive words from $w^{-1}T$ simply by intersecting it with other shifts of T , and so we instead will define a set based on T for which T_w is easy to find simply by intersecting its shifts. In order to do this, we will look at subsets of \mathbb{F}_∞ , the free group on countably many generators. We list the generators of \mathbb{F}_∞ as

$$\{a, b, x_a, x_b, x_{aa}, x_{ab}, x_{ba}, x_{bb}, x_{aaa}, \dots\}.$$

Using the two generators a, b we identify T with a subset of the group, to which we add the sets $x_w w T_w$ for $w \in T$. Call this new set T' . Note that for all $w \in T$, $w \frown T_w$ is a subset of T , and so $x_w w T_w \subseteq T'$. Then $T' \cap x_w^{-1} T' = w T_w$, since $w T_w$ is the set of positive words in $x_w^{-1} T'$. We can then multiply by $w T_w$ by w^{-1} to find T_w . However, the map sending T to T' is not a Borel reduction. Although we can now find T_w by intersecting shifts of T' , T_w maps to $(T_w)'$, so that is the set we need to find. The following proof addresses this issue.

THEOREM 4.2. $\preceq_{\mathbb{F}_\infty} \upharpoonright \Omega(\mathbb{F}_\infty)$ is a universal countable Borel quasi-order.

PROOF. We will construct the reduction in a few steps. We start with trees on $\{a, b\}$, which we then map to trees on $\{a, b, c, d\}$ for technical reasons. Next we define a map $f: \{a, b, c, d\}^{<\mathbb{N}} \rightarrow \mathcal{P}(\mathbb{F}_\infty)$, which will induce a map $F: Tr(\{a, b, c, d\}) \rightarrow \mathcal{P}(\mathbb{F}_\infty)$. The composition of these two maps will be our reduction.

If $T \in Tr(\{a, b\})$, define

$$t_a(T) = \{w \in T \mid w \frown a \notin T\}.$$

Similarly define $t_b(T)$. These sets are elements of T , which are “along the edge” of the tree, i.e., some immediate extension of these words is not in the tree. We define $S: Tr(\{a, b\}) \rightarrow Tr(\{a, b, c, d\})$ by

$$S(T) = T \cup (t_a(T) \frown c) \cup (t_b(T) \frown d), \tag{4.1}$$

where $X \frown z = \{x \frown z \mid x \in X\}$. Here, S “outlines” the tree using the letters c and d . The following property of S will be important later.

LEMMA 4.3. If $T, T' \in Tr(\{a, b\})$ and $S(T) \subseteq S(T')$, then $T = T'$.

PROOF. It is easily seen that $S(T) \subseteq S(T')$ implies $T \subseteq T'$, as

$$S(T) \cap \{a, b\}^{<\mathbb{N}} = T \text{ and } S(T') \cap \{a, b\}^{<\mathbb{N}} = T'.$$

Suppose $w \in \{a, b\}^{<\mathbb{N}} \setminus T$. Then there is some initial segment of $w' \subset w$ (possibly the empty string) and some $x \in \{a, b\}$ such that $w' \in t_x(T)$, i.e., $w = w' \frown x \frown t$, where $w' \in T$, $w' \frown x \notin T$, and $t \in \{a, b\}^{<\mathbb{N}}$. Then $w' \frown y \in S(T)$ for some $y \in \{c, d\}$, and so $w' \frown y \in S(T')$. This is only possible if $w' \frown x$ and all its extensions are not in T' , and in particular $w \notin T'$. \dashv

We list the generators of \mathbb{F}_∞ as

$$\{a, b, c, d, x_a, x_b, x_c, x_d, x_{aa}, x_{ab}, x_{ac} \dots\},$$

i.e., every string in $\{a, b, c, d\}^{<\mathbb{N}}$ (except the empty string) has a unique generator associated with it in addition to generators corresponding to the letters in our trees.

The empty string in $\{a, b, c, d\}^{<\mathbb{N}}$ and the identity element in \mathbb{F}_∞ will both be written as e . This should not cause confusion, although both uses will appear close to each other. Finally, we recall that if $A, B \in \mathcal{P}(\mathbb{F}_\infty)$, then $AB = \{ab \mid a \in A, b \in B\}$. We can now define $f: \{a, b, c, d\}^{<\mathbb{N}} \rightarrow \mathcal{P}(\mathbb{F}_\infty)$ inductively.

$$\begin{aligned} f(e) &= \{e\} \\ f(a) &= \{a, x_a a\} \\ f(b) &= \{b, x_b b\} \\ f(c) &= \{c, x_c c\} \\ f(d) &= \{d, x_d d\} \\ f(w) &= \left(\bigcup_{\substack{w=s\hat{\ }t \\ s,t \neq e}} f(s)f(t) \right) \cup \{x_w w\}. \end{aligned}$$

The idea here is that every set $f(w)$ contains elements, which encode the relation of w to its initial segments. Then define $F: Tr(\{a, b, c, d\}) \rightarrow \mathcal{P}(\mathbb{F}_\infty)$ by

$$F(T) = \bigcup_{w \in T} f(w).$$

There are a few helpful facts to record at this point. The simplest one is that $w \in f(w)$, which follows by a simple induction. The others we record as lemmas.

LEMMA 4.4. *If $u, v \in \{a, b, c, d\}^{<\mathbb{N}}$ are not equal, the sets $f(u)$ and $f(v)$ are disjoint.*

PROOF. Define the function $\Phi: \mathbb{F}_\infty \rightarrow \{a, b, c, d\}^{<\mathbb{N}}$ as

$\Phi(g)$ = the word in $\{a, b, c, d\}^{<\mathbb{N}}$ obtained by removing all other letters from the freely reduced representation of g .

By a simple inductive argument we see that for all $w \in \{a, b, c, d\}^{<\mathbb{N}}$, Φ is constant on $f(w)$ and equal to w . Thus the sets are disjoint. \dashv

LEMMA 4.5. *If a word starting with x_w is in $f(u)$, then $w \subset u$.*

PROOF. This follows from a straightforward induction on the length of u . \dashv

LEMMA 4.6. *If $\gamma \in f(u)$ starts with $x_w w$ and $u = w \hat{\ } t$, then $\gamma = x_w w \lambda$, with $\lambda \in f(t)$.*

PROOF. We prove this by an induction on the length of t . If $t = e$, then $\gamma = x_w w$. Otherwise, there must be some α, β such that $u = \alpha \hat{\ } \beta$ and $\gamma \in f(\alpha)f(\beta)$. We can then split γ into two words, $\gamma = \delta \lambda$, where δ starts with $x_w w$ and $\delta \in f(\alpha)$, while $\lambda \in f(\beta)$. By the Lemma 4.5, $w \subset \alpha$, say $\alpha = w \hat{\ } z$. Then $u = w \hat{\ } z \hat{\ } \beta$. We write $t = z \hat{\ } \beta$. As $|z| < |t|$, by assumption $\delta = x_w w \delta'$ with $\delta' \in f(z)$. Then $\gamma = x_w w \delta' \lambda$, and $\delta' \lambda \in f(z)f(\beta) \subseteq f(t)$ by definition. \dashv

We define the map $G: Tr(\{a, b\}) \rightarrow \mathcal{P}(\mathbb{F}_\infty)$ by

$$G(T) = F(S(T)),$$

where S is the map defined in (4.1).

LEMMA 4.7. For all $w \in \{a, b\}^{<\mathbb{N}}$ and all nonempty $T \in \text{Tr}(\{a, b, c, d\})$,

$$G(T) \cap x_w^{-1}G(T) = wG(T_w)$$

and hence

$$w^{-1}G(T) \cap (x_w w)^{-1}G(T) = G(T_w).$$

PROOF. First, we will show that $G(T) \cap x_w^{-1}G(T) \subseteq wG(T_w)$. Every element of $G(T)$ is a positive word in the generators of \mathbb{F}_∞ , so any word not starting with x_w will be freely reduced in $x_w^{-1}G(T)$ and so begin with x_w^{-1} , and thus not be in $G(T)$. So we need only focus on the words that start with x_w .

Suppose $g \in f(u) \subseteq G(T)$ and $g = x_w \alpha$ for some $\alpha \in \mathbb{F}_\infty$. By our inductive definition, this implies $g = x_w w \beta$ for some $\beta \in \mathbb{F}_\infty$. By Lemma 4.5, we must have $u = w \frown t$ for some $t \in \{a, b, c, d\}^{<\mathbb{N}}$. By Lemma 4.6, $\beta \in f(t)$. Also, $w\beta$ is in $f(w)f(t)$, so $w\beta \in G(T) \cap x_w^{-1}G(T)$. In addition, $\beta \in G(T_w)$, since $t \in (S(T))_w = S(T_w)$ (since $w \in \{a, b\}^{<\mathbb{N}}$) and so $f(t) \subseteq G(T_w)$. Thus $G(T) \cap x_w^{-1}G(T) \subseteq wG(T_w)$.

If $g \in G(T_w)$, then there is some $u \in S(T_w)$ such that $g \in f(u)$. Then

$$x_w w g, w g \in f(w)f(u) \subseteq G(T),$$

so $w g \in G(T) \cap x_w^{-1}G(T)$. Thus $G(T) \cap x_w^{-1}G(T) \supseteq wG(T_w)$. ⊖

Lemma 4.7 shows that for $T, S \in \Lambda(2)$, if $T \preceq_2^{tree} S$, then $G(T) \preceq_t^{\mathbb{F}_\infty} G(S)$. Next we check the other direction.

Suppose that $G(T) = g_1 G(T') \cap \dots \cap g_n G(T')$. We know that $e \in G(T)$ (since T is nonempty), which means that each g_i must be an inverse of an element in $G(T')$, say $g_i^{-1} = h_i \in G(T')$. Fix some $1 \leq i \leq n$ and suppose that $\Phi(h_i) = w$, i.e., $h_i \in f(w)$. If $u \in S(T)$, then $x_w u \in G(T)$. This implies $h_i x_w u \in G(T') \cap f(w \frown u)$, and, in particular, the intersection is nonempty, so $w \frown u \in S(T')$. Thus $S(T) \subseteq S(T')_w$.

If $w \notin \{a, b\}^{<\mathbb{N}}$, then $S(T')_w$ is either empty or a single element, but $S(T)$ is infinite. Thus $w \in \{a, b\}^{<\mathbb{N}}$, and so $S(T')_w = S(T'_w)$. It follows that $S(T) \subseteq S(T'_w)$, and so by Lemma 4.3, $T = T'_w$. Thus G is a Borel reduction. This completes the proof of Theorem 4.2. ⊖

COROLLARY 4.8. $\preceq_t^{\mathbb{F}_2} \upharpoonright \Omega(\mathbb{F}_2)$ is a universal countable Borel quasi-order, and so $\preceq_t^{\mathbb{F}_2}$ is a universal countable Borel quasi-order.

PROOF. Let $\phi: \mathbb{F}_\infty \rightarrow \mathbb{F}_2$ be an embedding. Then ϕ induces a map

$$\begin{aligned} \Phi: \Omega(\mathbb{F}_\infty) &\rightarrow \Omega(\mathbb{F}_2) \\ A &\mapsto \{\phi(a) \mid a \in A\}. \end{aligned}$$

If $A, B \in \Omega(\mathbb{F}_\infty)$ and there exist $g_1, \dots, g_n \in \mathbb{F}_\infty$ such that

$$A = g_1 B \cap \dots \cap g_n B,$$

then $\Phi(A) = \phi(g_1)\Phi(B) \cap \dots \cap \phi(g_n)\Phi(B)$.

Conversely, suppose that

$$\Phi(A) = h_1 \Phi(B) \cap \dots \cap h_n \Phi(B). \tag{*}$$

If some h_i is not in the image of ϕ , then $h_i \Phi(B)$ is disjoint from any set in the image of Φ , and so the right hand side cannot equal the left hand side unless $\Phi(A) = \emptyset$,

which is impossible. This implies that every h_i in $(*)$ is in the image of ϕ . It follows that $A = \phi^{-1}(h_1)B \cap \dots \cap \phi^{-1}(h_n)B$. ⊖

REMARK 4.9. The above proof shows that if G is any countable group containing \mathbb{F}_2 as a subgroup, then $\preceq_t^G \upharpoonright \Omega(G)$ is a universal countable Borel quasi-order.

Let $E_c(G)$ denote the conjugacy equivalence relation on the standard Borel space $\text{Sg}(G)$ of subgroups of G , i.e., for $A, B \in \text{Sg}(G)$,

$$A E_c(G) B \iff (\exists g \in G) A = gBg^{-1}.$$

In [7], Gao used a simple coding technique to prove the following result.

THEOREM 4.10 (Gao). *If $G = K * H$, where K has a nonabelian free subgroup and H is nontrivial cyclic, then $E_c(G)$ is a universal countable Borel equivalence relation.*

In light of the relationship between E_∞ and $\preceq_t^{\mathbb{F}_2}$, it is natural to consider the following countable Borel quasi-order:

DEFINITION 4.11. Let G be a countable group. Then \preceq_c^G is the countable Borel quasi-order on $\text{Sg}(G)$ defined by

$$A \preceq_c^G B \iff (\exists g_1, \dots, g_n \in G) A = g_1 B g_1^{-1} \cap \dots \cap g_n B g_n^{-1}.$$

Let $\Gamma(G)$ be the standard Borel space of infinite subgroups of G . Then the proof of the following result is a straightforward adaptation of Gao’s argument in [7].

THEOREM 4.12. *Suppose that G is a countable group containing a nonabelian free subgroup and that H is a nontrivial cyclic group. Then $\preceq_c^{G * H} \upharpoonright \Gamma(G * H)$ is a universal countable Borel quasi-order, and so $\preceq_c^{G * H}$ is a universal countable Borel quasi-order.*

PROOF. Let $h \in H$ be a generator of H . We define the map $K : \Omega(G) \rightarrow \text{Sg}(G * H)$ by

$$K(A) = \langle hxh^{-1} : x \in A \rangle.$$

This map is Borel, so we need only check that it is a reduction from $\preceq_t^G \upharpoonright \Omega(G)$ to $\preceq_c^{G * H} \upharpoonright \Gamma(G * H)$. We will make use of the observation that $K(A) = \bigstar_{g \in A} gHg^{-1}$.

If $A, B \in \mathcal{P}(G)$, then clearly $K(A \cap B) \subseteq K(A) \cap K(B)$. We will show that $K(A) \cap K(B) \subseteq K(A \cap B)$, so $K(A \cap B) = K(A) \cap K(B)$. Suppose that $g \in K(A) \cap K(B)$, and so can be written both as $g = x_1 h x_1^{-1} \dots x_n h x_n^{-1}$ with $x_1, \dots, x_n \in A$ and as $g = y_1 h y_1^{-1} \dots y_m h y_m^{-1}$ with $y_1, \dots, y_m \in B$. Then clearly $x_1 = y_1$, and so multiplying g on the left by $y_1 h^{-1} y_1^{-1} = x_1 h^{-1} x_1^{-1}$ we find that

$$x_2 h x_2^{-1} \dots x_n h x_n^{-1} = y_2 h y_2^{-1} \dots y_m h y_m^{-1}.$$

Thus $x_2 = y_2$, and repeating this argument we find that $x_i = y_i$ for $1 \leq i \leq \min\{n, m\}$. If for example $m < n$, then we would have the equation

$$x_{m+1} h x_{m+1}^{-1} \dots x_n h x_n^{-1} = e,$$

which is impossible. Similarly it cannot be that $n < m$. Thus $m = n$, and it follows that $g \in K(A \cap B)$.

Also note that if $g \in G$, then $K(gA) = gK(A)g^{-1}$. Thus

$$\begin{aligned} K(g_1 A \cap \dots \cap g_n A) &= K(g_1 A) \cap \dots \cap K(g_n A) \\ &= g_1 K(A) g_1^{-1} \cap \dots \cap g_n K(A) g_n^{-1}. \end{aligned}$$

Suppose that $A, B \in \mathcal{P}(G)$ and that $A \preceq_i^G B$, i.e., there exist $g_1, \dots, g_n \in G$ such that $A = g_1 B \cap \dots \cap g_n B$. Then

$$K(A) = K(g_1 B \cap \dots \cap g_n B) = g_1 K(B) g_1^{-1} \cap \dots \cap g_n K(B) g_n^{-1}.$$

Thus $A \preceq_i^G B$ implies that $K(A) \preceq_c^{G*H} K(B)$.

Next, suppose that $A, B \in \mathcal{P}(G)$ and that $K(A) \preceq_c^{G*H} K(B)$, so there exist $\gamma_1, \dots, \gamma_n \in G * H$ such that $K(A) = \gamma_1 K(B) \gamma_1^{-1} \cap \dots \cap \gamma_n K(B) \gamma_n^{-1}$. For each $x \in A$ and $1 \leq i \leq n$, let $w_{x,i} \in K(B)$ be the element such that $x h x^{-1} = \gamma_i w_{x,i} \gamma_i^{-1}$. Clearly for each $i = 1, \dots, n$ the map $x \mapsto w_{x,i}$ is an injection.

Note that $x h x^{-1}$ is a reduced word in $G * H$. For $1 \leq i \leq n$, we may assume that γ_i is a reduced word in $G * H$, and that $w_{x,i} \in K(B)$ can be written as

$$w_{x,i} = z_1 h^{m_1} z_1^{-1} \dots z_k h^{m_k} z_k^{-1} \quad (z_j \in B, m_k \in \mathbb{Z}).$$

We can rewrite this word as

$$w_{x,i} = u_1 h^{m_1} u_2 h^{m_2} \dots u_t h^{m_t} u_{t+1},$$

where $m_j \in \mathbb{Z} \setminus \{0\}$, $u_j \in G$ and the product $u_1 u_2 \dots u_j = z_j \in B$ for $1 \leq j \leq t + 1$. Furthermore, $w_{x,i}$ is never the trivial word.

The equation $x h x^{-1} = \gamma_i w_{x,i} \gamma_i^{-1}$ implies that starting with the right-hand side, there is a cancellation procedure which eventually leads to the left-hand side. In any such procedure, there must be some occurrence of h in the right-hand side, which is never cancelled. We call this the preserved occurrence of h . Let $\Delta_i \subseteq A$ be the set of elements $x \in A$ for which the preserved occurrence of h in some cancellation procedure is in the original expression for $w_{x,i}$.

We claim that $A \setminus \Delta_i$ is finite for each $1 \leq i \leq n$. If $x \in A \setminus \Delta_i$, then the preserved occurrence of h is in either γ_i or γ_i^{-1} . Suppose that $x_1, x_2 \in A \setminus \Delta_i$ are both words such that the preserved occurrence of h is in γ_i . Then the preserved occurrence of h must be the first h in γ_i , since γ_i is assumed to be reduced. Thus $\gamma_i = k h u$ for some $k \in G, u \in G * H$, and this gives us the two equations

$$\begin{aligned} x_1 h x_1^{-1} &= k h u w_{x_1,i} \gamma_i^{-1}, \\ x_2 h x_2^{-1} &= k h u w_{x_2,i} \gamma_i^{-1}, \end{aligned}$$

which implies that $x_1 = k = x_2$. Thus there is at most one element in $A \setminus \Delta_i$ such that the preserved occurrence of h is in γ_i . A similar argument shows that there is at most one element in $A \setminus \Delta_i$ such that the preserved occurrence of h is in γ_i^{-1} . So $|A \setminus \Delta_i| \leq 2$ for $1 \leq i \leq n$.

As A is infinite, this implies that $\Delta = \bigcap_{1 \leq i \leq n} \Delta_i$ is infinite. If we fix some $x_0 \in \Delta$, then for each $i = 1, \dots, n$, we can write

$$\begin{aligned} x_0 h x_0^{-1} &= \gamma_i w_{x_0,i} \gamma_i^{-1} \\ &= \gamma_i u_i (z_i h z_i^{-1}) v_i \gamma_i^{-1}, \end{aligned}$$

with $z_i \in B, u_i, v_i \in K(B)$, and the displayed h is the preserved occurrence in some cancellation procedure. This implies that $x_0 = \gamma_i u_i z_i$, and $x_0^{-1} = z_i^{-1} v_i \gamma_i^{-1}$.

Let $\beta_i = x_0 z_i^{-1} \in G$. Then $\gamma_i = \beta_i u_i^{-1}$. Thus

$$\begin{aligned} K(A) &= \beta_1 u_1^{-1} K(B) u_1 \beta_1^{-1} \cap \dots \cap \beta_n u_n^{-1} K(B) u_n \beta_n^{-1} \\ &= \beta_1 K(B) \beta_1^{-1} \cap \dots \cap \beta_n K(B) \beta_n^{-1} \\ &= K(\beta_1 B \cap \dots \cap \beta_n B) \end{aligned}$$

and so $A = \beta_1 B \cap \dots \cap \beta_n B$, with each $\beta_i \in G$. Thus $A \preceq_i^G B$, as desired. \dashv

The following result is an immediate consequence of Theorem 4.12.

COROLLARY 4.13. *If $n \geq 3$, then $\preceq_c^{\mathbb{F}_n} \upharpoonright \Gamma(\mathbb{F}_n)$ is a universal countable Borel quasi-order.*

Finally, the proof of the following result is a straightforward adaptation of the proof of Proposition 1 of Thomas-Velickovic [12].

COROLLARY 4.14. *$\preceq_c^{\mathbb{F}_2} \upharpoonright \Gamma(\mathbb{F}_2)$ is a universal countable Borel quasi-order.*

PROOF. Recall that a subgroup H of a group G is said to be malnormal if $gHg^{-1} \cap H = \{1\}$ for all $g \in G \setminus H$, and that \mathbb{F}_3 can be embedded as a malnormal subgroup of \mathbb{F}_2 . Arguing as in Corollary 4.8, we see this embedding induces a Borel reduction from $\preceq_c^{\mathbb{F}_3} \upharpoonright \Gamma(\mathbb{F}_3)$ to $\preceq_c^{\mathbb{F}_2} \upharpoonright \Gamma(\mathbb{F}_2)$. \dashv

§5. Embeddability of countable groups. Our ultimate goal is to show that embeddability of finitely generated groups is a universal countable Borel quasi-order. The techniques we will use in the proof are easier to understand in the more general setting of arbitrary countable groups. With this in mind, we first turn our attention to the embeddability relation for countable groups, \sqsubseteq_{Gp} . By removing the restriction that the groups we work with should be finitely generated, we are allowed more freedom with regards to how we construct groups for our Borel reduction. At the same time, removing this restriction means that \sqsubseteq_{Gp} is an analytic quasi-order, rather than a countable Borel quasi-order. We will prove the following result.

THEOREM 5.1. *\sqsubseteq_{Gp} is a universal analytic quasi-order.*

COROLLARY 5.2. *The biembeddability relation for countable groups \equiv_{Gp} is a universal analytic equivalence relation.*

This is in contrast with the isomorphism relation for countable groups \cong_{Gp} , which is known to be universal among all analytic equivalence relations induced by a Borel action of S_∞ . (This is due to Mekler in [11].) However, such equivalence relations are known not to be universal among all analytic equivalence relations.

Before we prove Theorem 5.1, we need to make a few definitions. We will write \mathcal{C} for the set of countable graphs whose vertex set is \mathbb{N} . By identifying each graph with its edge relation, we see that \mathcal{C} is a closed subset of $2^{\mathbb{N}^2}$ and so is a Polish space.

DEFINITION 5.3. If $S, T \in \mathcal{C}$, then we write $S \sqsubseteq_c T$ if S embeds into T , i.e., there exists $f : \mathbb{N} \rightarrow \mathbb{N}$ such that for all $m, n \in \mathbb{N}$, $(m, n) \in S \Leftrightarrow (f(m), f(n)) \in T$.

In [9], it was shown that \sqsubseteq_c is a universal analytic quasi-order. Thus to show that \sqsubseteq_{Gp} is universal, we only need to show that \sqsubseteq_c Borel reduces to it. To do this, we will use small cancellation techniques to create groups that encode the edge relations of graphs. We recall the following definitions and theorems from small cancellation theory. (See Chapter V of [10] for a fuller treatment of small cancellation theory.)

DEFINITION 5.4. Let R be a subset of a free group F . We say R is *symmetrized* if every element of R is cyclically reduced and whenever $r \in R$, all cyclic permutations of r and r^{-1} are in R .

THEOREM 5.5 (Theorem V.4.4 in [10]). *Let F be a free group. Let R be a symmetrized subset of F and N its normal closure. If R satisfies $C'(\lambda)$ for some $\lambda \leq 1/6$, then every nontrivial element $w \in N$ contains a subword s of some $r \in R$ with $|s| > (1 - 3\lambda)|r| \geq \frac{1}{2}|r|$.*

THEOREM 5.6 (Theorem V.10.1 in [10]). *Suppose that $G = \langle x_1, x_2, \dots \mid R \rangle$ is such that R is a symmetrized subset of $\langle x_1, x_2, \dots \rangle$ satisfying the $C'(1/6)$ small cancellation condition. If w represents an element of finite order in G , then there is some $r \in R$ of the form $r = v^n$ such that w is conjugate to a power of v .*

Note that this implies that if furthermore w is cyclically reduced, then w is a cyclic permutation of some power of v with $v^n \in R$. We will often refer to this consequence of Theorem 5.6.

Now we can proceed to the proof of Theorem 5.1.

PROOF OF THEOREM 5.1. Let $T \in \mathcal{C}$. Although the vertices of T are in fact elements of \mathbb{N} , to avoid confusion we will call the vertices of T v_0, v_1, v_2, \dots . Then G_T is defined to be the group with generators v_0, v_1, \dots and relators

- v_i^7 for all $i \in \mathbb{N}$
- $(v_i v_j)^{11}$ if $(v_i, v_j) \in T$
- $(v_i v_j)^{13}$ if $(v_i, v_j) \notin T$.

Let R_T be the symmetrization of the set of defining relations for G_T . Note that if T is any graph, then R_T satisfies the $C'(1/6)$ condition. Now suppose that $S, T \in \mathcal{C}$ are such that S embeds into T , say via the map f . Then f extends to a group homomorphism from G_S to G_T , as it sends the relations of G_S to relations of G_T .

To see that f is an embedding, let $\alpha = v_{i_1}^{k_1} v_{i_2}^{k_2} \dots v_{i_n}^{k_n}$ be a word in the generators of G_S , so that

$$f(\alpha) = f(v_{i_1})^{k_1} f(v_{i_2})^{k_2} \dots f(v_{i_n})^{k_n}$$

and suppose that $f(\alpha) = 1$ in G_T . Then by the $C'(1/6)$ condition, $f(\alpha)$ must contain more than $1/2$ of a relation in R_T . Note that any such relation must involve only generators in the image of the graph embedding $f: S \rightarrow T$. Suppose that $f(\alpha)$ contains more than half of a relation of the form $f(v_i)^{\pm 7}$. Since f is one-to-one, this cannot happen unless α already contained more than half of $v_i^{\pm 7}$.

Suppose that $f(\alpha)$ contains more than $1/2$ of the relation $(f(v_i) f(v_j))^k$, where the value of k depends on whether or not $(f(v_i), f(v_j)) \in T$. Since

$$(v_i, v_j) \in S \Leftrightarrow (f(v_i), f(v_j)) \in T,$$

it must be the case that $(v_i v_j)^k \in R_S$, and α already contained more than $1/2$ of $(v_i v_j)^k$. Thus $f(\alpha)$ does not contain more than $1/2$ of a relation in R_T unless α contains more than $1/2$ of the corresponding relation in R_S . Since every nontrivial element in G_S may be written as a word which does not contain more than $1/2$ of a relation in R_S , every nontrivial element in G_S maps to a nontrivial element in G_T . Thus if S embeds into T , then G_S embeds into G_T .

Conversely, suppose that $\theta: G_S \rightarrow G_T$ is an embedding. By Theorem 5.6, after adjusting the embedding θ by an inner automorphism of G_T if necessary, we may

assume $\theta(v_0) = t_0^k$ for some k such that $|k| < 7$, where t_0 is some vertex of T , since $\theta(v_0)$ must have order 7. Let $v_j \neq v_0$ be some vertex of S . Again by Theorem 5.6, we find that $\theta(v_j) = ut_j^l u^{-1}$ for some l such that $|l| < 7$, where $u \in G_T$ and t_j is some vertex of T . Unfortunately we cannot eliminate u by an inner automorphism without possibly changing the value of $\theta(v_0)$. We may assume that u is freely reduced and does not start with any power of t_0 . To see this, note that if u began with t_0^m , then we would be able to follow θ by the inner automorphism corresponding to t_0^m without changing the value of $\theta(v_0)$. Thus $\theta(v_0 v_j) = t_0^k u t_j^l u^{-1}$ is cyclically reduced. Since $v_0 v_j$ is a torsion element, so is $\theta(v_0 v_j)$. By Theorem 5.6, $\theta(v_0 v_j)$ must be a cyclic permutation of some power of $(v_m v_n)$, $m, n \in \mathbb{N}$. It immediately follows that $u = 1$, since no such words contain a mix of positive and negative powers. Thus $\theta(v_0 v_j) = t_0^k t_j^l$.

From this we find that $t_0 \neq t_j$, since otherwise $\theta(v_0 v_j)$ would have order 1 or 7, which is impossible since θ is an embedding and $v_0 v_j$ has order 11 or 13. Again, by Theorem 5.6, we find that $t_0^k t_j^l$ has finite order only if $k = l = \pm 1$. As the orders of $v_0 v_j$ and $\theta(v_0 v_j) = t_0^{\pm 1} t_j^{\pm 1}$ are equal, we see that

$$(v_0, v_j) \in S \iff (t_0, t_j) \in T.$$

Let $v_m \neq v_n$ be arbitrary vertices in S . Repeating the above argument with v_0 and v_m , as well as v_0 and v_n , we find there are inner automorphisms ψ_1, ψ_2 of G_T , corresponding to conjugating by suitable powers of t_0 , such that $\psi_1(\theta(v_m)) = t_m^{\pm 1}$ and $\psi_2(\theta(v_n)) = t_n^{\pm 1}$, where $t_m \neq t_0$ and $t_n \neq t_0$. A priori it may be the case that, for example, $\psi_1(\theta(v_n)) = t_0^k t_n^{\pm 1} t_0^{-k}$, with $k \neq 0$. But then

$$\psi_1(\theta(v_m v_n)) = t_m^{\pm 1} t_0^k t_n^{\pm 1} t_0^{-k}$$

has infinite order, which is impossible. Thus $\psi_1 = \psi_2$, and so $\psi_1(\theta(v_m v_n)) = t_m^{\pm 1} t_n^{\pm 1}$, and the above argument shows $t_m \neq t_n$ and that

$$(v_m, v_n) \in S \iff (t_m, t_n) \in T.$$

As v_m and v_n were arbitrary, the function $g : S \rightarrow T$ defined by $g(v_i) = t_i$ for all $i \in \mathbb{N}$ is an embedding. Thus $\sqsubseteq_G \leq_B \sqsubseteq_{G_P}$, which establishes the result. \dashv

§6. Embeddability of finitely generated groups. We now turn our attention to the embeddability relation for finitely generated groups.

DEFINITION 6.1. Let \mathcal{G} denote the Polish space of finitely generated groups. (See [2].) If $A, B \in \mathcal{G}$, then we write $A \preceq_{em} B$ if and only if there is a group embedding from A into B . We write \equiv_{em} for the associated equivalence relation.

It can be seen that \preceq_{em} is a countable Borel quasi-order, as any finitely generated group contains only countably many finitely generated subgroups, and isomorphism classes in \mathcal{G} are countable. We will show that in fact it is universal by reducing \preceq_2^{tree} to \preceq_{em} . Given a tree $T \in Tr(2)$, our general strategy is to define a finitely generated group G_T with subgroups corresponding to the trees T_w for $w \in 2^{<\mathbb{N}}$. We will start with two generators and then add relations to this group according to the nodes present in T . As in the section 5, these additional relations will allow us to control

the embeddings that exist between two of these groups and thus ensure that $T \mapsto G_T$ is a Borel reduction. Thus we will have shown:

THEOREM 6.2. \preceq_{em} is a universal countable Borel quasi-order.

In order to define the relations of G_T , we first define the following two homomorphisms:

$$\begin{array}{ll} f_0: \mathbb{F}_2 \rightarrow \mathbb{F}_2 & f_1: \mathbb{F}_2 \rightarrow \mathbb{F}_2 \\ x \mapsto x^5y & x \mapsto x^2yx^5yx \\ y \mapsto y^5x & y \mapsto y^2xyxy. \end{array}$$

We also define f_e to be the identity map. For any element $w \in 2^{<\mathbb{N}}$, we define f_w to be the corresponding composition of f_0 and f_1 , e.g., $f_{01} = f_0 \circ f_1$ and $f_{110} = f_1 \circ f_1 \circ f_0$. In other words, if we can write w as $u \frown v$, then $f_w = f_u \circ f_v$. The associativity of function composition ensures that this is well-defined.

One basic property of these maps is that for all $u \in 2^{<\mathbb{N}}$, the first letter of $f_u(a)$ is different for each $a \in \{x^{\pm 1}, y^{\pm 1}\}$, and the same is true for the last letter. This can be established through an easy induction on the length of u . If $u = e$, then this is immediate, and for $u = 0$ or $u = 1$, we quickly check that it holds. Now suppose that this is true for u . Then for $i \in \{0, 1\}$, consider $f_{u \frown i}(a) = f_u(f_i(a))$. We have already seen that the first and last letters of $f_i(a)$ are different for each a . By assumption, f_u takes the first and last letters of $f_i(a)$ to words with first and last letters different from those of $f_i(b)$ for any $b \neq a$ with $b \in \{x^{\pm 1}, y^{\pm 1}\}$, and this completes the induction.

With this established, a similar induction shows that every f_u takes freely reduced words to freely reduced words. In fact, every f_u takes cyclically reduced words to cyclically reduced words, since for $a, b \in \{x^{\pm 1}, y^{\pm 1}\}$, the first letter of $f_u(a)$ is the inverse of the last letter of $f_u(b)$ only if $b = a^{-1}$, by the uniqueness of the last letters.

If $\alpha \in \mathbb{F}_2$, then for any $w \in 2^{<\mathbb{N}}$, we can think of $f_w(\alpha)$ as a word on

$$\{f_w(a) \mid a \in \{x^{\pm 1}, y^{\pm 1}\}\}.$$

We refer to these special subwords as f_w -blocks. See figure 2.

Given $T \in Tr(2)$, we define

$$G_T = \langle x, y \mid \{(f_w(x))^{59}, (f_w(y))^{61} \mid w \in T\}, \{(f_w(x))^{67}, (f_w(y))^{71} \mid w \notin T\} \rangle.$$

The numbers in the exponents were chosen to be relatively prime and so that the relations satisfy small cancellation conditions, and they have no significance beyond that. We will eventually show that the map $T \mapsto G_T$ is a Borel reduction from \preceq_2^{free} to \preceq_{em} . To see that this is the case, we proceed by a series of lemmas.

$$\underbrace{\underbrace{x^5yx^5yx^5yx^5yx^5yy^5x}_{f_{00}(x)} \underbrace{x^5yx^5yx^5yx^5yx^5yy^5x}_{f_{00}(x)} \underbrace{x^5yx^5yx^5yx^5yx^5yy^5x}_{f_{00}(x)}}_{3 \text{ times}}$$

FIGURE 2. $f_{00}(x)^3$, with its f_{00} -blocks shown.

LEMMA 6.3. Let $u \in 2^{<\mathbb{N}}$, $a, b, c \in \{x^{\pm 1}, y^{\pm 1}\}$. Suppose $f_u(a)$ is a subword of $f_u(bc) = f_u(b)f_u(c)$. Then $f_u(a)$ does not contain letters from both $f_u(b)$ and $f_u(c)$. In other words, $f_u(a)$ must equal $f_u(b)$ or $f_u(c)$. It follows that if $\alpha, \beta \in \mathbb{F}_2$ are nontrivial and $f_u(\alpha)$ is a subword of $f_u(\beta)$, then α is a subword of β .

PROOF. We prove this inductively. It is easily checked to be true in the case that $u \in \{0, 1\}$. Now suppose that it is true for all u with $|u| < n$. Then if $u' = u \frown i$ with $|u'| = n$ and $i \in \{0, 1\}$ we may write $f_{u'}(a) = f_u(f_i(a))$ and $f_{u'}(bc) = f_u(f_i(bc))$. By assumption, the f_u -blocks in $f_{u'}(a)$ line up with the f_u -blocks in $f_{u'}(bc)$, and since $f_{u'}(a)$ is a subword of $f_{u'}(bc)$, it follows that $f_i(a)$ is a subword of $f_i(bc)$. This implies that $f_i(a)$ equals $f_i(b)$ or $f_i(c)$. Thus we find $f_{u'}(a)$ equals $f_{u'}(b)$ or $f_{u'}(c)$. \dashv

LEMMA 6.4. Let $w \in 2^{<\mathbb{N}}$, $\alpha, \beta \in \mathbb{F}_2$. If $f_w(\alpha)$ is a cyclic permutation of $f_w(\beta)$, then α is a cyclic permutation of β .

PROOF. We may write $\alpha = a_1 \dots a_n$ and $\beta = b_1 \dots b_n$ with $a_i, b_j \in \{x^{\pm 1}, y^{\pm 1}\}$, $1 \leq i, j \leq n$. Thus $f_w(\alpha) = f_w(a_1) \dots f_w(a_n)$ and $f_w(\beta) = f_w(b_1) \dots f_w(b_n)$. As $f_w(\alpha)$ is a cyclic permutation of $f_w(\beta)$, there is some $1 \leq k \leq n$ such that if we write $\alpha = \alpha_1 a_k \alpha_2$, then

$$f_w(b_1) \dots f_w(b_n) = g f_w(\alpha_2) f_w(\alpha_1) h,$$

where $hg = f_w(a_k)$. By Lemma 6.3, we must have $g = e$ or $h = e$. Suppose $g = e$. (The $h = e$ case is similar.) Then $f_w(\beta) = f_w(\alpha_2 \alpha_1 a_k)$, and so again by Lemma 6.3, $\beta = \alpha_2 \alpha_1 a_k$, which is a cyclic permutation of α . \dashv

LEMMA 6.5. Let $T \in Tr(2)$. If R_T denotes the symmetrization of the defining relations for G_T , then R_T satisfies the $C'(1/8)$ small cancellation condition.

PROOF. We only need to check the positive relations, since they satisfy the $C'(1/8)$ condition iff their inverses do, and there is no overlap between a positive word and a negative word.

We begin with an easy case. Suppose that $w \in 2^{<\mathbb{N}}$ and consider $f_w(x)^{n_x}$ and $f_w(y)^{n_y}$, where n_x and n_y denote the appropriate exponent, which depends on whether $w \in T$. As $f_w(x)$ and $f_w(y)$ do not start with the same letter, they do not have a common initial segment. We must also consider common initial segments of cyclic permutations of these two words, since we had to add the cyclic permutations of $f_w(x)^{n_x}$ and $f_w(y)^{n_y}$ to R_T to make sure that it was symmetrized.

A picture of sorts helps in the analysis. Before any sort of cyclic permutation, the two words can naturally be seen as being split into f_w -blocks. When a word is cyclically permuted a bit, the blocks at the beginning and end are truncated, as in figure 3. Now we cannot determine which word is a power of $f_w(x)$ and which is a power of $f_w(y)$ just by looking at the first letter of the words as before.

$$yx^5yx^5yx^5yy^5x \underbrace{x^5yx^5yx^5yx^5yx^5yy^5x}_{f_{00}(x)} \dots \underbrace{x^5yx^5yx^5yx^5yx^5yy^5x}_{f_{00}(x)} x^5yx^5$$

$n_x - 1$ times

FIGURE 3. $f_{00}(x)^{n_x}$ after being cyclically permuted.

Let r_1 be a cyclic permutation of $f_w(x)^{n_x}$ and r_2 be a cyclic permutation of $f_w(y)^{n_y}$ and let $M = \min\{|r_1|, |r_2|\}$. Suppose that $r_1 = st_1$ and $r_2 = st_2$ with s maximal. Permuting both r_1 and r_2 leftwards by $< |f_w(x)|$ letters, we get r_1^* and r_2^* , with $r_1^* = f_w(x)^{n_x}$. If the f_w -blocks in r_2^* also line up correctly, i.e., $r_2^* = f_w(y)^{n_y}$, then we know that r_1^* and r_2^* disagree at their first letter. Thus $|s| < |f_w(x)| < \frac{1}{8}M$ if $r_1 \neq r_2$.

Suppose that r_2^* is not in alignment, i.e., it is not $f_w(y)^{n_y}$. Then for r_1^* and r_2^* to agree for $\geq |f_w(x)|$ letters, $f_w(x)$ must be a subword of $f_w(y)^2$ containing letters from each copy of $f_w(y)$. But this is not possible, by Lemma 6.3. Thus r_1^* and r_2^* agree for $< |f_w(x)|$ letters if they are out of alignment and so $|s| < 2|f_w(x)| < \frac{1}{8}M$. In fact, the same reasoning shows that two different cyclic permutations of $f_w(x)^{n_x}$ or of $f_w(y)^{n_y}$ also overlap for less than $\frac{1}{8}M$ letters.

Now we consider the case when $w, v \in T$ are distinct and $a, b \in \{x, y\}$. Let r_1 be a cyclic permutation of $(f_w(a))^{n_a}$ and r_2 be a cyclic permutation of $(f_w(b))^{n_b}$, and let $M = \min\{|r_1|, |r_2|\}$. If $v = e$ and $w \neq e$, then we observe that no cyclic permutation of $(f_w(a))^{n_a}$ agrees with $(f_e(b))^{n_b} = b^{n_b}$ for more than six letters, which is less than $1/8$ of the length of either word. If w and v begin with different symbols, then one of r_1 and r_2 will be a cyclic permutation of a word in x^5y and y^5x , while the other will be a cyclic permutation of a word in x^2yxyx and y^2xyxy . Then the biggest possible common initial segment between r_1 and r_2 is xyx^3 or yx^3y , which is less than $1/8$ of the length of either word.

So we may assume that w and v start with the same symbols. Suppose that $w = u \hat{w}'$ and $v = u \hat{v}'$, with u maximal. Taking our cue from the notation for the greatest common divisor, we will write this as $u = (w, v)$. This should not cause confusion, as there are no ordered pairs (or greatest common divisors!) in what follows. Then up to some truncated bits at the beginning and end, r_1 and r_2 are both words in $f_u(x)$ and $f_u(y)$, and so we are in a situation very similar to our first case, except that now r_1 and r_2 contain a mix of $f_u(x)$ - and $f_u(y)$ -blocks, rather than just being conjugates of a power of one or the other. See figure 4 for a picture.

Suppose that r_1 and r_2 are both made up entirely of f_u -blocks, i.e., there are no truncated f_u -blocks at their beginning and end. Because $f_u(x)$ and $f_u(y)$ start with different letters, we see that if r_1 and r_2 agree on the beginning of a block, then they agree for the entire block. So the largest common initial segment s which r_1 and r_2 share is made up of entire f_u -blocks. We have $r_1 = f_u(\alpha)$, $r_2 = f_u(\beta)$, and $s = f_u(\gamma)$ with α, β, γ words in x and y , and so by Lemma 6.3, we find that γ is a common initial segment of α and β . Furthermore, $r_1 = f_u(\alpha)$ is a cyclic permutation of $f_w(a^{n_a}) = f_u(f_{w'}(a^{n_a}))$, and so by Lemma 6.4, α is a cyclic permutation of $f_{w'}(a^{n_a})$. Similarly, β is a cyclic permutation of $f_{v'}(b^{n_b})$.

So γ is a common initial segment of a cyclic permutation of $f_{w'}(a^{n_a})$ and a cyclic permutation of $f_{v'}(b^{n_b})$. This brings us back to the earlier cases. If w' and v' are

$$y \underbrace{x^5y}_{f_0(x)} \underbrace{x^5y}_{f_0(x)} \underbrace{x^5y}_{f_0(x)} \underbrace{y^5x}_{f_0(y)} \underbrace{x^5y}_{f_0(x)} \underbrace{x^5y}_{f_0(x)} \underbrace{x^5y}_{f_0(x)} \underbrace{x^5y}_{f_0(x)} \dots \underbrace{x^5y}_{f_0(x)} \underbrace{x^5y}_{f_0(x)} \underbrace{x^5y}_{f_0(x)} \underbrace{y^5x}_{f_0(y)} \underbrace{x^5y}_{f_0(x)} x^5$$

FIGURE 4. $f_{00}(x)^{n_x}$ after being cyclically permuted, with $f_0(x)$ - and $f_0(y)$ -blocks shown.

both nontrivial words, then they start with different symbols, which implies that $|\gamma| \leq 5$, and so s is made up of at most 5 f_u -blocks. On the other hand, r_1 and r_2 are made up of at least $6 \cdot \min\{|f_{w'}(a^{n_a})|, |f_{v'}(b^{n_b})|\}$ f_u -blocks, and so $|s| < \frac{1}{8}M$. If $w' = e$ and $v' \neq e$ or vice versa, then $|\gamma| \leq 6$ and we still find that $|s| < \frac{1}{8}M$. If $w' = e$ and $v' = e$, then γ is empty unless $a = b$, which implies that $r_1 = r_2$.

This leaves only the out of alignment cases to deal with. As before, we may permute r_1 and r_2 leftwards by $< |f_u(x)|$ letters to get r_1^* and r_2^* , with r_1^* a product of f_u -blocks. If r_2^* is not a product of f_u -blocks, then Lemma 6.3 tells us that r_1^* and r_2^* agree for $< |f_u(x)|$ letters, and so r_1 and r_2 agree for $< 2|f_u(x)|$ letters, which is $< 1/8$ of the length of each word. If r_2^* is a product of f_u -blocks, then we are in the previous case, and we have seen that either $r_1^* = r_2^*$ or the corresponding common initial segment between them consists of at most 6 f_u -blocks. This implies that $|s| < 7|f_u(a)| < \frac{1}{8}M$. We have finally exhausted all of the cases and have shown that R_T satisfies the $C'(1/8)$ condition, as desired. \dashv

LEMMA 6.6. *If $T, T' \in Tr(2)$ and there exists $w \in 2^{<\mathbb{N}}$ such that $T = T'_w$, then $G_T \hookrightarrow G_{T'}$.*

PROOF. This is obvious if $w = e$, and so we may assume that w is nontrivial. It is easy to see that f_w , viewed as a map from G_T to $G_{T'}$, is a homomorphism, since it will take defining relations in G_T to defining relations in $G_{T'}$. In more detail,

$$\begin{aligned} f_v(x)^{59}, f_v(y)^{61} \in R_T &\iff v \in T \\ &\iff w \hat{\ } v \in T' \\ &\iff f_{w \hat{\ } v}(x)^{59} = f_w(f_v(x)^{59}), \\ &\quad f_{w \hat{\ } v}(y)^{61} = f_w(f_v(y)^{61}) \in R_{T'} \end{aligned}$$

and similar equivalences hold for $f_v(x)^{67}$ and $f_v(y)^{71}$. It remains to show that f_w is an embedding.

We still need to show that nontrivial elements of G_T do not map to the identity in $G_{T'}$. As in section 5, we will show that if $\alpha \in \mathbb{F}_2$ is such that $f_w(\alpha)$ contains more than $1/2$ of a relation in $R_{T'}$, then α contains more than $1/2$ of a relation in R_T , which easily implies the result.

Suppose that $\alpha \in \mathbb{F}_2$ and that $f_w(\alpha) = 1$ in $G_{T'}$. Then $f_w(\alpha)$ contains more than half of a relation $r \in R_{T'}$. We know that r is a cyclic permutation of some $f_v(a^{n_a})$, where $v \in 2^{<\mathbb{N}}$, $a \in \{x^{\pm 1}, y^{\pm 1}\}$, and $n_a \in \{59, 61, 67, 71\}$. Let $u = (w, v)$, so that $w = u \hat{\ } w'$ and $v = u \hat{\ } v'$. Then $f_w(\alpha) = f_u(f_{w'}(\alpha))$, and r is a cyclic permutation of $f_u(f_{v'}(a^{n_a}))$. By assumption, the subword of r that both words contain must be big enough to contain an entire f_u -block. Thus Lemma 6.3 tells us that the f_u -blocks of r and $f_w(\alpha)$ must line up. The f_u -blocks are uniquely identified by their first or last letters, so once $f_w(\alpha)$ and r agree for part of an f_u -block, they agree on the whole thing, unless r begins and ends with a truncated f_u -block. In this case, cyclically permuting r until it is made up of f_u -blocks will “complete” the f_u -block at one end of r . This new word is also a relation which agrees with $f_w(\alpha)$ for at least as long as r did, since either only one end of r was in $f_w(\alpha)$ and cyclically permuting increases the length of the word the two agree on, or r was a subword of $f_w(\alpha)$ and this cyclic permutation is also a subword of $f_w(\alpha)$.

Thus we may assume that $r = f_u(\omega)$ for some $\omega \in \mathbb{F}_2$, and that r and $f_w(\alpha)$ share a subword of the form $f_u(\gamma)$, where $\gamma \in \mathbb{F}_2$. By Lemma 6.4, we know that ω is a cyclic permutation of $f_{v'}(a^{n_a})$. Then γ is a subword of a cyclic permutation of $f_{v'}(a^{n_a})$ and a subword of $f_{w'}(\alpha)$. If w' and v' are both nontrivial, then they begin with different symbols, and so $|\gamma| \leq 5$. But then

$$\begin{aligned} \frac{|f_u(\gamma)|}{|f_{v'}(a^{n_a})|} &= \frac{|\gamma|}{|f_{v'}(a^{n_a})|} \\ &\leq \frac{5}{|f_{v'}(a^{n_a})|} \\ &< 1/2, \end{aligned}$$

which is a contradiction. If $v' = e$ but $w' \neq e$, then virtually the same inequalities hold, since $f'_w(\alpha)$ does not contain any letter to a power greater than 6, and again we get a contradiction. Thus $w' = e$, meaning $w = u$, and so α contains $> 1/2$ of a cyclic permutation of $f_{v'}(a^{n_a})$. Further, since $w \hat{\ } v' \in T' \Leftrightarrow v' \in T$, it follows that $f_{v'}(a^{n_a}) \in R_T$. Thus if $f_w(\alpha) = 1$ in $G_{T'}$, then α contains $> 1/2$ of a word in R_T , as desired. \dashv

The proof of the converse of Lemma 6.6 will depend on the following two lemmas.

LEMMA 6.7. *Suppose $\alpha, \beta \in \mathbb{F}_2$ are cyclically reduced, and $w, v \in 2^{<\mathbb{N}}$. If r_1 is a cyclic permutation of $f_w(\alpha)$ and r_2 is both a cyclic permutation of $f_v(\beta)$ and a subword of r_1 , then $v \subset w$ or $w \subset v$.*

Moreover, if $w = v \hat{\ } w'$ then a cyclic permutation of $f_{w'}(\alpha)$ contains a cyclic permutation of β , and if $v = w \hat{\ } v'$, then a cyclic permutation of α contains a cyclic permutation of $f_{v'}(\beta)$.

PROOF. The result is trivial if $w = e$ or $v = e$, so we may assume that w and v are nontrivial. Let $u = (w, v)$, with $w = u \hat{\ } w'$ and $v = u \hat{\ } v'$. If $u = e$, then w and v begin with different symbols, which is impossible, since $|r_1|, |r_2| > 5$, the length of the longest possible agreement between r_1 and r_2 . So u is nontrivial, and r_1 is a cyclic permutation of $f_u(f_{w'}(\alpha))$, while r_2 is a cyclic permutation of $f_u(f_{v'}(\beta))$. The f_u -blocks of each word must line up, by Lemma 6.3. Further, since r_2 is a subword of r_1 , any truncated bits of f_u -blocks at the ends of r_2 are duplicated in r_1 . So we can permute r_1 and r_2 the same amount to get $r_1^* = f_u(\gamma)$ and $r_2^* = f_u(\omega)$, words composed entirely of f_u -blocks, with r_2^* contained in r_1^* . By Lemma 6.4, we know γ is a cyclic permutation of $f_{w'}(\alpha)$ and ω is a cyclic permutation of $f_{v'}(\beta)$, and that ω is a subword of γ .

If w' and v' are nontrivial, then they start with different symbols, and as above we reach a contradiction. Thus either $w' = e$, and so $w \subset v$ and a cyclic permutation of α contains a cyclic permutation of $f_{v'}(\beta)$, or $v' = e$, so $v \subset w$ and a cyclic permutation of $f_{w'}(\alpha)$ contains a cyclic permutation of β . \dashv

LEMMA 6.8. *Suppose that $t, u, v \in 2^{<\mathbb{N}}$, and some cyclic permutation of $f_t(x^k)$ is a product of a cyclic permutation of $f_u(x^l)$ and a cyclic permutation of $f_v(y^m)$, with $k, l, m \in \mathbb{Z} \setminus \{0\}$ all having the same sign. Then $u = v$, $t = u \hat{\ } 0$, $k = m = \pm 1$, $l = 5k$, and $f_t(x^k) = f_{u \hat{\ } 0}(x^{\pm 1})$ is either*

$$f_u(x^5)f_u(y)$$

or

$$f_u(y^{-1})f_u(x^{-5}).$$

PROOF. By Lemma 6.7, either $t \subset u$ or $u \subset t$. If $t \subset u$ and $u = t \frown u'$, then by the previous lemma, we find that a cyclic permutation of x^k contains a cyclic permutation of $f_{u'}(x^l)$. This is impossible unless $u' = e$. So we may assume that $u \subset t$ and $t = u \frown t'$. Similarly we find that $v \subset t$ and $t = v \frown t''$. It follows that $u \subset v$ or $v \subset u$.

Suppose that $u \subset v$ and $v = u \frown v'$. We know that the f_u -blocks in the cyclic permutation of $f_u(x^l)$ and the cyclic permutation of $f_u(f_{v'}(y^m))$ must line up with those in the cyclic permutation of $f_t(x^k)$. This means, in particular, that a truncated f_u -block at the end of the cyclic permutation of $f_u(x^l)$ must be completed by a truncated f_u -block at the beginning of the cyclic permutation of $f_u(f_{v'}(y^m))$, and vice versa. So we can assume that the cyclic permutations we are considering are made up of complete f_u -blocks. Then by Lemma 6.3 we obtain that a cyclic permutation of $f_{t'}(x^k)$ is a product of x^l and a cyclic permutation of $f_{v'}(y^m)$. Now, $f_{t'}(x^k)$ is composed of $f_{v'}$ -blocks, which must line up with those in the cyclic permutation of $f_{v'}(y^m)$. So it must be the case that x^l is a cyclic permutation of $f_{v'}$ -blocks. But this can only happen if $v' = e$, meaning $u = v$. Similar reasoning applies if $v \subset u$. Thus $u = v$.

It is not possible for a truncated $f_u(x)$ -block to be completed by a truncated $f_u(y)$ -block, or vice versa, and so we must have that the cyclic permutation of $f_t(x^k)$ we started with is either $f_u(x^l)f_u(y^m)$ or $f_u(y^m)f_u(x^l)$.

It follows that $x^l y^m$ is a cyclic permutation of $f_{t'}(x^k)$. This can only happen if $t' = 0$ and k, l, m are as in the statement of the lemma, since if $t' = e$, then x^k contains no occurrences of y , and if $t' \neq 0$ is nontrivial, then $f_{t'}(x^k)$ must contain at least two distinct blocks of x 's and y 's. ⊖

We now take up the converse of Lemma 6.6, which will complete the proof of Theorem 6.2.

LEMMA 6.9. *If $T, T' \in Tr(2)$ and $G_T \hookrightarrow G_{T'}$, then $\exists w \in 2^{<\mathbb{N}}$ such that $T = T'_w$.*

PROOF. Suppose that $\theta: G_T \rightarrow G_{T'}$ is a monomorphism. Our main goal is to prove that θ must actually be f_w for some $w \in 2^{<\mathbb{N}}$, up to an inner automorphism of $G_{T'}$. Once we know this, it is easy to recover the relations in each group, and thus to show that $T = T'_w$.

Since $x = f_e(x)$, x has some finite order n_x in G_T . Then $\theta(x)$ has order n_x , and so by Theorem 5.6, $\theta(x)$ must be conjugate to a power of some $f_w(x)$, where $w \in T' \Leftrightarrow e \in T$. If we follow θ by an inner automorphism of $G_{T'}$, we may assume that $\theta(x) = (f_w(x))^\delta$ for some nonzero integer δ .

Similarly, $\theta(y)$ is conjugate to a power of some $f_v(y)$ with $v \in T' \Leftrightarrow e \in T$. We find that $\theta(y) = u(f_v(y))^y u^{-1}$. We can assume that u does not contain more than half of an element of $R_{T'}$. We may also follow θ by the inner automorphism corresponding to $f_w(x)$ as necessary to ensure that u does not begin with a power of $f_w(x)$, and this will not change the value of $\theta(x)$. After freely reducing we get that $\theta(y) = u'ru'^{-1}$, where r is a cyclic permutation of $(f_v(y))^y$. To see this, suppose that $u = \alpha\beta^{-1}$, where $(f_v(y))^y = \beta\mu$ and β is the longest subword of u for which this is true. Then $u(f_v(y))^y u^{-1} = \alpha\beta^{-1}\beta\mu\beta\alpha^{-1} = \alpha\mu\beta\alpha^{-1}$. Then $u' = \alpha$ and $r = \mu\beta$. A similar argument works if $(f_v(y))^y$ cancels with u^{-1} . For example,

if we had $\theta(y) = xy^{-1}f_0(y)yx^{-1} = xy^{-1}(y^5x)yx^{-1}$, then after freely reducing we would find $\theta(y) = xy^4xyx^{-1} = xrx^{-1}$, where $r = y^4xy$.

We now proceed much as in the proof of Theorem 5.1. Let m_x be the order of $f_0(x)$ in G_T . Since θ is an embedding, it must take $f_0(x)$ to a torsion element of order m_x . We know that

$$\begin{aligned} \theta(f_0(x)) &= \theta(x^5y) = (f_w(x))^{5\delta}u'ru'^{-1} \\ &= (f_w(x))^{\delta'}u'ru'^{-1}, \end{aligned}$$

where $|\delta'| < \lfloor \frac{m_x}{2} \rfloor$, since we may eliminate any copies of $f_w(x)^{n_x}$ in $\theta(x^5y)$, and replace $f_w(x)^k$ with $f_w(x)^{k-n_x}$ for $k < n_x$. Note that, as written, this word may not be freely reduced, so we cannot necessarily use Theorem 5.5 yet. Let $z = \theta(f_0(x))$.

Suppose that z is cyclically reduced as written and that $u' \neq 1$. Then z is a cyclically reduced word with finite order in $G_{T'}$, which contains a mixture of positive and negative letters, which is impossible by Theorem 5.6, since the words in $R_{T'}$ are either entirely positive or entirely negative. So either $u' = 1$ or else z is not cyclically reduced.

First suppose that $u' = 1$, so that $z = (f_w(x))^{\delta'}r$ and r is cyclically reduced and is a cyclic permutation of $(f_v(y))^\gamma$. If δ' and γ have the same sign, then z is cyclically reduced as written. By Theorem 5.6, z is a cyclic permutation of some $f_t(x^k)$. Also, it is a product of a cyclic permutation of $f_w(x^k)$ and $f_v(y^\gamma)$, so by Lemma 6.8, $t = w \frown 0$, $v = w$, and $k = \pm 1$. Thus z is a cyclic permutation of $f_w(x^5)f_w(y)$ or $f_w(y^{-1})f_w(x^{-5})$. Because z begins with an $f_w(x^{\pm 1})$ block and cannot end with one, it follows that $\delta' = 5$, which implies that $\delta = 1$, and so $\theta(x) = f_w(x)^{\pm 1}$ and $\theta(y) = f_w(y)^{\pm 1}$.

Suppose that δ' and γ have opposite signs. There is a (possibly trivial) inner automorphism ψ such that $\psi(z) = sr'$, where s is a cyclic permutation of $f_w(x)^{\delta'}$, and r' is a cyclic permutation of $f_v(y)^\gamma$, and freely reducing sr' will leave a cyclically reduced word. Let $\psi(z) = z'$ with z' the result of freely reducing sr' . Since z' is a torsion element, it must be a cyclic permutation of some $f_t(x^k)$, and so its letters must all have the same sign.

Suppose z' and s have letters of the same sign. Then $z'r'^{-1}$ is cyclically reduced and so we have a cyclic permutation of $f_w(x)^{\delta'}$ written as a product of a cyclic permutation of $f_t(x)^k$ and a cyclic permutation of $f_v(y)^\gamma$ with k, γ having the same sign. By Lemma 6.8, we get that $\theta(x) = f_{v \frown 0}(x^{\pm 1}) = f_v((x^5y)^{\pm 1})$, and $\theta(y) = f_v(y^{\mp 1})$. But then either $\theta(xy) = f_v(x^5)$ or $\theta(xy) = f_v(y^{-1}x^{-5}y) = f_v(y)^{-1}f_v(x^{-5})f_v(y)$. Both of these are torsion elements, but xy is not a torsion element in G_T , which contradicts the fact that θ is an embedding. So suppose that z' and r' have letters of the same sign. Then $z's^{-1}$ is cyclically reduced and so we have a cyclic permutation of $f_v(y)^\gamma$ written as a product of a cyclic permutation of $f_t(x)^k$ and a cyclic permutation of $f_w(x)^{\delta'}$ with k, δ' having the same sign. Arguing as in the proof of Lemma 6.8, we find that $w = t$ and that $w \subset v$. Let $v = w \frown v'$. We obtain that $f_{v'}(y^\gamma) = x^{-\delta'+k}$, which is absurd.

We still must address the case where $u' \neq 1$ and z is not cyclically reduced. This can happen for two reasons. It may be that u' and $f_w(x)$ begin in the same way. We know that u' does not begin with an entire copy of $f_w(x)$, and we have assumed that u' does not further cancel with r , so there is an inner automorphism ψ such that

$\psi(\theta(f_0(x))) = su''ru''^{-1}$, a cyclically reduced word with s a cyclic permutation of $(f_w(x))^{\delta'}$. If $u'' \neq 1$, then $\psi(\theta(f_0(x)))$ contains positive and negative letters, and we have already seen that this is impossible. Thus we must have that $\psi(\theta(f_0(x))) = sr$, and as before we see that $\psi(\theta(x)) = f_w(x)^{\pm 1}$ and $\psi(\theta(y)) = f_w(y)^{\pm 1}$.

The other possibility is that u' cancels with the end of $f_w(x)$. It cannot cancel with the whole of $f_w(x)$, and so again after following θ by an appropriate inner automorphism ψ we get that $\psi(\theta(f_0(x))) = su''ru''^{-1}$, a cyclically reduced word with s a cyclic permutation of $(f_w(x))^{\delta'}$. This case is treated exactly as in the previous paragraph.

So we have shown that, up to an inner automorphism, $\theta(x) = f_w(x)^{\pm 1}$ and $\theta(y) = f_w(y)^{\pm 1}$ with the signs matching. If $\theta(x) = f_w(x)$ and $\theta(y) = f_w(y)$, then $\theta = f_w$, and hence

$$\begin{aligned} u \in T &\Leftrightarrow f_u(x^{53}) \in R_T \\ &\Leftrightarrow f_w(f_u(x^{53})) \in R_{T'} \\ &\Leftrightarrow w \frown u \in T'. \end{aligned}$$

Thus $T = T'_w$, as desired. Thus it only remains is to eliminate the undesirable case when $\theta(x) = f_w(x^{-1})$ and $\theta(y) = f_w(y^{-1})$. In this case we have that

$$\begin{aligned} \theta(f_{00}(x)) &= \theta(x^5yx^5yx^5yx^5yx^5y^5x) \\ &= f_w(x^{-5}y^{-1}x^{-5}y^{-1}x^{-5}y^{-1}x^{-5}y^{-1}x^{-5}y^{-1}x^{-5}y^{-1}y^{-5}x^{-1}) \\ &= f_w((xy^5yx^5yx^5yx^5yx^5y^5x)^{-1}). \end{aligned}$$

We will show this is not a torsion element in $G_{T'}$. Since $f_{00}(x)$ is a torsion element in G_T , this implies that θ is not a homomorphism, which is a contradiction.

Let $\alpha = xy^5yx^5yx^5yx^5yx^5y^5x$. It is easy to see that $f_{00}(x)$ is the only torsion element that has length 36 and that contains 26 x s. However, α is not a cyclic permutation of $f_{00}(x)$. It follows that $f_w(\alpha)$ (and hence $\theta(f_{00}(x))$) is not a torsion element, since otherwise it would have to be a cyclic permutation of some $f_t(x^k)$ with $t \in T'$. Thus by Lemma 6.7, either $w \subset t$ or $t \subset w$. Suppose that $w \subset t$ and $t = w \frown t'$. Then α must be a cyclic permutation of $f_{t'}(x^k)$, which we have already seen is impossible. If $t \subset w$ and $w = t \frown w'$, then x^k must be a cyclic permutation of $f_{w'}(\alpha)$, which is also impossible. Thus we reach a contradiction, eliminating the final undesirable case. ⊣

PROOF OF THEOREM 6.2. Lemmas 6.6 and 6.9 establish that the map $T \mapsto G_T$ is a Borel reduction from \preceq_2^{tree} to \preceq_{em} . ⊣

COROLLARY 6.10. \equiv_{em} is a universal countable Borel equivalence relation.

It may have been possible to prove this corollary without any reference to quasi-orders, by reducing some known universal countable Borel equivalence relation to \equiv_{em} . However, it seems that the most natural and direct route to this result is through Theorem 6.2. A closer look at the above proof also leads to the following result, which tells us that the biembeddability relation on the groups constructed above is much more complicated than the isomorphism relation on these groups.

COROLLARY 6.11. With the above notation, $G_T \cong G_S \Leftrightarrow T = S$.

PROOF. One direction is trivial. For the other, suppose that $T, S \in Tr(2)$ are such that $G_T \cong G_S$, via $\phi: G_T \rightarrow G_S$. Then, in particular, ϕ is an embedding, and by

the proof of Lemma 6.9, there exists $w \in 2^{<\mathbb{N}}$ such that $T = S_w$. Furthermore, after adjusting by an inner automorphism of G_S if necessary, we can suppose that $\phi = f_w$. We will show that if $w \neq e$, then $f_w: G_T \rightarrow G_S$ is not surjective. Hence $w = e$ and $G_T = G_S$.

Suppose that $w \in 2^{<\mathbb{N}}$ is nontrivial and that $f_w: G_T \rightarrow G_S$ is surjective. Then there is some word $\alpha \in \mathbb{F}_2$, which we may assume does not contain more than $1/2$ of a relation in R_T , such that $f_w(\alpha) = x$ in G_S , where x is one of the generators of G_S . This means that $f_w(\alpha)x^{-1} = 1$ in G_S . By the proof of Lemma 6.6, we know that $f_w(\alpha)$ does not contain more than $1/2$ of a relation in G_S . By Theorem 5.5, for $f_w(\alpha)x^{-1}$ to represent the identity in G_S , it must contain more than $(1 - 3/8) = 5/8$ of a relation in R_T . But $f_w(\alpha)x^{-1}$ has at most one more letter in common with a relation than $f_w(\alpha)$ does. Since $f_w(\alpha)$ contains less than $1/2$ of a relation in R_T , $f_w(\alpha)x^{-1}$ contains less than $5/8$ of a relation in R_T . This is a contradiction. \dashv

§7. Quasi-orders that symmetrize to universal countable Borel equivalence relations. We have seen that symmetrizing universal countable Borel quasi-orders produces universal countable Borel equivalence relations. It would be desirable to be able to strengthen this and say that symmetrizing a non-universal countable Borel quasi-order produces a nonuniversal countable Borel equivalence relation. However, it is easy to see that this is not the case. Let E be a universal countable Borel equivalence relation. Then E is not universal as a quasi-order, as symmetry is preserved downwards under \leq_B , but obviously E symmetrizes to a universal equivalence relation, i.e., E itself.

If this were the extent of the phenomenon, then it would still be possible to use negative results about the universality of a given countable Borel quasi-order in order to prove negative results about the universality of its associated equivalence relation, so long as the quasi-order was asymmetric somewhere. Unfortunately, things are as bad as they could be.

THEOREM 7.1. *There are 2^{\aleph_0} countable Borel quasi-orders Q , distinct up to Borel bireducibility, for which E_Q is a universal countable Borel equivalence relation.*

PROOF. Recall that Adams and Kechris showed in [1] that there are 2^{\aleph_0} countable Borel equivalence relations up to Borel bireducibility. The equivalence relations they produced are all defined on different spaces; we will write X_E for the space on which E is defined. Further, these equivalence relations have all equivalence classes of size at least 2, a minor technical point that will be used later.

Given a countable Borel equivalence relation E on a standard Borel space X with a Borel linear order \leq , define $E(\leq)$ to be $E \cap \leq$. We see that $E(\leq)$ is an asymmetric countable Borel quasi-order, unless E is just equality on X , written $\Delta(X)$. Note that

$$E(\leq) \leq_B F(\leq) \implies E \leq_B F$$

and so if $E(\leq)$ and $F(\leq)$ are Borel bireducible, then so are E and F . Thus by the result of Adams and Kechris we find that there are 2^{\aleph_0} quasi-orders of the form $E(\leq)$ up to Borel bireducibility.

Now we define the family of quasi-orders $\{E(\leq) \vee E_\infty\}$, where E ranges over the equivalence relations from the proof of Adams and Kechris and \vee denotes the

disjoint union of two equivalence relations. Now suppose that

$$E(\leq) \vee E_\infty \leq_B F(\leq) \vee E_\infty.$$

Then $E(\leq) \leq_B F(\leq)$, since $E(\leq)$ is completely asymmetric, and so any two $E(\leq)$ -comparable elements in X_E must map to X_F . Since the equivalence classes of E are all of size at least 2, every element of X_E is $E(\leq)$ -comparable with some other element of X_E , and so every element of X_E is mapped to X_F .

Thus there are 2^{\aleph_0} countable Borel quasi-orders of the form $E(\leq) \vee E_\infty$. Each of these symmetrizes to $\Delta(X_E) \vee E_\infty$, which is universal. \dashv

Most of the reducibilities from computability theory, such as Turing reducibility \leq_T or 1-reducibility \leq_1 , are countable Borel quasi-orders. The equivalence relations associated with them have been the subject of a great deal of work in descriptive set theory; for example, see [4]. It remains open whether \equiv_T or \equiv_1 are universal countable Borel equivalence relations. Thus the following question naturally arises.

OPEN PROBLEM 7.2. *Are any of \leq_1, \leq_T , etc. universal countable Borel quasi-orders?*

In light of theorem 7.1, showing that \leq_T or \leq_1 are not universal countable Borel quasi-orders would not settle the question of whether \equiv_T or \equiv_1 are universal, although it may be taken as “evidence” that the answer is no.

Acknowledgements. This research was partially supported by NSF Grant DMS 1101597. The results in this paper (except for Theorem 7.1) are from my thesis, which was done under the direction of Simon Thomas, whose help was invaluable. I would also like to thank Arthur Apter, Justin Bush, Gregory Cherlin, David Duncan, and Charles Weibel for many helpful mathematical discussions.

REFERENCES

- [1] SCOT ADAMS and ALEXANDER S. KECHRIS, *Linear algebraic groups and countable Borel equivalence relations*. *Journal of American Mathematical Society*, vol. 13 (2000), no. 4, pp. 909–943.
- [2] CHRISTOPHE CHAMPETIER, *L'espace des groupes de type fini*. *Topology*, vol. 39 (2000), no. 4, pp. 657–680.
- [3] R. DOUGHERTY, S. JACKSON, and A. S. KECHRIS, *The structure of hyperfinite Borel equivalence relations*. *Transactions of the American Mathematical Society*, vol. 341 (1994), no. 1, pp. 193–225.
- [4] RANDALL DOUGHERTY and ALEXANDER S. KECHRIS, *How many Turing degrees are there? Computability theory and its applications* (Boulder, CO, 1999), Contemporary Mathematics, vol. 257, *American Mathematical Society*, Providence, RI, 2000, pp. 83–94.
- [5] JACOB FELDMAN and CALVIN C. MOORE, *Ergodic equivalence relations, cohomology, and von Neumann algebras. I*. *Transactions of the American Mathematical Society*, vol. 234 (1977), no. 2, pp. 289–324.
- [6] SY-DAVID FRIEDMAN and LUCA MOTTO ROS, *Analytic equivalence relations and bi-embeddability*, this JOURNAL, vol. 76 (2011), no. 1, pp. 243–266.
- [7] SU GAO, *Coding subset shift by subgroup conjugacy*. *Bulletin of the London Mathematical Society*, vol. 32 (2000), no. 6, pp. 653–657.
- [8] ALEXANDER S. KECHRIS, *Classical descriptive set theory*. *Graduate Texts in Mathematics*, vol. 156, Springer-Verlag, New York, 1995.
- [9] ALAIN LOUVEAU and CHRISTIAN ROSENDAL, *Complete analytic equivalence relations*. *Transactions of the American Mathematical Society*, vol. 357 (2005), no. 12, pp. 4839–4866.
- [10] ROGER C. LYNDON and PAUL E. SCHUPP, *Combinatorial group theory*. *Classics in Mathematics*, Springer-Verlag, Berlin, 2001. Reprint of the 1977 edition.

[11] ALAN H. MEKLER, *Stability of nilpotent groups of class 2 and prime exponent*. this JOURNAL, vol. 46 (1981), no. 4, pp. 781–788.

[12] SIMON THOMAS and BOBAN VELICKOVIC, *On the complexity of the isomorphism relation for finitely generated groups*. *Journal of Algebra*, vol. 217 (1999), no. 1, pp. 352–373.

DEPARTMENT OF MATHEMATICS
CALIFORNIA INSTITUTE OF TECHNOLOGY
PASADENA, CA 91125, USA
E-mail: jaywill@caltech.edu