

Headaches, shoulder pains, and upper back pain also decreased. Patients reported high levels of benefit from the acupuncture and encouraged other patients to continue to come and try the acupuncture. Usually, the needles could be placed within the first third of the group.

Conclusions: Acupuncture improved anxiety ratings for people in group psychotherapy for anxiety over group alone, though the possibility of a placebo effect cannot be eliminated. Patients chose acupuncture, which could also present a potential bias.

Disclosure of Interest: None Declared

O0003

Using Virtual Reality Assisted Therapy to Reduce Cognitive Test Anxiety and Dysfunctional Metacognitions

F. Obuća^{1*}, P. Ünal-Aydın¹ and O. Aydın¹

¹Psychology, International University of Sarajevo, Sarajevo, Bosnia and Herzegovina

*Corresponding author.

doi: 10.1192/j.eurpsy.2024.884

Introduction: Cognitive test anxiety and dysfunctional metacognitions can significantly impact an individual's performance and overall mental health. However, the effectiveness of various treatment strategies, including Virtual Reality (VR) therapy, is yet to be fully explored.

Objectives: This study aimed to examine the effectiveness of VR therapy in reducing cognitive test anxiety and dysfunctional metacognitions in adults.

Methods: A total of 64 participants were enrolled in the study, with 40 in the treatment group and 24 in the control group. Data were collected using the Metacognition Questionnaire-30, Cognitive Test Anxiety Scale, and a sociodemographic questionnaire. Paired samples t-tests were used to compare pretest and posttest scores, while independent samples t-tests were used to compare the means between the groups.

Results: The findings suggest that the treatment group experienced a significant reduction in cognitive test anxiety and negative metacognition scores following VR therapy. No significant changes were observed in the control group, and there were no significant differences in pretest scores between the treatment and control groups.

Conclusions: The study indicates that VR therapy may be an effective treatment strategy for reducing cognitive test anxiety and dysfunctional metacognitions. Further research is recommended to validate these findings and explore the potential of VR therapy in treating other psychological disorders.

Disclosure of Interest: None Declared

EPP0553

Perfectionism, self-efficacy and mindfulness as predictors of test anxiety among university students

S. Tvrtkovic

Psychology, International University of Sarajevo, Sarajevo, Bosnia and Herzegovina

doi: 10.1192/j.eurpsy.2024.885

Introduction: Test anxiety includes subjective experience of intense physiological, cognitive and/or behavioral symptoms

during test-taking situations such as pacing, headaches, excessive feelings of fear, anger, troubles concentrating, sudden forgetfulness and negative self-talk. Especially students who are striving for flawlessness, have overly critical self-evaluations and beliefs that other expect perfection are sensitive to experiencing these feelings. On the contrary, individuals who believe in their ability, are present in the moment and are open to experiences tend to be more resilient to stressors and anxiety symptoms.

Objectives: The aim of this study was to investigate perfectionism, self-efficacy, and mindfulness as predictors of test-anxiety among undergraduate and postgraduate students of different study fields.

Methods: 525 undergraduate and postgraduate students from the fields of Natural, Medical Sciences and Engineering, Social Sciences, Humanities and Art, and Economics, Business and Administration Studies participated in the study. A sociodemographic form, the Test Anxiety Inventory (TAI), 15-Item Five Facet Mindfulness Questionnaire (FFMQ-15), Scale of General Self-efficacy (GSES) and Frost Multidimensional Perfectionism Scale-Brief (FMPS-Brief) were used. Descriptive statistics were used to show the sociodemographics of our sample, while correlational analyses were performed to assess the associations between the variables. To further validate the findings, multiple linear regression analyses were performed.

Results: Higher perfectionistic evaluative concerns and lower self-efficacy showed associations with test-anxiety and were proven as predictors among undergraduate and postgraduate students. In addition, being younger and female in postgraduates, and having perfectionistic strivings and being Mindful-Observe in undergraduate students proved to be significant predictors of test-anxiety.

Conclusions: The found associations of perfectionism and self-efficacy, as well as its predictor roles further validate the information we have in literature, but widen the sample to postgraduate students and various study fields which help us generalize the findings more. What adds the most value in theoretical and practical aspects is the findings associated with mindfulness. Mindfulness techniques are very popular intervention methods for anxiety related symptoms, and the positive relationship of Mindful-Observe with test anxiety opens new viewpoints to mindfulness-based interventions. Particularly focusing on Mindful-Observe while treating test-anxiety may yield better outcomes in alleviation of symptoms.

Disclosure of Interest: None Declared

Bipolar Disorders

EPV0093

Lithium withdrawal and relapse in bipolar disorder when kidney function deteriorates

C. De Andrés-Lobo, M. D. C. Vallecillo Adame, T. Jiménez Aparicio, A. Rodríguez Campos*, N. Navarro Barriga, M. J. Mateos Sexmero, B. Rodríguez Rodríguez, M. Fernández Lozano, M. Calvo Valcárcel, M. Andreo Vidal, P. Martínez Gimeno, M. P. Pando Fernández, L. Rojas Vázquez, G. Lorenzo Chapatte, M. Ríos Vaquero and A. Monllor Lazarraga
Psychiatry, HCUV, Valladolid, Spain

*Corresponding author.

doi: 10.1192/j.eurpsy.2024.886