

Range expansion of the whitenose shark, *Nasolamia velox*, and migratory movements to the oceanic Revillagigedo Archipelago (west Mexico)

FRIDA LARA-LIZARDI^{1,2}, MAURICIO HOYOS-PADILLA^{2,3}, JAMES T. KETCHUM^{2,4}
AND FELIPE GALVÁN-MAGAÑA¹

¹Instituto Politécnico Nacional, Centro Interdisciplinario de Ciencias Marinas, Av. IPN s/n. C.P. 23096. La Paz, B.C.S, Mexico,

²Pelagios-Kakunjá A. C. 1540 Sinaloa, C.P. 23070, La Paz, B.C.S., Mexico, ³Fins Attached, 19675 Still Glen Way, Colorado Springs, CO 80908, USA, ⁴Centro de Investigaciones Biológicas del Noroeste, Playa Palo de Santa Rita Sur, 23096 La Paz, B.C.S, Mexico

*Current literature considers that *Nasolamia velox* has a limited distribution along the coastline of the Eastern Pacific with sporadic sightings in the Galapagos Archipelago. This study provides evidence of the occurrence of this species at the Revillagigedo Archipelago (18°99'186"N 112°08'44"W), Mexico, using acoustic telemetry and videos taken from 2014 to 2016. We report here movements from a coastal location (National Park Cabo Pulmo) to a group of oceanic islands (Revillagigedo Archipelago) by one single individual, supporting the idea of the potential connectivity of sharks between the Gulf of California and the Revillagigedo Archipelago. This report extends the known distribution of *N. velox* to 400 km off the mainland coast of the Americas, thereby increasing the knowledge of the distribution of a species commonly reported in fishery landings of the Eastern Pacific.*

Keywords: Revillagigedo Archipelago, acoustic telemetry, visual census, range extension, shark distribution, connectivity

Submitted 26 May 2016; accepted 19 January 2017; first published online 27 February 2017

INTRODUCTION

The whitenose shark (*Nasolamia velox*) is an endemic species of the Eastern Tropical Pacific (ETP), however very little is known about the biology and distribution of this species. The whitenose shark was previously known as *Carcharhinus velox*. Compagno & Garrick (1983) determined that the species differs externally from all other carcharhinids by very wide, transversely oriented nostrils placed close together so that the internarial width is only slightly greater than the width of each nostril, hence, they proposed a new genus *Nasolamia*. Recent molecular phylogenetic studies suggest that this species might be closely related to *Carcharhinus acronotus* (Naylor *et al.*, 2012).

Nasolamia velox (Gilbert, 1898) is a medium-sized shark that reaches a maximum of 165 cm total length (TL) (Ruiz *et al.*, 2009). The species is viviparous, with a yolk-sac placenta (Ruiz-Alvarado & Mijangos-López, 1999; Compagno, 2001). It feeds mainly on anchovies and crabs. The size at birth is about 53 cm TL. Size at maturity in the male is 114 cm and in the female 130–162 cm (Compagno, 2001). The usual reproduction, mating and birth season for *N. velox* is May–

July, although recent births have been observed toward the end of March (Ruiz-Alvarado & Mijangos-López, 1999; Mendizábal *et al.*, 2000; Villavicencio, 2000; Bizzarro *et al.*, 2007). The species *N. velox* is classified as highly vulnerable and its conservation is critical as it requires very specific nursery areas and has a low fecundity rate (four individuals per female with 9 months' gestation period; Soriano-Velásquez *et al.*, 2006).

Nasolamia velox is commonly reported among fishery landings of the Eastern Pacific Coast, such as Mexico (Saucedo-Barrón, 1982; Cabrera, 2000; Soriano *et al.*, 2006; Bizzarro *et al.*, 2009; Walther-Mendoza *et al.*, 2013), Panama (Compagno & Garrick, 1983), Costa Rica (Garro *et al.*, 2011), Peru (Kato *et al.*, 1967), Colombia (Mantilla, 1998; Mejía-Falla *et al.*, 2010) and Ecuador (Bearez, 1996). In Mexico, the distribution of *N. velox* ranges from Baja California (Walther-Mendoza *et al.*, 2013), south of Sinaloa (van der Heiden & Findley, 1988), east coast of Baja California Sur (Bizzarro *et al.*, 2007), Michoacan (Madrid-Vera *et al.*, 1998), Oaxaca (Alavez-Jiménez, 2006) to the Gulf of Tehuantepec, Chiapas (Soriano-Velásquez & Acal-Sánchez, 2003). In the Gulf of California, *N. velox* has also been reported in the outer Gorda Banks, Cabo San Lucas, and Santa Maria Bay in Baja California Sur, San Felipe, Bahía Las Animas in Baja California, Guaymas in Sonora, Mazatlán and Bahía Topolobampo in Sinaloa (Compagno & Garrick, 1983).

Corresponding author:

F. Galván-Magaña

Emails: galvan.felipe@gmail.com and fgalvan@ipn.mx

Some reports have been published about the presence of this species in insular areas of the ETP. In Galapagos, reports show evidence of records at the northern end of Isabela (1987) and Baltra Island (1989) (Grove & Lavenberg, 1997). In Guadalupe Island, tourists and fishermen have reported the presence of *N. velox*, however there are no records in the formal literature (Walther-Mendoza *et al.*, 2013). The goals of this study were to describe an extension in the range of *N. velox* and examine their movements at the Revillagigedo Archipelago, a group of oceanic islands off the west coast of Mexico.

MATERIALS AND METHODS

Study site

The Revillagigedo Archipelago, located 400 km south of Baja California, Mexico, is a Marine Protected Area composed of four islands: Socorro, Clarion, San Benedicto and Roca Partida (Figure 1). Prevailing currents in the Gulf of California favour chance dispersal of organisms to these islands, particularly of reef fauna (Bizzarro *et al.*, 2009). Due to the large percentage of endemic biota and high biodiversity, the Revillagigedo Biosphere Reserve was listed as a UNESCO World Heritage site in July 2016. The MPA includes a 9.5 nautical mile no-take zone from the coast of the islands to the surrounding waters. The oceanographic conditions of the Revillagigedo Archipelago support one of the most diverse shark communities in the world and the greatest diversity of these species in the ETP (IUCN, 2016), where nine species of sharks are frequently observed (*Galeocerdo cuvier*, *Carcharhinus falciformis*, *Carcharhinus albimarginatus*, *Carcharhinus galapagensis*, *Carcharhinus limbatus*, *Carcharhinus obscurus*, *Sphyrna lewini*, *Rhincodon typus*, *Triaenodon obesus*) and another eight species of sharks are occasionally found (*Alopias pelagicus*, *Alopias superciliosus*, *Alopias vulpinus*, *Echinorhinus cookei*, *Isurus oxyrinchus*, *Nasolamia velox*, *Prionace glauca*) (CONANP, 2005; M. Hoyos, unpublished data).




Fig. 1. Locations of acoustic receivers (grey dots) and the sighting during a shark survey (El Boiler, SB) in the Revillagigedo Archipelago (black dot).

Shark tagging and acoustic receivers

During a study on shark movements and residency in Cabo Pulmo National Park, Mexico in March 2013, a *N. velox* was captured using hand-lines baited with fish. The individual shark was fitted with a coded acoustic transmitter (V16, 69.0 kHz, Vemco, Halifax), which emitted a uniquely coded acoustic signal at random intervals between 60–120 s. The shark was brought onto a small fishing boat, immobilized on the deck, and a hose placed in the mouth with running salt-water to pump through the gills while the shark was manipulated. The coded transmitter was implanted surgically into the body cavity of the shark through a small 2 cm incision and then the wound was closed with three sutures. In addition, total, fork and precaudal lengths of the shark were measured, sex determined by the presence of claspers, and location recorded with a GPS. The identification of *N. velox* was determined on the basis of videos and pictures that highlighted the diagnostic characters (Figure 2, described by Compagno & Garrick, 1983).

Since 2009 Pelagios Kakunja and University of California-Davis have set an array of eight autonomous acoustic receivers (VR2W, Vemco, Halifax) in different sites around the Revillagigedo Archipelago at depths easily reached by scuba divers (average 25 m below the surface). These receivers were designed to listen for coded transmitters and to record the date and time of arrival and departure of individual sharks. The acoustic range of each receiver varied depending on water depth, tide and neighbouring reef structure. Range tests at other sites indicated transmitter detection ranges of 200–300 m. Therefore, as long as a shark tagged (with a unique acoustic signal code) was within the detection range of one of the acoustic receivers, we were able to determine




Fig. 2. *Nasolamia velox* tagged in Cabo Pulmo National Park, Mexico (March 2013). (a) Dorsal view showing the white colouration in the tip of the snout, (b) Ventral view showing the particular nostrils position that differentiated the genus *Nasolamia* from other *Carcharhinidae*.

the presence of this shark. If the shark was recorded in two or more receivers, we knew that the individual was moving between two or more of the monitored sites.

The sighting recorded on video

During a shark survey performed on 5–13 March 2014, visual censusing was conducted by scuba divers. A total of 22 observations were taken at six different areas of the Archipelago: Socorro Island (Punta Tosca and Cabo Pearce), Roca Partida and San Benedicto (Cuevitas, El Boiler and El Canyon). Each survey lasted ~45 minutes. Information on depth and temperature were recorded and a number of images and videos of the sharks were taken using a GOPRO 3+ camera.

The identification of the whitnose shark was made on the basis of photographs and videos that documented diagnostic evidence of this species. Once the species was confirmed, we verified its geographic range using specialized literature and revised data on collected specimens from worldwide museums in order to confirm that there was no existing record of this species from the Revillagigedo Archipelago. This specialized literature and revised data were available on the International Centre for Living Aquatic Resources Management website (<http://www.fishbase.org>; Froese & Pauly, 2016).

RESULTS

The presence of *N. velox* was recorded by acoustic telemetry and video in the Revillagigedo Archipelago. The first record occurred at the receiver located at the east side of Roca Partida Island (18°99'186"N 112°08'44"W) at a depth of 33 m. The tagged individual was first detected on 12 May 2014 and last detection was on 22 February 2016, hence total duration of monitoring was 646 days, however this individual was only present 14% of the time (96 days; Figure 3). The second record occurred at San Benedicto Island at 24 m on 6 March 2015, at the dive site known as 'El Boiler' (19°19'57"N 110°49'211"W). Water temperature was 25.5°C. The identified specimen was ~120 cm in length. In both cases (telemetry and video records) the sharks were not collected, but positive identification was based on a




Fig. 3. Map showing the movement from where the shark was tagged (National Park Cabo Pulmo, March 2013) to where it was first detected (Roca Partida, Revillagigedo Archipelago, May 2014).

single high-definition video of one of the sharks by observing diagnostic characteristics (Figures 2 & 4; Compagno, 2001).

DISCUSSION

We report here for the first time, the occurrence of *N. velox* in Cabo Pulmo and the Revillagigedo Archipelago, and movements from a coastal location (Cabo Pulmo reef) to a group of oceanic islands (Revillagigedo). We provide evidence that the movement patterns of *N. velox* are longer and more complex than what was previously considered for this species, and its distribution range is extended over 487 km.

According to the literature and museum records, the distribution range of *N. velox* is restricted to the Eastern Pacific, from Mexico (Saucedo-Barrón, 1982; van der Heiden & Findley, 1988; Gilbert, 1898; Cabrera 2000; CONANP 2005; Soriano *et al.*, 2006; Bizzarro *et al.*, 2009; Walther-Mendoza *et al.*, 2013; Madrid-Vera *et al.*, 2015) to Panama (Compagno & Garrick, 1983), Costa Rica (Garro *et al.*, 2011), Peru (Kato *et al.*, 1967), Colombia (Mejia-Falla *et al.*, 2010) and Ecuador (Bearez, 1996). Previous records at insular locations of the Eastern Pacific included the Galapagos Archipelago, 3150 km south of Revillagigedo Archipelago. There were no previous records from the Revillagigedo Archipelago, and the closest records were from Cabo San Lucas, 487 km north-east of the Revillagigedo Archipelago (Bizzarro *et al.*, 2007).

The whitnose shark is frequently classified as a tropical inshore and offshore shark, normally found over the continental shelves in shallow coastal waters at depths of 15–24 m, but occasionally it can be found down to 192 m (Compagno, 2001). In Guatemala, this species has been reported 30–100 km off the coastline on the continental slope (Porras, 1997; Ruiz-Alvarado & Mijangos-López, 1999). In Costa Rica, *N. velox* is found in offshore fisheries 80–120 km off the coast and in demersal fisheries on the slopes of the continental shelf (Garro *et al.*, 2011).

Our findings show evidence that there may be potential connectivity of sharks between the Gulf of California and the Revillagigedo Archipelago, however population level observations are needed to confirm this idea. Previous studies have also found other species such as tiger shark, *Galeocerdo cuvier*, moving between Cabo Pulmo and Revillagigedo (J. Ketchum, unpublished data), and the giant manta, *Manta birostris*, using both areas as part of their life



Fig. 4. The shark *N. velox* sighting during a survey at El Boiler, San Benedicto Island (6 March 2015). Individual of ~120 cm TL.

cycle (R. Rubin, Pacific Manta Research Group personal communication, 2015).

Very little is known about the biology, reproduction and ecology of *N. velox*, therefore, future research on this species is necessary. It has been reported that the species has a low fecundity rate and very specific nursery areas (Compagno, 1988; Ruiz-Alvarado & Mijangos-López, 1999). Because of these characteristics this species has been classified as vulnerable to habitat degradation and marine pollution (Alavez-Jiménez, 2006). Responses to the El Niño–Southern Oscillation (ENSO) were documented with high catch per unit effort (CPUE) in landings off the Pacific Coast of Mexico during 1998, prompting the dispersal of a larger number of whitnose sharks (Soriano-Velásquez *et al.*, 2004). It is necessary to increase our understanding of this species and to generate effective management and conservation strategies for vulnerable shark species in the region.

ACKNOWLEDGEMENTS

Thanks to National Geographic and Fischer Productions.

FINANCIAL SUPPORT

We thank the World Wildlife Fund–Gulf of California Program for providing funds to carry out the first shark-tagging project in Cabo Pulmo. We would also like to acknowledge The International Community Foundation and UC Davis for supporting the deployment of the first array of acoustic receivers at Cabo Pulmo and the Revillagigedo Islands. Finally, we thank the Instituto Politécnico Nacional (COFAA, EDI) and the Comisión Nacional para la Ciencia y Tecnología (CONACYT) of Mexico that supported FLL to carry out this work.

REFERENCES

- Alavez-Jiménez C.** (2006) Aspectos de la biología y pesquería de los tiburones sedoso (*Carcharhinus falciformis*) y pico blanco (*Nasolamia velox*), capturados por la flota artesanal pesquera de ensenada Chipehua, Oaxaca, durante el periodo 2004–2005. Bachelor thesis. Instituto Tecnológico de Salina Cruz, Mexico.
- Bearez P.** (1996) Lista de los peces marinos del Ecuador continental. *Revista de Biología Tropical* 44, 731–741.
- Bizzarro J., Smith W., Hueter R., Tyminski J., Márquez-Farías J.F., Castillo-Géniz J.L., Cailliet G.M. and Villavicencio-Garayzar C.J.** (2007) The status of shark and ray fishery resources in the Gulf of California: Applied research to improve management and conservation. Report to the David and Lucille Packard Foundation, 237 pp.
- Cabrera C.C.** (2000) Determinación de los hábitos alimenticios durante las estaciones primavera y verano de *Carcharhinus falciformis*, *Sphyrna lewini* y *Nasolamia velox* (Carcharhiniformes: Carcharhinidae) a partir del análisis de su contenido estomacal en el Golfo de Tehuantepec. México Bachelor Thesis. UNAM, México.
- Compagno L.J.** (1988) *Sharks of the order Carcharhiniformes*. Princeton, NJ: Princeton University Press.
- Compagno L.J.** (2001) *Sharks of the world: an annotated and illustrated catalogue of shark species known to date*. Rome: FAO.
- Compagno L.J. and Garrick J.A.F.** (1983) *Nasolamia*, new genus, for the shark *Carcharhinus velox* Gilbert, 1898 (Elasmobranchii: Carcharhinidae). Wellington: Zoology Publications from Victoria University of Wellington, pp. 76–77.
- CONANP (Comisión Nacional de Áreas Naturales Protegidas)** (2005) *Programa de conservación y manejo. Reserva de la Biosfera Archipiélago de Revillagigedo*. México, DF: CONANP.
- Froese R. and Pauly D.** (eds) (2016) *FishBase* (version 10.2015). Available at: <http://www.fishbase.org/>.
- Garro A.L., Vargas R.A., Zanella I. and Le Foulgo L.** (2011) Análisis de las capturas de tiburones y rayas en las pesquerías artesanales de Tárcoles, Pacífico Central de Costa Rica. *Revista Ciencias Marinas y Costeras* 1, 145–157.
- Grove J.S. and Lavenberg J.** (1997) *The fishes of the Galapagos islands*. Stanford, CA: Stanford University Press.
- IUCN** (2016) IUCN report for the World Heritage Committee, 40th session, Istanbul, Turkey. WHC/16/40.COM/INF.8B2.
- Kato S., Springer S. and Wagner M.H.** (1967) *Field guide of Eastern Pacific and Hawaiian sharks*. U.S. Fish and Wildlife Service, Circular 271, 47 pp.
- Madrid Vera J., Luna A.R. and Bravo I.R.** (2015) Peces de la plataforma continental de Michoacán y sus relaciones regionales en el Pacífico mexicano. *International Journal of Tropical Biology and Conservation* 46, 267–276.
- Mantilla L.** (1998) Lista de especies elasmobranchios de Colombia. *Rev. Fen. Anat.* Vol I. 19/08/2006. Available at <http://www.bioaquaticresearch.com/Sharks/1-2-01-Tib.html>.
- Mejía-Falla P.A., Tobón-López A., Navia A.F., Narváez K. and Lozano R.A.** (2010) Avistamiento de elasmobranchios en aguas colombianas. In Mejía-Falla P.A., Narváez K., Bohórquez J., Osaer F., Ramírez V. and Hleap J.S. (eds) *Libro de resúmenes II Encuentro colombiano sobre condricios*. Cali, Colombia: Fundación SQUALUS, p. 29.
- Mendizábal O.D., Vélez-Marín R., Soriano-Velásquez S.R. and Castillo-Géniz J.L.** (2000) *Tiburones oceánicos del Pacífico. Sustentabilidad y Pesca Responsable en México: Evaluación y Manejo*. Instituto Nacional de la Pesca, SEMARNAP, pp. 155–195.
- Naylor G.J., Caira J.N., Jensen K., Rosana K.A.M., White W.T., and Last P.R.** (2012) A DNA sequence-based approach to the identification of shark and ray species and its implications for global elasmobranch diversity and parasitology. *Bulletin of the American Museum of Natural History* 367, 1–262.
- Porras G.** (1997) *Contribución al conocimiento de la biología y pesquería del tiburón punta de zapato, Nasolamia velox, capturado por la pesca artesanal en el Pacífico de Guatemala*. Problema Especial. Guatemala: CEMA-USAC/DIGESEPE-DITEPESCA.
- Ruiz-Alvarado C.L. and Mijangos-López N.** (1999) *Estudio sobre la pesquería del tiburón en Guatemala*. [Case Studies for the Management of Elasmobranch Fisheries.] Rome: FAO, Unidad Especial de Ejecución para la Pesca y la Acuicultura.
- Ruiz C., Arauz R., Pérez-Jiménez J.C., Castillo-Geniz J.L. and Soriano-Velásquez S.** (2009) *Nasolamia velox*. The IUCN Red List of Threatened Species. Version 2014.3.
- Saucedo-Barrón C.J.** (1982) El tiburón: contribución al estudio de la pesquería en la zona sur de Sinaloa, México. *Ciencias del Mar, Universidad Autónoma de Sinaloa. Epoca* 1, 14–29.
- Soriano-Velásquez S. and Acal Sánchez D.** (2003) La pesquería de tiburón en Chiapas y su entorno socioeconómico. La Jornada Ecológica. UNAM. Available at <http://www.jornada.unam.mx/2003/01/27/eco-f.html>.

- Soriano-Velásquez S.R., Acal D.J.L. and Castillo-Géniz J.L.** (2004) *Aspectos reproductivos del tiburón coyotito Nasolamia velox* (Compagno y Garrick, 1983), capturado en el Golfo de Tehuantepec, México. Reporte de Investigación. México: INAPESCA.
- Soriano-Velásquez S.R., Acal-Sánchez D.E., Castillo-Géniz J.L., Vázquez-Gómez N.Y. and Ramírez-Santiago C.E.** (2006) Tiburón del Golfo de Tehuantepec. Sustentabilidad y Pesca Responsable en México. Arreguín-Sánchez F., Beléndez-Moreno L.F., Méndez Gómez-Humarán I., Solana-Sansores R. and Rangel-Dávalos C. (eds). México: Instituto Nacional de la Pesca, SAGARPA, pp. 323–360.
- Van der Heiden A.M. and Findley L.T.** (1988) Lista de los peces marinos del sur de Sinaloa, México. *Anales del Instituto de Ciencias del Mar y Limnología, Universidad Nacional Autónoma de México* 15, 209–224.
- Villavicencio-Garáyzar C.J.** (2000) *Áreas de crianza de tiburones en el Golfo de California*. Universidad Autónoma de Baja California Sur, Área Interdisciplinaria de Ciencias del Mar, Informe final. México, DF: SNIBCONABIO.
- and
- Walther-Mendoza M., Ayala-Bocos A., Hoyos-Padilla M. and Reyes-Bonilla H.** (2013) New records of fishes from Guadalupe Island, northwest Mexico. *Hidrobiológica* 23, 410–414.

Correspondence should be addressed to:

F. Galván-Magaña
Instituto Politécnico Nacional,
Centro Interdisciplinario de Ciencias Marinas, Av. IPN s/n.
C.P. 23096. La Paz, B.C.S, Mexico
Email: galvan.felipe@gmail.com; fgalvan@ipn.mx