

Extra-hepatopulmonary cystic echinococcosis in Bulgaria: frequency, management and outcome of the disease

Research Article

Cite this article: Harizanov RN, Rainova IG, Kaftandjiev IT (2011). Extra-hepatopulmonary cystic echinococcosis in Bulgaria: frequency, management and outcome of the disease. *Parasitology* **148**, 562–565. <https://doi.org/10.1017/S0031182020002206>


Received: 31 July 2020
Revised: 26 October 2020
Accepted: 16 November 2020
First published online: 20 November 2020

Key words:

Cystic echinococcosis; diagnosis; extra-hepatopulmonary localization; outcome

Author for correspondence:

Rumen N. Harizanov,
E-mail: harizanov@ncipd.org

Rumen N. Harizanov , Iskra G. Rainova and Iskren T. Kaftandjiev

Department of Parasitology and Tropical Medicine, National Centre of Infectious and Parasitic Diseases, 26 Yanko Sakazov Blvd., 1504 Sofia, Bulgaria

Abstract

In the endemic countries, human cystic echinococcosis (CE) poses a serious medical and social problem. Because it most often affects the liver and lungs we aimed to define the proportion of cases with different organ localization, the diagnostic and therapeutic approaches in such cases, and the outcome of them. For a period 2010–2019, a total of 2863 cases of CE were registered in Bulgaria, of which 148 (5.17%) with organ localization other than liver and/or lung. The majority of patients with extra-hepatopulmonary localization of CE were adults. The distribution by gender showed predominance of female patients (57.43%) over those of males and primary cases (85.14%) exceeded the cases of recurrence. According to our study most common is the spleen involvement, followed by involvement of the abdominal cavity, kidneys and muscle/subcutaneous tissue. Other extra-hepatopulmonary organ localizations are significantly less common. This study shows that the extra-hepatopulmonary localization of CE is not so rare, and in most cases it is a primary disease. In respect of this, clinicians should consider hydatidosis in the differential diagnosis when cystic formation(s) is found, regardless of the organ involved.

Introduction

According to the WHO definition, human cystic echinococcosis (CE) is a zoonotic disease caused by species complex centred on *Echinococcus granulosus*. The disease is widespread and cases among humans are recorded all over the world except for Antarctica (WHO, 2020). In high-endemic countries, the disease poses a serious medical and social problem and a challenge to the systems of public health.

According to a various literature sources, the disease most often affects the liver with the lungs being in second place, and these organ localizations are found in at least 90% of all cases (Eckert *et al.*, 2001; Cöl *et al.*, 2003; Ahmadi and Hamidi, 2010). Often the course of the disease is without symptoms and if any are present these are not pathognomonic and depend on the affected organ (Eckert *et al.*, 2001).

Among the member states of the European Union (EU), Bulgaria holds a leading position by the number of cases of CE among its population. Regardless of the steady trend of decreasing annual number of cases by 43.5% for the period 2008–2017 (i.e. from 386 in 2008 to 218 in 2017), the country is still with highest annual incidence of CE in the EU (EFSA and ECDC, 2018). Because of the obvious importance of the disease, and because it most often affects the liver and lungs we aimed to define the proportion of cases with different organ localization, the diagnostic and therapeutic approaches in such cases, and the outcome of them.

Patients and methods

Study design

The study is a retrospective analysis of cases of CE with extra-hepatopulmonary localization recorded for a 10-year period from January 2010 to December 2019. Cases involving the liver and/or lungs alone, or in combination with other affected organs were excluded from the study to determine the frequency of cases in which only other organs and systems of the human body were affected. Data from epidemiological survey cards of cases with CE registered in the country and containing medical information about the organ localization and number of hydatid cysts, the method of diagnosis and confirmation of disease, the therapeutic approach and clinical outcome were processed. According to the existing legislation in the country, each case of CE is subject to a mandatory notification and epidemiological investigation. At a local level, this is performed by the Regional Health Inspectorates (RHI) who every month sends the collected information to the National Center for Infectious and Parasitic Diseases (NCIPD), where it is analysed and on its grounds are prepared proposals to the Ministry of Health for improvement of disease surveillance and control.

Table 1. Reported cases of extra-hepatopulmonary CE by year

Year	Total cases of CE	Cases of extra-hepatopulmonary CE	Percent
2010	335	13	3.88
2011	345	15	4.35
2012	346	18	5.20
2013	306	12	3.92
2014	333	13	3.90
2015	313	20	6.39
2016	269	15	5.58
2017	218	11	5.05
2018	206	13	6.31
2019	192	18	9.38
Total	2863	148	5.17
Median	309.5	14	5.13
Minimum	192	11	3.88
Maximum	346	20	9.38

Ethical standards

The current study is not part of a funded research project, and the data were collected by routine diagnostic activities and epidemiological investigations carried out under Bulgarian legislation. As the presented data are retrospective, de-identified and depersonalized an ethical approval and consent to participate were not applicable.

Statistical analysis

Some indicators such as median, min–max and percent have been identified.

Results

A total of 2863 cases of CE were registered in the country, of which 148 with organ localization other than liver and/or lung. The proportion of these cases is variable over the years, averaging 5.17% over the period (Table 1). The organ localization of hydatid disease is presented in Table 2.

The distribution of cases by age, gender and type (primary or relapses) is presented in Table 3. Primary cases prevail among persons over 18 years of age and among females.

Only in 9.46% of the cases ($n = 14$) the disease proceeded without clinical symptoms, as in six of the patients it was a recurrence found during the follow-up observation, and in eight of them it was diagnosed accidentally in imaging studies performed on another occasion. In the remaining 90.54% ($n = 134$) there were symptoms most often expressed as: heaviness, discomfort and dull pain in the affected organ ($n = 86$, 58.11%) and detection of swelling or tumour formation ($n = 27$, 18.24%).

Imaging methods [ultrasound, radiography, computerized tomography scan and magnetic resonance imaging (MRI)] were used in all patients with extra-hepatopulmonary CE. The diagnosis was confirmed by serological and histological methods. Both methods were used in 29.05% of the cases ($n = 43$), only serological methods in 8.11% ($n = 12$), and only by histological study were confirmed 62.84% of the cases ($n = 93$). In eight cases, the serological results were negative or borderline and diagnosis was confirmed histologically.

Table 2. Organ involvement in the cases of extra-hepatopulmonary CE

Organ localization	Number of cases	Percent
Spleen	40	27.03
Abdominal cavity	33	22.29
Kidneys	21	14.19
Muscles/subcutaneous tissue	21	14.19
Bones	11	7.43
Heart	7	4.73
Uterus, ovaries, adnexa	5	3.38
Pancreas	3	2.03
Pelvis	3	2.03
CNS	3	2.03
Spinal cord	1	0.67
Total	148	100

Regarding the therapeutic approach, 95.95% ($n = 142$) of the patients underwent surgical treatment, 3.38% ($n = 5$) underwent medical treatment and one patient (0.67%) was treated by a puncture–aspiration–injection–reaspiration (PAIR) technique. With exception of one case that ended with death caused by a spontaneous rupture of a hydatid cyst in the right atrium of the heart, in all other cases the outcome was favourable.

Discussion

Literature data on extra-hepatopulmonary echinococcosis are scarce and most are based on descriptions of clinical cases or small series of patients (Çakır *et al.*, 2016). Studies based on larger groups of patients are relatively rare and are mostly from Turkey and Iran, where the disease is endemic (Cöl *et al.*, 2003; Ahmadi and Hamidi, 2010; Akcam *et al.*, 2014; Çakır *et al.*, 2016).

CE is one of the parasitic diseases of greatest medical importance in Bulgaria (Harizanov *et al.*, 2019). Several hundred cases are registered annually in the country. The relative proportion of patients with extra-hepatopulmonary echinococcosis is just over 5%, which is consistent with the literature (Eckert *et al.*, 2001). The majority of patients with extra-hepatopulmonary localization of CE were persons over 18 years of age (86.49%). This can be explained by the slow, chronic course of the disease that in many cases started long time ago, i.e. the infection can occur in childhood but it can be diagnosed much later. The distribution by gender showed a predominance of female patients ($n = 85$, 57.43%) over those of males ($n = 63$, 42.57) and primary cases ($n = 126$, 85.14%) exceeded the cases of recurrence ($n = 22$, 14.86%). In 81.08% ($n = 120$) of the patients, the presence of a solitary hydatid cyst was detected, and in 18.92% ($n = 28$) two or more were diagnosed.

According to our study, most common is the spleen involvement: 1.4% from all reported cases of CE for the studied period and 27.03% from the cases with extra-hepatopulmonary localization. Regarding this organ localization, our data are similar to those of some authors (Eris *et al.*, 2013; Belli *et al.*, 2014; Çakır *et al.*, 2016; Bartın, 2019) and differ significantly from others (Akcam *et al.*, 2014). In 92.5% of the cases with splenic localization the presence of a solitary hydatid cyst in the organ was found, and only in 7.5% ($n = 3$) the presence of two or more cysts was found. From the total of 40 patients with splenic hydatid cyst (SHC), only three of them (7.5%) had secondary SHC. In this respect, our data differ somewhat from those in the studies of Akbulut *et al.* (2013).

Table 3. Reported cases of extra-hepatopulmonary CE by age, gender and type of case

Year	Age groups		Sex		Type of case	
	Children	Adults	Male	Female	Primary	Relapse
2010	3	15	8	10	16	2
2011	1	12	5	8	12	1
2012	0	11	3	8	11	0
2013	2	13	6	9	12	3
2014	4	16	9	11	16	4
2015	2	11	5	8	11	2
2016	2	10	5	7	11	1
2017	4	14	9	9	14	4
2018	1	14	7	8	12	3
2019	1	12	6	7	11	2
Total	20	128	63	85	126	22
Percent	13.51	86.49	42.57	57.43	85.14	14.86

The incidence of CE of the peritoneal cavity and retroperitoneal space varies from 6 to 16% according to various studies (Engin *et al.*, 2000; Akcam *et al.*, 2014). Our data are for 22.2% involvement of the abdominal cavity (retroperitoneum and mesentery), and in these cases the percentage of multiple echinococcosis is much higher – 18.18% ($n = 6$).

Our study found that the frequency of kidney and muscle/subcutaneous tissue involvement was identical (14.19% each). All cases of renal echinococcosis were with unilateral involvement and had a solitary cyst in 90.48% of the patients ($n = 19$). Regarding CE affecting muscles and/or subcutaneous tissue, we found that the most common site of involvement is the cervical ($n = 8$), and gluteal and femoral regions (three cases, respectively). In this respect, our data differ greatly from those of the authors, according to whom muscle involvement alone is very rare (García-Alvarez *et al.*, 2002; Burgazli *et al.*, 2013).

Other organs and systems are affected with a relatively lower frequency. It is noteworthy that the relative share of multiple echinococcosis is higher in bone involvement (54.55%, six out of 11 cases). We found that the spine was most often affected ($n = 10$), and more frequently the thoracic vertebrae ($n = 7$) were involved. Common to all of these cases was that they had symptoms with acute pain in the affected area, and in one of them spontaneous fracture of a thoracic vertebra occurred.

Cardiac involvement was found in 4.73% of all cases with extra-hepatopulmonary localization of CE, while in published studies the incidence of cardiac involvement ranged from 0.2 to 2% (Prousalidis *et al.*, 1998; Engin *et al.*, 2000; Ibn Elhadj *et al.*, 2014). In 42.86% of these cases the cysts were found in the pericardium ($n = 3$), and in the rest, echinococcal cysts were localized in the epicardium, right atrium, right and left ventricles.

Although the study covers a considerable period of time, the identified cases of CE involving the central (CNS) and peripheral nervous system (spinal cord) are relatively small, and their frequency is similar to that presented in the literature (Altinörs *et al.*, 2000; Engin *et al.*, 2000). In all patients the disease progressed with pronounced neurological symptoms (paraplegia, headache, vomiting, epileptiform seizures and paresthesia of the lower extremities).

Interestingly, in all three cases of pancreatic involvement, reported during the study period, the main clinical symptom was development of mechanical jaundice, one of most serious

complications in pancreatic hydatid cyst disease. The frequency of PMS (2.03%) in the current study was similar to that described by other authors (Akbulut *et al.*, 2014).

The diagnostic algorithm for extra-hepatopulmonary CE does not differ from that in liver or lung involvement. Imaging examinations play a major role, with more than half of the patients using more than one method. The primary imaging diagnosis includes ultrasound or radiography, depending on the nature of the patients' complaints, and subsequently some of them underwent additional computed tomography in the course of diagnostic clarification.

MRI was used only in cases where patients had neurological symptoms and suspected volume-occupying lesions in the brain and spinal cord. Confirmatory diagnosis most often was histological, which is probably due to the fact that over 95% of patients were treated surgically. However, 37.16% of patients were tested by serological methods. A negative or borderline result was observed in 14.55% ($n = 8$) of a total of 55 patients. In the diagnostic practice in Bulgaria, of all immunological techniques, enzyme-linked immunosorbent assay is most often used as a screening and western blot as a confirmatory test. Commercial kits are used according to the manufacturer's protocol. However, serological tests still play a complementary role to the imaging examinations in the diagnosis of CE. The relatively low sensitivity (up to 30% of false-negative results) of available commercial kits, especially in cases of extrahepatic localization of CE, makes serological results difficult for interpretation (Sarkari and Rezaei, 2015). Akbulut *et al.* (2014) reported that 15 of the 40 patients with pancreatic echinococcosis described in the literature had negative serological tests for CE.

In the patients included in our study, main therapeutic approach was the surgical removal of the hydatid cyst/cysts. This can be explained by the extra-hepatopulmonary localization of the disease and is in line with the currently established guidelines for treatment of the disease (Eckert *et al.*, 2001). PAIR was administered to only one patient with solitary hydatid cyst of the kidney, and five individuals were treated with medications only, three of whom had recurrence and multiple involvement. The other two were inoperable due to concomitant pathology, which did not allow surgical intervention. According to the treatment algorithm of CE accepted in Bulgaria, each patient treated surgically or with PAIR is subject to a drug anti-relapse

prophylaxis with albendazole, and the duration of the course is determined individually, according to severity of the disease (from 1 to more of 12 months). A decision on antiparasitic medical treatments given to patients before surgery or PAIR is made by the treating physician and lasts from 1 day to 1 week before the scheduled procedure, but this prophylaxis is not precisely regulated and does not cover all patients with CE. Because, recurrences are registered annually, this gives us reason to believe that not all patients treated for CE with invasive techniques are covered by anti-relapse drug prophylaxis.

For the period studied by us, except for one case that ended with death due to spontaneous rupture of an echinococcal cyst located in the right atrium, all others had a favourable outcome.

This study shows that the extra-hepatopulmonary localization of CE is not so rare, and in most cases it is a primary disease. In this aspect, clinicians should consider hydatidosis in the differential diagnosis when cystic formation(s) is found, regardless of the organ involved.

Author contribution. RH conceived and designed the study. IK and IR conducted data gathering, RH, IR and IK wrote the manuscript.

Conflict of interest. The authors declare that they have no conflict of interest.

Ethical standards. The current study is not part of a funded research project, and the data were collected by routine diagnostic activities and epidemiological investigations carried out under Bulgarian legislation. As the presented data are retrospective, de-identified and depersonalized an ethical approval and consent to participate were not applicable.

References

- Ahmadi NA and Hamidi M** (2010) Unusual localizations of human hydatid disease in Hamedan province, west of Iran. *Helminthologia* **47**, 94–98.
- Akbulut S, Sogutcu N and Eris C** (2013) Hydatid disease of the spleen: single-center experience and a brief literature review. *Journal of Gastrointestinal Surgery* **17**, 1784–1795.
- Akbulut S, Yavuz R, Sogutcu N, Kaya B, Hatipoglu S, Senol A and Demircan F** (2014) Hydatid cyst of the pancreas: report of an undiagnosed case of pancreatic hydatid cyst and brief literature review. *World Journal of Gastrointestinal Surgery* **6**, 190–200.
- Akcam AT, Ulku A, Koltas IS, Izol V, Bicer OS, Kilicbagir E, Sakman G, Poyrazoglu H, Erman T, Aridogan IA, Parsak CK, Inal M and Iskiti S** (2014) Clinical characterization of unusual cystic echinococcosis in southern part of Turkey. *Annals of Saudi Medicine* **34**, 508–516.
- Altinörs N, Baybek M, Caner HH and Erdogan B** (2000) Central nervous system hydatidosis in Turkey: a cooperative study and literature survey analysis of 458 cases. *Journal of Neurosurgery* **93**, 1–8.
- Bartın MK** (2019) Hydatid cyst disease with extra hepatic localizations. *Biomedical Journal of Scientific & Technical Research* **19**, 14625–14628. doi: 10.26717/BJSTR.2019.19.003367
- Belli S, Akbulut S, Erbay G and Koçer NE** (2014) Spontaneous giant splenic hydatid cyst rupture causing fatal anaphylactic shock: a case report and brief literature review. *Turkish Journal of Gastroenterology* **25**, 88–91.
- Burgazli KM, Ozdemir CS, Beken Ozdemir E, Mericiler M and Polat ZP** (2013) Unusual localization of a primary hydatid cyst: a subcutaneous mass in the paraumbilical region. *European Review for Medical and Pharmacological Sciences* **17**, 1766–1768.
- Çakır M, Balasar M, Küçükkartallar T, et al.** (2016) Management of extra-hepatopulmonary hydatid cysts (157 cases). *Turkiye Parazitoloji Dergisi* **40**, 72–76.
- Cöl C, Cöl M and Lafçi H** (2003) Unusual localizations of hydatid disease. *Acta Medica Austriaca* **30**, 61–64.
- Eckert J, Gemmell MA, Meslin F-X and Pawlowski ZS and World Health Organization** (2001) *WHO/OIE Manual on Echinococcosis in Humans and Animals: A Public Health Problem of Global Concern*. Paris, France: World Organisation for Animal Health.
- Engin G, Acunaş B, Rozanes I and Acunaş G** (2000) Hydatid disease with unusual localization. *European Radiology* **10**, 1904–1912.
- Eris C, Akbulut S, Yildiz MK, Abuoglu H, Odabasi M, Ozkan E, Atalay S and Gunay E** (2013) Surgical approach to splenic hydatid cyst: single center experience. *International Surgery* **98**, 346–353.
- European Food Safety Authority & European Centre for Disease Prevention and Control** (2018) The European Union summary report on trends and sources of zoonoses, zoonotic agents and food-borne outbreaks in 2017. *EFSA Journal* **16**, 5500.
- García-Alvarez F, Torcal J, Salinas JC, et al.** (2002) Musculoskeletal hydatid disease: a report of 13 cases. *Acta Orthopaedica Scandinavica* **73**, 227–231.
- Harizanov R, Rainova I and Kaftandjiev I** (2019) Human cystic echinococcosis, trichinellosis, and toxocariasis in Bulgaria: an update of data for 2015–2017. *International Medicine* **1**, 43–50.
- Ibn Elhadj Z, Boukhris M, Kammoun I, Halima AB, Addad F and Kachboursa S** (2014) Cardiac hydatid cyst revealed by ventricular tachycardia. *Journal of the Saudi Heart Association* **26**, 47–50.
- Prousalidis J, Tzardinoglou K, Sgouradis L, Katsohis C and Aletras H** (1998) Uncommon sites of hydatid disease. *World Journal of Surgery* **22**, 17–22.
- Sarkari B and Rezaei Z** (2015) Immunodiagnosis of human hydatid disease: where do we stand? *World Journal of Methodology* **5**, 185–195.
- World Health Organization** (2020) Echinococcosis fact sheet. Retrieved from WHO website: <https://www.who.int/news-room/fact-sheets/detail/echinococcosis> (Accessed 29 June 2020).