

Is There a Functional Relation Between Set Shifting and Hyperactivity in Children with Attention-Deficit/Hyperactivity Disorder (ADHD)?

Lauren N. Irwin, Nicole B. Groves, Elia F. Soto and Michael J. Kofler* 
Department of Psychology, Florida State University, Tallahassee, FL 32306, USA

(RECEIVED November 19, 2019; FINAL REVISION March 6, 2020; ACCEPTED April 28, 2020; FIRST PUBLISHED ONLINE May 27, 2020)

Abstract

Objective: Replicated evidence indicates that children with attention-deficit/hyperactivity disorder (ADHD) show disproportionate increases in hyperactivity/physical movement when their underdeveloped executive functions are taxed. However, our understanding of hyperactivity's relation with set shifting is limited, which is surprising given set shifting's importance as the third core executive function alongside working memory and inhibition. The aim of this study was to experimentally examine the effect of imposing set shifting and inhibition demands on objectively measured activity level in children with and without ADHD. **Method:** The current study used a validated experimental manipulation to differentially evoke set shifting, inhibition, and general cognitive demands in a carefully phenotyped sample of children aged 8–13 years with ADHD ($n = 43$) and without ADHD ($n = 34$). Activity level was sampled during each task using multiple, high-precision actigraphs; total hyperactivity scores (THS) were calculated. **Results:** Results of the 2×5 Bayesian ANOVA for hyperactivity revealed strong support for a main effect of task ($BF_{10} = 1.79 \times 10^{18}$, $p < .001$, $\omega^2 = .20$), such that children upregulated their physical movement in response to general cognitive demands and set shifting demands specifically, but not in response to increased inhibition demands. Importantly, however, this manipulation did not disproportionately increase hyperactivity in ADHD as demonstrated by significant evidence *against* the task \times group interaction ($BF_{01} = 18.21$, $p = .48$, $\omega^2 = .002$). **Conclusions:** Inhibition demands do not cause children to upregulate their physical activity. Set shifting produces reliable increases in children's physical movement/hyperactivity over and above the effects of general cognitive demands but cannot specifically explain hyperactivity in children with ADHD.

Keywords: ADHD, Set shifting, Cognitive flexibility, Hyperactivity, Global–local, Executive function

INTRODUCTION

Attention-deficit/hyperactivity disorder (ADHD) is a chronic and heterogeneous neurodevelopmental disorder that affects approximately 5% of school-age children (Polanczyk, de Lima, Horta, Biederman, & Rohde, 2007; Polanczyk, Willcutt, Salum, Kieling, & Rohde, 2014). It has been proposed that underlying deficits in executive function(s) may drive ADHD's phenotypic behavioral presentation for many, if not most, children with ADHD (Barkley, 1997; Chacko, Kofler, & Jarrett, 2014; Kasper, Alderson, & Hudec, 2012; Mahone, 2011; Rapport et al., 2009; Sonuga-Barke, Bitsakou, & Thompson, 2010). Hyperactivity, or excess physical movement, is considered a core and impairing deficit in the clinical

model of ADHD (APA, 2013). Recent meta-analytic evidence suggests that children with ADHD exhibit elevated motor activity compared to children without ADHD, regardless of subtype/current presentation, particularly during activities that challenge their underdeveloped executive functions (Kofler, Raiker, Sarver, Wells, & Soto, 2016; Mahone, 2011). However, the extent to which hyperactivity in ADHD is evoked by cognitively challenging tasks in general or by demands on specific executive functions remains unclear. The current study focuses primarily on one of the three core executive functions, set shifting (Miyake et al., 2000), and uses a carefully controlled, counterbalanced experimental manipulation to examine the effects of imposing set shifting demands on objectively measured activity level in children with and without ADHD. In addition, the experimental design included explicit manipulation of inhibitory control given prior evidence implicating this core executive function in performance on set shifting tasks (Irwin, Kofler, Soto, & Groves, 2019).

*Correspondence and reprint requests to: Michael J. Kofler, Department of Psychology, Florida State University, 1107 W. Call Street, Tallahassee, FL 32306-4301, USA. Phone: +(850) 645-0656, Fax: +(850) 644-7739. E-mail: kofler@psy.fsu.edu

Executive functions and hyperactivity

Meta-analytic and experimental evidence indicates that children with ADHD show disproportionate increases in hyperactivity/physical movement when environmental demands challenge their underdeveloped executive functions (Kofler et al., 2016, 2018; Rapport et al., 2009). This pattern may suggest that hyperactivity in ADHD is a response to any type of cognitive/executive function demand. However, emerging experimental results suggest the potential for specificity in this link. For example, multiple studies have reported causal links between working memory demands and hyperactivity (e.g., Hudec et al., 2015; Kofler, Sarver, & Wells, 2015; Patros, Alderson, Hudec, Tarle, & Lea, 2017; Rapport et al., 2009), such that children with ADHD show disproportionate increases in activity level as working memory demands are experimentally evoked. In contrast, emerging evidence suggests that experimentally increasing inhibition does not impact activity level for children with or without ADHD (Alderson, Rapport, Kasper, Sarver, & Kofler, 2012). Thus, the available evidence suggests that hyperactivity in ADHD may be linked with specific executive functions rather than cognitive demands in general.

Importantly, however, no study to date has experimentally examined this relation with set shifting – a critical omission given set shifting's importance as the third core executive function alongside working memory and inhibition (Karr et al., 2018; Miyake et al., 2000). Correlational studies suggest limited, mixed evidence for an association between set shifting abilities and hyperactivity in ADHD with evidence both supporting (McCandless & O'Laughlin, 2007) and failing to support this relation (Toplak, Bucciarelli, Jain, & Tannock, 2008). However, the few studies that have examined this relation have been limited by the use of correlational methods and informant ratings of children's hyperactive behavior that occurred in settings disconnected from the measurement of set shifting abilities (Mahone & Hoffman, 2007). To our knowledge, no study to date has conducted the experimental manipulations of set shifting demands and concurrent measurement of hyperactivity needed to provide evidence for or against a causal link between set shifting and ADHD-related hyperactivity.

Current study

The current study combined a validated experimental manipulation of set shifting and inhibitory control (Irwin et al., 2019) with concurrent and objective measurement of hyperactivity (actigraphy) to test for a causal role of these executive functions on ADHD-related hyperactivity. We hypothesized that set shifting demands would be functionally related to hyperactivity as evidenced by significant increases in activity level for both ADHD and non-ADHD children during shifting relative to nonshifting task conditions. Evidence to support a causal role of set shifting demands on ADHD-related hyperactivity would require disproportionate increases in activity level when set shifting demands are

increased (i.e., group \times task interaction), indicating that set shifting demands specifically evoke or exacerbate differences in hyperactive behavior between children with and without ADHD. In contrast, we did not expect evidence to support main or interaction effects of the inhibitory control manipulation given previous evidence that experimentally increasing inhibition demands does not produce changes in actigraph-measured activity level for children with or without ADHD (Alderson et al., 2012).

METHOD

Participants

The sample included 77 children aged 8–13 years ($M = 10.46$, $SD = 1.54$; 32 girls) from the Southeastern US recruited through community resources from 2015 to 2018 for participation in a clinical research study of the neurocognitive mechanisms underlying pediatric attention and behavioral problems. Psychoeducational evaluations were provided to all caregivers. All parents and children gave informed consent/assent, and IRB approval was obtained/maintained. Sample ethnicity was mixed with 51 White/non-Hispanic (66.2%), 10 Hispanic/English-speaking (13.0%), 9 African-American (11.7%), 3 Asian (3.9%), and 4 multiracial children (5.2%; Table 1).

All children and caregivers completed an identical, comprehensive evaluation that included detailed, semi-structured clinical interviewing and multiple norm-referenced parent and teacher questionnaires. Please refer to Irwin et al. (2019) for a detailed account of the comprehensive psychoeducational evaluation. The final sample included 43 children with ADHD and 34 children without ADHD (Table 1). Psychostimulants ($N_{prescribed}=13$) were withheld ≥ 24 hr for neurocognitive testing. Children were excluded for gross neurological, sensory, or motor impairment; history of seizure disorder, psychosis, autism spectrum disorder, or intellectual disability; or nonstimulant medications that could not be withheld for testing.

Procedures

Children participated in two research sessions (3 hr each) following the baseline psychoeducational assessment. The set shifting and control tasks were administered as part of a larger battery of executive and nonexecutive laboratory tasks. The tasks were counterbalanced across participants to minimize order effects. Children were seated in a caster wheel swivel chair approximately .66 m from the computer monitor for all tasks. Performance was monitored at all times by the examiner, who was stationed just out of the child's view to provide a structured setting while minimizing performance improvements associated with examiner demand characteristics (Gomez & Sanson, 1994). All children received brief (2–3 min) breaks after each task and preset longer (10–15 min) breaks after every 2–3 tasks to minimize fatigue.

Table 1. Descriptive statistics

Demographics	ADHD (<i>N</i> = 43)	Non-ADHD (<i>N</i> = 34)	$\chi^2(4, N = 77)$	<i>P</i>	Phi	BF ₀₁
Gender (boys/girls)	25/18	20/14	.004	.95, ns	.007	2.29
Ethnicity (AA/A/C/H/M)	5/0/32/3/3	4/3/19/7/1	11.13	.09, ns	.38	1.28
	<i>M</i> (<i>SD</i>)	<i>M</i> (<i>SD</i>)	<i>t</i> (75)	<i>P</i>	Cohen's <i>d</i>	BF ₀₁
Age	10.26(1.51)	10.71(1.56)	1.27	.21, ns	.29	2.75
SES	46.71(11.65)	51.59(11.48)	1.84	.07, ns	.42	2.10
VCI (IQ)	102.67(14.16)	109.21(10.30)	2.26	.03	.53	.24*
BASC-2/3 attention problems						
Teacher	63.49(8.69)	52.62(10.76)	-4.91	<.001	1.13	<.01***
Parent	65.98(7.10)	56.47(11.46)	-4.46	<.001	1.03	<.01***
BASC-2/3 hyperactive problems						
Teacher	62.62(15.16)	54.15(12.75)	-2.60	<.01	.60	.11*
Parent	68.00(13.46)	54.62(11.40)	-4.63	<.001	1.06	<.01***
Total hyperactivity scores						
Beginning point	36.60(28.32)	28.45(21.76)	1.38	.17, ns	.32	2.38
Global–global	116.26(88.21)	97.97(100.09)	.85	.40, ns	.19	4.01*
Local–local	97.70(111.19)	69.38(53.94)	1.36	.18, ns	.32	2.44
Global–local	159.82(142.55)	115.84(108.19)	1.49	.14, ns	.35	2.07
Ending point	58.03(46.76)	43.47(43.49)	1.40	.17, ns	.32	2.33

Note. AA, African-American; A, Asian; BASC, Behavior Assessment System for Children; C, Caucasian/non-Hispanic; H, Hispanic/English-speaking; M, multiracial; SES, socioeconomic status; VCI (IQ), Verbal Comprehension Index (Intelligence Quotient).

*Moderate = BF₀₁ < .33 or > 3.00; **strong = BF₀₁ < .10 or > 10.00; ***extreme = BF₀₁ < .01 or > 100.00.

Table 2. Bayesian mixed-model ANOVA examining hyperactivity in both ADHD and non-ADHD groups across conditions

Dependent variable	Task main effect				Group main effect				Group* task interaction effect			
	<i>F</i>	<i>p</i>	ω^2	BF ₁₀	<i>F</i>	<i>p</i>	ω^2	BF ₀₁	<i>F</i>	<i>p</i>	ω^2	BF ₀₁
Total hyperactivity score	28.78	<.001	.20	1.79 × 10 ¹⁸ ***	2.82	.10, ns	.02	1.60	.79	.48, ns	.002	18.21**

Note. ω^2 = omega-squared (small = .01, medium = .06, large = .16); *moderate = BF₀₁ < .33 or > 3.00; **strong = BF₀₁ < .10 or > 10.00; ***extreme = BF₀₁ < .01 or > 100.00.

Experimental manipulation of set shifting

As described in Irwin et al. (2019), the current study adapted the Miyake et al. (2000) global–local set shifting task for use with children. Three task variants were created to be identical in all aspects except our primary dependent variable (set shifting demands). In addition to the global–local set shifting task and two baseline (low cognitive demand) control conditions, we administered both global–global and local–local nonshifting variants to provide more precise control for both higher- and lower-order processes involved in successful performance on the global–local task. These computerized tasks use Navon (1977) figures, which feature a “global” shape (e.g., a circle) constructed using smaller, “local” figures (e.g., squares; Figure 1). To minimize memory demands, on-screen cues (“big shape”, “small shapes”) were positioned next to each quadrant (Figure 1). Sixty trials were administered following three blocks of 6–8 practice trials (100% correct required). Children responded via mouse click. Technical details regarding task parameters and administration are reported in Irwin et al. (2019). Importantly, there was strong evidence to support the integrity of the

experimental manipulation for increasing set shifting demands with this sample, including large increases in speed shift costs relative to both control conditions ($\omega^2 = .12, .14$, both $p < .001$), which did not differ ($\omega^2 = -.002, p = .62$; Irwin et al., 2019).

Set shifting condition: global–local

As shown in Figure 1, children were required to shift their response between global and local features depending on which quadrant the figures appeared (top quadrants: global; bottom quadrants: local). Trials with stimuli in the top left or bottom right quadrants involved set shifting (shift trials) because responses required a different rule than the previous trial; trials with stimuli in the top right or bottom left quadrants did not require shifting because they featured the same rule as the previous trial (nonshift trials).

Choice-response control: global–global

The global–global task was identical to the global–local task described above except that children always responded to the prepotent global figure (i.e., no explicit shifting or inhibition




Fig. 1. A sample trial from the global–local task. Children are instructed to click a response button (bottom) based on the presented stimulus (top) and rule set. Navon figures are presented sequentially in each quadrant in clockwise rotation. In this example, the Navon figure is a circle (global feature) comprised of squares (local features). Shift trials require children to inhibit the rule set from the previous trial and cognitively shift to the alternate rule set (top left and bottom right quadrants). Nonshift trials require children to apply to same rule set from the previous trial (top right and bottom left quadrants). The first trial of each task was excluded from analysis because it was neither a shift nor a nonshift trial. Figure adapted from Irwin et al. (2019).

demands). This control condition accounted for the well-established finding that children with ADHD make more errors and show slower/more variable reaction times on choice-response tasks compared to children without ADHD (Kofler et al., 2013; Klein, Wendling, Huettner, Ruder, & Peper, 2006).

Inhibition control: local–local

The local–local task was identical to the two previously described tasks except that children always responded to the local features (i.e., no explicit shifting demands). This control condition accounted for action-restraint inhibition demands required for children in the target age range to ignore a prepotent global figure and respond to the smaller (local) figures (i.e., Stroop effect; Lansbergen, Kenemans, & Van Engeland, 2007; Poirel et al., 2011).

Baseline (low cognitive demand) controls: computerized painting tasks

Children used Microsoft® Paint for five consecutive minutes at the beginning (C1) and end (C2) of both research sessions. The Paint program served as pre- and post-conditions to assess and control for potential within-day fluctuations in activity level (e.g., fatigue effects). Children sat in the same chair and interacted with the same computer used for the set

shifting tasks while interacting with a program that placed relatively modest demands on cognitive processes (i.e., the Paint program allows children to draw/paint anything they like on the monitor using a variety of interactive tools¹). Following Rapport et al. (2009), the two C1 and two C2 control conditions were separately averaged to create beginning and end of session composite scores.

Activity level

Actigraph

An actigraph is an acceleration-sensitive device that measures motor activity. The estimated reliability for actigraphs placed at the same site on the same person ranges from .90 to .99 (Tryon, Pinto, & Morrison, 1991). Basic Motionlogger® (Ambulatory Monitoring, 2004) actigraphs were used to measure children’s activity level. The acceleration-sensitive devices resemble wristwatches and were set to Proportional Integrating Measure (low-PIM) mode, which measures the intensity of movement (i.e., quantifies gross activity level). Movement was sampled 16 times per second (16 Hz) and collapsed into 1 s epochs. Children were told that the actigraphs were “special watches” that let them play the computer learning games. Observer XT (Noldus, 2012) software was used to code start and stop times for each task, which were matched to the time stamps from the actigraphs. Actigraphs were placed on the child’s nondominant wrist and both ankles (i.e., three actigraph scores per child per task).

Dependent variables

Following Rapport et al. (2009), we computed total hyperactivity scores (THS) by summing activity level across the three actigraph sites to index overall movement, separately for each of the three tasks.

Intellectual functioning (IQ) and socioeconomic status

IQ was estimated using the Wechsler Intelligence Scale for Children (WISC–V; Wechsler, 2014) Verbal Comprehension Index. Socioeconomic status (SES) was estimated using the Hollingshead (1975) scoring based on caregiver(s)’ education and occupation.

Bayesian analyses

Bayesian analyses were selected because they allow stronger conclusions by estimating the magnitude of support for both the alternative and null hypotheses (Rouder, Morey, Speckman, & Province, 2012). That is, Bayesian methods can confirm the null hypothesis rather than just fail to reject it (Wagenmakers et al., 2016b). Bayes factor mixed-model ANOVAs with JZS default prior scales (Rouder et al., 2012; Wagenmakers et al., 2016a) were conducted using JASP 0.8.5 (JASP Team, 2017). Instead of a *p*-value, these

¹As noted by Rapport et al. (2009), successful interaction with the Paint program requires some cognitive processes such as focused attention and interaction with long-term memory, as well as limited short-term storage/rehearsal processes. As such, we refer to these conditions as having “low” rather than “no” cognitive demands.

analyses provide BF_{10} , which is the Bayes factor of the alternative hypothesis (H_1) against the null hypothesis (H_0). BF_{10} is an odds ratio, where values above 3.0 are considered moderate evidence supporting the alternative hypothesis (conceptual equivalent of $p < .05$). BF_{10} values above 10 are considered strong (>30 = very strong, >100 = decisive/extreme support; Wagenmakers et al., 2016b). A finding of $BF_{10}=10$, for example, would indicate that the data are 10 times more likely under the alternative hypothesis than under the null hypothesis (i.e., strong support for an effect).

Conversely, BF_{01} is the Bayes factor of the null hypothesis (H_0) against the alternative hypothesis (H_1). BF_{01} is the inverse of BF_{10} (i.e., $BF_{01}=1/BF_{10}$) and is reported when the evidence indicates a lack of an effect (i.e., favors the null hypothesis; Rouder et al., 2012). BF_{01} values are interpreted identically to BF_{10} (>3 = moderate, >10 = strong, >100 = decisive/extreme support for the null hypothesis that the ADHD and non-ADHD groups are *equivalent* on an outcome; Rouder et al., 2012).

Notably, interpretation of results was unchanged when examining frequentist p -values instead of Bayes factors, except that nonsignificant p -values cannot be interpreted as evidence of between-group equivalence.

Data analysis overview

The evidence for a functional role of set shifting in ADHD-related hyperactivity was evaluated via a 2 (group: ADHD, non-ADHD) \times 5 (task: C1 paint, global–global, local–local, global–local, C2 paint) Bayesian mixed-model ANOVA for actigraph-measured THS. A main effect of group in the absence of an interaction would indicate that any between-group differences in activity level during the set shifting condition cannot be specifically attributable to set shifting. A significant group \times task interaction with *post hoc* tests indicating disproportionate increases in activity level in the ADHD group during the shift versus both nonshift cognitive task conditions would provide support for a causal role of set shifting demands in ADHD-related hyperactivity. Similarly, *post hoc* tests for this interaction indicating disproportionate hyperactivity increases for the ADHD group during the shift and nonshift cognitive tasks (but not the set shifting task specifically) would indicate that ADHD-related hyperactivity is caused at least in part by general cognitive demands rather than by shifting demands specifically. Finally, in the absence of an interaction, a main effect of task would provide preliminary support for a functional relation between hyperactivity and general cognitive, inhibition, and/or set shifting demands, depending on the pattern of *post hoc* results. Specifically, significant hyperactivity increases during the shift and nonshift cognitive tasks relative to the baseline painting tasks would indicate a functional link between cognitive demands in general and children's activity level. An additional increase in hyperactivity between the choice-response control and inhibition control tasks would provide strong evidence for a functional relation between inhibition and children's activity level, whereas an

additional increase from the choice-response and inhibition control tasks to the set shifting task would support a functional relation between hyperactivity and set shifting specifically (albeit not ADHD-related hyperactivity specifically).

RESULTS

Preliminary analyses

Outliers beyond 3.00 SD were winsorized relative to the within-group distribution (ADHD, non-ADHD). This process affected 1.5% (ADHD group) to 0.6% (non-ADHD group) of all data points. Task performance data for the current sample was previously reported in Irwin et al. (2019). Hyperactivity data during these tasks have not been reported previously for any children in the current sample. There was no significant evidence to support ADHD versus non-ADHD group differences in age ($BF_{01}=2.75$, $p=.21$), gender ($BF_{01}=2.29$, $p=.69$), or SES ($BF_{01}=2.10$, $p=.14$). All parent and teacher ADHD symptom ratings were higher for the ADHD than non-ADHD group as expected (all $BF_{10}>8.00$, all $p<.01$; Table 1). There was moderate evidence suggesting a difference in IQ ($BF_{10}=4.13$, $p=.01$). However, IQ was not included as a covariate based on compelling statistical, methodological, and conceptual rationale against covarying IQ when investigating cognitive processes in ADHD (Dennis et al., 2009; Irwin et al., 2019).

Experimental manipulation of set shifting demands

Results of the 2 (group: ADHD versus Non-ADHD) \times 5 (task: C1 paint, global–global, local–local, global–local, C2 paint) Bayesian mixed-model ANOVA provided decisive support for a main effect of task ($BF_{10}=1.79 \times 10^{18}$, $p<.001$, $\omega^2=.20$) but no evidence to support a main effect of group ($BF_{01}=1.60$, $p=.10$, $\omega^2=.02$; Figure 2). *Post hoc* comparisons for the main effect of task (Table 2) indicated that children's activity level significantly increased from both baseline control conditions (i.e., paint) to all three cognitive task conditions (i.e., choice response, inhibitory control, set shifting; all $BF_{10}>9.85$, all $p<.01$, all $d=.47$ – 1.15), indicating that children upregulated their physical movement in response to the general cognitive demands common across the shift and nonshift cognitive tasks. However, children's activity level did not increase significantly during the inhibition control (i.e., local–local) as compared to the choice-response control condition (i.e., global–global; $BF_{01}=1.30$, $p=.05$, $d=-.25$), indicating that children's physical activity level was not significantly affected when inhibitory control demands were evoked.² In contrast, there was decisive evidence indicating an additional increase between the inhibition control

²As shown in Figure 2, the null hypothesis testing 'trend' toward differences between the choice response and inhibitory control conditions reflects an atheoretical decrease rather than increase in movement when inhibitory control demands are evoked. A one-tailed directional test of the hypothesis that inhibition demands evoke greater levels of hyperactivity provided strong evidence against a role of inhibitory control demands for increasing children's physical movement/hyperactivity ($BF_{01}=22.53$, $p=.97$).




Fig. 2. Graph depicting group mean differences in total hyperactivity scores (THS) across the five task conditions. Error bars reflect 95% confidence intervals. The unit of measure on the y-axis is called Proportional Integrating Measure (PIM) and is a measure of movement intensity.

and set shifting conditions ($BF_{10} = 20543.66$, $p < .001$, $d = .49$), indicating that children specifically upregulated their motor movement in response to set shifting demands in their environment. This increase was also significant between the choice-response control and set shifting conditions based on null hypothesis testing but failed to demonstrate sufficient Bayesian evidentiary value ($BF_{10} = 2.65$, $p = .02$, $d = .29$).

Importantly, there was strong evidence *against* the group \times task interaction ($BF_{01} = 18.21$, $p = .48$, $\omega^2 = .002$; Table 2). This finding indicated that the functional links between hyperactivity and both general cognitive demands and specific set shifting demands occurred equivalently for children with ADHD and without ADHD. In other words, all children upregulate their physical activity in response to environmental demands that challenge their cognitive/set shifting abilities. However, set shifting, inhibitory control, and general cognitive demands in the environment do not appear to be viable explanations for ADHD-related hyperactivity because these increases occurred equivalently for children with and without ADHD in response to both general cognitive demands and specific cognitive processes required to shift between rule sets.

DISCUSSION

The current study was the first to test for a causal role of set shifting demands on ADHD-related hyperactivity by combining a validated, experimental manipulation of set shifting (Irwin et al., 2019) with concurrent and objective measurement of hyperactivity (actigraphy) in a carefully phenotyped sample of children with and without ADHD. Results from the current study indicate that hyperactivity in children is functionally related to cognitive demands in general and set shifting demands specifically while providing strong evidence *against* a functional link between inhibitory control and increased hyperactivity in children (Alderson et al., 2012).

Importantly, the current results indicate a functional if not causal role of set shifting demands on children's physical movement, while providing strong evidence that set shifting demands cannot explain hyperactivity in ADHD given that children with and without ADHD increased their physical activity equivalently when set shifting demands were experimentally evoked.

The current findings provided strong evidence *against* our hypothesis that hyperactivity in ADHD would be functionally related to set shifting demands. Rather, the findings provided decisive evidence to indicate that children with and without ADHD increase their activity level equivalently in response to set shifting demands. The use of Bayesian statistics allowed stronger conclusions by providing significant support for the null hypothesis rather than just failing to reject it. These findings suggest that set shifting is not a viable explanation for ADHD-related hyperactivity and help to clarify the limited, mixed evidence by indicating that hyperactivity in ADHD is not linked specifically to set shifting demands (McCandless & O'Laughlin, 2007; Toplak et al., 2008). The incongruence between our *a priori* hypothesis and our findings may be because the former was developed from mixed findings across correlational studies rather than from experimentally manipulated cognitive demands and objectively measured behavior. Additionally, these findings coincide with recent evidence, based on part on the current sample, indicating that children with ADHD likely do not have unique set shifting deficits (Irwin et al., 2019), and that set shifting abilities do not covary with ADHD-related hyperactive/impulsive symptom severity (Kofler, Irwin et al., 2019).

Although of secondary interest in the current study, the current findings were consistent with recent experimental evidence indicating that inhibitory control demands do not evoke hyperactive behavior in children with or without ADHD (Alderson et al., 2012). That is, there was strong

evidence *against* an incremental increase in children's physical activity from the choice-response control (i.e., global–global task) to the inhibition control (i.e., local–local task) condition and as such increasing inhibition demands did not increase hyperactivity in children. This pattern of results is consistent with the only other experimental study to date examining the functional role of inhibition on hyperactivity (Alderson et al., 2012) and extends previous findings by examining this pattern using an experimental manipulation of general cognitive demands, inhibition demands, and set shifting demands within a single study.

Despite clear evidence linking hyperactivity in ADHD with cognitive demands in the environment (for meta-analytic review, see Kofler et al., 2016), it was unclear whether hyperactivity in ADHD is evoked or exacerbated by any cognitive demands or whether this occurs specifically in response to specific executive function abilities. The current study helps clarify this emerging literature, while providing evidence for both possibilities. That is, the current findings indicate that children increase their activity level both in response to general cognitive demands as well as specifically in response to set shifting demands. Taken together, the literature at this time appears to indicate that of the three core executive functions (Miyake et al., 2000), inhibition demands do not cause children to upregulate their physical activity (current study; Alderson et al., 2012), whereas working memory (e.g., Kofler et al., 2015) and set shifting demands (current study) both evoke hyperactive behavior in children. However, it appears that only working memory does so disproportionately for children with ADHD (Hudec et al., 2015; Patros et al., 2017; Rapport et al., 2009). In other words, there is consistent evidence that children become more physically active when faced with cognitively challenging tasks (Kofler et al., 2016), and emerging evidence that these increases may be linked specifically to two of the three core executive functions (i.e., working memory and set shifting). However, as shown in the current study, the functional link between set shifting and increased physical movement/hyperactivity is not likely to be a viable explanation for ADHD-related hyperactivity because these demands do not disproportionately affect children with ADHD. Taken together, the most parsimonious conclusion appears to be that set shifting affects hyperactivity in children over and above the effects of general cognitive demands, but cannot specifically explain why hyperactivity in children with ADHD appears to occur primarily in the context of environmental demands that challenge them cognitively (Kofler et al., 2016).

Limitations

Despite the experimental methodology and carefully phenotyped sample, the following limitations must be considered when interpreting results. Children with all ADHD current presentation specifiers were included in this sample based on meta-analytic evidence that children with

ADHD inattentive and combined/hyperactive subtypes/current presentations do not differ in terms of actigraph-measured movement despite both subtypes/specifiers demonstrating elevated activity level versus neurotypical and clinical controls at the group and individual levels (Kofler et al., 2016). However, meta-analytic evidence suggests that ADHD symptom domains may differentially relate to neurocognitive functions and as such future work should collect larger samples of each ADHD presentation to investigate whether associations with hyperactivity remain the same or differ as a function of ADHD presentation (Willcutt et al., 2012).

Despite the use of a carefully phenotyped sample and strong evidence for group differences in parent and teacher ratings of ADHD hyperactivity/impulsivity symptoms, there was no evidence to support a main effect of group differences in objectively assessed hyperactivity across the five experimental conditions. Although this finding was initially surprising, on close inspection it is consistent with meta-analytic findings indicating minimal differences between ADHD and non-ADHD groups during tasks with relatively low cognitive demands ($d = .36$; Kofler et al., 2016). Thus, it may be that the set shifting task, despite being cognitively challenging enough to evoke higher levels of hyperactivity versus the cognitive task controls, does not place sufficient demands on the specific cognitive processes known to disproportionately give rise to hyperactive behavior in ADHD (e.g., working memory; Hudec et al., 2015; Kofler et al., 2015; Patros et al., 2017; Rapport et al., 2009). Additional research is needed to test this hypothesis, as to our knowledge no study to date has concurrently examined hyperactivity across conditions that systematically differ in all three of the core executive functions.

Finally, the current study evoked set shifting demands and observed effects on children's hyperactive behavior. However, further clarifying the directionality of the set shifting/hyperactivity relation will require the opposite, namely, experimentally manipulating children's activity level while concurrently measuring children's set shifting performance. This line of work will be critical given evidence that increasing physical movement may facilitate performance on cognitive tests for children with ADHD (Hartanto, Krafft, Iosif, & Schweitzer, 2016; Sarver, Rapport, Kofler, Raiker, & Friedman, 2015), presumably because the physical activity serves as a physiological arousal mechanism that actuates cortical networks implicated in attention and executive function (Rapport et al., 2009). Thus, despite the evidence for a functional relation detected herein, more work is needed to conclude that hyperactivity serves as a compensatory mechanism that actuates and augments cognitive functioning for children with ADHD while they are engaged in tasks that tax those underdeveloped cognitive processes (Rapport et al., 2009).

Clinical and research implications

The current results indicate a functional relation between set shifting and hyperactivity, while also providing evidence

against set shifting as a viable cause of hyperactive behavior in children with ADHD. Given the additional finding that general cognitive demands evoked higher levels of hyperactive behavior for both groups, it will be critical for future studies to include cognitive task control conditions to ensure specificity in their results. For example, if we had not included the choice-response control task, we would likely have erroneously concluded that there was a functional link between inhibitory control and hyperactivity. However, because we included a choice-response task with minimal inhibition demands, we were able to see that the increased hyperactivity during the inhibition task relative to baseline was not attributable specifically to the task's inhibition demands (Alderson et al., 2012). Moreover, additional work is needed to test our hypothesis above that only working memory appears to disproportionately evoke hyperactivity in ADHD because to our knowledge no ADHD hyperactivity studies have included manipulations of all three core executive functions in a single experiment (working memory, inhibitory control, set shifting; Karr et al., 2018).

ACKNOWLEDGMENTS

This work was supported in part by an NIH grant (R34 MH102499-01, PI: Kofler). The sponsor had no role in design and conduct of the study; collection, management, analysis, and interpretation of the data; or preparation, review, or approval of the manuscript.

CONFLICT OF INTEREST

The authors have no conflicts of interest to report.

References

- Alderson, R.M., Rapport, M.D., Kasper, L.J., Sarver, D.E., & Kofler, M.J. (2012). Hyperactivity in boys with attention deficit/hyperactivity disorder (ADHD): The association between deficient behavioral inhibition, attentional processes, and objectively measured activity. *Child Neuropsychology*, 18(5), 487–505.
- Ambulatory Monitoring. (2004). *The Micromini Motionlogger Actigraph and Family of Recorders user's Manual* (3rd ed.). New York, NY: Author.
- American Psychiatric Association. (2013). *Diagnostic and Statistical Manual of Mental Disorders (DSM-5®)*. American Psychiatric Pub.
- Barkley, R.A. (1997). Behavioral inhibition, sustained attention, and executive functions: Constructing a unifying theory of ADHD. *Psychological Bulletin*, 121, 65–94.
- Chacko, A., Kofler, M., & Jarrett, M. (2014). Improving outcomes for youth with ADHD: A conceptual framework for combined neurocognitive and skill-based treatment approaches. *Clinical Child and Family Psychology Review*, 17(4), 368–384.
- Dennis, M., Francis, D.J., Cirino, P.T., Schachar, R., Barnes, M. A., & Fletcher, J. M. (2009). Why IQ is not a covariate in cognitive studies of neurodevelopmental disorders. *Journal of the International Neuropsychological Society*, 15(3), 331–343.
- Gomez, R., & Sanson, A. (1994). Effects of experimenter and mother presence on the attentional performance and activity of hyperactive boys. *Journal of Abnormal Child Psychology*, 22, 517–529.
- Hartanto, T.A., Krafft, C.E., Iosif, A.M., & Schweitzer, J.B. (2016). A trial-by-trial analysis reveals more intense physical activity is associated with better cognitive control performance in attention-deficit/hyperactivity disorder. *Child Neuropsychology*, 22(5), 618–626.
- Hollingshead, A.B. (1975). *Four Factor Index of Social Status*. New Haven, CT: Yale.
- Hudec, K.L., Alderson, R.M., Patros, C.H., Lea, S.E., Tarle, S.J., & Kasper, L.J. (2015). Hyperactivity in boys with attention-deficit/hyperactivity disorder (ADHD): The role of executive and non-executive functions. *Research in Developmental Disabilities*, 45, 103–109.
- Irwin, L.N., Kofler, M.J., Soto, E.F., & Groves, N.B. (2019). Do children with attention-deficit/hyperactivity disorder (ADHD) have set shifting deficits?. *Neuropsychology*, 33(4), 470.
- JASP Team. (2017). JASP. Version 0.8.1. [computer software]
- Karr, J.E., Areshenkoff, C.N., Rast, P., Hofer, S.M., Iverson, G.L., & Garcia-Barrera, M.A. (2018). The unity and diversity of executive functions: A systematic review and re-analysis of latent variable studies. *Psychological Bulletin*, 144(11), 1147.
- Kasper, L.J., Alderson, R.M., & Hudec, K.L. (2012). Moderators of working memory deficits in children with attention-deficit/hyperactivity disorder (ADHD): A meta-analytic review. *Clinical Psychology Review*, 32(7), 605–617.
- Klein, C., Wendling, K., Huettner, P., Ruder, H., & Peper, M. (2006). Intra-subject variability in attention-deficit hyperactivity disorder. *Biological Psychiatry*, 60(10), 1088–1097.
- Kofler, M.J., Irwin, L.N., Soto, E.F., Groves, N.B., Harmon, S.L., & Sarver, D.E. (2019). Executive functioning heterogeneity in pediatric ADHD. *Journal of Abnormal Child Psychology*, 47(2), 273–286.
- Kofler, M.J., Raiker, J.S., Sarver, D.E., Wells, E.L., & Soto, E.F. (2016). Is hyperactivity ubiquitous in ADHD or dependent on environmental demands? Evidence from meta analysis. *Clinical Psychology Review*, 46, 12–24.
- Kofler, M.J., Rapport, M.D., Sarver, D.E., Raiker, J.S., Orban, S.A., Friedman, L.M., & Kolomeyer, E.G. (2013). Reaction time variability in ADHD: a meta-analytic review of 319 studies. *Clinical Psychology Review*, 33(6), 795–811.
- Kofler, M.J., Sarver, D.E., & Wells, E.L. (2015). Working memory and increased activity level (hyperactivity) in ADHD: Experimental evidence for a functional relation. *Journal of Attention Disorders*, 1087054715608439.
- Kofler, M.J., Spiegel, J.A., Austin, K.E., Irwin, L.N., Soto, E.F., & Sarver, D.E. (2018). Are episodic buffer processes intact in ADHD? Experimental evidence and linkage with hyperactive behavior. *Journal of Abnormal Child Psychology*, 46(6), 1171–1185.
- Lansbergen, M.M., Kenemans, J.L., & Van Engeland, H. (2007). Stroop interference and attention-deficit/hyperactivity disorder: A review and meta-analysis. *Neuropsychology*, 21(2), 251.
- Mahone, E.M. (2011). ADHD: Volumetry, motor, and oculomotor functions. In *Behavioral Neuroscience of Attention Deficit Hyperactivity Disorder and Its Treatment* (pp. 17–47). Berlin, Heidelberg: Springer.
- Mahone, E. M. & Hoffman, J. (2007). Behavior ratings of executive function among preschoolers with ADHD. *The Clinical Neuropsychologist*, 21(4), 569–586.
- McCandless, S. & O'Laughlin, L. (2007). The clinical utility of the Behavior Rating Inventory of Executive Function (BRIEF) in

- the diagnosis of ADHD. *Journal of Attention Disorders*, 10(4), 381–389.
- Miyake, A., Friedman, N.P., Emerson, M. J., Witzki, A. H., Howerter, A., & Wager, T. D. (2000). The unity and diversity of executive functions and their contributions to complex “frontal lobe” tasks: A latent variable analysis. *Cognitive Psychology*, 41, 49–100.
- Navon, D. (1977). Forest before trees: The precedence of global features in visual perception. *Cognitive Psychology*, 9(3), 353–383.
- Noldus Information Technology. (2012). *The Observer XT Reference Manual*. Wageningen, The Netherlands: Author.
- Patros, C.H., Alderson, R.M., Hudec, K.L., Tarle, S.J., & Lea, S.E. (2017). Hyperactivity in boys with attention-deficit/hyperactivity disorder: The influence of underlying visuospatial working memory and self-control processes. *Journal of Experimental Child Psychology*, 154, 1–12.
- Poirel, N., Simon, G., Cassotti, M., Leroux, G., Perchey, G., Lanoe, C., . . . Houde, O. (2011). The shift from local to global visual processing in 6-year-old children is associated with grey matter loss. *PLoS ONE* 6(6), e20879.
- Polanczyk, G., de Lima, M.S., Horta, B.L., Biederman, J., & Rohde, L.A. (2007). The worldwide prevalence of ADHD: A systematic review and metaregression analysis. *American Journal of Psychiatry*, 164(6), 942–948.
- Polanczyk, G.V., Willcutt, E.G., Salum, G.A., Kieling, C., & Rohde, L.A. (2014). ADHD prevalence estimates across three decades: an updated systematic review and meta-regression analysis. *International Journal of Epidemiology*, 43(2), 434–442.
- Rapport, M.D., Bolden, J., Kofler, M.J., Sarver, D.E., Raiker, J.S., & Alderson, R.M. (2009). Hyperactivity in boys with attention-deficit/hyperactivity disorder (ADHD): A ubiquitous core symptom or manifestation of working memory deficits?. *Journal of Abnormal Child Psychology*, 37(4), 521–534.
- Rouder, J.N., Morey, R.D., Speckman, P.L., & Province, J.M. (2012). Default Bayes factors for ANOVA designs. *Journal of Mathematical Psychology*, 56(5), 356–374.
- Sarver, D.E., Rapport, M.D., Kofler, M.J., Raiker, J.S., & Friedman, L.M. (2015). Hyperactivity in attention-deficit/hyperactivity disorder (ADHD): Impairing deficit or compensatory behavior?. *Journal of Abnormal Child Psychology*, 43(7), 1219–1232.
- Sonuga-Barke, E., Bitsakou, P., & Thompson, M. (2010). Beyond the dual pathway model: Evidence for the dissociation of timing, inhibitory, and delay-related impairments in attention-deficit/hyperactivity disorder. *Journal of the American Academy of Child & Adolescent Psychiatry*, 49, 345–355.
- Toplak, M.E., Bucciarelli, S.M., Jain, U., & Tannock, R. (2008). Executive functions: Performance-based measures and the behavior rating inventory of executive function (BRIEF) in adolescents with attention deficit/hyperactivity disorder (ADHD). *Child Neuropsychology*, 15(1), 53–72.
- Tryon, W.W., Pinto, L.P., & Morrison, D.F. (1991). Reliability assessment of pedometer activity measurements. *Journal of Psychopathology and Behavioral Assessment*, 13(1), 27–44.
- Wagenmakers, E.J., Love, J., Marsman, M., Jamil, T., Ly, A., Verhagen, J., . . . Meerhoff, F. (2016a). Bayesian inference for psychology. part II: Example applications with JASP. *Psychonomic Bulletin & Review*, 1–19.
- Wagenmakers, E.J., Morey, R.D., & Lee, M.D. (2016b). Bayesian benefits for the pragmatic researcher. *Current Directions in Psychological Science*, 25(3), 169–176.
- Wechsler, D. (2014). *Wechsler Intelligence Scale for Children-Fourth/Fifth Edition*. San Antonio: Pearson.
- Willcutt, E.G., Nigg, J.T., Pennington, B.F., Solanto, M.V., Rohde, L.A., Tannock, R., . . . Lahey, B.B. (2012). Validity of DSM-IV attention deficit/hyperactivity disorder symptom dimensions and subtypes. *Journal of Abnormal Psychology*, 121(4), 991.