

## Regular Article

# Adverse childhood experiences and transcriptional response in school-age children

A. Marie-Mitchell<sup>1</sup>  and S.W. Cole<sup>2</sup>

<sup>1</sup>Preventive Medicine & Pediatrics Departments, Loma Linda University, Loma Linda, CA, USA and <sup>2</sup>Department of Psychiatry and Behavioral Sciences & Department of Medicine, Division of Hematology-Oncology, UCLA School of Medicine, Los Angeles, CA, USA

### Abstract

This study evaluated whether children with higher adverse childhood experiences (ACE) scores had alterations in immune cell gene expression profiles. RNA sequencing was conducted on dried blood spot samples from 37 generally healthy English-speaking children (age 5–11) who were recruited from well-child visits at a university-affiliated pediatric practice. The Whole Child Assessment was used to assess ACE exposure. Primary analyses examined an a priori-specified composite of 19 pro-inflammatory gene transcripts. Secondary analyses examined a 34-gene composite assessing Type I interferon response, and used Transcript Origin Analyses to identify cellular mechanisms. After controlling for age, body mass index percentile, sex, race/ethnicity, current insurance status, and household smoking exposure, pro-inflammatory gene expression was elevated by 0.094 log<sub>2</sub> RNA expression units with each Child-ACE total score point ( $p = .019$ ). Type I interferon gene expression was similarly upregulated (0.103;  $p = .008$ ). Transcript origin analyses implicated CD8+ T cell as the primary sources of gene transcripts upregulated, and nonclassical (CD16+) monocytes as sources of downregulated transcripts. These preliminary analyses suggest that parent-reported ACE exposures are associated with increased expression of both inflammatory and interferon gene transcripts in children's circulating blood cells.

**Keywords:** adverse childhood experiences, adversity, biomarkers, immunology, inflammation

(Received 5 February 2020; revised 4 July 2020; accepted 13 October 2020; First Published online 28 January 2021)

Adverse childhood experiences (ACEs) include child maltreatment, domestic violence, substance abuse, mental illness, incarceration, and single parenting (Felitti et al., 1998). Extensive research demonstrates that ACEs double risk for chronic diseases in adults such as heart disease, cancer, liver disease, lung disease, and autoimmune diseases, and quadruple risk for mental health problems including suicide, alcoholism, illicit drug use, depression, and dementia (Anda et al., 2006; Brown et al., 2009; Felitti et al., 1998). While there is evidence that ACEs are linked to adult pathology through adoption of unhealthy lifestyles, there is also evidence that ACEs are associated with physiologic changes even in the presence of healthy lifestyles (Shonkoff et al., 2012). These physiologic changes include activation of the hypothalamus–pituitary–adrenal (HPA) axis and sympathetic nervous system (SNS) with co-occurring alterations in immune system function (Chiang, Taylor, & Bower, 2015; Repetti, Taylor, & Seeman, 2002; Shonkoff et al., 2012).

Innate (or natural) immunity is the first line of defense against injury, infection, or necrosis. Harmful stimuli are eliminated and tissue repair is initiated through inflammation which involves granulocytes, including neutrophils, and macrophages, and

the release of communication molecules called cytokines (Segerstrom & Miller, 2004). Adaptive (or acquired) immunity is a slower but more specific response that involves three-types of lymphocytes: T helper (CD4+) cells, T cytotoxic (CD8+) cells, and B cells. T helper cells release cytokines that amplify the activity of CD8+ T cells and B cells (Segerstrom & Miller, 2004). One subset of cytokines known as Type I interferons play a key role in protecting cells against viral infections (Trinchieri, 2010). Prolonged stress can stimulate immune cell production of the cytokines involved in inflammation while reducing the expression of Type I interferons and cytokines that support B cell antibody production, such that there is an increase in vulnerability to both infection and neoplastic disease, as well as autoimmune and allergic disease (Irwin & Cole, 2011; Miller, Chen, & Cole, 2009). Inflammation is protective as an acute response, but when prolonged can become a toxic precursor to variety of diseases including depression, dementia, diabetes, cardiovascular disease, and some cancers (Murphy, Slavich, Chen, & Miller, 2015).

The psychological stress response begins with a perception of threat or uncertainty by the central nervous system that triggers the HPA axis and/or SNS, and results in a transient reconfiguration of the immune system that involves both inflammation and suppression of some adaptive immune responses (Adamao, 2014). A broad literature demonstrates that stressors activate proinflammatory pathways, including laboratory studies of stressors like giving a speech or arguing with a spouse, and naturalistic

**Author for Correspondence:** Ariane Marie-Mitchell, 24785 Stewart St, Suite 206, Loma Linda, CA 92354, USA; E-mail [amariemitchell@llu.edu](mailto:amariemitchell@llu.edu)

**Cite this article:** Marie-Mitchell A, Cole SW (2022). Adverse childhood experiences and transcriptional response in school-age children. *Development and Psychopathology* 34: 875–881. <https://doi.org/10.1017/S095457942000187X>

studies of social difficulties, loneliness, and targeted rejection (Murphy et al., 2015; Murphy, Slavich, Rohleder, & Miller, 2013). The presence of a proinflammatory phenotype in response to early-life stressors has been observed in infants (David, Measelle, Ostlund, & Ablow, 2017; Jiang et al., 2017; Measelle & Ablow, 2018), young children (Cicchetti, Handley, & Rogosch, 2015; Slopen, Kubzansky, McLaughlin, & Koenen, 2013; Tyrka, Parade, Valentine, Eslinger, & Seifer, 2015), and adolescents (Chiang et al., 2019; Miller & Chen, 2010; Miller & Cole, 2012). Several studies have demonstrated that adults with a history of child maltreatment or other ACEs have increased levels of inflammation as shown by elevations in white blood cell counts and circulating proinflammatory cytokines (Danese et al., 2009; Danese, Pariante, Caspi, Taylor, & Poulton, 2007; Howren, Lamkin, & Suls, 2009; Surtees et al., 2003). Studies of chronic stressors (such as family stressors and trauma) suggest that inflammation persists, but is accompanied by alterations in cytokines and reductions in lymphocyte proliferation (Ehrlich, Miller, & Chen, 2015; Segerstrom & Miller, 2004).

Studies of gene expression offer a complementary approach to examining the impact of stress on cellular processes. “Gene expression” refers to the transcription of a gene’s DNA blueprint into RNA molecules that subsequently serve as the template for producing the biologically active proteins that mediate cell function (including the cytokines noted above) (Ehrlich et al., 2015; Irwin & Cole, 2011; Segerstrom & Miller, 2004). Gene expression studies suggest the presence of a common biological pathway that is activated in response to stressful social environments and can increase risk for various chronic diseases (Cole, 2013, 2014; Schwaiger et al., 2016). This common biological pathway is called the conserved transcriptional response to adversity (CTRA) and refers to SNS signaling that upregulates gene expression of proinflammatory cytokines, downregulates gene expression for Type 1 interferons, and stimulates hematopoietic output of classical monocytes (Cole, 2019; Schwaiger et al., 2016). CTRA profiles in blood are predictive of inflammation-related disease outcomes in cancer and general chronic disease (Antoni et al., 2016; Knight et al., 2016, 2019; Simons et al., 2017), and complement blood cytokine measures by quantifying cellular mediators of inflammation that traffic through blood to diseased tissues in atherosclerosis, cancer, infections, and other localized disease processes (Irwin & Cole, 2011; Miller et al., 2009).

Studies of loneliness, poverty, bereavement, and chronic stress have all shown activation of the CTRA, which provides a biological mechanism linking adverse social environments with the development of inflammation-related diseases and reductions in antiviral defenses (Cole, 2019). Two studies of pediatric asthma patients demonstrated CTRA activation in response to targeted rejection or harsh parent–child conflict (Ehrlich et al., 2015; Murphy et al., 2015). One study of adolescents at risk for depression showed increased transcription factor activation related to CTRA (Murphy et al., 2013). These studies suggest that CTRA activation may be a key component to biological mechanisms linking ACEs to poor health outcomes, but in order to translate these findings into preventive interventions, studies are needed on correlations with the assessment of ACEs and potential differences in biological response to ACEs by age group. Studying interferon and cell type profiles is particularly important to providing a more comprehensive understanding of immune reconfiguration, beyond the inflammatory response that is observed in response to both acute and chronic stressors. Evaluations of the immune system can be done at different levels, from gene expression to

circulating cell counts. Studies of gene expression have the advantage of avoiding some of the challenges of interpreting circulating cell counts (Zhoua, Fragalaa, McElhaney, & Kuchel, 2010), and the added benefit of being feasible based upon the minimally invasive approach of dried blood spot (DBS) sampling (McDade, Williams, & Snodgrass, 2007).

The purpose of this study was to test the hypothesis that children with higher ACE scores would have alterations in blood gene expression profiles that resemble the CTRA in showing increased expression of pro-inflammatory genes. As exploratory hypotheses, we also tested for differential expression of Type I interferon genes due to their typical downregulation in the adult CTRA profile, and examined potential cellular origins of differences in gene expression. If children with higher ACE scores do have alterations in immune cell gene expression profiles, then these alterations could serve as a predictive biomarker of risk for adult pathology and an outcome biomarker to assess the potential impact of pediatric interventions.

## Method

### *Participants and recruitment*

Participants were recruited to attend a single follow-up research visit after receiving well-child care at one of two university-affiliated pediatric primary care practices. Pediatric patients were eligible for the research study if they were age 5–11 years old, English-speaking, and generally healthy based upon medical record review. Patients were excluded if they had a neurodevelopmental disability that would prohibit their participation in psychological testing or if they had a sibling already enrolled in the study. All eligible patients were mailed an information letter about the research study and called by phone within 1–2 weeks after their well-child visit. A total of 131 eligible patients were successfully contacted by phone, of whom 100 agreed to participate. The study was powered to detect the direct effect of ACEs on the total Pediatric Symptom Checklist score, a measure of psychosocial dysfunction that correlates with common mental disorders (target  $N = 89$ ) (Gardner, Lucas, Kolko, & Campo, 2007; Jellinek et al., 1999; Murphy, Reed, Jellinek, & Bishop, 1992). The collection of DBS samples was added as a secondary measure about halfway through the study, resulting in a total of 40 samples. The Institutional Review Board at our university approved this research study (IRB# 5140273).

### *Adversity measure*

The Whole Child Assessment (WCA) was used to assess child adversity at the research visit (Marie-Mitchell et al., 2019; Marie-Mitchell, Watkins, Copado, & Distelberg, 2020). Ten questions were about child exposure to ACEs (Child-ACEs) based upon the categories described in the original ACE study, including child physical abuse, child sexual abuse, child verbal abuse, child physical neglect, child emotional neglect, domestic violence, substance abuse by a household member, mental illness of a household member, incarceration of a household member, and single parenting (Felitti et al., 1998). Six questions were about risk of Child-ACEs, including risk of emotional abuse, physical abuse, caregiver mental illness, caregiver alcohol abuse, and family dysfunction (Centers for Disease Control and Prevention & National Center for Chronic Disease Prevention and Health Promotion, 2009; Dubowitz, Lane, Semiatin, & Magder, 2012;

Feigelman, Dubowitz, Lane, Grube, & Kim, 2011; Kroenke, Spitzer, & Williams, 2003; Kroenke, Spitzer, Williams, Monahan, & Löwe, 2007; Taj, Devera-Sales, & Vinson, 1998). Although most ACE scales count exposure to adversity only, the WCA includes questions about risk of adversity in order to both increase reporting of family stressors and opportunities for pediatric providers to prevent ACEs. Prior research suggests that adding questions about risk to the Child-ACE total score increases associations with child health outcomes (Marie-Mitchell et al., 2020). The Child-ACE exposure score was the sum of the 10 exposure questions, and the Child-ACE total score was the sum of the 16 questions about exposure and risk.

At the end of the research visit, research assistants reviewed questionnaires for caregiver disclosure of domestic violence, child maltreatment, or risk of child maltreatment. Further history was obtained to assess current safety and community resources were provided when indicated. Research assistants were prepared to report new cases of maltreatment to Child Protective Services, but all cases in this sample were previously reported.

### Gene expression measure

Caregivers were notified that children could not participate in the research visit if they had been sick in the last 3 days, and if necessary rescheduled. In addition, after providing written consent and prior to beginning the research protocol, caregivers were asked whether the child had a fever in the last 3 days and whether they had used any medication within the last 24 hours. All participants responded negatively to these screening questions and all children had normal vital signs.

A trained research assistant collected DBS samples using a sterile disposable lancet and filter paper cards (Whatman #903, GE Healthcare, Piscataway, NJ) (McDade et al., 2007). The cards were allowed to dry at room temperature and then stored in a freezer until batch processing. RNA was extracted from 40 available DBS samples and subject to genome-wide transcriptional profiling by RNA sequencing as previously described (Ross, Carroll, Dunkel Schetter, Hobel, & Cole, 2019; Ross, Cole, Carroll, & Dunkel Schetter, 2019). Briefly, total RNA was extracted (Qiagen RNeasy) and polyadenylated RNA was converted to cDNA (Lexogen QuantSeq 3' FWD with low mass buffer) and sequenced on an Illumina HiSeq4000 instrument in the UCLA Neuroscience Genomics Core Laboratory, all following the manufacturers' standard protocols for low-mass samples. Samples yielded an average of 14.3 million sequencing reads, each of which was mapped to the reference human transcriptome using the STAR aligner (94% average mapping rate) (Dobin et al., 2013). The total number of reads for each human gene was normalized to transcripts per million total mapped reads, floored at 1 read per million to suppress spurious variability, and log<sub>2</sub> transformed for analysis by linear statistical models as described below. Three samples failed to pass standard endpoint quality control criteria for DBS RNA sequencing (mean profile correlation with each other sample >.55) and were omitted from subsequent analyses, leaving a total of 37 RNA samples for analysis.

### Statistical analysis

For the purpose of describing the study participants only, the sample was divided into those with a lower Child-ACE score (0–1) and higher Child-ACE score (2+) which categorization is consistent with our prior exploration of clinical outcomes

associated with Child-ACE scores (Marie-Mitchell et al., 2020). The Mann–Whitney test was used to compare continuous variables and the Fisher's Exact test was used to compare categorical variables.

Primary analyses examined an a priori-specified 19-gene composite of pro-inflammatory gene transcripts (e.g., Interleukin-1B (IL1B), Interleukin-6 (IL6), Interleukin-8 (IL8), Tumor Necrosis Factor (TNF), Prostaglandin-Endoperoxide Synthase 2 (PTGS2)) that has been used in previous CTRA research (Cole, 2019), with statistical analyses testing for association of their average expression level with the continuous Child-ACE scores while controlling for continuous age, continuous body mass index percentile, and indicator variables for participant sex, race/ethnicity (White/Black/Latino/Other), current insurance coverage, and household smoking exposure. Statistical significance of average association coefficients was derived from standard errors estimated by bootstrap re-sampling of linear model residual vectors to control for any potential correlation across genes. Secondary analyses examined Type I interferon gene expression by parallel analysis of an a priori-specified 34-gene composite (e.g., *IFI*-, *MX*-, and *OAS*-family genes) used in previous CTRA research (Cole, 2019).

To explore the cellular mechanisms that might potentially be involved in ACE-related differences in gene expression, we also carried out Transcript Origin Analyses (TOA) as previously described using cell type-specific reference profiles from isolated leukocyte subsets (Black, Cole, Christodoulou, & Figueiredo, 2018; Cole, Hawkey, Arevalo, & Cacioppo, 2011). In these exploratory analyses, all genes showing >20% differential expression per ACE count (after adjustment for covariates) served as inputs into the higher-order TOA bioinformatics analysis, and statistical significance of TOA cell type-specific diagnosticity scores was derived from standard errors estimated by bootstrap re-sampling of linear model residual vectors to control for any potential correlation across genes. Because these analyses were exploratory and involved multiple cell types, statistical significance was based on Bonferroni-corrected *p* values adjusting for the 10 cell types examined (i.e., CD4+ T cells, CD8+ T cells, B cells, NK cells, DC1 dendritic cells, DC2 dendritic cells, DC3 dendritic cells, CD16-classical monocytes, and CD16+ nonclassical monocytes). Note that TOA does not attempt to quantify the relative prevalence of different cell types within the circulating blood cell pool, but rather attempts the more modest objective of determining whether the genes that are found to be differentially expressed (i.e., associated with Child-ACE scores) are predominately expressed by one or more specific subset of leukocytes (e.g., CD4+ T cells, CD8+ T cell, etc.) based on previous RNA profiling analyses of physically isolated reference cell samples (for more detail, see Black et al., 2018; Cole et al., 2011). TOA findings may reflect either differences in leukocyte subset prevalence (e.g., increased or decreased numbers of a given cell type) or differences in leukocyte activation (e.g., per-cell changes in RNA transcription rates and activation of a given cell type), or a combination of both processes.

## Results

### Sample characteristics

As shown in Table 1, higher Child-ACE exposure scores were associated with female gender, Medicaid insurance, and smoke exposure. Higher Child-ACE total score was only associated

**Table 1.** Characteristics of children in research sample by Child-ACE score

Child-ACE exposure only	Score 0–1	Score 2+	<i>p</i> value
Age – Median (SD)	9 (2.2)	9 (2.0)	0.437
BMI% – Median (SD)	69 (22.2)	71 (21.8)	0.851
Female – N (%)	9 (39%)	12 (86%)	0.007*
Male – N (%)	14 (61%)	2 (14%)	
White – N (%)	5 (22%)	2 (14%)	0.697
Latino/Hispanic – N (%)	11 (48%)	7 (50%)	
Black – N (%)	2 (9%)	0 (0%)	
Other – N (%)	5 (22%)	5 (36%)	
Medicaid insurance – N (%)	14 (61%)	14 (100%)	0.007*
Private insurance – N (%)	9 (39%)	0 (0%)	
No smoke exposure – N (%)	22 (96%)	9 (64%)	0.021*
Smoke exposure – N (%)	1 (4%)	5 (36%)	
PSC total – median (SD)	5 (4.9)	9 (8.8)	0.162
Child-ACE total	Score 0–1	Score 2+	<i>p</i> value
Age – Median (SD)	9.5 (1.9)	7 (2.2)	0.094
BMI% – Median (SD)	69 (20.4)	73 (23.3)	0.724
Female – N (%)	8 (50%)	13 (62%)	0.520
Male – N (%)	8 (50%)	8 (38%)	
White – N (%)	5 (31%)	2 (10%)	0.378
Latino/Hispanic – N (%)	6 (38%)	12 (57%)	
Black – N (%)	1 (6%)	1 (5%)	
Other – N (%)	4 (25%)	6 (29%)	
Medicaid insurance – N (%)	8 (50%)	20 (95%)	0.002*
Private insurance – N (%)	8 (50%)	1 (5%)	
No smoke exposure – N (%)	15 (94%)	16 (76%)	0.206
Smoke exposure – N (%)	1 (6%)	5 (24%)	
PSC Total – Median (SD)	4.5 (3.6)	8 (8.2)	0.166

Note: \*  $p < 0.05$ ; ACE = adverse childhood experiences; BMI = body mass index; PSC = Pediatric Symptom Checklist; SD = standard deviation.

with Medicaid insurance. The median Pediatric Symptom Checklist total score was higher for higher Child-ACE scores, but this difference was nonsignificant.

### Inflammatory gene expression

Primary analyses examined whether a pre-specified composite of 19 pro-inflammatory gene transcripts would show elevated levels of expression in DBS samples in proportion to the child's ACE total score (i.e., both exposure and risk). Results showed that was indeed the case (Figure 1a), with average pro-inflammatory gene expression elevated by 0.094 log<sub>2</sub> RNA expression units with each Child-ACE total score point ( $\pm 0.036$  standard error,  $p = .019$ ) in analyses controlling for age, body mass index percentile, sex, race/ethnicity, current insurance status, and household smoking exposure. Similar effects emerged for the Child-ACE exposure measure that omitted measures based on risk and scored solely on reported exposures ( $0.121 \pm 0.057$  standard error,  $p = .049$ ; Figure 1b).

### Interferon gene expression

Secondary analyses examined whether a composite of 34 Type I interferon-related gene transcripts might also show differential expression as a function of Child-ACE scores. Average expression of the Type I interferon composite was elevated by  $0.103 \pm 0.037$  log<sub>2</sub> RNA expression units with each Child-ACE total score point ( $p = .008$ ), and similar effects emerged for the Child-ACE exposure score ( $0.175 \pm 0.051$ ,  $p = .002$ ).

### Cellular origins

To identify potential cellular mechanisms of the observed differences in DBS whole blood gene expression, we conducted TOA testing for differential cellular origins of the 796 gene transcripts that showed >20% difference in average expression per Child-ACE total score point. Results implicated CD8+ T cell as the primary sources of the 759 gene transcripts upregulated in association with Child-ACE scores, and nonclassical (CD16+) monocytes as the source of the 37 downregulated genes (Figure 1c).






Figure 1. Childhood adversity and gene expression profiles

## Discussion

The present study documents elevated levels of inflammatory gene expression in whole blood RNA samples from children in proportion to caregiver-reported exposure to ACEs early in life. These results are independent of major demographic confounders (age, body mass index percentile, sex, race/ethnicity), independent of parental risk factors (health insurance status, household smoking), robust to inclusion versus exclusion of risk factors (i.e., total vs. exposure Child-ACE scores) and are consistent with previous research linking ACEs to circulating inflammatory protein biomarkers. These results document the correlation between the Child-ACE score reported by caregivers completing the WCA and inflammatory gene expression in the child. As such, the present results suggest that alterations in immune cell gene regulation might represent one pathway by which ACE exposures come to be associated with greater levels of inflammation and elevated health risk later in life.

Results of the primary analysis linking ACE exposure to increased pro-inflammatory gene expression are consistent with previous research linking adverse life circumstances in adolescence and adulthood to increased expression of pro-inflammatory genes in circulating immune cells (Cole, 2013, 2014, 2019; Schwaiger et al., 2016). However, the results of secondary analyses examining Type I interferon-related gene expression and immune cell origins suggest that the transcriptomic correlates of ACE exposure in this sample of children does not resemble the classical CTRA profile observed in adolescents and adults. In the classical CTRA profile, elevated expression of pro-inflammatory genes is

accompanied by decreased expression of Type I interferon genes, and involves upregulation of classical monocytes (Cole, 2019; Schwaiger et al., 2016). By contrast, the present analyses link ACE exposure to increased Type I interferon-related gene expression, decreased non-classical monocyte activity, and increased activity of CD8<sup>+</sup> T cells (which is uncharacteristic of the classical CTRA profile).

Future research in larger samples is needed to confirm these results, and to further define the cellular and molecular mechanisms involved. Upregulation of Type I interferon-related gene expression along with increased activity of CD8<sup>+</sup> T cells may reflect a pattern of generalized activation within the T cell compartment of the adaptive immune system, in contrast to the suppression observed in most previous CTRA studies. Three hypotheses may explain why our results differ from the classical CTRA pattern. (a) The anti-viral response of the immune system may vary for infants, young children, adolescents, and adults. Our sample only included young children (age 5–11 years old), in contrast to the older age samples of other studies that have evaluated CTRA in youth (Ehrlich et al., 2015; Miller et al., 2018; Murphy et al., 2013, 2015). Recent experience with differences in susceptibility to the coronavirus supports the merit of considering differences in immune response by age groups (Carsetti et al., 2020). (b) The natural history of toxic stress may change with the temporal proximity of exposure to ACEs. This is consistent with literature on the normal function of Type I interferon in protection from disease by increased production in response to acute stimuli and decreased production when stimuli are prolonged (Trinchieri, 2010), as well as with studies of dysfunctional

states (i.e., autoimmune disorders which are marked by persistent upregulation of basal T cell activity) (Trinchieri, 2010; Wahadat et al., 2018) and a broader literature on the influence of temporal parameters on immune response (Segerstrom & Miller, 2004). (c) ACE exposure may be correlated with increased exposure to or activity of viral infections, which could result in upregulation of both Type I interferon activity and CD8+ T cell activity as observed here. If subsequent research supports this hypothesis, then the basis for differential viral activity would become an important question for future ACE research.

Some limitations should be considered when interpreting these results. First, although we were able to detect statistically significant associations for the relationships examined, the sample size for this study was small and may have failed to detect other associations. Thus, our results should be considered preliminary and in need of replication. Second, although the results are consistent with the causal role of ACEs in health risks via chronic inflammation, the direction of the association cannot be determined from this study since it is cross-sectional. However, this study does have an advantage over adult studies of ACEs in that ACE exposure was measured based upon caregiver report, whereas the immune response was measured in the child and not likely to have influenced the caregiver report. Third, although associations with health outcomes is plausible based upon our prior exploration of Child-ACE scores (Marie-Mitchell et al., 2020), we were unable to conduct a broad examination of health outcomes in this sample so the health significance of the present results remain to be clarified in future research. Finally, future studies should extend the current RNA-based findings to include direct measures of inflammatory proteins (e.g., plasma cytokines) and cell populations (e.g., flow cytometry enumeration) that were not feasible in the context of the present study's DBS sampling method.

Preliminary analyses suggest that a brief questionnaire with caregiver-reported responses to questions about childhood adversity can distinguish young children at lower and higher risk for elevated levels of inflammatory gene expression. Our results also suggest that patterns of immune response may vary by age and temporal proximity to stressors. Of note to general pediatricians and population health researchers, our assessment of the immune system used DBS methods, which offers a minimally invasive and pragmatic approach to advancing precision medicine. While it is humbling to see this biopsychosocial process in children as young as 5–11 years old, it also underscores the need for early interventions to protect against the potential sequelae of adversity.

**Acknowledgment.** We are grateful for the support of the Loma Linda University Pediatrics Department.

**Financial Support.** This work was supported by the Loma Linda University Pediatrics Department Research Initiative (PRI #558502). The sponsor did not have a role in the study design, analysis, or preparation of this manuscript for publication.

**Conflict of Interest.** None.

## References

Adamao, S. (2014). The effects of stress hormones on immune function may be vital for the adaptive reconfiguration of the immune system during fight-or-flight behavior. *Integrative and Comparative Biology*, *54*, 419–426. doi:10.1093/icb/icu005

Anda, R., Felitti, V., Bremner, J., Walker, J., Whitfield, C., Perry, B., ... Giles, W. (2006). The enduring effects of abuse and related adverse experiences in childhood. A convergence of evidence from neurobiology and

epidemiology. *European Archives of Psychiatry and Clinical Neuroscience*, *256*, 174–186.

Antoni, M., Bouchard, L., Jacobs, J., Lechner, S., Jutagir, D., Gudenkauf, L., ... Blomberg, B. (2016). Stress management, leukocyte transcriptional changes and breast cancer recurrence in a randomized trial: An exploratory analysis. *Psychoneuroendocrinology*, *74*, 269–277.

Black, D., Cole, S., Christodoulou, G., & Figueiredo, J. (2018). Genomic mechanisms of fatigue in survivors of colorectal cancer. *Cancer*, *124*, 2637–2644. doi:10.1002/cncr.31356

Brown, D., Anda, R., Tiemeier, H., Felitti, V., Edwards, V., Croft, J., & Giles, W. (2009). Adverse childhood experiences and the risk of premature mortality. *American Journal of Preventive Medicine*, *37*, 389–396.

Carsetti, R., Quintarelli, C., Quinti, I., Mortari, E., Zumla, A., Ippolito, G., & Locatelli, F. (2020). The immune system of children: The key to understanding SARS-CoV-2 susceptibility? *Lancet Child & Adolescent Health*, *4*, 414–416.

Centers for Disease Control and Prevention, & National Center for Chronic Disease Prevention and Health Promotion. (2009). *The Power of Prevention: Chronic Disease...The Public Health Challenge of The 21st Century*. Retrieved from Atlanta, GA.

Chiang, J. J., Ko, A., Bower, J. E., Taylor, S. E., Irwin, M. R., & Fuligni, A. J. (2019). Stress, psychological resources, and HPA and inflammatory reactivity during late adolescence. *Development and Psychopathology*, *31*, 699–712. doi:10.1017/s0954579418000287

Chiang, J. J., Taylor, S. E., & Bower, J. E. (2015). Early adversity, neural development, and inflammation. *Developmental Psychobiology*, *57*, 887–907. doi:10.1002/dev.21329

Cicchetti, D., Handley, E. D., & Rogosch, F. A. (2015). Child maltreatment, inflammation, and internalizing symptoms: Investigating the roles of C-reactive protein, gene variation, and neuroendocrine regulation. *Development and Psychopathology*, *27*, 553–566. doi:10.1017/s0954579415000152

Cole, S. (2013). Social regulation of human gene expression: Mechanisms and implications for public health. *American Journal of Public Health*, *103*, S84–S92.

Cole, S. (2014). Human social genomics. *PLOS Genetics*, *10*, e1004601.

Cole, S. (2019). The conserved transcriptional response to adversity. *Current Opinion in Behavioral Sciences*, *28*, 31–37.

Cole, S.W., Hawkey, L.C., Arevalo, J. M., & Cacioppo, J. T. (2011). Transcript origin analysis identifies antigen-presenting cells as primary targets of socially regulated gene expression in leukocytes. *Proceedings of the National Academy of Sciences of the United States of America*, *108*, 3080–3085. doi:10.1073/pnas.1014218108

Danese, A., Moffitt, T., Harrington, H., Milne, B., Polanczyk, G., Pariante, C., ... Caspi, A. (2009). Adverse childhood experiences and adult risk factors for age-related disease: Depression, inflammation, and clustering of metabolic risk markers. *Archives of Pediatrics and Adolescent Medicine*, *163*, 1135–1143.

Danese, A., Pariante, C., Caspi, A., Taylor, A., & Poulton, R. (2007). Childhood maltreatment predicts adult inflammation in a life-course study. *Proceedings of the National Academy of Sciences of the United States of America*, *104*, 1319–1324.

David, J., Measelle, J., Ostlund, B., & Ablow, J. (2017). Association between early life adversity and inflammation during infancy. *Developmental Psychobiology*, *59*, 696–702. doi:10.1002/dev.21538

Dobin, A., Davis, C. A., Schlesinger, F., Drenkow, J., Zaleski, C., Jha, S., ... Gingeras, T. R. (2013). STAR: Ultrafast universal RNA-seq aligner. *Bioinformatics*, *29*, 15–21.

Dubowitz, H., Lane, W., Semiati, J., & Magder, L. (2012). The SEEK model of pediatric primary care: Can child maltreatment be prevented in a low-risk population? *Academic Pediatrics*, *12*, 259–268.

Ehrlich, K., Miller, G., & Chen, E. (2015). Harsh parent-child conflict is associated with decreased anti-inflammatory gene expression and increased symptom severity in children with asthma. *Development and Psychopathology*, *27*, 1547–1554.

Fiegelman, S., Dubowitz, H., Lane, W., Grube, L., & Kim, J. (2011). Training pediatric residents in a primary care clinic to help address psychosocial problems and prevent child maltreatment. *Academic Pediatrics*, *11*, 474–480. doi:10.1016/j.acap.2011.07.005

- Felitti, V. J., Anda, R. F., Nordenberg, D., Williamson, D. F., Spitz, A. M., Edwards, V., ... Marks, J. S. (1998). Relationship of childhood abuse and household dysfunction to many of the leading causes of death in adults. The Adverse Childhood Experiences (ACE) Study. *American Journal of Preventive Medicine*, *14*, 245–258.
- Gardner, W., Lucas, A., Kolko, D. J., & Campo, J. V. (2007). Comparison of the PSC-17 and alternative mental health screens in an at-risk primary care sample. *Journal of the American Academy of Child & Adolescent Psychiatry*, *46*, 611–618.
- Howren, M., Lamkin, D., & Suls, J. (2009). Associations of depression with C-reactive protein, IL-1, and IL-6: A meta-analysis. *Psychosomatic Medicine*, *71*, 171–186.
- Irwin, M., & Cole, S. (2011). Reciprocal regulation of the neural and innate immune systems. *Nature Reviews Immunology*, *11*, 625–632. doi:10.1038/nri3042
- Jellinek, M., Murphy, J., Little, M., Pagano, M., Comer, D., & Kelleher, K. (1999). Use of the Pediatric Symptom Checklist (PSC) to screen for psychosocial problems in pediatric primary care: A national feasibility study. *Archives of Pediatrics & Adolescent Medicine*, *153*, 254–260.
- Jiang, N. M., Tofail, F., Ma, J. Z., Haque, R., Kirkpatrick, B., Nelson, C. A., & Petri, W. A. (2017). Early life inflammation and neurodevelopmental outcome in Bangladeshi infants growing up in adversity. *American Journal of Tropical Medicine and Hygiene*, *97*, 974–979. doi:10.4269/ajtmh.17-0083
- Knight, J., Rizzo, J., Logan, B., Wang, T., Arevalo, J., Ma, J., & Cole, S. (2016). Low Socioeconomic status, adverse gene expression profiles, and clinical outcomes in hematopoietic stem cell transplant recipients. *Clinical Cancer Research*, *22*, 69–78. doi:10.1158/1078-0432.CCR-15-1344
- Knight, J., Rizzo, J., Wang, T., He, N., Logan, B., Spellman, S., ... Cole, S. (2019). Molecular correlates of socioeconomic status and clinical outcomes following hematopoietic cell transplantation for leukemia. *JNCI Cancer Spectrum*, *3*, pkz073. doi:10.1093/jncics/pkz073
- Kroenke, K., Spitzer, R., & Williams, J. (2003). The patient health questionnaire-2: Validity of a two-item depression screener. *Medical Care*, *41*, 1284–1292.
- Kroenke, K., Spitzer, R., Williams, J., Monahan, P., & Löwe, B. (2007). Anxiety disorders in primary care: Prevalence, impairment, comorbidity, and detection. *Annals of Internal Medicine*, *146*, 317–325.
- Marie-Mitchell, A., Lee, J., Siplon, C., Chan, F., Riesen, S., & Vercio, C. (2019). Implementation of the Whole Child Assessment to screen for adverse childhood experiences. *Global Pediatric Health*, *6*, 1–9.
- Marie-Mitchell, A., Watkins, H., Copado, I., & Distelberg, B. (2020). Use of the Whole Child Assessment to identify children at risk of poor outcomes. *Child Abuse and Neglect*, *104*, 104489. doi:10.1016/j.chiabu.2020.104489
- McDade, T., Williams, S., & Snodgrass, J. (2007). What a drop can do: Dried blood spots as a minimally invasive method for integrating biomarkers into population-based research. *Demography*, *44*, 899–925.
- Measelle, J., & Ablow, J. (2018). Contributions of early adversity to pro-inflammatory phenotype in infancy: The buffer provided by attachment security. *Attachment & Human Development*, *20*, 1–23. doi:10.1080/14616734.2017.1362657
- Miller, G., & Chen, E. (2010). Harsh family climate in early life presages the emergence of a proinflammatory phenotype in adolescence. *Psychological Science*, *21*, 848–856.
- Miller, G., Chen, E., & Cole, S. (2009). Health psychology: Developing biologically plausible models linking the social world and physical health. *Annual Review of Psychology*, *60*, 501–524.
- Miller, G., Chen, E., Shalowitz, M., Story, R., Leigh, A., Ham, P., ... Cole, S. (2018). Divergent transcriptional profiles in pediatric asthma patients of low and high socioeconomic status. *Pediatric Pulmonology*, *53*, 710–719.
- Miller, G., & Cole, S. (2012). Clustering of depression and inflammation in adolescents previously exposed to childhood adversity. *Biol Psychiatry*, *72*, 34–40. doi:10.1016/j.biopsych.2012.02.034
- Murphy, J., Reed, J., Jellinek, M., & Bishop, S. (1992). Screening for psychosocial dysfunction in inner-city children: Further validation of the pediatric symptom checklist. *Journal of the American Academy of Child & Adolescent Psychiatry*, *31*, 1105–1111.
- Murphy, M., Slavich, G., Chen, E., & Miller, G. (2015). Targeted rejection predicts decreased anti-inflammatory gene expression and increased symptom severity in youth with asthma. *Psychological Science*, *26*, 111–121.
- Murphy, M., Slavich, G., Rohleder, N., & Miller, G. (2013). Targeted rejection triggers differential pro- and anti-inflammatory gene expression in adolescents as a function of social status. *Clinical Psychological Science*, *1*, 30–40.
- Repetti, R., Taylor, S., & Seeman, T. (2002). Risky families: Family social environments and the mental and physical health of offspring. *Psychological Bulletin*, *128*, 330–366.
- Ross, K., Carroll, J., Dunkel Schetter, C., Hobel, C., & Cole, S. (2019). Pro-inflammatory immune cell gene expression during the third trimester of pregnancy is associated with shorter gestational length and lower birthweight. *American Journal of Reproductive Immunology*, *82*, e13190. doi:10.1111/aji.13190
- Ross, K., Cole, S., Carroll, J., & Dunkel Schetter, C. (2019). Elevated pro-inflammatory gene expression in the third trimester of pregnancy in mothers who experienced stressful life events. *Brain, Behavior and Immunity*, *76*, 97–103. doi:10.1016/j.bbi.2018.11.009
- Schwaiger, M., Grinberg, M., Moser, D., Zang, J., Heinrichs, M., Hengstler, J., ... Kumsta, R. (2016). Altered stress-induced regulation of genes in monocytes in adults with a history of childhood adversity. *Neuropsychopharmacology*, *41*, 2530–2540.
- Seegerstrom, S., & Miller, G. (2004). Psychological Stress and the Human Immune System: A MetaAnalytic Study of 30 Years of Inquiry. *Psychological Bulletin*, *130*, 601–630.
- Shonkoff, J., Garner, A., The Committee on Psychosocial Aspects of Child and Family Health, Committee on Early Childhood Adoption and Dependent Care, Section on Developmental and Behavioral Pediatrics, Siegel, B., ... Wood, D. (2012). The lifelong effects of early childhood adversity and toxic stress. *Pediatrics*, *129*, e232.
- Simons, R., Lei, M., Beach, S., Barr, A., Cutrona, C., Gibbons, F., & Philibert, R. (2017). An index of the ratio of inflammatory to antiviral cell types mediates the effects of social adversity and age on chronic illness. *Social Science & Medicine*, *185*, 158–165.
- Sloven, N., Kubzansky, L. D., McLaughlin, K. A., & Koenen, K. C. (2013). Childhood adversity and inflammatory processes in youth: A prospective study. *Psychoneuroendocrinology*, *38*, 188–200. doi:10.1016/j.psyneuen.2012.05.013
- Surtees, P., Wainwright, N., Day, N., Brayne, C., Luben, R., Khaw, K. T., ... Khaw, K.-T. (2003). Adverse experience in childhood as a developmental risk factor for altered immune status in adulthood. *International Journal of Behavioral Medicine*, *10*, 251–268.
- Taj, N., Devera-Sales, A., & Vinson, D. (1998). Screening for problem drinking: Does a single question work? *Journal of Family Practice*, *46*, 328–335.
- Trinchieri, G. (2010). Type I Interferon: Friend or foe? *Journal of Experimental Medicine*, *207*, 2053–2063.
- Tyrka, A. R., Parade, S. H., Valentine, T. R., Eslinger, N. M., & Seifer, R. (2015). Adversity in preschool-aged children: Effects on salivary interleukin-1beta. *Development and Psychopathology*, *27*, 567–576. doi:10.1017/S0954579415000164
- Wahadat, M., Bodewes, I., Maria, N., van Helden-Meeuwse, C., van Dijk-Hummelman, A., Steenwijk, E., ... Versnel, M. (2018). Type I IFN signature in childhood-onset systemic lupus erythematosus: A conspiracy of DNA- and RNA-sensing receptors? *Arthritis Research & Therapy*, *20*, doi:10.1186/s13075-017-1501-z
- Zhou, X., Fragalaa, M., McElhaney, J., & Kuchel, G. (2010). Conceptual and methodological issues relevant to cytokine and inflammatory marker measurements in clinical research. *Current Opinion in Clinical Nutrition and Metabolic Care*, *13*, 541–547.