

Original Article

*These authors shared first authorship.

Cite this article: Knies AK, Jutagir DR, Ercolano E, Pasacreta N, Lazenby M, McCorkle R (2019). Barriers and facilitators to implementing the commission on cancer's distress screening program standard. *Palliative and Supportive Care* **17**, 253–261. <https://doi.org/10.1017/S1478951518000378>

Received: 1 April 2018

Revised: 9 April 2018

Accepted: 13 April 2018

Key words:

Quality of life; distress screening; cancer; anxiety/depression; interventions

Author of correspondence: Andrea K. Knies, Department of Neurology, Division of Neurocritical Care & Emergency Medicine, Yale School of Medicine, 15 York Street, New Haven, CT 06510. E-mail: andrea.knies@yale.edu

Barriers and facilitators to implementing the commission on cancer's distress screening program standard

Andrea K. Knies, DR. RER. MEDIC.^{1*}, Devika R. Jutagir, PH.D.^{2*}, Elizabeth Ercolano, D.N.S.¹, Nicholas Pasacreta, B.A.¹, Mark Lazenby, PH.D.¹ and Ruth McCorkle, PH.D.¹

¹Yale University School of Nursing, West Haven, CT and ²Yale University School of Medicine, New Haven, CT

Abstract

Objective. Many cancer centers struggle to implement standardized distress screening despite the American College of Surgeons' Commission on Cancer 2012 mandate for a distress screening program standard of care by 2015. This paper presents outcomes for the first cohort of participants ($n = 36$) of a Screening for Psychosocial Distress Program (SPDP), a 2-year training program designed to assist clinicians in implementing routine distress screening as mandated by the American College of Surgeons Commission on Cancer. Specifically, participants' success with distress screening implementation, institutional barriers and facilitators to implementation, and the role of the SPDP are described.

Method. This research followed a longitudinal pre- and posttest mixed methods design. An investigator-developed questionnaire collected qualitative (distress screening goals, institutional barriers and facilitators, facilitators associated with participation in the SPDP) and quantitative (level of goal achievement) data at 6, 12, and 24 months of participation in the SPDP. Conventional content analysis was applied to qualitative data. Mixed methods data analysis in Dedoose evaluated (1) types and number of distress screening goals, barriers, and facilitators, and (2) goal achievement at 6, 12, and 24 months of participation.

Result. Ninety-five percent of distress screening implementation goals were completed after 2 years of participation. Most common institutional barriers to distress screening implementation were "lack of staff," "competing demands," and "staff turn-over." Most common institutional facilitators were "buy-in," "institutional support," and "recognition of participants' expertise." The number of reported facilitators associated with SPDP participation was higher than the number associated with any institutional factor, and increased over time of participation.

Significance of results. Participating in training programs to implement distress screening may facilitate successful achievement of the Commission on Cancer's distress screening standard, and benefits seem to increase with time of participation. Training programs are needed to promote facilitators and overcome barriers to distress screening.

Introduction

In 2012, the American College of Surgeons' Commission on Cancer (CoC) mandated that, for cancer centers to be accredited, a distress screening program standard of care be achieved by 2015 (Commission on Cancer, 2012). This mandate followed a 1997 National Comprehensive Cancer Network recommendation for universal distress screening at all cancer care centers (Holland et al., 2013). These guidelines were intended to improve uptake of distress screening, yet the CoC deadline has passed and many institutions continue to struggle to meet these requirements. In 2007, only 15% of oncologists reported adhering to standardized distress screening (Pirl et al., 2007), and in a 2015 survey, only 54.6% of 467 oncology social workers screened for distress using a standardized instrument (BrintzenhofeSzoc et al., 2015). Given that up to 43% of patients report clinically significant distress over the course of cancer diagnosis and treatment (Zabora et al., 2001), widespread adoption of comprehensive distress screening has the potential to connect a large proportion of cancer patients with psychosocial intervention and other needed services that reduce distress (Jacobsen & Jim, 2008). Insight is needed into why some cancer centers can implement standardized distress screening, whereas others cannot, and whether training programs can facilitate adoption of the program standard.

Comprehensive distress screening is a multistep process designed to provide initial detection of distress then documented referral and follow-up. It is recommended that distress screening not only detect distress, but also further evaluate distressed patients and provide referrals for appropriate services (Pirl et al., 2014). The five steps of comprehensive psychosocial distress screening include: (1) screening patients for distress using a validated screening tool, (2) evaluating the screen for clinically significant distress, (3) referring distressed patients to psychosocial healthcare services, (4) following up to reevaluate distress, and (5)

documenting distress screening steps in the health record and conducting Quality Improvement by auditing health records (Lazenby et al., 2015a). Such procedures enhance patient communication with clinicians and increase referrals for psychosocial services (Carlson et al., 2012).

However, at each step of distress screening, barriers and facilitators may determine whether a comprehensive screening program is successfully implemented. Protected time, engaged leadership, and a culture that values distress screening have all been identified as facilitators of distress screening implementation (Riblet et al., 2014). In contrast, several reports of attempts to implement distress screening at teaching hospitals (Lo et al., 2016; O'Connor et al., 2017) and across outpatient settings nationwide (Tavernier et al., 2013) show that lack of time on the part of clinicians is an implementation barrier. Review papers have indicated that lack of staff buy-in and lack of staff knowledge about the need for screening interfere with implementation of evidence-based psychosocial care in oncology (Schofield et al., 2006) and with implementation of distress screening (Pearman et al., 2015). A significant body of literature outlines common barriers to and facilitators of implementing distress screening. However, to support institutions in developing screening programs, more research is needed that clarifies how barriers and facilitators correspond to specific steps of distress screening.

For this reason, we designed a 2-year Screening for Psychosocial Distress Program (SPDP) to teach and facilitate routine comprehensive psychosocial distress screening to US cancer care professionals. In this paper, we report data available for clinicians from 18 cancer care institutions participating in the SPDP. To our knowledge, this is the first program to delineate barriers and facilitators to achieving distress screening goals in a cohort of participants enrolled in a training program to implement distress screening over time. Specifically, this article describes (1) participants' achievement of distress screening related goals at 6, 12, and 24 months postenrollment; (2) barriers and facilitators to goal achievement experienced by participants in their institutions; and (3) the helpfulness of the SPDP for goal achievement.

Methods

Design

The SPDP used a longitudinal pre- and posttest design. Mixed methods research data including both data on process (barriers and facilitators) and outcomes (goal achievement) were collected at 6, 12, and 24 months. A cohort of two cancer-care clinicians (dyads) from 18 institutions (36 clinicians total) completed the 2-year SPDP (2013–2015). The program consisted of an introductory workshop in early 2014; bimonthly teleconferences in the first year and quarterly in the second year for information, problem-solving, and support; and an advanced workshop at 12 months of participation. The workshops, described previously (Lazenby et al., 2018), included faculty presentations on content relevant to distress screening and participants' group work and presentations on their distress screening activities.

Participants

Participants were recruited through advertisements online (<http://apos-society.org/screening/>) and in trade periodicals as described previously (Lazenby et al., 2015b). Applications were screened for inclusion and exclusion criteria. Inclusion criteria included (1)

two people from one institution applying as a dyad, who (2) had direct access to patients to screen, and (3) had strong letters of support from the institution's administrators. Excluded were applicants who applied individually and who had noncommittal letters of support. From the pool of applicants, we selected those institutions that provided the strongest evidence for (1) a well-grounded expectation that the dyads would work well together, and (2) continuous strong administration buy-in and support for distress screening efforts of the dyad over the full 2 years of training (purposive sampling).

Formation of goals relevant to distress screening

Participants were required to state three distress screening goals on their applications. During the introductory workshop, participants were supported by faculty in refining these goals into the Specific, Measurable, Achievable, Realistic, Time-bound (SMART) goal format (Doran, 1981). Exercises were driven by the Reach, Effectiveness, Adoption, Implementation, Maintenance framework as previously described (Lazenby et al., 2015b) to help them develop goals that they would be able to work on over the 2 years of their participation in the SPDP.

Data collection

At 6, 12, and 24 months after baseline, we collected data on dyads' progress on implementing distress screening in their institutions using an investigator-developed Goal Evaluation Form (Grant et al., 1999, 2007) sent by e-mail to be completed online. The Goal Evaluation Form collected both quantitative and qualitative data: For quantitative data, dyads rated the level of achievement of their SMART goals refined during the introductory workshop at baseline (1 = never started; 2 = stopped/cancelled; 3 = stalled; 4 = in process; 5 = completed). Open-ended questions prompted the provision of qualitative data on institutional barriers and facilitators to goal achievement and the helpfulness of materials provided during the SPDP (Table 1). One institution (18) did

Table 1. Open-ended questions used in the Goal Evaluation Form to elicit institutional barriers and facilitators to goal achievement and the helpfulness of materials provided during the SPDP

<p>Questions to elicit barriers and facilitators experienced within institutions</p> <ul style="list-style-type: none"> • What happened in your setting either positively or negatively to support your goals at the institutional level? • What happened in your setting either positively or negatively to support your goals at the administrative level? • What happened among your colleagues in your setting either positively or negatively to support your goals? • What resources in your setting either positively or negatively to support your goals? • What activities have occurred that demonstrated respect of your agency in relation to your expertise in distress screening? • What changes in your positions have occurred since you completed the workshop? How have these changes affected your ability to achieve your goals? <p>Question to elicit the helpfulness of materials provided during the SPDP</p> <ul style="list-style-type: none"> • What materials received during the workshop have been most helpful to you in achieving your goals?

SPDP, Screening for Psychosocial Distress Program.

not provide 24-month data. Raw data were imported into Dedoose, a web application for managing, analyzing, and presenting qualitative and mixed method research data (SocioCultural Research Consultants, LLC, 2016).

Data analysis

Goals specified by dyads were categorized according to the five steps of distress screening: (1) screening, (2) evaluating, (3) referring if needed, (4) following up, and (5) documenting (Lazenby et al., 2015a). Of the original 54 goals, three goals were recategorized into two different distress screening steps and one goal was divided into two separate goals, which resulted in a total of 58 individual coded goals. Goals were defined as unfinished when a level of achievement was reported as “never started,” “stopped/cancelled,” or “stalled” at 6 and 12 months, or reported as “never started,” “stopped/cancelled,” “stalled,” or “in process” at 24 months.

Qualitative data on institutional barriers and facilitators were analyzed with conventional content analysis (Hsieh & Shannon, 2005) by identifying categories directly from the text. Categories were organized into a coding system consisting of main categories (e.g., “lack of buy-in”) and, if applicable, ancillary subcategories specifying more nuanced aspects of the main category (e.g., “lack of administrator buy-in”). Three investigators (RM, ML, EE), who are experts in psychosocial distress screening, independently applied the codes to subjects’ self-reported barriers and facilitators. Overall, there were minimal discrepancies noted; these investigators discussed each coding discrepancy and reached consensus on the classification. Data saturation was reached; therefore, no additional coding was needed. In Dedoose, goals’ categories and levels of achievement at 6, 12, and 24 months were specified, and codes yielded during content analysis were applied to qualitative data collected for the open-ended questions. As 24 months, data were missing for institution 18, their goal achievement levels provided for 12 months (“completed,” $n = 3$) were used for 24 months evaluation. However, the identification of institutional barriers, facilitators, and SPDP materials for institution 18 at 24 months had to be omitted. Data for all evaluation time points were downloaded in an Excel spreadsheet and frequencies of (1a) goal type (as pertaining to the five steps of psychosocial distress screening), (1b) goal achievement levels, (2a) types of institutional barriers, (2b) types of institutional facilitators, and (3) facilitative components of the SPDP were descriptively analyzed.

Ethical approval

The program was exempt from review by Yale University’s Human Investigation Committee.

Results

Participants’ characteristics

Of 18 dyads (36 participants total), 11 identified their institutions as National Cancer Institute Designated/Comprehensive Cancer Centers; the remaining seven institutions identified as community/general hospitals. Size of institutions (measured by number of cancer patients treated annually) ranged from 263 to 13,683 patients/year, with a median size of 3,300 patients/year ($M = 4,716.66$, $SD = 4,343.37$). Participants’ disciplines were social

work (58%), nursing (25%), psychiatry/psychology (14%), and other (3%).

Goal achievement

Table 2 displays the 58 SMART goals pertaining to the five steps of psychosocial distress screening along with their level of goal achievement at 6, 12, and 24 months, and direct quotes of goal examples. The most common goals were creating “stakeholder buy-in” ($n = 12$), establishing a “referral network” ($n = 9$), and starting “brief screening” ($n = 8$).

Over the course of the 2-year training program, participants reported 15 (25.9%), 38 (65.5%), and 55 (94.8%) of the 58 goals as completed at 6, 12, and 24 months, respectively. Of the remaining three incomplete goals at 24 months, two goals, categorized as “referral network” (“create a list of current psychosocial programs, including svcs provided by our behavioral health dept”) (institution 25) and “piloting & beginning” (“pilot distress screening process in 4 identified clinics and record data in spreadsheet”) (institution 13) were in process. Another goal, categorized as “stakeholder buy-in” (“engage key stakeholders at each hospital cancer institute to complete current state assessment across the system”) (institution 51) was stopped/cancelled.

At 6 or 12 months, 18 goals (31.0%) were unfinished, but completed at 12 or 24 months. Specifically, at 6 months, 16 goals (27.6%) were unfinished, but completed at 12 months (“referral network” [$n = 2$], “education on buy-in & sustainability” [$n = 2$], “piloting & beginning” [$n = 1$], “follow-up” [$n = 1$], “electronic health record” [$n = 1$], “stakeholder buy-in” [$n = 1$], and “clinical evaluation” [$n = 1$]), or 24 months (“piloting & beginning” [$n = 3$], “education on process” [$n = 2$], “follow-up” [$n = 1$], and “electronic health record” [$n = 1$]). At 12 months, two goals (3.4%) were unfinished (after being “in process” at 6 months), but completed at 24 months (“brief screening” and “referral network” [each $n = 1$]).

Institutional barriers

Over the course of the 2-year program, dyads reported a total of 65 barriers occurring during their distress screening implementation efforts in their institutions. The top three barriers were “lack of staff” ($n = 15$), “staff turnover” and “competing demands” (each $n = 11$), and “screening process mechanics” ($n = 7$). Screening process mechanics related barriers refer to the selection, utility/validity of the screening tool, and the documentation of screening activities. Examples of participants’ direct quotes of barriers are shown in Table 3.

Twenty-one barriers (32.3%) were reported for goals, which at any one evaluation time point (6, 12, or 24 months) were reported as unfinished ($n = 25$, including goals that were repeatedly reported as unfinished at various time points). Analogously to the frequencies found for barriers reported for all goals, among these 21 barriers, the top three were: “staff turnover” and “lack of staff” (each $n = 5$), “screening process mechanics” ($n = 3$), and “competing demands” and “lack of buy-in in institution” (each $n = 2$) (Table 3.). Among barriers reported for the 25 unfinished goals, percentages for staff and screening process mechanics-related barriers were higher (47.6% vs 40%, respectively) than among barriers reported for all goals, including completed goals (14.3% vs. 10.8%, respectively). Percentages for barriers related to competing demands, however, were lower among barriers reported with unfinished goals (9.5%) than among barriers for all goals (16.9%). Barriers related to a lack

Table 2. Level of achievement of goals by category at 6, 12, and 24 months (*N* = 58 goals) and goal example quotations

Goals (<i>N</i> = 58) Direct quotes of goal example	#(%)	Levels of goal achievement														
		6 months					12 months					24 months				
		1	2	3	4	5	1	2	3	4	5	1	2	3	4	5
0. Preplanning process																
Stakeholder buy-in “Submit proposal to cancer committee for review, revision and authorization” (institution 31)	12 (20.7)	0	0	1	5	6	0	0	0	2	10	0	1	0	0	11
Policy “Develop pathways and protocols for assessment, referrals and follow-up based on distress screen cut-off score” (institution 28)	3 (5.2)	0	0	0	2	1	0	0	0	1	2	0	0	0	0	3
Education on process “Educate cancer center of excellence staff about implementation of the psychosocial distress tool” (institution 25)	7 (12.1)	1	0	1	4	1	0	0	0	3	4	0	0	0	0	7
Education on buy-in & sustainability “To educate and cultivate ‘buy-in’ from clinical trials/research dept to begin screening all patients entering clinical trials” (institution 53)	5 (8.6)	1	0	1	2	1	0	0	0	1	4	0	0	0	0	5
1. Screening																
Brief screening “The distress thermometer will be given to all patients at the chemo class by the nurse doing the education” (institution 65)	8 (13.8)	0	0	0	4	4	0	1	0	1	6	0	0	0	0	8
Piloting & beginning “Pilot distress screening process in four identified clinics and record data in spreadsheet” (institution 13)	6 (10.3)	5	0	0	1	0	3	0	0	1	2	0	0	0	1	5
2. Evaluation of screen																
Clinical evaluation “Develop the process and procedure for connecting information of patients with a distress screening score greater than 4” (institution 65)	3 (5.2)	1	0	0	2	0	0	0	0	1	2	0	0	0	0	3

(Continued)

Table 2. (Continued.)

Goals (N = 58) Direct quotes of goal example	#(%)	Levels of goal achievement														
		6 months					12 months					24 months				
		1	2	3	4	5	1	2	3	4	5	1	2	3	4	5
#	#	#	#	#	#	#	#	#	#	#	#	#	#	#		
3. Referral																
Referral network "Update and connect with outside community resource referrals" (institution 64)	9 (15.5)	1	0	1	5	2	0	0	1	3	5	0	0	0	1	8
4. Follow-up																
Patient follow-up "Identify and follow up with oncology patients referrals (those given the assessment tool) to assess session follow through of appt." (institution 25)	2 (3.4)	1	0	1	0	0	0	0	0	1	1	0	0	0	0	2
5. Documentation & quality improvement																
Electronic health record "Determine and create a database to evaluate distress screen" (institution 37)	2 (3.4)	1	0	1	0	0	0	0	0	1	1	0	0	0	0	2
Documentation & quality improvement "Analyze baseline and follow up data for all patients screened to date by July 2014 and summarize results for cancer committee" (institution 51)	1 (1.7)	0	0	0	1	0	0	0	0	0	1	0	0	0	0	1
Total	58 (100)	11	0	6	26	15	3	1	1	15	38	0	1	0	2	55

Levels of achievement: 1 = never started; 2 = stopped/cancelled; 3 = stalled; 4 = in process; 5 = completed .

of buy-in within institutions were reported equally frequently for unfinished versus all goals (9.5% vs. 9.2%, respectively).

Institutional facilitators

Dyads reported a total of 189 facilitators that they experienced during their distress screening implementation efforts within their institutions. Table 4 shows the frequencies of all facilitator categories at 6, 12, and 24 months along with examples of participants' direct quotes. By far the most frequent facilitator reported was "buy-in" ($n = 76$, 40.2%), followed by "institution support" ($n = 28$, 14.8%), and "dyad viewed as knowledgeable & a resource" ($n = 23$, 12.2%).

With regard to the 18 goals that were unfinished at 6 or 12 months, 35 (17.7%) facilitators were reported concurrently with completion of these goals at 12 or 24 months. Among those 35 facilitators, "buy-in" ($n = 12$), "education efforts in institution," and "institution support" (each: $n = 6$) were most frequent.

Whereas the frequency for "buy-in" facilitators slightly decreased from 12 ($n = 7$) to 24 months ($n = 5$), the frequencies for facilitators related to "education efforts in institution" and "institution support" both increased from 12 months ($n = 1$) to 24 months ($n = 5$) (data not shown).

Helpfulness of the screening for psychosocial distress training program

Overall, across all evaluation time points, 65 SPDP-related facilitators were reported. Beyond the materials provided by the SPDP, dyads reported conference calls and interactions among peers from other institutions as instrumental in helping them achieve their goals.

Table 5 shows frequencies reported for components of the training program that were useful at 6, 12, and 24 months along with examples of direct quotes reflective of the component reported by dyads.

Table 3. Frequencies and quotes of barrier categories reported by dyads at 6, 12, and 24 months (*N* = 65)

Barriers (<i>N</i> = 65)	All goals #	Unfinished goals #	Direct quote of barrier example
Lack of resources:			
Lack of staff	15	5	“Clinic staff is burdened with high acuity and numbers of patients” (institution 72)
Lack of finances	2	–	“[...], fiscal pressures from the overall institution present a barrier” (institution 5)
Staff attitudes & fears	4	1	“[...] the only push back is from some cancer center staff who are concerned about the added burden of the tool in their already busy roles” (institution 51)
Staff turnover	11	5	“A more challenging aspect is changes in personnel [...]” (institution 53)
Lack of staff education	3	1	“A more challenging aspect is [...] the ongoing need for education” (institution 53)
Multisite coordination	2	1	“The offsite locations are more difficult to connect to resources” (institution 1)
Electronic health record*	4	1	“Attempting to get a new assessment in the EMR was a time consuming process” (institution 28)
Lack of buy-in within institution	6	2	“Some of the front line staff didn’t have the buy in” (institution 18)
Screening process mechanics†	7	3	“There has been some challenges regarding the current form, and patients understanding of the thermometer and if their distress is truly about cancer concerns” (institution 64)
Competing demands	11	2	“The difficulty comes in with competing time demands for staff on the oncology unit- it is challenging to fulfill distress screening in addition to existing requirements” (institution 8)
Total	65	21	

*Barriers related to electronic health record were “lack of IT support” and “limited usefulness.”

†Barriers related to screening process mechanics were “selection of tool,” “utility and validity of tool,” and “incomplete documentation process.”
EMR, electronic medical record; IT, information technology.

Table 4. Frequencies and quotes of facilitator categories reported by dyads at 6, 12, and 24 months (*N* = 189)

Categories of facilitators (<i>N</i> = 189)	Total # (%)	6 months # (%)*	12 months # (%)*	24 months # (%)*	Direct quote of facilitator example
Buy-in	76 (40.2)	32 (42.1)	31 (40.8)	13 (17.1)	“Our staff was very supportive of the program” (institution 32)
Institution support	28 (14.8)	2 (7.14)	11 (39.3)	15 (53.6)	“The administrative support we’ve received has been extremely helpful” (institution 66)
Dyad viewed as knowledgeable & a resource	23 (12.2)	7 (30.4)	8 (34.8)	8 (34.8)	“I was asked to write an article on screening for distress for our cancer committee’s annual report” (institution 53)
Education efforts in institution	22 (11.6)	7 (31.8)	8 (36.4)	7 (31.8)	“Staff education and training on how to talk about distress with patients” (institution 42)
Resources	13 (6.9)	6 (46.2)	1 (7.7)	6 (46.2)	“We are very fortunate to have so many resources available internally and externally” (institution 18)
Electronic health record (screening integrated in EHR)	11 (5.8)	8 (72.7)	2 (18.2)	1 (9.1)	“Having the distress tool loaded into Epic has been helpful” (institution 13)
Psychosocial committee	8 (4.2)	3 (37.5)	3 (37.5)	2 (25)	“Active participants of staff on committee” (institution 25)
Multisite coordination	4 (2.1)	2 (50)	1 (25)	1 (25)	“[...] we get monthly reports/check ins on how each site is doing on the project” (institution 51)
Dyad support (dyad receives support from dyad or from others)	2 (1.1)	1 (50)	1 (50)	0	“My colleague, [...], and I work very closely and well together in this process” (institution 31)
Successful screening at diagnosis: improving and adding resources	2 (1.1)	0	0	2 (100)	“Full support allowed me to get added FTEs and to make system-wide policies/protocols” (institution 1)

*The relative frequencies at 6, 12, and 24 months refer to the absolute frequencies of each barrier, not to the absolute frequency of all barriers (*N* = 189).
FTE, full-time equivalent; EHR, electronic health record.

Table 5. Frequencies of facilitators associated with the SPDP to meet the mandate at 6, 12, and 24 months with examples of quotes ($N = 65$)

Facilitators ($N = 65$)	Total # (%)	6 months # (%)	12 months # (%)	24 months # (%)	Direct quote of facilitator example
Printed materials	20 (30.8)	10 (50)	6 (30)	4 (20)	"I have utilized many of the slides in the presentations to use in my presentation to staff [...]. [They] were very helpful in helping us to develop our education program. [...]" (institution 65)
Workshops	15 (23.1)	6 (40)	4 (26.7)	5 (33.3)	"Everything, the workbooks with all of the presentations were vital. Also poster presentations are key" (institution 1)
Intragroup correspondence	12 (18.5)	3 (25)	5 (41.7)	4 (33.3)	"Networking with fellow program participants, particularly when sharing policies and tools they've developed - gives a broader perspective and new insights" (institution 66)
Conference calls	10 (15.4)	2 (20)	6 (60)	4 (40)	"We benefited from the support on the conference calls throughout the cycle" (institution 13)
Slides for education	3 (4.6)	0	2 (66.7)	1 (33.3)	"The slide deck used for talk points with other providers" (institution 72)
Mandate	2 (3.1)	0	2 (100)	0	"Inpatient distress screening will strengthen our standing about meeting CoC standards for psychosocial care" (institution 8)
Selection of tool	2 (3.1)	0	0	2 (100)	"Initiation of PHQ2 screening for depression" (institution 42)
Affiliation with program helpful	1 (2.3)	0	0	1 (100)	"This training and expertise that we have with the YSN distress screen program has just elevated the respect we have in the field" (institution 7)

CoC, Commission on Cancer; SPDP, Screening for Psychosocial Distress Program; PHQ2, Patient Health Questionnaire - 2; YSN, Yale School of Nursing.

Concurrently with the 18 goals that were unfinished at 6 or 12 months, but finished at 12 or 24 months, SPDP facilitators were reported 13 times (i.e., more often than the top facilitator "buy-in" that dyads experienced within their institutions). Whereas the frequency for "buy-in" facilitators decreased from 12 months ($n = 7$) to 24 months ($n = 5$), facilitators reported as related to the SPDP increased from 12 months ($n = 4$) to 24 months ($n = 9$).

Discussion

The 18 dyads who participated in the 2-year SPDP were successful in achieving their goals related to implementing the CoC's distress screening standard, with nearly all goals attained by the end of the 2-year training program.

During the implementation of distress screening programs in their institutions, our participants primarily faced staff-related barriers (lack of staff, staff turnover), as well as barriers related to competing demands and screening process mechanics. Barriers related to screening process mechanics and, particularly, staff-related barriers were not only most frequently reported for all goals, but also most frequently reported together with unfinished goals. Therefore, staff-related barriers and barriers related to screening process mechanics are not only most prevalent, but may impede goal achievement. In contrast, although barriers related to competing demands were relatively frequently reported overall, they were less frequently reported with unfinished goals and therefore do not appear to impede goal completion. Rather, competing demands may be an inherent barrier when already busy clinicians take on the responsibility of implementing distress screening. Although it has previously been documented that lack of time is an implementation barrier (Mitchell et al., 2012) and that conveying lack of time to patients may discourage them from reporting their distress (Biddle et al., 2016), our study suggests that competing demands may not ultimately hinder goal achievement. Although each of these barriers should be

anticipated, these results suggest that a sufficient and steady staff base, as well as the piloting and fine-tuning of screening process mechanics tailored to the individual institution are the facilitators that are crucial for the success of a distress screening program.

Most frequent facilitators reported by our participants were buy-in, institutional support, and dyad viewed as knowledgeable and a resource. Whereas buy-in was a frequent facilitator for dyads' distress screening efforts during the first year of our program, institution support increased over time and particularly during the second year of training. It is possible that it takes the buy-in of key stakeholders within institutions to subsequently unlock institutional support. Both buy-in and institutional support were also the most frequent facilitators to be reported concurrently with unfinished, but eventually completed, goals. These findings support prior literature that emphasizes the importance of enhancing buy-in of stakeholders for successful implementation of distress screening (Butz et al., 2011; Groff et al., 2018; Lozcalzo et al., 2011). Education efforts in institutions were also one of the most prevalent facilitators reported with previously unfinished and eventually completed goals. It is possible that dyads recognized the need for education as crucial for achieving their goals and enhanced their education efforts accordingly. This interpretation may be supported by the trend that the frequency of Education Efforts facilitators reported concurrently with previously unfinished, but eventually completed goals increased from 12 to 24 months, which is a not a pattern seen for Education Efforts facilitators reported for all goals (Table 4).

The distress screening program appeared to be particularly valuable for the completion of unfinished goals: the SPDP was reported concurrently with previously unfinished, but eventually completed goals more often than any of these other facilitators, with increasing frequencies from 12 to 24 months. Therefore, the SPDP may have been critical for the ultimate achievement of these more challenging goals that seemed to require facilitators beyond those that dyads were able to activate within their

institutions. Elements of the SPDP that were reported as particularly helpful were attendance at both workshops and printed materials provided by the faculty. Printed materials such as the slide deck were reported as helpful in implementing education efforts that seemed crucial for success. Participants also reported the training program served as an opportunity to exchange information and resources about distress screening with dyads at other sites. Access to distress screening experts through scheduled conference calls tailored to the needs of the dyads was also considered an effective strategy in promoting goal achievement. Although previous research has demonstrated that conducting a distress screening educational program for oncology staff members is feasible and may enhance distress screening implementation (Grassi *et al.*, 2001), our findings indicate specific components that might be particularly helpful to include in future distress screening training programs, particularly in-person workshops, printed educational materials, and regular opportunities to exchange information with experts and staff members implementing distress screening at other institutions.

Strengths

The program contributes to the literature by identifying barriers and facilitators that seem crucial to the successful implementation of a routine comprehensive distress screening program in community healthcare settings. As such, the method of analysis using a type of software that allowed for analyzing the frequencies of barriers and facilitators alongside goal key phrases was a strength. Literature on distress screening attempts in community settings is scarce, because community settings are less likely to implement and document distress screening attempts than academic medical centers (Pearman *et al.*, 2015). Furthermore, given the diversity of the types of medical centers included in the sample, the results may be generalizable to a variety of types of similar medical institutions.

Limitations

Some limitations of the program should be considered when interpreting the results. The program consisted of one intervention group and did not include a comparison group. It is possible that participating dyads would have achieved distress screening implementation success without participating in the program. Size of institutions varied and three dyad members were reassigned to different locations; therefore, working together was not always possible. Given that participants self-selected to apply to the program and were admitted only if two staff members were committed to participating and demonstrated strong administrative support, results may not generalize to less motivated and less resourced institutions. However, barriers such as lack of staff may be more wide spread at institutions that would not meet criteria to participate in this program.

The data reported are limited to self-report and no objective patient data were audited or recorded; therefore, outcomes are limited to the participants' perceptions. Furthermore, phrasing of the open-ended questions on the goal form did not allow identifying relationships for barriers and facilitators with specific goals. However, the temporal overlap of goal achievement and the concurrent reporting of barriers and facilitators may suggest connections.

Future research

Given that our program consisted of a small sample size of 18 dyads, future evaluation should include a greater number of participants to either replicate or challenge the present findings. Evaluation should be conducted that directly links specific barriers and facilitators to distress screening goals. Furthermore, assessing patient perspectives on barriers and facilitators to patient participation in distress screening and provider-reported barriers and facilitators to distress screening may shed light on other factors related to distress screening goal achievement (Lambert *et al.*, 2014). Future research should also examine whether and how the CoC mandate itself acts as a facilitator for distress screening implementation.

Given that lack of staff and competing demands were reported as barriers, more efforts are needed to understand how to conduct distress screening in a time-efficient way, and how distress screening can be used to maximize time during diagnostic and treatment visits (O'Connor *et al.*, 2017). Future research should examine strategies for convincing administrators of cancer care institutions to provide adequate resources to establish and maintain distress screening programs. Our results suggest that gaining institutional support and leveraging its benefits for implementing distress screening takes time. To support staff efforts to secure resources for distress screening, studies are needed that establish cost-effective distress screening policies and identify mechanisms through which distress screening may contribute to reduced health care spending (Pearman *et al.*, 2015). These efforts will be dependent on dyads' abilities to analyze patient data on distress screening from electronic medical records and demonstrate linkages to patient outcomes such as treatment adherence and completion, and cost savings.

Conclusion

Since 2015, cancer centers are mandated by the American College of Surgeons' CoC to have a routine comprehensive distress screening program in place to be accredited. Our 2-year SPDP was designed to support cancer care clinicians with distress screening implementation. Results gained from the evaluation of the first cohort of participants indicate that successful implementation of a distress screening program requires time, buy-in and support from key stakeholders, a solid and steady staff base, and education efforts within institutions. Participation in distress screening training programs may help cancer care clinicians achieve distress screening implementation goals that are particularly challenging and require support of external experts, or otherwise may be left unfinished. Future research is needed to develop training programs that assist clinicians in overcoming barriers and activating facilitators that are crucial to meeting the distress screening mandate.

Acknowledgments. This research was supported by the National Cancer Institute grant R25 CA177553. D.R.J. was supported by T32 CA009461. The authors declare that there is no conflict of interest.

A.K.K. and D.R.J. assumed equal lead responsibility for this paper.

References

- Biddle L *et al.* (2016) Patients' and clinicians' experiences of holistic needs assessment using a cancer distress thermometer and problem list: a qualitative study. *European Journal of Oncology Nursing* 22, 59–65.

- BrintzenhofeSzoc K et al.** (2015) Screening for psychosocial distress: a national survey for oncology social workers. *Journal of Psychosocial Oncology* **33**, 34–47.
- Butz BD et al.** (2011). Implementing screening for distress, the 6th vital sign: A Canadian strategy for changing practice. *Psycho-Oncology* **20**, 463–469.
- Carlson LE, Walker A and Mitchell AJ** (2012). Screening for distress and unmet needs in patients with cancer: Review and recommendations. *Journal of Clinical Oncology* **30**, 1160–1177.
- Commission on Cancer** (2012) Cancer Program Standards 2012 Version 1.1: Ensuring patient-centered care. Available from <http://www.facs.org/cancer/coc/programstandards2012.html>
- Doran GT** (1981) There's a S.M.A.R.T. way to write management's goals and objectives. *Management Review* **70**, 35–36.
- Grant M et al.** (2007) Disseminating end-of-life-education to cancer centers: Overview of program and of evaluation. *Journal of Cancer Education* **22**, 140–148.
- Grant M et al.** (1999) Improving cancer pain management using a performance improvement framework. *Journal of Nursing Care Quality* **13**, 60–72.
- Grassi L et al.** (2001) Educational intervention in cancer outpatient clinics on routine screening for emotional distress: An observational study. *Psycho-Oncology* **20**, 669–674.
- Groff S et al.** (2018) Examining the sustainability of Screening for Distress, the sixth vital sign, in two outpatient oncology clinics: A mixed-methods study. *Psycho-Oncology* **27**, 141–147.
- Holland JC et al.** (2013) Distress management. *Journal of the National Comprehensive Cancer Network* **11**, 190–209.
- Hsieh HF and Shannon SE** (2005) Three approaches to qualitative content analysis. *Qualitative Health Research* **15**, 1277–1288.
- Jacobsen PB and Jim HS** (2008) Psychosocial interventions for anxiety and depression in adult cancer patients: Achievements and challenges. *CA: A Cancer Journal for Clinicians* **58**, 214–230.
- Lambert SD et al.** (2014) Psycho-oncology services among women with gynecologic cancer following distress screening. *Journal of the National Comprehensive Cancer Network* **12**, 899–906.
- Lazenby M et al.** (2015a) Supporting commission on cancer-mandated psychosocial distress screening with implementation strategies. *Journal of Oncology Practice* **11**, e413–e420.
- Lazenby M et al.** (2015b) The five steps of comprehensive psychosocial distress screening. *Current Oncology Reports* **17**, 447.
- Lazenby M et al.** (2018) An educational program's effect on participants' goals for implementing psychosocial distress screening in routine cancer care. *Clinical Journal of Oncology Nursing* **22**.
- Lo SB et al.** (2016) Experience implementing distress screening using the National Comprehensive Cancer Network distress thermometer at an urban safety-net hospital. *Psycho-Oncology* **25**, 1113–1115.
- Lozalzo M, Clark KL and Holland J** (2011) Successful strategies for implementing biopsychosocial screening. *Psycho-Oncology* **20**, 455–462.
- Mitchell AJ et al.** (2012) How feasible is implementation of distress screening by cancer clinicians in routine clinical care? *Cancer* **118**, 6260–6269.
- O'Connor M et al.** (2017) Detecting distress: Introducing routine screening in a gynecologic cancer setting. *Clinical Journal of Oncology Nursing* **21**, 79–85.
- Pearman T et al.** (2015) Implementation of distress screening in an oncology setting. *The Journal of Community and Supportive Oncology* **13**, 423–428.
- Pirl WF et al.** (2014) Recommendations for the implementation of distress screening programs in cancer centers. *Cancer* **120**, 2946–2954.
- Pirl WF et al.** (2007) Screening for psychosocial distress: A national survey of oncologists. *Journal of Supportive Oncology* **5**, 499–504.
- Riblet N et al.** (2014) Addressing distress in patients with head and neck cancers: A mental health quality improvement project. *Journal of the National Comprehensive Cancer Network* **12**, 1005–1013.
- Schofield P et al.** (2006) Barriers to the provision of evidence-based psychosocial care in oncology. *Psycho-Oncology* **15**, 863–872.
- SocioCultural Research Consultants, LLC** (2016) Dedoose version 7.0.23. Available from <http://www.dedoose.com/>
- Tavernier SS, Beck SL and Dudley WN** (2013) Diffusion of a distress management guideline into practice. *Psycho-Oncology* **22**, 2332–2338.
- Zabora J et al.** (2001) The prevalence of psychological distress by cancer site. *Psycho-Oncology* **10**, 19–28.