

Self-sustaining process of minimal attached eddies in turbulent channel flow

Yongyun Hwang^{1,†} and Yacine Bengana²

¹Department of Aeronautics, Imperial College London, South Kensington, London SW7 2AZ, UK

²Département de Mécanique, Université de Pierre et Marie Curie, 4 Place Jussieu, 75005 Paris, France

(Received 15 July 2015; revised 17 March 2016; accepted 28 March 2016;
first published online 20 April 2016)

It has been recently shown that the energy-containing motions (i.e. coherent structures) in turbulent channel flow exist in the form of Townsend's attached eddies by a numerical experiment which simulates the energy-containing motions only at a prescribed spanwise length scale using their self-sustaining nature (Hwang, *J. Fluid Mech.*, vol. 767, 2015, pp. 254–289). In the present study, a detailed investigation of the self-sustaining process of the energy-containing motions at each spanwise length scale (i.e. the attached eddies) in the logarithmic and outer regions is carried out with an emphasis on its relevance to 'bursting', which refers to an energetic temporal oscillation of the motions (Flores & Jiménez, *Phys. Fluids*, vol. 22, 2010, 071704). It is shown that the attached eddies in the logarithmic and outer regions, composed of streaks and quasi-streamwise vortical structures, bear the self-sustaining process remarkably similar to that in the near-wall region: i.e. the streaks are significantly amplified by the quasi-streamwise vortices via the lift-up effect; the amplified streaks subsequently undergo a 'rapid streamwise meandering motion', reminiscent of streak instability or transient growth, which eventually results in breakdown of the streaks and regeneration of new quasi-streamwise vortices. For the attached eddies at a given spanwise length scale λ_z between $\lambda_z^+ \simeq 100$ and $\lambda_z \simeq 1.5h$, the single turnover time period of the self-sustaining process is found to be $Tu_\tau/\lambda_z \simeq 2$ (u_τ is the friction velocity), which corresponds well to the time scale of the bursting. Two additional numerical experiments, designed to artificially suppress the lift-up effect and the streak meandering motions, respectively, reveal that these processes are essential ingredients of the self-sustaining process of the attached eddies in the logarithmic and outer regions, consistent with several previous theoretical studies. It is also shown that the artificial suppression of the lift-up effect of the attached eddies in the logarithmic and outer regions leads to substantial amounts of turbulent skin-friction reduction.

Key words: turbulence simulation, turbulence theory, turbulent boundary layers

1. Introduction

Townsend (1961, 1976) originally introduced the concept of the 'attached eddy' to describe the energy-containing motions (i.e. coherent structures) populating the logarithmic region of wall-bounded turbulent flows. Given the length scale of

† Email address for correspondence: y.hwang@imperial.ac.uk

the logarithmic region, the size of the energy-containing motions there would be proportional to the distance of their centre from the wall. This implies that, at least some parts of the energy-containing motions would reach the wall, and, in this sense, they are expected to be ‘attached’ to the wall. Under the assumption that these energy-containing motions in the logarithmic region (i.e. attached eddies) are statistically self-similar to one another, Townsend (1976) theoretically predicted that turbulence intensity of the wall-parallel velocity components would exhibit a logarithmic wall-normal dependence as the mean-velocity profile does. The original theory of Townsend (1976) has been significantly refined over a number of years by Perry and coworkers (e.g. Perry & Chong 1982; Perry, Henbest & Chong 1986; Perry & Marusic 1995; Nickels *et al.* 2005, among many others), who have elaborated to develop a self-consistent model predicting the statistics of wall-bounded turbulent flows with a structural model of the attached eddy in the form of a hairpin or Λ vortex. Especially, these authors further predicted and verified the emergence of k_x^{-1} law in the spectra of wall-parallel velocity components (k_x is the streamwise wavenumber), which would be linked to the logarithmic wall-normal dependence of turbulence intensity of these velocity components (Perry & Chong 1982).

A growing body of evidence, which supports the theoretical predictions made with the attached eddy hypothesis, has emerged, especially over the last decade. The logarithmic growth of the near-wall streamwise turbulence intensity with the Reynolds number is an example of this, as it would be given by extending Townsend’s prediction to the near-wall region (Marusic & Kunkel 2003). The linear growth of the spanwise integral length scale with the distance from the wall has also been understood as another important evidence (Tomkins & Adrian 2003; del Álamo *et al.* 2004; Monty *et al.* 2007), as this indicates that the spanwise size of the energy-containing motions in the logarithmic region is proportional to the distance from the wall. Lastly, the logarithmic dependence of turbulence intensity of the wall-parallel velocity components has recently been indeed confirmed at sufficiently high Reynolds numbers (Jiménez & Hoyas 2008; Marusic *et al.* 2013; Orlandi, Bernardini & Pirozzoli 2015), suggesting that the energy-containing motions in wall-bounded turbulent flow would indeed be organised in the form of Townsend’s attached eddies.

Very recently, the first author of the present study managed to compute the statistics of the attached eddies at a given length scale with a numerical experiment performed at moderate Reynolds numbers ($Re_\tau \simeq 1000\text{--}2000$ where Re_τ is the friction Reynolds number) (Hwang 2015). Built upon the linearly growing nature of the spanwise integral length scale with the distance from the wall, the numerical experiment is designed to isolate the energy-containing motions only at a given spanwise length scale using their ‘self-sustaining nature’ (Hwang & Cossu 2010c, 2011). The isolated self-sustaining energy-containing motions were found to be self-similar with respect to their spanwise length scale, and their statistical structures were remarkably similar to those of the attached eddies, demonstrating the existence of Townsend’s attached eddies as energy-containing motions in wall-bounded turbulent flows. It was also shown that the single attached eddy is composed of two distinct elements, one of which is a long streaky motion and the other is a relatively compact streamwise vortical structure. At the considered Reynolds numbers, the size and the wall-normal location of the former streaky motion, the major carrier of the streamwise turbulence intensity, are roughly self-similar along

$$y \simeq 0.1\lambda_z \quad \text{and} \quad \lambda_x \simeq 10\lambda_z, \quad (1.1a,b)$$

where y is the wall-normal direction, λ_x the streamwise length scale and λ_z the spanwise length scale, while those of the latter vortical structure, carrying all the velocity components, also self-similarly scale with

$$y \simeq 0.5 \sim 0.7\lambda_z \quad \text{and} \quad \lambda_x \simeq 2 \sim 3\lambda_z. \quad (1.1c,d)$$

The scaling also reveals that the smallest attached eddy, given with $\lambda_z^+ \simeq 100$, is a near-wall coherent motion in the form of a near-wall streak and with several quasi-streamwise vortices aligned to it (Hwang 2013), while the largest one, given with $\lambda_z \simeq 1.5h$ in a turbulence channel with the half-height h , is an outer motion composed of a very-large-scale motion (VLSM) and with several large-scale motions (LSMs) aligned to it (Hwang 2015). The attached eddies at intermediate length scales, the size of which is proportional to their distance from the wall, are also in the form of a long streak and several quasi-streamwise vortices, and contribute to the logarithmic region by filling the gap caused by the separation between the inner and outer length scales. It should be mentioned that this scenario incorporates all the coherent structures currently known and exhibits the behaviour consistent with the spectra of all the velocities and Reynolds stress (including k_x^{-1} spectra), providing an integrated description for the coherent structures within the attached eddy scenario (for further details, see also Hwang 2015).

The present study is an extension of the work by Hwang (2015), and, in particular, it is aimed to explore the ‘dynamics’ of the self-sustaining attached eddies and its relevance to the corresponding motions in a fully turbulent environment. For this purpose, in the present study we consider the smallest computational box which allows for the self-sustaining mechanism of each of the attached eddies (i.e. the minimal unit). We then examine the detailed physical processes in the minimal unit by comparing them with those under fully developed turbulence. Particular emphasis of the present study is given to relating the self-sustaining process of the attached eddies with the dynamical behaviour often referred to as ‘bursting’ in the logarithmic and outer regions (Flores & Jiménez 2010). To this end, we introduce two additional numerical experiments, designed to examine the physical processes composing the self-sustaining process of the attached eddies in the logarithmic and outer regions.

The paper is organised as follows. In §2, we briefly introduce the numerical methods of extracting self-sustaining attached eddies at a given spanwise length scale (see also Hwang 2015, for further details). A careful observation of the instantaneous flow field is then followed in §3 with examination of auto- and cross-correlation functions of several flow variables of interest. Especially in this section, we show that the bursting in the logarithmic and outer regions is likely a consequence of the self-sustaining process of the attached eddies. In §4, a comprehensive discussion on the self-sustaining mechanism of the attached eddies is given with the two numerical experiments mentioned. The paper finally concludes in §5.

2. Numerical experiment

2.1. Computation of self-sustaining attached eddies

We consider a turbulent channel, in which we denote x , y and z as the streamwise, wall-normal and spanwise directions, respectively. The upper and lower walls are set to be at $y = 0$ and $y = 2h$, respectively, where h is the half-height of the channel. The numerical experiment in the present study is performed using the Navier–Stokes solver in our previous studies (Hwang & Cossu 2010c, 2011; Hwang

2013, 2015). In this solver, the streamwise and spanwise directions are discretised using the Fourier–Galerkin method with the 2/3 rule, and the wall-normal direction is discretised using a second-order central difference scheme. Time integration is performed semi-implicitly using the Crank–Nicolson method for the terms with wall-normal derivatives and a third-order low-storage Runge–Kutta method for the rest of the terms. The computations are performed by imposing a constant volume flux across the channel.

As mentioned, the present numerical experiment is largely based on the previous work by Hwang (2015) (the reader is strongly recommended to refer to this paper in which the numerical technique introduced in this section is fully verified). In turbulent channel flow, the spanwise length scale exhibits a linear growth with the distance from the wall (i.e. y) in the range from $\lambda_z^+ \simeq 100$ to $\lambda_z \simeq 1.5h$ (del Álamo *et al.* 2004). The spanwise spectra of the streamwise velocity clearly reveal this feature, and are approximately aligned along $y \sim 0.1\lambda_z$ (del Álamo *et al.* 2004; Hwang 2015). The key idea of Hwang (2015) is to design a numerical simulation only for the energy-containing motions at a prescribed spanwise length $\lambda_z = \lambda_{z,0}$ between $\lambda_z^+ \simeq 100$ and $\lambda_z \simeq 1.5h$ by artificially removing the motions at the other spanwise length scales (i.e. $\lambda_z \neq \lambda_{z,0}$).

First, the removal of the motions, the spanwise size of which is larger than the prescribed spanwise length $\lambda_{z,0}$ (i.e. $\lambda_z > \lambda_{z,0}$) is implemented by setting the size of the spanwise computational domain L_z to be identical to the prescribed spanwise length scale $\lambda_{z,0}$ (i.e. $L_z = \lambda_{z,0}$), while keeping a sufficiently long streamwise domain L_x . However, even in such a computational domain, the motions at $\lambda_z > \lambda_{z,0}$ are not completely removed, as a spurious motion, uniform along the spanwise direction, would still survive (Hwang 2013). This spurious motion is subsequently removed by taking the approach in Hwang (2013) which explicitly filters out the spanwise uniform components by setting the right-hand side of the discretised momentum equation at each Runge–Kutta substep as follows:

$$\left. \begin{aligned} \widehat{\text{RHS}}_x(y; k_x \neq 0, k_z = 0) &= 0, \\ \widehat{\text{RHS}}_y(y; k_x \neq 0, k_z = 0) &= 0, \end{aligned} \right\} \quad (2.1)$$

where the RHS_i the right-hand side of i -component of the discretized momentum equation, $\widehat{\cdot}$ denotes Fourier-transformed state in x and z directions and k_x and k_z are the streamwise and spanwise wavenumbers, respectively. We note that this technique is only aimed at removing the two-dimensional spurious motion populating the x – y plane, thus no action is taken for $\widehat{\text{RHS}}_z$ (for further details, see also Hwang 2013).

Once the motions at $\lambda_z > \lambda_{z,0}$ are removed, the motions at $\lambda_z < \lambda_{z,0}$ are subsequently quenched using an over-damped large-eddy simulation (LES) (Hwang & Cossu 2010c; Hwang 2013, 2015). For the given scope of the present study, it is very important for the residual stress model of the LES not to generate any possible energy transfer to the resolved quantity. Therefore, as in Hwang (2015), we consider the static Smagorinsky model in which the residual stress is modelled in a purely diffusive form:

$$\tilde{\tau}_{ij} - \frac{\delta_{ij}}{3} \tilde{\tau}_{kk} = -2\nu_t \tilde{\mathcal{S}}_{ij}, \quad (2.2a)$$

with

$$\nu_t = (C_s \tilde{\Delta})^2 \tilde{\mathcal{S}} \mathcal{D}, \quad (2.2b)$$

where $\tilde{\cdot}$ denotes the filtered quantity, S_{ij} the strain rate tensor, C_s the Smagorinsky constant, $\tilde{\Delta} = (\tilde{\Delta}_1 \tilde{\Delta}_2 \tilde{\Delta}_3)^{1/3}$ the nominal filter width, $\tilde{\mathcal{S}} = (2\tilde{S}_{ij}\tilde{S}_{ij})^{1/2}$ the norm of the strain rate tensor and $\mathcal{D} = 1 - \exp[-(y^+/A^+)^3]$ is the van Driest damping function (Härtel & Kleiser 1998). The removal of the motions at $\lambda_z < \lambda_{z,0}$ is subsequently carried out by increasing C_s with the increment $\Delta C_s = 0.05$ until we isolate the self-sustaining motions at $\lambda_z = \lambda_{z,0}$. We note that this approach is equivalent to increasing the filter width of LES without losing actual resolution (Mason & Cullen 1986). Finally, it should be mentioned that this approach itself does not strongly depend on the choice of the eddy viscosity model (Hwang & Cossu 2010c, 2011).

The smallest computational domain, which allows for survival of the isolated self-sustaining attached eddies at a given spanwise length scale $\lambda_{z,0}$ (i.e. the minimal unit), is then sought. Since the spanwise size of the target attached eddies determines the spanwise computational domain L_z (i.e. $L_z = \lambda_{z,0}$), only the size of the streamwise computational domain L_x needs to be examined by gradually reducing it from a sufficiently large value. Such a search of the minimal unit of the self-sustaining attached eddies at a given spanwise length scale was previously performed in Hwang & Cossu (2010c) with the present over-damped LES, but without the filtering (2.1). In the present study, we therefore repeat this search again to check any possible effect of (2.1). It has been found that the filtering (2.1) does not significantly affect the size of the minimal unit, resulting in the following streamwise computational domain size for the self-sustaining attached eddies at $\lambda_{z,0} = L_z$:

$$L_x = 2L_z. \quad (2.3)$$

In the present study, all the simulations are performed by keeping with this aspect ratio of the computational domain (see also table 1). We note that the use of this minimal streamwise computational domain is not a great limitation of the present study, as the minimal domain is found not to significantly change the second-order statistics of the isolated self-sustaining attached eddies at $\lambda_{z,0} = L_z$. A detailed discussion on this issue is given in appendix B where the second-order statistics of the attached eddies with the minimal unit are compared with those with a long streamwise domain in Hwang (2015). Finally, it should be mentioned that the filtering of the uniform motion along the streamwise direction, such as (2.1) for the spanwise direction, is not implemented. As we shall see in §§ 3 and 4, the streamwise uniform Fourier mode in the minimal unit mainly resolves the long streaky motion observed in a long computational domain. Indeed, we will see that the streamwise uniform Fourier mode is strongly correlated with the streaky motion (see § 3), and inhibition of the related physical process destroys the self-sustaining process in the minimal unit (see § 4).

The simulation parameters of the present study are summarised in table 1, where six different simulations are considered at two different Reynolds numbers ($Re_\tau \simeq 950$ and $Re_\tau \simeq 1800$). For each computational domain, we consider a pair of simulations for the purpose of comparison. One is with $C_s = 0.05$, the value known to provide the best statistics compared to that of direct numerical simulations (DNS) (Härtel & Kleiser 1998) and is performed to track the attached eddies with background turbulence, and the other is with an increased C_s and is to study the attached eddies surviving only through the self-sustaining mechanism. Typical spatial grid spacing of the simulations is set to be in the range of $\Delta x^+ = 50\text{--}70$, $\Delta y_{\min}^+ = 1\text{--}2$ and $\Delta z^+ = 20\text{--}30$, in order to properly resolve the near-wall motions. We note that this grid spacing is finer than that in Hwang (2015) where the numerical technique applied here is fully verified

Case	Re_m	Re_τ	L_x/h	L_z/h	$N_x \times N_y \times N_z$	C_s	$T_{avg}u_\tau/h$
<i>L950a</i>	38 133	941	1.5	0.75	$24 \times 81 \times 24$	0.05	247
<i>SL950a</i>	38 133	936	1.5	0.75	$24 \times 81 \times 24$	0.20	246
<i>L950b</i>	38 133	976	2.0	1.0	$32 \times 81 \times 32$	0.05	256
<i>SL950b</i>	38 133	1004	2.0	1.0	$32 \times 81 \times 32$	0.25	263
<i>O950</i>	38 133	997	3.0	1.5	$48 \times 81 \times 48$	0.05	262
<i>SO950</i>	38 133	1152	3.0	1.5	$48 \times 81 \times 48$	0.40	302
<i>L1800a</i>	73 333	1446	0.75	0.375	$24 \times 129 \times 24$	0.05	84
<i>SL1800a</i>	73 333	1438	0.75	0.375	$24 \times 129 \times 24$	0.20	84
<i>L1800b</i>	73 333	1606	1.0	0.5	$32 \times 129 \times 32$	0.05	93
<i>SL1800b</i>	73 333	1685	1.0	0.5	$32 \times 129 \times 32$	0.30	98
<i>L1800c</i>	73 333	1745	1.5	0.75	$48 \times 129 \times 48$	0.05	102
<i>SL1800c</i>	73 333	1954	1.5	0.75	$48 \times 129 \times 48$	0.40	114

TABLE 1. Simulation parameters in the present study. Here, Re_m is the Reynolds number based on the bulk velocity and T_{avg} is the time interval for the average. In the names of the simulations, L indicates ‘logarithmic’, O ‘outer’ and S ‘self-sustaining’.

with very long L_x . Finally, it should be mentioned that significantly long average times (T_{avg}) are considered to ensure sufficient convergence of the time correlation functions introduced in § 3. The convergence of the correlation functions have also been checked by doubling T_{avg} , but it is not significantly improved.

2.2. Validation and preliminary test

To verify the present LES, a reference simulation for $C_s = 0.05$ without applying (2.1) is first performed with the domain of *O950*: i.e. the minimal unit for the outer attached eddies (Hwang & Cossu 2010c). Figure 1 shows the mean-velocity profile and turbulent velocity fluctuations of the reference simulation. The statistics of the simulation show reasonable agreement with those of DNS by del Álamo *et al.* (2004), consistent with Lozano-Durán & Jiménez (2014) who reported the effect of the computational domain size on the statistics. As discussed in § 2.1, the present numerical experiment also implements the filtering action (2.1) unlike the ordinary minimal box simulation which only considers a spatially confined computation box. Therefore, the statistics of the simulation for $C_s = 0.05$ with (2.1) is also checked (the dashed line in figure 1). Although the filtering (2.1) a little distorts the mean-velocity profile and the velocity fluctuations roughly for $y^+ > 150$ ($y > 0.15h$), it does not yield significant change overall in the statistics. Further inspection is performed by computing the time correlation functions of interest defined in § 3, and reveals that the filtering (2.1) does not significantly change the dynamical process in the considered computational domain. For further details, the reader refers to appendix A where a detailed comparison of the computed time correlation functions is made.

The only difference between the present numerical experiment and that in Hwang (2015) is the streamwise size of the computational domain. To ensure that the change of the streamwise computational domain does not affect the self-similar nature of the computed attached eddies, the statistics of the attached eddies, obtained by implementing the technique in § 2.1, are also reported. Figure 2 shows the normalised second-order statistics of the simulations in which the attached eddies at the given




FIGURE 1. (a) Mean-velocity profile and (b) turbulent velocity fluctuations of the reference simulation: —, O950 without the filtering (2.1); ---, O950; - · - · -, DNS at $Re_\tau = 934$ (del Álamo *et al.* 2004).




FIGURE 2. Normalised second-order statistics of the attached eddies, scaled with the given spanwise length L_z : (a) streamwise velocity, $y = 0.3L_z$; (b) wall-normal velocity, $y = 0.4L_z$; (c) spanwise velocity, $y = 0.4L_z$; (d) Reynolds stress, $y = 0.2L_z$. Here, ---, from $Re_\tau \simeq 950$ (SL950a, SL950b); —, from $Re_\tau \simeq 1800$ (SL1800a, SL1800b, SL1800c).

spanwise length scale are isolated (i.e. the simulations tagged with ‘S’ in table 1). The computed statistics of the attached eddies are approximately self-similar with respect to their spanwise length scale L_z below certain wall-normal locations: the streamwise velocity fluctuation is self-similar roughly below $y \simeq 0.3L_z$ (figure 2a), the

wall-normal and the spanwise velocity fluctuations are below $y \simeq 0.4L_z$ (figure 2*b,c*) and the Reynolds stress is below $y \simeq 0.2L_z$ (figure 2*d*). Above these wall-normal locations, the second-order statistics depend on the box size, as in Hwang (2015). However, we note that the non-self-similar part of the fluctuations above these wall-normal locations is merely a fluctuation induced by the self-similar part in the wall-normal locations supposed to be ‘empty’, due to the absence of the motions larger than the attached eddies of interest (for further discussion, see also Hwang 2013, 2015). Finally, it should be mentioned that the second-order statistics of the wall-parallel velocity components show non-negligibly large contributions to the region close to the wall (figure 2*a,c*), whereas those of the wall-normal velocity and the Reynolds stress are fairly small (figure 2*b,d*): for example, at $y = 0.05L_z$, the normalised streamwise and the spanwise velocity fluctuations are respectively approximately 0.98 and 0.68 times their maxima, whereas the wall-normal velocity and Reynolds stress are respectively found to be approximately 0.28 and 0.45 times their maxima. This is essentially due to the impermeability condition caused by the presence of the wall, and is exactly as in the statistical structure of the single attached eddy hypothesised by Townsend (1976).

3. Results

3.1. The largest attached eddies: large-scale and very-large-scale motions

We first explore the temporal evolution of the largest attached eddies (i.e. the energy-containing motions at $\lambda_z = 1.5h$) in the minimal unit (O950 and SO950 in table 1). As discussed in Hwang (2015), the largest attached eddy is composed of a VLSM (outer streaky motion) and LSMs (outer streamwise vortical structures) aligned with it. In a large computational domain, the VLSM is typically featured to be a long structure mainly carrying a large amount of the streamwise turbulent kinetic energy, and this is quite similar to the behaviour of the streaks in the near-wall region (Hwang 2015). Therefore, in the outer minimal unit, in which only single VLSM would be resolved mainly by zero streamwise wavenumber due to its long length (i.e. $k_x = 0$; see also figure 6), its temporal evolution would be well tracked by

$$E_u = \frac{1}{2V_h} \int_{\Omega_h} u^2 dV, \quad (3.1a)$$

where u is the streamwise velocity fluctuation, Ω_h the lower (or upper) half of the computational domain of interest and V_h the volume of the half of the computational domain.

On the contrary, the LSM is a relatively short structure given with its streamwise length scale at $\lambda_x \simeq 3\text{--}4h$. The structure at this length is identical to that often postulated as a hairpin vortex packet by Adrian and coworkers (Adrian 2007). This structure is relatively isotropic, compared to the VLSM which mostly carries the streamwise velocity fluctuation, in the sense that it contains all the velocity components (Hwang 2015). It is important to mention that this structure was found to share a number of statistical similarities with quasi-streamwise vortex in the near-wall region (Hwang 2015). This suggests that the LSMs would likely to be quasi-streamwise vortical structures in the outer region, although they are much more disorganised due to small-scale turbulence associated with cascade and turbulent dissipation (Jiménez 2012). It should be mentioned that this view is consistent with del Álamo *et al.* (2006) who called this structure the ‘tall-attached vortex cluster’.




FIGURE 3. Time evolution of the flow field of O950: (a) time trace of E_u^+ (solid) and E_v^+ (dashed); (b) magnification of (a) for $tu_\tau/h \in [158, 164]$; (c–f) the corresponding flow visualisation at $tu_\tau/h = 159, 160.2, 160.9, 162$. In (c–f), the red and blue iso surfaces indicate $u^+ = -3.2$ and $v^+ = 1.4$, respectively.

Since the VLSM carries very little wall-normal or spanwise turbulent kinetic energy, in the minimal unit, the LSMs would be well characterised by either

$$E_v = \frac{1}{2V_h} \int_{\Omega_h} v^2 \, dV, \quad (3.1b)$$

or

$$E_w = \frac{1}{2V_h} \int_{\Omega_h} w^2 \, dV, \quad (3.1c)$$

where v and w are the wall-normal and spanwise velocity fluctuations, respectively. Later, we shall see that these two variables are indeed strongly correlated to each other (figure 5c,d). Here, it should also be pointed out that the use of the primitive variables in characterising the quasi-streamwise vortical structure instead of enstrophy or streamwise vorticity is intentionally introduced to track their behaviour more precisely. Given the significant amount of small-scale turbulence associated with energy cascade and turbulent dissipation in the logarithmic and the outer regions, the variables such as enstrophy or streamwise vorticity would not faithfully represent the large-scale organised vortical structure. Indeed, Jiménez (2012) recently showed that such variables carry dominant energy around turbulent dissipation length scales.

A set of the flow fields of O950, in which the largest attached eddies are contained with other smaller-scale turbulent motions, are first examined. Figure 3 shows the time trace of E_u and E_v , and the corresponding evolution of the flow field in time. Both E_u and E_v show large-scale temporal oscillations with a time scale roughly at $Tu_\tau/h \simeq 2-5$ (figure 3a). This is exactly the feature known as ‘bursting’ by Flores &




FIGURE 4. Time evolution of the flow field of *SO950*: (a) time trace of E_u^+ (solid) and E_v^+ (dashed); (b) magnification of (a) for $tu_\tau/h \in [564, 572]$; (c–f) the corresponding flow visualisation at $tu_\tau/h = 565, 568.5, 569.3, 570$. In (c–f), the red and blue iso surfaces indicate $u^+ = -4$ and $v^+ = 1.5$, respectively.

Jiménez (2010). A careful observation reveals that E_u and E_v oscillate with a certain phase difference: for example, at $tu_\tau/h = 159$, E_v is around the local maximum while E_u being at fairly low-energy state (figure 3*b*), and, at $tu_\tau/h = 160.2$, E_v reaches the local minimum whereas E_u becomes considerably larger (figure 3*b*). Visualisation of the corresponding flow fields in figure 3(c–f) suggests that this is likely due to the interactive dynamics between the VLSM (streak) and the LSMs (streamwise vortical structures). At $tu_\tau/h = 159$ (figure 3*c*), the flow field exhibits a few fairly strong large-scale v structures, which would be a part composing the LSMs in the minimal unit. On the other hand, u structures in the flow field are fairly weak at this time, and they appear to be collectively gathered around the v structures. At $tu_\tau/h = 160.2$ (figure 3*d*), the v structures become considerably weakened whereas the u structures are significantly amplified, forming a strong streaky structure extending over the entire streamwise domain. This amplification of the streaky structure appears to be a consequence of the ejection of the streamwise momentum by the strong v structures observed at $tu_\tau/h = 159$ (figure 3*c*), clearly reminiscent of the ‘lift-up’ effect predicted by previous theoretical studies (del Álamo & Jiménez 2006; Cosu, Pujals & Depardon 2009; Pujals *et al.* 2009; Hwang & Cosu 2010*b*). The amplified streaky structure subsequently meanders along the streamwise direction (figure 3*e*). This process eventually leads to regeneration of new v structures with the breakdown of the streaky motion (figure 3*f*).

Qualitatively the same behaviour is observed in the over-damped simulation *SO950*, in which only the energy-containing motions at $\lambda_z = 1.5h$ are isolated by replacing all the smaller-scale motions with a crude eddy viscosity given with the increased C_s . Figure 4 shows the time trace of E_u and E_v , and the corresponding evolution of the flow field of *SO950*. As in *O950* (figure 3*a*), both E_u and E_v exhibit temporal

oscillations with the time scale at $Tu_\tau/h = 2-5$ (figure 4a). In this case, a more prominent phase difference is observed between the oscillations of E_u and E_v (figure 4a,b), presumably because the smaller-scale surrounding turbulent motions are removed in *SO950*. However, this could also be partially due to the increased oscillation amplitude of E_u in *SO950*, which has previously been shown to be an artefact caused either by the crude eddy viscosity itself or by the lack of the energy-containing motions in the near-wall and logarithmic regions (Hwang 2015). Not surprisingly, the flow fields of *SO950* are composed of much smoother structures than those of *O950* (figure 4c-f). However, the temporal evolution of the u and v structures of *SO950* are remarkably similar to that of *O950*: the v structures significantly amplify the streaky u structure via the lift-up effect (figure 4c,d); the amplified streaky u motion subsequently meanders along the streamwise direction (figure 4e); the v structures are finally regenerated with breakdown of the streaky u motion (figure 4f).

To statistically quantify this cyclic dynamical process, auto- and cross-correlations of several variables of interest are computed. It is also useful to introduce two additional variables, such that:

$$E_0 = \int_{\Omega_{y,h}} |\hat{u}(y; k_x, k_z)|^2 + |\hat{v}(y; k_x, k_z)|^2 + |\hat{w}(y; k_x, k_z)|^2 dy, \quad (3.2a)$$

for $k_x = 0$ and $k_z = 2\pi/L_z$, and

$$E_1 = \int_{\Omega_{y,h}} |\hat{u}(y; k_x, k_z)|^2 + |\hat{v}(y; k_x, k_z)|^2 + |\hat{w}(y; k_x, k_z)|^2 dy, \quad (3.2b)$$

for $k_x = 2\pi/L_x$ and $k_z = 2\pi/L_z$. Here, $\hat{\cdot}$ denotes the Fourier-transformed state in the x and z directions, and $\Omega_{y,h}$ is the lower (or upper) half of the wall-normal domain. We note that, in (3.2), E_0 is introduced to compute the energy of the streamwise uniform component of the motions at $\lambda_z = 1.5h$, while E_1 is to compute the energy of the first streamwise Fourier component, which would measure the meandering motion observed in figures 3 and 4. The correlation functions are defined as

$$C_{ij}(\tau) = \frac{\langle E_i(t + \tau)E_j(t) \rangle}{\sqrt{\langle E_i^2(t) \rangle} \sqrt{\langle E_j^2(t) \rangle}}, \quad (3.3)$$

where $i, j = u, v, w, 0, 1$ and $\langle \cdot \rangle$ indicates average in time. In the present study, all the correlation functions are computed by averaging the lower and upper half-channels.

Figure 5 shows the correlation functions computed with E_u , E_v and E_w from *O950* and *SO950*. In the case of *O950*, all the auto-correlations decay to zero at $|\tau u_\tau/h| \simeq 1-2$, although C_{uu} tends to drop a little more slowly than C_{vv} and C_{ww} (figure 5a). The size of the time interval, in which the auto-correlations remain positive, is $Tu_\tau/h \simeq 2-4$, and this roughly corresponds to the single period of the temporal oscillation of E_u and E_v (figure 3a). The cross-correlation C_{uv} of *O950* also reveals that there is indeed a phase difference between E_u and E_v (figure 5c): the peak of C_{uv} is located at $\tau u_\tau/h \simeq -0.6$, indicating that E_u statistically reaches its local extremum approximately $\Delta\tau u_\tau/h \simeq 0.6$ before E_v reaches its local extremum. On the other hand, E_v and E_w are found to be strongly correlated to each other: $C_{vw}(\tau = 0) \simeq 0.7$ and C_{uw} also exhibits a peak at $\tau u_\tau/h \simeq -0.6$ as C_{uv} does (figure 5c).




FIGURE 5. Time correlation functions of (a,c) O950 and (b,d) SO950. In (a,b), —, $C_{uu}(\tau)$; ----, $C_{vv}(\tau)$; - · - · -, $C_{wv}(\tau)$. In (c,d), —, $C_{uv}(\tau)$; ----, $C_{uw}(\tau)$; - · - · -, $C_{vw}(\tau)$.

Qualitatively the same behaviour is observed in the correlation functions of SO950 in which only the energy-containing motions at $\lambda_z \simeq 1.5h$ are isolated (figure 5b,d). All the auto-correlations drop to zero near $|\tau u_\tau/h| \simeq 1.5-2$ (figure 5b), resulting in the time interval of positive auto-correlation to be $Tu_\tau/h \simeq 4$. As for O950, this time scale reasonably well quantifies a single period of the temporal oscillations of E_u and E_v (figure 4a). The phase difference between E_u and E_v is also seen in C_{uv} of SO950 (figure 5d) which shows the peak at $\tau u_\tau/h \simeq -1.7$. Finally, E_v and E_w are also found to be strongly correlated to each other, as seen in C_{vw} (figure 5d).

The correlation functions with E_0 and E_1 are also presented in figure 6. The auto- and cross-correlations obtained only with E_0 and E_1 (i.e. C_{00} , C_{11} and C_{01}) are given in figure 6(a,b). The correlation functions of O950 show fairly similar behaviour to those of SO950: C_{00} and C_{11} of O950 exhibit similar time scales to those of SO950, and C_{01} of both O950 and SO950 has a phase difference between E_0 and E_1 with $\Delta\tau u_\tau/h = 1$. The phase difference between E_0 and E_1 implies that the streamwise uniform motion occurs first and the meandering motion follows, as in the instantaneous fields shown in figures 3 and 4. Here, it is interesting to note that the time scale of C_{11} ($Tu_\tau/h \simeq 2$) appears to be much smaller than that of C_{00} as well as that of C_{uu} , C_{vv} and C_{wv} ($Tu_\tau/h \simeq 4$; see also figure 5a,b). This suggests that the meandering motion of the streak (figures 5c and 6c), which would be characterised by E_1 , is a rapid process which persists only for roughly half of the bursting period.

Cross-correlations are further computed by also considering E_u , E_v and E_w , as shown in figure 6(c,d). The correlation C_{0u} shows that E_0 and E_u are strongly




FIGURE 6. Time correlation functions of (a,c) *O950* and (b,d) *SO950*. In (a,b), —, $C_{00}(\tau)$; - - - - , $C_{11}(\tau)$; - - - - - , $C_{01}(\tau)$. In (c,d), —, $C_{0u}(\tau)$; - - - - , $C_{0v}(\tau)$; - - - - - , $C_{1v}(\tau)$.

correlated to each other in both *O950* and *SO950*, indicating that the streamwise uniform mode ($k_x = 0$) at $\lambda_z = 1.5h$ well represents the streaky motion (i.e. VLSM) in the minimal unit. This feature can also be confirmed from C_{0v} of *O950* and *SO950*, which behaves very similarly to C_{uv} (figure 5c,d). Finally, the peak of C_{1v} for both *O950* and *SO950* is found roughly at $\tau u_\tau/h \simeq -1$. This indicates that the meandering motion of the streak statistically appears before the vortical structures (LSMs) are fully amplified, consistent with the instantaneous flow fields given in figures 3 and 4.

Despite many qualitative similarities between the dynamical behaviours of the energy-containing motions in *O950* and *SO950*, it should be pointed out that the correlation functions of the two simulations (*O950* and *SO950*) are not quantitatively the same, although this is not very surprising given the aggressive nature of the present numerical experiment: for example, the time intervals of $C_{vv} > 0$ and $C_{ww} > 0$ of *O950* are smaller than those of *SO950* (figure 5a,b), and the peak locations of C_{uw} , C_{uv} and C_{0v} of *O950* are a little different from those of *SO950* (figures 5c,d, 6c,d). Apparently, these differences could stem from two possible origins: one is that E_u , E_v and E_w of *O950* contain the effect of the surrounding smaller-scale turbulent motions, and the other is that the dynamical behaviour of the motions in *SO950* is probably a little distorted by the artificially increased eddy viscosity of *SO950*. The issue of which of the two more dominantly yields the differences could be clarified by further inspecting C_{00} , C_{11} and C_{01} (figure 6a,b), as they are, in a way, obtained by applying cutoff filters to both of *O950* and *SO950*. It appears that the correlation functions of *O950* are not exactly the same as those of *SO950*, suggesting that

increasing C_s a little distorts the motions at $\lambda_z = 1.5h$. However, the times scales of these correlation functions for $O950$ do not appear to be significantly different from those for $SO950$. This implies that the correlation functions obtained with E_u , E_v and E_w of $O950$ do not precisely represent the dynamical behaviour of the largest attached eddies at $\lambda_z = 1.5h$ due to the contribution by the surrounding smaller-scale turbulent fluctuations.

3.2. The attached eddies in the logarithmic region

Now, we consider the attached eddies in the logarithmic region. As shown in §2.2, the attached eddies in the logarithmic region generate a self-similar statistical structure with respect to the spanwise length scale $\lambda_{z,0}$ ($= L_z$). This self-similar part, which contains the essential physical process of the energy-containing motions in the logarithmic region, appears mainly below $y \simeq 0.3\text{--}0.4L_z$ (figure 2). To trace the dynamical behaviour of this self-similar part more precisely, the definitions of E_u , E_v , E_w , E_0 and E_1 given in (3.1) and (3.2) are a little modified, so that the wall-normal domain of the integration becomes only $[0, 2/3L_z]$. We note that this new definition is still consistent with (3.1), as the wall-normal domain of the integration becomes $[0, 1]$ for $L_z = 1.5h$, which is the spanwise length scale of the VLSM and the LSM.

Inspection of the instantaneous flow fields of all the simulations in the logarithmic region (see also table 1) reveals that basically the same dynamical process occurs in the attached eddies, given with streaks and quasi-streamwise vortical structures, at each of the spanwise length scales belonging to the logarithmic region: i.e. amplification of the streaks by the quasi-streamwise vortical structures, subsequent meandering of the streaks along the streamwise direction and breakdown of the streaks with regeneration of the quasi-streamwise vortical structures. To avoid repetition of the same discussion given in §3.1, here we only report auto- and cross-correlation functions of the simulations concerning the logarithmic region with a focus on the scaling of the computed correlation functions.

Figure 7 shows auto-correlations obtained from all the simulations concerning the energy-containing motions in the logarithmic region. All the correlation functions from both full (figure 7*a,c,e*) and over-damped simulations (figure 7*b,d,f*) scale fairly well with the spanwise size of the attached eddies L_z ($= \lambda_{z,0}$), suggesting that the attached eddies in the minimal unit are ‘dynamically’ self-similar with respect to the spanwise size. The auto-correlation functions C_{uu} and C_{vv} in figure 7(*a–d*) reach zero at $\tau u_\tau/L_z \simeq \pm 1$, indicating that the single period of the self-sustaining process would be roughly given by $Tu_\tau/L_z \simeq 2$. As in the case of the outer attached eddies, C_{11} , which characterises the time scale of the meandering motion of the streak, is found to have much shorter time scale $Tu_\tau/L_z \simeq 1$ than that of C_{uu} and C_{vv} .

Figure 8 shows several cross-correlations obtained from the same simulations. The cross-correlation C_{uw} of both full (figure 8*a*) and over-damped simulations (figure 8*b*) exhibits a phase difference between E_u and E_v (or E_w), similar to that of the simulations concerning the outer-scaling motions (figure 5*c,d*). A strong correlation between E_v and E_w is also observed in the motions in the logarithmic region, similarly to those in the outer region (figure 8*c,d*). Finally, the cross-correlation C_{1v} also confirms that amplification of E_v appears a little after E_1 is amplified, indicating that the vortical structures in the logarithmic region are regenerated after the meandering motion (figure 8*e,f*).




FIGURE 7. Auto-correlation functions: (a,b) $C_{uu}(\tau)$; (c,d) $C_{vv}(\tau)$; (e,f) $C_{11}(\tau)$. In (a,c,e), — $L950a$; - - - $L950b$; - - - - $L1800a$; - · - · - $L1800b$; · · · · · $L1800c$, while, in (b,d,f), — $LS950a$; - - - $LS950b$; - - - - $LS1800a$; - · - · - $LS1800b$; · · · · · $LS1800c$.

4. Discussion

Thus far, we have explored the dynamical behaviour of the attached eddies in the logarithmic and outer regions. The attached eddies in the minimal unit exhibit relatively large-scale temporal oscillations, ‘bursting’ as reported by Flores & Jiménez (2010). The bursting is also observed in the over-damped simulations, in which the attached eddies only at the given spanwise length scale survive through their self-sustaining mechanisms. The qualitatively good comparison of instantaneous flow fields and correlation functions between the full and the over-damped simulations suggests that the bursting is presumably the reflection of the self-sustaining process of the attached eddies with the characteristic turnover time scale given by:

$$\frac{Tu_\tau}{\lambda_z} \simeq 2, \quad (4.1)$$




FIGURE 8. Cross-correlation functions: (a,b) $C_{uv}(\tau)$; (c,d) $C_{vw}(\tau)$; (e,f) $C_{lv}(\tau)$. In (a,c,e), — $L950a$; - - - $L950b$; - - - - $L1800a$; - · - · - $L1800b$; · · · · · $L1800c$, while, in (b,d,f), — $SL950a$; - - - $SL950b$; - - - - $SL1800a$; - · - · - $SL1800b$; · · · · · $SL1800c$.

where λ_z is the spanwise size of the given attached eddies, which corresponds to the spanwise domain size of the minimal unit L_z . Here, we note that if the spanwise length scale is chosen as $\lambda_z^+ \simeq 100$, (4.1) gives $T^+ \simeq 200$, the well-known bursting period of the near-wall coherent structures in the minimal unit (Hamilton, Kim & Waleffe 1995; Jiménez *et al.* 2005). On the other hand, if $\lambda_z = 1.5h$ is considered, (4.1) yields $Tu_\tau/h \simeq 3$, which roughly corresponds to the bursting time scale of the largest attached eddies composed of VLSMs and LSMs (see § 3.1). The attached eddies the spanwise size of which is between $\lambda_z^+ = 100$ and $\lambda_z = 1.5h$ burst in a self-similar manner with the time scale given by (4.1) (see § 3.2). We note that if $\lambda_z = 3y$ is chosen, the time scale given in (4.1) becomes consistent with the bursting time scale of $Tu_\tau/y \simeq 6$ reported by Flores & Jiménez (2010) who extracted this time scale




FIGURE 9. A schematic diagram of self-sustaining process of the attached eddies (originally proposed by Hamilton *et al.* (1995) for the near-wall motions). The diagram incorporates the streak transient growth proposed by Schoppa & Hussain (2002) for the near-wall motions as well as the Orr mechanism proposed by Jiménez (2013, 2015). Here, u_s denotes the streamwise velocity given with the amplified streaks.

from a direct numerical simulation, and, interestingly, $\lambda_z = 3y$ corresponds well to the scaling $y \simeq 0.3\text{--}0.4\lambda_z$, below which the computed statistics of the isolated attached eddies are found to be self-similar (see also figure 2).

In the present study, it is also shown that the self-sustaining process of the attached eddies involves its two dynamically interconnected structural elements: one is the long streaky motion extending over the entire streamwise domain of the minimal unit (figure 6*c,d*), and the other is the vortical structure which would be statistically in the form of quasi-streamwise vortices given the strong correlation between the wall-normal and spanwise velocities (figures 5*c,d* and 8*c,d*). The interactive dynamics between the two elements appears to be the backbone of the self-sustaining process of the attached eddies in the logarithmic and the outer regions, and is remarkably similar to that in the near-wall region (Hamilton *et al.* 1995; Schoppa & Hussain 2002). It should be pointed out, however, that this should not be very surprising, given the fact that the near-wall motion in the form of the near-wall streak and the quasi-streamwise vortices would simply be the smallest attached eddy (Hwang 2015).

Figure 9 is a schematic diagram of the self-sustaining process of the attached eddies at a given spanwise length scale proposed in the present study based on the observations made in the previous section. This schematic diagram is originally from Hamilton *et al.* (1995), and, here, we further incorporate several previous contributions in this context (Schoppa & Hussain 2002; Jiménez 2013, 2015). The streaks are amplified via the ‘lift-up’ effect by which the streamwise vortical structures transfer energy of the mean shear to streaks (figures 3*c,d* and 4*c,d*). The amplified streaks then undergo ‘rapid’ meandering or oscillatory motion in the streamwise direction via streak instability or transient growth (see also § 3.2; Hamilton *et al.* 1995; Schoppa & Hussain 2002; Park, Hwang & Cossu 2011). The quasi-streamwise vortical structures

are finally regenerated with breakdown of the streak via linear and/or nonlinear mechanisms (Hamilton *et al.* 1995; Schoppa & Hussain 2002; Jiménez 2013, 2015). In order to provide more solid evidence on the existence of these physical processes, we now introduce two additional numerical experiments, designed to examine the lift-up effect (the left branch from streamwise vortical structures to streaks in figure 9) and the streak meandering and regeneration of the streamwise vortical structures (the right and lower branches from streaks to streamwise vortical structures in figure 9), respectively.

4.1. Streak amplification: lift-up effect

The ‘lift-up’ effect has been very well understood as the robust mechanism for generation of the streaky motions both in transitional (Ellingsen & Palm 1975; Landahl 1980; Gustavsson 1991; Butler & Farrell 1992; Reddy & Henningson 1993; Schmid & Henningson 2001) and turbulent flows (Landahl 1990; Butler & Farrell 1993; Kim & Lim 2000; Chernyshenko & Baig 2005; del Álamo & Jiménez 2006; Cossu *et al.* 2009; Pujals *et al.* 2009; Hwang & Cossu 2010*a,b*; Willis, Hwang & Cossu 2010). From a vortex dynamical viewpoint, it simply represents tilting of the streamwise vortices by mean shear: i.e.

$$\frac{D\omega_y}{Dt} \sim \frac{dU}{dy}\omega_x, \tag{4.2}$$

where D/Dt is the material time derivative, ω_x and ω_y are the streamwise and wall-normal vorticities, respectively. The ‘lift-up’ effect is an important origin of the ‘non-normality’ of the linearised Navier–Stokes operator, leading to a large amplification of an initial condition as well as a body forcing containing a significant amount of the wall-normal velocity component. Therefore, the non-modal stability analysis (Schmid & Henningson 2001; Schmid 2007), which quantifies the amplification mechanism of the stable linearised Navier–Stokes operator, has been a popular methodology of most of the previous investigations. However, direct relevance of the ‘lift-up’ effect to fully developed turbulence has been very rarely shown, except by Kim & Lim (2000) who demonstrated its importance on the near-wall turbulence at a low Reynolds number by performing a direct numerical simulation without the off-diagonal term of the Orr–Sommerfeld–Squire operator representing (4.2).

The goal of this section is to demonstrate that the ‘lift-up’ effect is the essential part of the self-sustaining process of the attached eddies in the logarithmic and outer regions with a similar approach of Kim & Lim (2000). To this end, we modify their approach to suppress the lift-up effect only at a prescribed spanwise length scale. In the minimal unit, the streak amplification by the lift-up effect is found to be dominant at infinitely long streamwise wavelength (figure 6*c,d*). To artificially suppress this lift-up effect at zero streamwise wavenumber ($k_x = 0$), the following modified streamwise momentum equation is solved with the minimal unit:

$$\frac{\partial \bar{u}}{\partial t} + (\mathbf{u} \cdot \nabla)\bar{u} - ([\hat{v}e^{ik_z z} + \hat{v}^*e^{-ik_z z}]|_{(k_x, k_z)=(0, 2\pi/L_z)}) \frac{d\bar{U}}{dy} = -\frac{1}{\rho} \frac{\partial p}{\partial x} + (\nu + \nu_T)\nabla^2 \bar{u}, \tag{4.3}$$

where the superscript * denotes the complex conjugate, \bar{u} is the streamwise velocity and \bar{U} is the streamwise velocity averaged along the streamwise and spanwise directions at each time. In (4.3), the energy extraction mechanism from the mean shear \bar{U} by the wall-normal velocity (i.e. lift-up effect) is now artificially eliminated for the streamwise uniform streaky motions ($k_x = 0$) at $\lambda_z = L_z$ by adding the third




FIGURE 10. Time trace of E_u and E_v of *SO950*. At $tU_e/h = 7500$ (U_e is the centreline velocity of the laminar velocity profile with the same volume flux), the streamwise momentum equation is replaced by (4.3). Here, E_u and E_v are normalised with U_e .

term on the left-hand side. We also note that this is equivalent to removing the off-diagonal term of the Orr–Sommerfeld–Squire operator for $k_x = 0$ and $k_z = \pm 2\pi/L_z$, as in the approach by Kim & Lim (2000) whose numerical solver is written in the wall-normal velocity and vorticity form.

The designed numerical experiment is first performed with *SO950* in which only the largest attached eddies sustain themselves without any smaller-scale turbulent motions. Figure 10 shows time trace of E_u and E_v . Initially, a normal *SO950* simulation is performed, yielding the non-trivial fluctuations of E_u and E_v via the self-sustaining process. At $tU_e/h = 7500$ (U_e is the centreline velocity of the laminar velocity profile with the same volume flux), the streamwise momentum equation is replaced by (4.3). Immediately after this time, E_u , which characterises the temporal evolution of the streak (the VLSM in this case), very rapidly decays, indicating that the streaky motion is very quickly destroyed by implementing (4.3) (figure 10*a*). We note that, in $tU_e/h \in [7500, 7600]$, this decay of E_u appears to be even faster than E_v , finally resulting in complete suppression of the self-sustaining process in *SO950*. The very rapid destruction of the streaky motion by implementing (4.3) is consistently observed with a few other initial conditions. This clearly suggests that the amplification of the long streaky motion is indeed governed by the ‘lift-up’ effect, consistent with a number of previous theoretical predictions along this line (del Álamo & Jiménez 2006; Pujals *et al.* 2009; Hwang & Cosu 2010*b*).

The numerical experimentation with (4.3) is also applied to *O950* in which the largest attached eddies are resolved with turbulent motions at other length scales. Not surprisingly, *O950* is not completely suppressed by (4.3), as the self-sustaining mechanisms at other scales would still be undisturbed. In figure 11, we compare




FIGURE 11. One-dimensional spanwise wavenumber spectra of (a,b) the streamwise and (c,d) the wall-normal velocities: (a,c) *O950*; (b,d) *O950* with (4.3). Here, the spectra are normalised with U_e for the purpose of comparison.

the spanwise wavenumber spectra of *O950* (figure 11a,c) with those of *O950* with (4.3) (figure 11b,d). The streamwise velocity spectra of *O950* with (4.3) clearly show a significant amount of reduction of the streamwise turbulent kinetic energy at $\lambda_z = 1.5h$ compared to those of *O950* (figure 11a,b), indicating that the streaky motion at $\lambda_z = 1.5h$ (VLSM) under full turbulence is also destroyed by inhibiting the lift-up effect. This also appears to yield reduction of the wall-normal turbulent kinetic energy at $\lambda_z = 1.5h$ for $y > 0.5h$ (figure 11c,d), as this length scale corresponds to that of the streamwise vortical structures involved in the self-sustaining process at $\lambda_z = 1.5h$ (see also (1.1c,d) with $\lambda_z = 1.5h$). However, it is quite interesting to note that the wall-normal turbulent kinetic energy at other spanwise wavelengths, particularly the ones near $\lambda_z \simeq 0.75h$, appears to be significantly amplified with the removal of the streaky motions at $\lambda_z = 1.5h$, and generates additional energy even at $\lambda_z = 1.5h$. This suggests that the non-trivial scale interactions exist between the attached eddies at neighbouring length scales, although the detailed investigation of this issue is beyond the scope of the present study. Inspection of the streamwise wavenumber spectra shown in figure 12 also allows us to reach the same conclusion. Since (4.3) only applies to the streamwise uniform streaky motion (i.e. $k_x = 0$), the spectra of the streamwise velocity show only a little change (figure 12a,b). However, with the destruction of the streaky motions uniform along the streamwise direction, the spectra of the wall-normal velocity at $\lambda_x \simeq 3h$ show a considerable amount of reduction of energy for $y > 0.5h$ (figure 12c,d), indicating that the streamwise vortical structures associated with the self-sustaining process at $\lambda_z = 1.5h$ are destroyed. We




FIGURE 12. One-dimensional streamwise wavenumber spectra of (a,b) the streamwise and (c,d) the wall-normal velocities: (a,c) O950; (b,d) O950 with (4.3). Here, the spectra are normalised with U_e for the purpose of comparison.

note that the increased energy at $\lambda_x \simeq 1.5h$ in the streamwise wavenumber spectra of the wall-normal velocity should be associated with the increased energy of the wall-normal velocity near $\lambda_z \simeq 0.75h$ (figure 12d).

Finally, it has been robustly found that suppressing the lift-up effect through (4.3) yields a considerable amount of turbulent skin-friction reduction, as reported in table 2 showing that the smaller the computational box is, the larger the drag reduction achieved. A considerable amount of turbulent skin-friction reduction by suppressing the lift-up effect here suggests that generation of turbulent skin-friction at high Reynolds numbers would be associated with the momentum transfer to the wall by the self-sustaining processes of the attached eddies in the logarithmic and outer region. This result also indicates that applying a linear flow control, e.g. with the approaches in Kim & Bewley (2007), to relatively large-scale attached eddies would be a promising strategy for turbulent drag reduction. Despite many interesting further issues, a detailed investigation on the relation between turbulent skin friction and the self-sustaining process (or lift-up effect) in the logarithmic and outer regions is beyond the scope of the present study and is currently a subject of on-going investigation.

4.2. Streak breakdown and regeneration of streamwise vortical structures

In contrast to the ‘lift-up’ effect which has been an issue in a number of previous studies, the breakdown of amplified streaks and the subsequent regeneration of the

Case	$\Delta C_f(\%)$	Case	$\Delta C_f(\%)$
L950a	20	L1800a	30
L950b	13	L1800b	25
O950	8	L1800c	18

TABLE 2. Skin-friction reduction ΔC_f by suppressing the lift-up effect with (4.3).

streamwise vortical structures (see figure 9) have been much less studied, even for the near-wall motions. In the case of the near-wall motions, which would be the smallest attached eddies (Hwang 2015), the amplified streak typically experiences a rapid sinuous meandering or oscillatory motion before the streamwise vortices are generated (Hamilton *et al.* 1995; Schoppa & Hussain 2002). This sinuous meandering motion is a consequence of the streak instability or transient growth (Hamilton *et al.* 1995; Schoppa & Hussain 2002), although, in practice, distinguishing one from another appears to be almost impossible, given the fact that both of the processes are basically a consequence of the interaction with the same sinuous-mode instability (Hœpffner, Brandt & Henningson 2005). It should also be mentioned that the sinuous mode instability essentially originates from the ‘spanwise shear’ generated by the amplified streak (i.e. $\partial u_s / \partial z$ where u_s is the streamwise velocity with the amplified streak; see also Park & Huerre (1995), Cossu & Brandt (2002), Hœpffner *et al.* (2005), Park *et al.* (2011)). It is therefore physically not meaningful to analyse this process using the approaches based on the linearised Navier–Stokes equation around ‘mean shear’, as the equation does not carry any relevant physics relating to this mechanism.

The sinuous meandering motion of the amplified streak feeds a small amount of the streamwise vortical structures, and they are subsequently amplified via nonlinear mechanisms (Hamilton *et al.* 1995; Schoppa & Hussain 2002). Hamilton *et al.* (1995) proposed that wall-normal advection of the streamwise vortices would be the leading nonlinear regeneration mechanism based on a minimal Couette flow simulation ($D\omega_x/Dt \sim v\partial\omega_x/\partial y$; figure 9). On the other hand, with a minimal channel flow simulation, Schoppa & Hussain (2002) showed that stretching of the streamwise vortices by the streamwise meandering (or wavy) motion caused by the streak instability or transient growth would play an important role in this process ($D\omega_x/Dt \sim \omega_x\partial u/\partial x$; figure 9). More recently, Jiménez (2013, 2015) proposed that the Orr mechanism, a process by which the wall-normal velocity takes energy from the mean shear, also appears to play a role in this process especially for the motions in the logarithmic and outer regions. At this moment, it is quite difficult to convincingly argue which of the mechanisms would be dominant or most important among the three, especially for the generation of the streamwise vortical structures in the logarithmic and outer regions. However, it does not appear that solely the Orr mechanism (Jiménez 2013, 2015), which merely concerns the amplification of the wall-normal velocity only, would play a dominant role, given the strong correlation between the wall-normal and spanwise velocities (figures 5c,d and 8c,d).

In the present study, it has been found that the rapid sinuous meandering motion of the amplified streak and the subsequent generation of the streamwise vortical structures are also robust features of the self-sustaining processes in the logarithmic and outer regions (figures 3c, 4c, 6c,d, and 8e,f). This suggests that the streak instability, which was previously analysed by Park *et al.* (2011) for the outer




FIGURE 13. Time trace of E_u and E_v of SO950. At $tU_e/h = 7500$, (4.4) is applied. Here, E_u and E_v are normalised with U_e .

attached eddies using the Floquet theory, or transient growth would also be an important process associated with the subsequent generation of the streamwise vortical structures. It is worth mentioning that this process is intricately tangled with the subsequent nonlinear amplification of the streamwise vortical structures, especially via the vortex stretching mechanism proposed by Schoppa & Hussain (2002), as the vortical structures take the energy required for the amplification from the meandering (or wavy) streaky motion (i.e. $\partial u_s/\partial x$).

Given this observation, it is very tempting to postulate that the streak meandering motion, presumably caused by the streak instability or transient growth, would play a crucial role in regeneration of the streamwise vortical structures. The goal of this section is to examine this mechanism. For this purpose, here, we introduce a numerical experiment which artificially damps the sinuous meandering motion of the amplified streak. Since the streak meandering motion is dominant at $\lambda_x \simeq L_x$ in the minimal unit (figures 3c and 4d), we implement a damping of this wave component by modifying the right-hand side of the discretised streamwise momentum equation at each Runge–Kutta substep, such that:

$$\widehat{\text{RHS}}_x \left(y; k_x = \frac{2\pi}{L_x}, k_z = \pm \frac{2\pi}{L_z} \right) \rightarrow \mu \widehat{\text{RHS}}_x \left(y; k_x = \frac{2\pi}{L_x}, k_z = \pm \frac{2\pi}{L_z} \right), \quad (4.4)$$

where μ is a factor which should be $\mu < 1$ for damping. Here, we again stress that this technique is implemented only to the streamwise momentum equation to only suppress the meandering motion of u .

The damping technique (4.4) is applied to both O950 and SO950 by considering a few different μ ($= 0.85, 0.90, 0.95$). However, the results are found to be qualitatively independent of the choice of μ , thus the results only for $\mu = 0.95$ are reported in




FIGURE 14. One-dimensional spanwise wavenumber spectra of (a,b) the streamwise and (c,d) the wall-normal velocities: (a,c) *O950*; (b,d) *O950* with (4.4). Here, the spectra are normalised with U_e for the purpose of comparison.

the present study. Figure 13 shows time trace of E_u and E_v of *SO950* in which (4.4) is applied at $tU_e/h = 7500$. Immediately after (4.4) is applied, E_v decays very quickly. Examination with a few other initial conditions reveals that this behaviour is qualitatively independent of the initial condition, suggesting that the streak meandering motion observed in *SO950* is indeed directly involved in generation of the streamwise vortical structures. In *SO950*, both E_u and E_v eventually decay to zero after application of (4.4), but, in this case, it is quite interesting to note that E_u persists for a substantially long time. This certainly differs from the behaviour of E_u observed after applying (4.3) (compare figure 13a with 10a), implying that the lift-up effect is a very strong amplification process of the streamwise velocity (or streak) even with a small amplitude of the vortical structures.

The damping technique (4.4) is also applied to *O950*. The streamwise and spanwise wavenumber spectra of the streamwise and the wall-normal velocity are compared with those of *O950*, as respectively reported in figures 14 and 15. Consistent with the case of *SO950*, application of (4.4) to *O950* yields significant reduction of the wall-normal turbulent kinetic energy at $\lambda_z = 1.5h$ (figure 14c,d), despite the increase of the streamwise turbulent kinetic energy in the spanwise wavenumber spectra (figure 14a,b; see also the next paragraph for a further discussion). The streamwise wavenumber spectra of the wall-normal turbulent kinetic energy also clearly exhibits a significant amount of reduction of the energy at $\lambda_x = 3h$ (figure 15c,d), indicating that the artificial damping of the streak meandering motion significantly destroys the generation mechanism of the streamwise vortical structures at this wavelength.




FIGURE 15. One-dimensional streamwise wavenumber spectra of (a,b) the streamwise and (c,d) the wall-normal velocities: (a,c) *O950*; (b,d) *O950* with (4.4). Here, the spectra are normalised with U_e for the purpose of comparison.

Despite the consistent result with *SO950*, in *O950*, application of (4.4) does not appear to completely destroy the structures at $\lambda_z = 1.5h$. Although the streamwise wavenumber spectra of the streamwise velocity show a large diminishing of the energy at $\lambda_z = 1.5h$ (figure 15a,b), confirming that (4.4) indeed inhibits the meandering motion of the streak at this wavelength, the increase of the streamwise turbulent kinetic energy in the spanwise wavenumber spectra (figure 14b) imply that the streamwise uniform streaky motion (i.e. $k_x = 0$ mode in the minimal unit) is amplified on average. This also appears to be consistent with little change of the skin-friction drag, despite the significant destruction of the vortical structures at $\lambda_x = 3h$.

The intensification of the energy of the streamwise uniform motion is surprising, at least to us, given the complete suppression of the self-sustaining process in *SO950*. The increase of the energy of the streamwise uniform motion might be explained by the lack of a proper breakdown mechanism through the streak instability or transient growth artificially suppressed, as weaker vortices could also sustain stronger streaks which can breakdown at shorter streamwise wavelengths with the elevated breakdown threshold. However, the persisting streaky motion even with the damaged breakdown mechanism can also suggest that there may exist some additional feeding mechanisms of the streamwise vortical structures. Currently, we do not have a precise understanding of these mechanisms, although they might be from scale interaction or from the Orr-mechanism proposed by Jiménez (2013, 2015). However, in any case, the result of the present numerical experiment directly supports the notion that the

streak instability (or transient growth) and the following nonlinear vortex stretching of the streamwise vortical structures play a crucial role in the formation of the vortical structures in the logarithmic and the outer regions as well as in the determination of their streamwise length scale (Schoppa & Hussain 2002; Park *et al.* 2011; Hwang 2015).

5. Concluding remarks

Thus far, we have investigated the self-sustaining process of the energy-containing motions in the logarithmic and outer regions (the attached eddies) by examining their dynamical behaviour in the minimal unit. The present study is summarised as follow:

- (i) The attached eddies at a given spanwise length scale in the minimal unit exhibit a statistically recurrent dynamical behaviour which has been called ‘bursting’ (Flores & Jiménez 2010). The bursting is also observed in the over-damped simulations in which the attached eddies survive only through their self-sustaining mechanisms, and is remarkably similar to that in the full simulations, indicating that the bursting is the reflection of the self-sustaining process of the attached eddies. For the attached eddies at the given spanwise length scale λ_z , the time period of the bursting (i.e. the self-sustaining process) is found to scale with $Tu_\tau/\lambda_z \simeq 2$, suggesting that the statistically self-similar attached eddies in the logarithmic region (Hwang 2015) are also ‘dynamically’ self-similar.
- (ii) It is shown that the self-sustaining process of the attached eddies in the logarithmic and outer regions is very similar to that of the near-wall motions (Hamilton *et al.* 1995; Schoppa & Hussain 2002), which would be the smallest attached eddies (Hwang 2015). The attached eddies in the minimal unit in the logarithmic and outer regions, composed of two elements, the streak and the streamwise vortical structures, sustain themselves through the interactive dynamics between the two: (a) the streak is significantly amplified by the streamwise vortical structures via the lift-up effect; (b) the amplified streak subsequently experiences very rapid meandering motion along the streamwise direction; and (c) the meandering streaks breakdown with regeneration of new streamwise vortical structures.
- (iii) To provide more convincing evidence on the existence of the proposed self-sustaining process in the logarithmic and outer regions, two numerical experiments are further performed. One is to artificially suppress the lift-up effect by modifying the technique introduced by Kim & Lim (2000), and the other is to artificially damp out the streak meandering motion, which would probably be a consequence of the streak instability or transient growth (Park *et al.* 2011). It is shown that the artificial suppression of the lift-up effect inhibits the amplification of the streak and subsequently suppresses the self-sustaining process of the attached eddies. Also, the artificial inhibition of the streak meandering motions destroy the generation of the streamwise vortical structures, significantly affecting the self-sustaining process. The numerical experiments also reveal that there are non-trivial scale interactions among the attached eddies.

The importance of the present study probably lies in the identification of the self-sustaining process of the energy-containing motions in the logarithmic and outer regions, given in the form of Townsend’s attached eddies. The existence of the self-sustaining process in these regions, which appears to be basically the same as that of the near-wall region, would provide some evidence on the relevance of

the so-called ‘exact coherent structures’ at high Reynolds numbers: i.e. the exact solutions of the Navier–Stokes equation in the form of unstable stationary/travelling waves and relative periodic orbits (e.g., Nagata 1990; Kawahara & Kida 2001; Waleffe 2001; Faisst & Eckhardt 2003; Wedin & Kerswell 2004; Jiménez *et al.* 2005). At low Reynolds numbers, these solutions have been shown to form a skeleton of the basin of attraction of turbulent solutions in phase space (e.g., Gibson, Halcrow & Cvitanovic 2008), and their discovery has played a crucial role in recent advancement on understanding bypass transition and low Reynolds number turbulence. It should be mentioned that the exact coherent structure, typically given in the form of a wavy streak and flanked streamwise vortices, physically represents the self-sustaining process given in figure 9 (e.g., Waleffe 2003). This therefore suggests that the attached eddies in the logarithmic and outer regions, which bear such a self-sustaining process, would be linked to the exact coherent structures at high Reynolds numbers at least to some extent.

Nevertheless, it is not yet clear how much such dynamical-system-based approaches could be extended further especially for describing the dynamics of coherent structures in high Reynolds number wall turbulence. Perhaps, one of the most important challenges would be description of the complex scale interactions between the self-sustaining structures at different length scales. In fact, the dynamical-system-based approaches implicitly assume that turbulence is a chaos occurring in a finite dimensional nonlinear system. Unfortunately, turbulence is, however, a chaos occurring in a spatially extended system, the dimension of which is virtually infinite, and is inherently a multi-scale phenomenon involving daunting scale interactions. Given this difficulty, the full suitability of the dynamical system approach especially to high Reynolds number turbulence is yet an open question, although the approach may enlighten at least some important aspects of turbulence.

Acknowledgements

Y.H. gratefully acknowledges Professor J. Jiménez for sharing his paper (Jiménez 2015) before its publication. This work was partially supported by the Engineering and Physical Science Research Council (EPSRC) in the UK (EP/N019342/1).

Appendix A. Effect of the filtering (2.1) on time correlation functions

As mentioned, all the simulations of the present study are performed with the filtering technique (2.1) which removes the two-dimensional spurious motions populating the x – y plane in the spanwise minimal unit simulations; for further details, the reader is also referred to Hwang (2013) in which a detailed discussion on this spurious motion is given. In figure 16, a set of time correlation functions of $O950$ without (2.1) are compared with those of $O950$. All the presented correlation functions of $O950$ without (2.1) show good agreement with those of $O950$, except $C_{uv}(\tau)$. It appears that $C_{uv}(\tau)$ of $O950$ without (2.1) is a little larger than that of $O950$, although the peak location of $C_{uv}(\tau)$ itself shows good agreement. However, this may have been expected in a way, given the nature of the filtering technique (2.1) which removes the streamwise and wall-normal motions in the x – y plane. Overall good agreement of the correlation functions from the two simulations suggests that application of (2.1) does not significantly affect the motions in $O950$, as in our previous studies (Hwang 2013, 2015).




FIGURE 16. Time correlation functions of (a) *O950* without (2.1) and (b) *O950*. Here, —, $C_{uu}(\tau)$; ----, $C_{vv}(\tau)$; - · - · -, $C_{uv}(\tau)$; · · · · ·, $C_{vw}(\tau)$.




FIGURE 17. Comparison of the second-order statistics of the self-sustaining attached eddies in the minimal unit (*SL1800c* in the present study) with those in the long streamwise domain (*L1800c* in Hwang 2015): (a) the streamwise, wall-normal and spanwise velocities; (b) Reynolds stress. Here, —, *SL1800c* in the present study; ----, *L1800c* in Hwang (2015).

Appendix B. Comparison of statistics between attached eddies in the minimal and long computational domains

Here, we compare the statistics of the self-sustaining attached eddies in the minimal streamwise domain ($L_x = 2L_z$; *SL1800c* in the present study) with those in a long streamwise domain ($L_x = 8\pi h$; *L1800c* in Hwang 2015). We note that the two simulations share the same spanwise computational domain size, $L_z = 0.75h$, although the resolution of the present study is a little finer than that in Hwang (2015). Except this, all the simulation parameters of the two simulations are identical to each other. Figure 17 shows the second-order statistics of *SL1800c* in the present study and those of *L1800c* in Hwang (2015). The second-order statistics obtained with the streamwise minimal domain tend to generate larger velocity fluctuations and Reynolds stress in the region relatively close to the wall, except the streamwise velocity fluctuation showing a non-trivial difference. However, overall, the second-order statistics with the streamwise minimal domain do not considerably deviate from those with the long domain, indicating that the minimal domain does not significantly distort the

second-order statistics of the interest. It should be mentioned that qualitatively the same behaviour is observed in all the other minimal unit simulations in the present study in comparison to those with sufficiently long computational domains in Hwang (2015).

REFERENCES

- ADRIAN, R. J. 2007 Hairpin vortex organization in wall turbulence. *Phys. Fluids* **19**, 041301.
- DEL ÁLAMO, J. C. & JIMÉNEZ, J. 2006 Linear energy amplification in turbulent channels. *J. Fluid Mech.* **559**, 205–213.
- DEL ÁLAMO, J. C., JIMÉNEZ, J., ZANDONADE, P. & MOSER, R. D. 2004 Scaling of the energy spectra of turbulent channels. *J. Fluid Mech.* **500**, 135–144.
- DEL ÁLAMO, J. C., JIMÉNEZ, J., ZANDONADE, P. & MOSER, R. D. 2006 Self-similar vortex clusters in the turbulent logarithmic region. *J. Fluid Mech.* **561**, 329–358.
- BUTLER, K. M. & FARRELL, B. F. 1992 Three-dimensional optimal perturbations in viscous shear flow. *Phys. Fluids A* **4**, 1637–1650.
- BUTLER, K. M. & FARRELL, B. F. 1993 Optimal perturbations and streak spacing in wall-bounded turbulent shear flow. *Phys. Fluids* **5**, 774–777.
- CHERNYSHENKO, S. I. & BAIG, M. F. 2005 The mechanism of streak formation in near-wall turbulence. *J. Fluid Mech.* **544**, 99–131.
- COSSU, C. & BRANDT, L. 2002 Stabilization of Tollmien–Schlichting waves by finite amplitude optimal streaks in the Blasius boundary layer. *Phys. Fluids* **14**, L57–L60.
- COSSU, C., PUJALS, G. & DEPARDON, S. 2009 Optimal transient growth and very large scale structures in turbulent boundary layers. *J. Fluid Mech.* **619**, 79–94.
- ELLINGSEN, T. & PALM, E. 1975 Stability of linear flow. *Phys. Fluids* **18**, 487–488.
- FAISST, H. & ECKHARDT, B. 2003 Travelling waves in pipe flow. *Phys. Rev. Lett.* **91**, 224502.
- FLORES, O. & JIMÉNEZ, J. 2010 Hierarchy of minimal flow units in the logarithmic layer. *Phys. Fluids* **22**, 071704.
- GIBSON, J. F., HALCROW, J. & CVITANOVIC, P. 2008 Visualizing the geometry of state space in plane Couette flow. *J. Fluid Mech.* **611**, 107–130.
- GUSTAVSSON, L. H. 1991 Energy growth of three-dimensional disturbances in plane Poiseuille flow. *J. Fluid Mech.* **224**, 241–260.
- HAMILTON, J. M., KIM, J. & WALEFFE, F. 1995 Regeneration mechanisms of near-wall turbulence structures. *J. Fluid Mech.* **287**, 317–348.
- HÄRTEL, C. & KLEISER, L. 1998 Analysis and modelling of subgrid-scale motions in near-wall turbulence. *J. Fluid Mech.* **356**, 327–352.
- HÖPPFNER, J., BRANDT, L. & HENNINGSON, D. S. 2005 Transient growth on boundary layer streaks. *J. Fluid Mech.* **537**, 91–100.
- HWANG, Y. 2013 Near-wall turbulent fluctuations in the absence of wide outer motions. *J. Fluid Mech.* **723**, 264–288.
- HWANG, Y. 2015 Statistical structure of self-sustaining attached eddies in turbulent channel flow. *J. Fluid Mech.* **767**, 254–289.
- HWANG, Y. & COSSU, C. 2010a Amplification of coherent streaks in the turbulent Couette flow: an input–output analysis at low Reynolds number. *J. Fluid Mech.* **643**, 333–348.
- HWANG, Y. & COSSU, C. 2010b Linear non-normal energy amplification of harmonic and stochastic forcing in the turbulent channel flow. *J. Fluid Mech.* **664**, 51–73.
- HWANG, Y. & COSSU, C. 2010c Self-sustained process at large scales in turbulent channel flow. *Phys. Rev. Lett.* **105**, 044505.
- HWANG, Y. & COSSU, C. 2011 Self-sustained processes in the logarithmic layer of turbulent channel flows. *Phys. Fluids* **23**, 061702.
- JIMÉNEZ, J. 2012 Cascades in wall-bounded turbulence. *Annu. Rev. Fluid Mech.* **44**, 27–45.
- JIMÉNEZ, J. 2013 How linear is wall-bounded turbulence. *Phys. Fluids* **25**, 110814.

- JIMÉNEZ, J. 2015 Direct detection of linearized bursts in turbulence. *Phys. Fluids* **27**, 065102.
- JIMÉNEZ, J. & HOYAS, S. 2008 Turbulent fluctuations above the buffer layer of wall-bounded flows. *J. Fluid Mech.* **611**, 215–236.
- JIMÉNEZ, J., KAWAHARA, G., SIMENS, M. P., NAGATA, M. & SHIBA, M. 2005 *Phys. Fluids* **17**, 015105.
- KAWAHARA, G. & KIDA, S. 2001 Periodic motion embedded in plane Couette turbulence: regeneration cycle and burst. *J. Fluid Mech.* **449**, 291–300.
- KIM, J. & BEWLEY, T. R. 2007 A linear systems approach to flow control. *Annu. Rev. Fluid Mech.* **39**, 383–417.
- KIM, J. & LIM, J. 2000 A linear process in wall-bounded turbulent shear flows. *Phys. Fluids* **12** (8), 1885–1888.
- LANDAHL, M. T. 1980 A note on an algebraic instability of inviscid parallel shear flows. *J. Fluid Mech.* **98**, 243–251.
- LANDAHL, M. T. 1990 On sublayer streaks. *J. Fluid Mech.* **212**, 593–614.
- LOZANO-DURÁN, A. & JIMÉNEZ, J. 2014 Effect of the computational domain on direct simulations of turbulent channels up to $Re = 4200$. *Phys. Fluids* **26**, 011702.
- MARUSIC, I. & KUNKEL, G. J. 2003 Streamwise turbulent intensity formulation for flat-plate boundary layers. *Phys. Fluids* **15** (8), 2461.
- MARUSIC, I., MONTY, J. P., HULTMARK, M. & SMITS, A. J. 2013 On the logarithmic region in wall turbulence. *J. Fluid Mech.* **716**, R3.
- MASON, P. J. & CULLEN, N. J. 1986 On the magnitude of the subgrid-scale eddy coefficient in large-eddy simulations of turbulent channel flow. *J. Fluid Mech.* **162**, 439–462.
- MONTY, J. P., STEWART, J. A., WILLIAMS, R. C. & CHONG, M. S. 2007 Large-scale features in turbulent pipe and channel flows. *J. Fluid Mech.* **589**, 147–156.
- NAGATA, M. 1990 Three-dimensional finite-amplitude solutions in plane Couette flow: bifurcation from infinity. *J. Fluid Mech.* **217**, 519–527.
- NICKELS, T. B., MARUSIC, I., HAFEZ, S. & CHONG, M. S. 2005 Evidence of the $k - 1$ law in a high Reynolds number turbulent boundary layer. *Phys. Rev. Lett.* **95**, 074501.
- ORLANDI, P., BERNARDINI, M. & PIROZZOLI, S. 2015 Poiseuille and couette flows in the transitional and fully turbulent regime. *J. Fluid Mech.* **770**, 424–441.
- PARK, D. S. & HUERRE, P. 1995 Primary and secondary instabilities of the asymptotic suction boundary layer on a curved plate. *J. Fluid Mech.* **283**, 249–272.
- PARK, J., HWANG, Y. & COSSU, C. 2011 On the stability of large-scale streaks in the turbulent Couette and Poiseuille flows. *C. R. Méc.* **339** (1), 1–5.
- PERRY, A. E. & CHONG, M. S. 1982 On the mechanism of turbulence. *J. Fluid Mech.* **119**, 173–217.
- PERRY, A. E., HENBEST, S. & CHONG, M. S. 1986 A theoretical and experimental study of wall turbulence. *J. Fluid Mech.* **165**, 163–199.
- PERRY, A. E. & MARUSIC, I. 1995 A wall-wake model for the turbulence structure of boundary layers. Part I: extension of the attached eddy hypothesis. *J. Fluid Mech.* **298**, 361–388.
- PUJALS, G., GARCÍA-VILLALBA, M., COSSU, C. & DEPARDON, S. 2009 A note on optimal transient growth in turbulent channel flows. *Phys. Fluids* **21**, 015109.
- REDDY, S. C. & HENNINGSON, D. S. 1993 Energy growth in viscous channel flows. *J. Fluid Mech.* **252**, 209–238.
- SCHMID, P. J. 2007 Nonmodal stability theory. *Annu. Rev. Fluid Mech.* **39**, 129–162.
- SCHMID, P. J. & HENNINGSON, D. S. 2001 *Stability and Transition in Shear Flows*. Springer.
- SCHOPPA, W. & HUSSAIN, F. 2002 Coherent structure generation in near-wall turbulence. *J. Fluid Mech.* **453**, 57–108.
- TOMKINS, C. D. & ADRIAN, R. J. 2003 Spanwise structure and scale growth in turbulent boundary layers. *J. Fluid Mech.* **490**, 37–74.
- TOWNSEND, A. A. 1961 Equilibrium layers and wall turbulence. *J. Fluid Mech.* **11** (1), 97–120.
- TOWNSEND, A. A. 1976 *The Structure of Turbulent Shear Flow*. Cambridge University Press.
- WALEFFE, F. 2001 Exact coherent structures in channel flow. *J. Fluid Mech.* **435**, 93–102.

- WALEFFE, F. 2003 Homotopy of exact coherent structures in plane shear flows. *Phys. Fluids* **15**, 1517–1534.
- WEDIN, H. & KERSWELL, R. R. 2004 Exact coherent structures in pipe flow: travelling wave solutions. *J. Fluid Mech.* **508**, 333–371.
- WILLIS, A. P., HWANG, Y. & COSSU, C. 2010 Optimally amplified large-scale streaks and drag reduction in the turbulent pipe flow. *Phys. Rev. E* **82**, 036321.