

Genetic heterogeneity and phylogenetic status of *Leishmania (Leishmania) infantum* zymodeme MON-1: epidemiological implications

M. HIDE*, A. L. BAÑULS and M. TIBAYRENC

Institut de Recherche IRD, Centre d'Etudes sur le Polymorphisme des Micro-organismes (CEPM),
Unité Mixte de recherche CNRS-IRD 9926, BP 5045, 34032 Montpellier Cedex 1, France

(Received 1 March 2001; revised 23 May 2001; accepted 30 May 2001)

SUMMARY

Leishmania (Leishmania) infantum zymodeme MON-1 is responsible for the majority of visceral leishmaniasis cases around the Mediterranean basin, albeit that it causes also cutaneous forms. The MON classification is based on starch gel multilocus enzyme electrophoresis (MLEE) typing. The aim of this work was to explore further the genetic diversity and phylogenetic status of this zymodeme by alternative typing techniques. Fourteen *L. (L.) infantum/L. (L.) chagasi* stocks identified as MON-1 by MLEE in reference laboratories, 3 *L. infantum* stocks attributed to other zymodemes (MON-24, MON-29, MON-33) and reference standard stocks belonging to other species (*L. (L.) major*, *L. (L.) tropica* and *L. (L.) donovani*) were characterized by 2 different markers: MLEE on cellulose acetate plates and Random Amplified Polymorphic DNA (RAPD). We have obtained 10 different genotypes with RAPD and 6 different genotypes with MLEE on cellulose acetate plates for the 14 *L. infantum/L. chagasi* MON-1 stocks studied. MLEE and RAPD data gave quite congruent phylogenetic results: *L. infantum* zymodeme MON-1 was shown to be polyphyletic and genetically heterogeneous. This work confirms the necessity of using different markers to build up a robust phylogeny. Finally the epidemiological and clinical implications of these results are discussed.

Key words: *Leishmania (Leishmania) infantum*, zymodeme MON-1, Random Amplified Polymorphic DNA (RAPD), Multilocus Enzyme Electrophoresis (MLEE), Discrete Typing Unit (DTU), phylogeny.

INTRODUCTION

The parasites belonging to the *Leishmania* genus are the causative agents of leishmaniasis. These diseases constitute a priority for WHO in terms of public health. With regard to *L. (L.) infantum* (Nicolle, 1908), many new cases are now appearing in Southern Europe, due to the frequent association between this species and HIV infection (Alvar *et al.* 1992). This species belongs, together with *L. (L.) chagasi* (Cunha and Chagas, 1937), *L. (L.) donovani* (Laveran and Mesnil, 1903) and *L. (L.) archibaldi* (Castellani and Chalmers, 1919), to the *L. donovani* complex. These 4 species are described as responsible for visceral leishmaniasis, although they are often isolated from cutaneous lesions too. *L. infantum* has been subdivided into zymodemes MON with starch gel Multilocus Enzyme Electrophoresis technique (MLEE) by the French WHO reference laboratory of Montpellier (Laboratoire d'Ecologie Médicale et Pathologie Parasitaire, France) (Rioux *et al.* 1990), (Lanotte *et al.* 1981). The MON classification is based on 15 enzymatic systems (DIA (NADH diaphorase), FH (fumarate hydratase),

G6PD (glucose-6-phosphate dehydrogenase), GLUD (glutamate dehydrogenase), GOT1 and GOT2 (glutamate oxaloacetate transaminase), GPI (glucose phosphate isomerase), IDH (isocitrate dehydrogenase), MDH (malate dehydrogenase), ME (malic enzyme), MPI (mannose-phosphate isomerase), NHI and NHD (nucleoside hydrolase), 6PGD (6-phosphogluconate dehydrogenase) and PGM (phosphoglucomutase). A zymodeme is defined as a collection of stocks that have the same isozyme profile (Godfrey & Kilgour, 1976).

The present study focused on *L. infantum* species and especially on the zymodeme MON-1. The zymodeme MON-1 is the more common and the more widespread zymodeme in the Mediterranean basin (Moreno *et al.* 1986). Being responsible for the majority of visceral leishmaniasis cases, together with the zymodeme MON-80 (Aoun *et al.* 1999; Marty *et al.* 1994) and the zymodeme MON-34 (Harrat *et al.* 1996). The zymodeme MON-1, however, can also produce cutaneous leishmaniasis cases (Marty *et al.* 1998) as do other zymodemes of the *L. donovani* complex.

The aim of this study was to re-examine the phylogenetic individualization and genetic homogeneity of the *L. infantum* zymodeme MON-1. Two genetic markers were used for phylogenetic analyses:

* Corresponding author: Tel: +33 467 41 62 72. Fax: +33 467 41 62 99. E-mail: hide@cepm.mpl.ird.fr

MLEE on cellulose acetate plates (Godfrey & Kilgour, 1976) and random amplified polymorphic DNA (RAPD) technique (Williams *et al.* 1990).

MATERIALS AND METHODS

Leishmania stocks

The origin of the stocks is detailed in Table 1. The stocks were cultivated at 26 °C in RPMI 1640 medium supplemented with 10% heat-inactivated foetal calf serum, 1% glutamine and 0.5% gentamycin in disposable flasks. Parasites were harvested in logarithmic phase by centrifugation (5000 g for 30 min at 4 °C) washed twice in PBS (phosphate-buffered saline: 0.01 M Na₂HPO₃; 0.01 M Na₂H₂PO₄; 0.15 M NaCl, pH 7.3). Parasite pellets were kept at -80 °C.

MLEE analysis

MLEE experiments were performed on cellulose acetate plates (Helena Laboratories) according to the method described by Godfrey & Kilgour (1976) and Ben Abderrazak *et al.* (1993) with slight modifications (Bañuls, 1998). Cellular pellets were lysed in an equal volume of enzyme stabilizer (EDTA, dithiothreitol, ϵ -aminocaproic, at 2 mM) on an ice bed for 15 min. The lysates were centrifuged at 13000 g for 10 min at 4 °C and the water-soluble fraction was removed and stored at -80 °C. Fifteen enzyme systems were studied; they were as follows. Aconitase (ACON, EC4.2.1.3), alanine aminotransferase (ALAT, EC2.6.1.2), glucose-phosphate isomerase (GPI, EC5.3.1.9), glucose-6-phosphate dehydrogenase (G6PD, EC1.1.1.49), glutamate oxaloacetate transaminase (GOT, EC2.6.1.1), isocitrate dehydrogenase (IDH, EC1.1.1.42), malate dehydrogenase NAD⁺ (MDH, EC1.1.1.37), malate dehydrogenase (NADP⁺) or malic enzyme (ME, EC1.1.1.40), mannose-phosphate isomerase (MPI, EC5.3.1.8), nucleoside hydrolase NHI (inosine substrate) and NHD (deoxyinosine substrate) (EC2.4.2.*), 6-phosphogluconate dehydrogenase (6PGD, EC1.1.1.44), phosphoglucomutase (PGM, EC2.7.5.1), peptidase 1, substrate 1-leucyl-leucine-leucine (PEP1, EC3.4.11) and peptidase 2, substrate 1-leucyl-1-alanine (PEP2, EC3.4.11), superoxide dismutase (SOD, EC1.15.1.1). This gives a total of 16 loci since NHI system shows an activity at 2 distinct loci (*Nhi1* and *Nhi2*, *Nhi1* being the fastest migrating locus).

The difference between the panels of enzymatic systems for MLEE on starch gel (MON classification) and for MLEE on acetate plates is based on: ACON, ALAT, PEP (1 and 2) and SOD for acetate plates and GLUD, DIA and FH for starch gel.

Random amplified polymorphic DNA

The technique was performed according to the

method described by Williams *et al.* (1990). DNA was extracted by phenol/chloroform (Sambrook, Fritsch & Maniatis, 1989) and DNA concentration was estimated by spectrophotometry at 260 nm. Primers giving reproducible patterns have been selected according to a previous screening performed in our laboratory on 120 decameric primers (Bañuls *et al.* 1999a). The following 16 primers from Operon Technologies (Alameda, California) have been used to study the sample (5'-3': A1-CAGGCCCTTC, A7-GAAACGGGTG, A10-GTGATCGCAG, B10-CTGCTGGGAC, B11-GTAGACCCGT, F10-GG-AAGCTTGG, F13-GGCTGCAGAA, N20-GGT-GCTCCGT, R2-CACAGCTGCC, R7-ACTGGC-CTGA, R8-CCCGTTGCCT, R15-GGACAACG-AG, U2-CTGAGGTCTC, U3-CTATGCCGAC, B4-GGACTGGAGT, B15-GGAGGGTGTT). Each RAPD reaction (60 μ l) contained: 20 ng template DNA, 100 μ M each dNTP, 6 μ l 10X *Taq* polymerase buffer, 12 pmol decameric primer, 0.9U *Taq* polymerase (Roche, Germany), 49.4 μ l nuclease-free water. Forty-five cycles (denaturation: 1 min at 94 °C; annealing: 1 min at 36 °C; elongation: 1 min at 72 °C) were followed by a final step of 7 min at 72 °C in a PTC-100 Thermocycler (MJ Research, Inc., Massachusetts, USA). RAPD products were visualized on 1.6% agarose gels and revealed with ethidium bromide under UV.

Data analysis

All bands obtained on RAPD and MLEE gels were numbered and scored as presence or absence data. For MLEE data, band number 1 is the fastest and for RAPD, band number 1 is the slowest. MLEE and RAPD data were computed with the *Genetics Toolbox* software designed in our laboratory (S. Noël and F. Chevenet). Genotypic diversity (number of multilocus genotypes on the total number of stocks) and mean genetic diversity H were calculated according to Selander & Levin (1980). Mean genetic diversity H is computed with the formula:

$$H = \sum h/r,$$

where $h = 1 - \sum q_i^2$, q is the frequency of each allele for a locus i and r the total number of loci.

Estimation of genetic differences among the stocks was performed using the Jaccard's distance (Jaccard, 1908), computed with the following formula:

$$D_{ij} = 1 - [a/(a+b+c)],$$

where a is the number of common bands between the stocks i and j , b is the number of bands present in i and absent in j and c is the number of bands present in j and absent in i . All the distances were regrouped in 2 distance matrices (one for RAPD and one for MLEE data). Dendrograms have been constructed from these matrices with the

Table 1. *Leishmania* spp. stocks

International Code	Zymodeme	Locality of isolation	Locality of infection*
<i>L. infantum</i>			
MHOM/FR/78/LEM75	1	Beziers	–
MHOM/FR/92/LEM2572	1	Toulouse	Milan (IT)
MHOM/FR/92/LEM2301	1	Toulouse	Marseille (FR)
MHOM/TN/80/IPT1	1	Monastir	–
MHOM/MA/67/ITMAP263	1	–	–
MHOM/FR/92/LPN83	1	Nice	–
MHOM/FR/93/LPN92	1	Nice	–
MHOM/FR/94/LPN94	1	Nice	–
MHOM/FR/94/LPN105	1	Nice	Bastia (FR)
MHOM/FR/95/LPN119	1	Nice	Cannes (FR)
MHOM/FR/96/LPN129	1	Nice	Vallauris (FR)
MHOM/FR/96/LPN132	1	Nice	Mouns Sartoux (FR)
MHOM/DZ/82/LIPA59†	24	El Asnam	–
MHOM/ES/81/BCN1†	29	Monserrat	–
MHOM/FR/82/LEM356†	33	Amélie-les bains	–
<i>L. chagasi</i>			
MHOM/BR/79/L101	1	–	–
MHOM/BR/74/PP75	1	Bahia	–
<i>L. donovani</i>			
MHOM/ET/67/HU3	18	Humera	–
<i>L. tropica</i>			
MHOM/SU/74/K27	60	–	–
<i>L. major</i>			
MHOM/SU/73/5ASKH	4	Askhabad	–

* FR, France; TN, Tunisia; MA, Morocco; DZ, Algeria; ES, Spain; BR, Brazil; ET, Ethiopia; SU, former Soviet Union.

† *L. infantum* LIPA59, BCN1, LEM356 were isolated from cutaneous lesions, the other stocks pertaining to the *L. donovani* complex were isolated from visceral lesions.




Fig. 1. MLEE profile revealed with the MDH enzymatic system. From left to right: Lane 1: *Leishmania major*, 5ASKH; lane 2: *L. tropica*, K27; lane 3: *L. infantum*, LEM75 (MON-1); lane 4: *L. chagasi*, L101 (MON-1); lane 5: *L. donovani*, HU3; lanes 6 and 7: *L. infantum*, BCN1 (MON-29), LIPA59 (MON-24). Lanes 3 and 4 reveal polymorphic patterns within zymodeme MON-1.

Neighbour-joining method (Saitou & Nei, 1987). Parsimony analysis of Wagner (Felsenstein, 1985) (with 100 replicates) was also used to build phylogenetic trees based on MLEE data compiled with RAPD data and the robustness of the nodes was tested by a bootstrap analysis. Dendrograms were analysed with the Treedyn software (Chevenet *et al.* 2000). Correlation between the results obtained with MLEE and RAPD was evaluated by a non-parametric test (Mantel, 1967) with 10^4 repetitions.




Fig. 2. MLEE profiles obtained for the PGM enzymatic system. From left to right: Lane 1: *Leishmania tropica*, K27; lane 2: *L. major*, 5ASKH; lane 3: *L. donovani*, HU3; lanes 4 and 5: *L. infantum*, LEM75, BCN1; lanes 6 and 7: *L. chagasi*, PP75, L101. This system is a specific tag of the *L. donovani* complex. The 5 stocks pertaining of the *L. donovani* complex have the same band and this band is absent for *L. tropica* and *L. major*.

RESULTS

MLEE analysis

We have obtained 47 different electrophoretic bands with the 16 enzyme loci. The genotypic diversity was 0.43 within *L. infantum/L. chagasi* MON-1 and 0.5 within the whole *L. donovani* complex. The mean genetic diversity (H) was 0.083, 0.095 and 0.21 for *L. infantum/L. chagasi* MON-1, the whole *L. donovani* complex and the whole sample, respectively. Only 5 enzyme loci were polymorphic within *L. infantum/L. chagasi*: MON-1: *6pgdh*, *Mdh*, *Nhi2*, *Got*, *Pep1* (see Fig. 1), which corresponds to a polymorphism rate




Fig. 3. RAPD patterns obtained with the primer F10. From left to right: Lane 1: *Leishmania tropica*, K-27; lane 2: *L. major*, 5ASKH; lane 3: *L. donovani*, HU3; lanes 4–8: *L. infantum*, LEM2301 (MON-1), LEM2572 (MON-1), LPN119 (MON-1), IPT1 (MON-1), ITMAP263 (MON-1); lane 9: *L. chagasi*, PP75 (MON-1); lanes 10–18: *L. infantum*, LIPA59 (MON-24), BCN1 (MON-29), LEM356 (MON-33), LPN132 (MON-1), LPN92 (MON-1), LPN94 (MON-1), LPN105 (MON-1), LPN129 (MON-1), LPN83 (MON-1); MW, Molecular weight. Note the polymorphic patterns within *L. infantum*/*L. chagasi* MON-1.




Fig. 4. RAPD patterns obtained with the primer A10. All the patterns are identical for the stocks pertaining to the *Leishmania donovani* complex, A10 primer is specific of the complex (tag). From left to right: Lanes 1–3: *L. infantum*, LIPA59 (MON-24), BCN1 (MON-29), LEM356 (MON-33); Lane 4: *L. chagasi*, PP75; Lane 5–16: *L. infantum*, LEM2301 (MON-1), LEM2572 (MON-1), LPN119 (MON-1), IPT1 (MON-1), ITMAP263 (MON-1), LPN132 (MON-1), LPN92 (MON-1), LPN94 (MON-1), LPN105 (MON-1), LPN129 (MON-1), LPN83 (MON-1), LEM75 (MON-1); Lane 17: *L. donovani*, HU3; Lane 18: *L. major*, 5ASKH; Lane 19: *L. tropica*, K-27; MW, Molecular weight.

of 0.312. For the *L. donovani* complex, we found 4 loci giving specific electromorphs: *G6pdh*, *Gpi*, *Pgm*, *Sod* (see Fig. 2), though there were no loci giving specific electromorphs either for *L. infantum* or for MON-1. Considering all these data, we have observed that (i) in some cases MLEE loci on acetate plates gave more discriminated patterns: some MON-1 stocks showed different patterns for 6PGDH, GOT, MDH, NHI2, PEP1 on acetate plates; and (ii) in some other cases MLEE on starch gel was more discriminated than MLEE on acetate plates: the 3 non-MON-1 stocks had the same pattern as several MON-1 stocks with the MDH system on acetate plates but were different from MON-1 stocks on starch gels.

RAPD analysis

We have obtained 117 different electrophoretic bands with the 16 primers used. The genotypic

diversity was 0.71 within *L. infantum*/*L. chagasi* MON-1. RAPD analysis revealed a more important genetic diversity within the MON-1 (0.205) compared to the MLEE data. The mean genetic diversity was 0.37 in the whole sample and 0.24 within the *L. donovani* complex. Six primers gave polymorphic patterns within *L. infantum*/*L. chagasi* MON-1 (U2, R15, F10, B4, B10, A1) (see Fig. 3). The corresponding polymorphism rate was 0.5. There were 5 primers giving specific amplicons of the *L. donovani* complex (A7, A10, B15, F13, U3) (see Fig. 4) but we did not find specific amplicons for MON-1.

Phylogenetic analysis

The 2 neighbour-joining trees (Fig. 5) designed from MLEE and RAPD distance matrices showed similar patterns. All stocks belonging to the *L. donovani* complex fell into 1 cluster. In both trees,




Fig. 5. Two Neighbour-joining dendrograms obtained from RAPD (left) and MLEE (right) data. The mentioned loci correspond to specific loci (tag) of the *Leishmania donovani* complex. In order to align the two dendrograms, the branch corresponding to the LPN83 stock in MLEE tree has been rotated.

the MON-1 stocks were not grouped into a specific cluster and were mixed among stocks from other zymodemes (Fig. 5). For example, the *L. donovani* stock was associated with the 2 *L. chagasi* MON-1 stocks. On the Wagner parsimony tree (Fig. 6), a strong bootstrap value (100%) supported the node of the whole *L. donovani* complex. This complex was characterized by several specific markers (5 RAPD amplicons and 4 MLEE electromorphs). Within this complex, no relevant additional subdivision was revealed by the bootstrap analysis.

Correlation test

Overall, strong agreement between the MLEE and RAPD trees was ascertained by the results of the nonparametric correlation test (Mantel, 1967). All *P* values were $<10^{-4}$, either for the whole sample ($r: 0.95$), for the *L. donovani* complex ($r: 0.73$) or for the MON-1 stocks ($r: 0.80$) considered separately.

DISCUSSION

The parasites of the *L. donovani* complex are responsible for very different clinical expressions and present different epidemiological features. In this framework, the aim of this study was to explore in depth the genetic structure of the zymodeme MON-1 at the micro-evolutionary level. It is considered that MON-1 has a considerable epidemiological relevance. The main goal was therefore to

verify the phylogenetic individualization of MON-1 within the *L. donovani* complex and its genetic homogeneity. The epidemiological and clinical implications of the genetic and phylogenetic results are discussed.

Individualization of the *L. donovani* complex

Our results confirmed the genetic individualization of the *L. donovani* complex against the other species (*L. tropica* and *L. major*). The phylogenetic construction and the 4 specific MLEE and 5 specific RAPD markers (tags) show that the *L. donovani* complex can be considered as a DTU (Discrete Typing Unit) according to the definition of Tibayrenc (Tibayrenc, 1998). These data confirmed those published by Mauricio & Howard (1999); indeed they postulated that the *L. donovani* complex has a monophyletic origin on the basis of RAPD data and *mspC* sequence.

Phylogenetic relationship between *L. donovani* complex species

The controversy still exists around the species and zymodeme classification belonging to the *L. donovani* complex. Indeed, it was widely accepted that the *L. donovani* complex is divided in 4 different species (*L. infantum*, *L. chagasi*, *L. donovani*, *L. archibaldi*). Moreover, recent data raise the question of the genetic individualization of *L. donovani* and *L.*




Fig. 6. Phylogenetic tree constructed by bootstrapped (100 replicates) parsimony analysis of Wagner. Bootstrap values for all nodes are given. The stocks pertaining to the zymodeme MON-1 fall into different groups and these groups are not supported by strong nodes. The node of the DTU “*L. donovani* complex” is supported by a strong bootstrap value (100%). * LNP 92, LPN94 and LPN105 have the same genotype with MLEE and RAPD techniques.

infantum (Rioux, J. A., personal communication; Bañuls, Hide & Tibayrenc, 1999b) and of *L. archibaldi*. It is difficult to know clearly the real phylogenetic status of each species. Mauricio & Howard (1999) suggest that *L. infantum*/*L. chagasi* is a monophyletic group (with the RAPD primer H1 5′–3′:CGCGCCCGCT as specific tag). The same authors showed by PCR-RFLP (ITS and mini-exon) and RAPD that *L. chagasi* could not be distinguished from *L. infantum* (Mauricio, Stothard & Miles, 2000). In 1986, other authors considered that *L. infantum* and *L. chagasi* pertained to the complex *L. infantum* and *L. donovani* was a distinct complex

(Moreno *et al.* 1986). Our results cannot distinguish the 3 species: *L. donovani*, *L. chagasi* and *L. infantum*. By the use of more resolutive molecular tools such as microsatellites and larger samples of *L. donovani*, *L. chagasi* and *L. archibaldi*, this fundamental point should be resolved.

Genetic heterogeneity and polyphyletism of MON-1 within the L. donovani complex

Analysis of the *L. infantum* species by other molecular methods than MLEE on starch gel allowed study of the genetic diversity and the

phylogenetic status of the zymodeme MON-1. The different calculations showed a comparable polymorphism in MON-1 and in the whole sample of *L. infantum*, as well as a significant genotypic diversity. The use of 2 different markers (better resolution than MLEE on starch gel) revealed an additional and non-negligible genetic variability within the zymodeme MON-1. From a phylogenetic point of view, MLEE and RAPD data gave congruent results confirmed by the Mantel test ($r = 0.80$ within MON-1; $P < 10^{-4}$). Analysis by phenetic and phylogenetic methods underscored that zymodeme MON-1 did not correspond to a DTU. The present study did not allow separation of the zymodeme MON-1 from the other zymodemes (MON-24, MON-33, MON-29) responsible for cutaneous forms in competent hosts (Rioux *et al.* 1985; Harrat *et al.* 1996; Aoun *et al.* 2000). MON-1 stocks are not grouped into a specific cluster, indeed other MON zymodemes are inserted among them. Moreover, the *L. chagasi* stock characterized as MON-1 is more distant from *L. infantum* than *L. infantum* stocks belonging to other zymodemes. These data suggest that, in spite of the MLEE on starch gel classification, zymodeme MON-1 does not have a monophyletic status for the studied markers, which is probably due to insufficient resolution of the 15 enzyme loci used for MON identification.

Clinical and epidemiological implications

The *L. donovani* species is described as producing, in the majority of cases, a visceral clinical form and evolving only in an anthroponotic cycle. The 2 other species, *L. chagasi* and *L. infantum* (among which MON-1), evolve in an anthrozoönotic cycle and have the potential to produce asymptomatic infections, benign cutaneous forms as well as lethal visceral forms of the disease. The additional genetic polymorphism within MON-1 obtained in this study, suggests the possibility of testing for statistical association between the clinical variability observed in patients and the parasite genotypes. This kind of work must be performed on a large *Leishmania* sample representative of the clinical variability, and information about the patient (immunological status, disease symptoms) must be taken into account. Moreover, the polyphyletic nature of MON-1 and the additional genetic variability revealed in this work also raise questions about the reservoir implicated in the parasite's transmission, the origin of transmission of the disease and the vector species incriminated. The general conclusion from our data is that MON-1 is not the result of a recent, common clonal descent, which has to be taken into account in all epidemiological surveys. The common ancestor of all MON-1 genotypes may be ancient, and it is common to other strains pertaining to different zymodemes. For such epidemiological surveys,

markers with a higher discriminative power are preferable (Tibayrenc, 1999). This is the case for microsatellites or multi-gene sequencing (Rossi *et al.* 1994; Vickerman, 1994; Minodier *et al.* 1997), which showed also their usefulness for the intraspecific analysis because of their fast molecular clock.

The authors gratefully acknowledge P. Marty (Department of Internal Medicine-Haematology, Sophia-Antipolis University, Nice, France) and J. P. Dedet and F. Pratlong (Laboratoire d'Ecologie Médicale et Pathologie Parasitaire, Montpellier, France) who kindly provided some *L. infantum* stocks.

REFERENCES

- ALVAR, J., GUTIERREZ-SOLAR, B., MOLINA, R., LOPEZ-VELEZ, R., GARCIA-CAMACHO, A., MARTINEZ, P., LAGUNA, F., CERCENADO, E. & GALMES, A. (1992). Prevalence of *Leishmania* infection among AIDS patients. *Lancet* **339**, 1427.
- AOUN, K., BOURATBINE, A., HARRAT, Z., GUIZANI, I., MOKNI, M., BEL HADJ ALI, S., BEN OSMAN, A., BELKAID, M., DELLAGI, K. & BEN ISMAIL, R. (2000). Epidemiologic and parasitologic data concerning sporadic cutaneous leishmaniasis in northern Tunisia. *Bulletin de la Société de Pathologie Exotique* **93**, 101–103.
- AOUN, K., BOURATBINE, A., HARRAT, Z., MAHERZI, A., BELKAID, M., BOUSNINA, S. & BEN ISMAIL, R. (1999). Confirmation of the presence of *Leishmania infantum* MON-80 in Tunisia. *Bulletin de la Société de Pathologie Exotique* **92**, 29–30.
- BAÑULS, A. L. (1998). Apport de la génétique évolutive à l'épidémiologie et à la taxonomie du genre *Leishmania*. Ph.D. thesis, University of Montpellier II, France.
- BAÑULS, A. L., BRISSE, A., SIDIBE, I., NOËL, S. & TIBAYRENC, M. (1999a). A phylogenetic analysis by multilocus enzyme electrophoresis and multiprimer random amplified polymorphic DNA fingerprinting of the *Leishmania* genome project Friedlin reference strain. *Folia Parasitologica* **46**, 10–14.
- BAÑULS, A. L., HIDE, M. & TIBAYRENC, M. (1999b). Molecular epidemiology and evolutionary genetics of *Leishmania* parasites. *International Journal for Parasitology* **29**, 1137–1147.
- BEN ABDERRAZAK, S., GUERRINI, F., MATHIEU-DAUDE, F., TRUC, P., NEUBAUER, K., LEWICKA, K., BARNABE, C. & TIBAYRENC, M. (1993). Isoenzyme electrophoresis for parasite characterization. *Methods in Molecular Biology* **21**, 361–382.
- CHEVENET, F., BAÑULS, A. L. & BARNABE, C. (2000). TreeDyn: un éditeur interactif d'arbres phylogénétiques. In *Actes des Premières Journées Ouvertes Biologie, Informatique et Mathématiques* (ed. Caraux, G., Gascuel, O. & Sagot, M. F.), pp. 87–90. ENSAM/LIRMM, Montpellier.
- FELSENSTEIN, J. (1985). Confidence limits on phylogenies: an approach using bootstrap. *Evolution* **39**, 783–791.
- GODFREY, D. G. & KILGOUR, V. (1976). Enzyme electrophoresis in characterizing the causative organism of Gambian trypanosomiasis. *Transaction of the Royal Society of Tropical Medicine and Hygiene* **70**, 219–224.

- HARRAT, Z., PRATLONG, F., BELAZZOU, S., DEREURE, J., DENIAU, M., RIOUX, J. A., BELKAID, M. & DEDET, J. P. (1996). *Leishmania infantum* and *L. major* in Algeria. *Transaction of the Royal Society of Tropical Medicine and Hygiene* **90**, 625–629.
- JACCARD, P. (1908). Nouvelles recherches sur la distribution florale. *Bulletin de la Société vaudoise de Sciences Naturelles* **44**, 223–270.
- LANOTTE, G., RIOUX, J. A., MAAZOUN, R., PASTEUR, N., PRATLONG, F. & LEPART, J. (1981). The application of a numerical method to the taxonomy of the genus *Leishmania* Ross, 1903. The recognition of 146 original lines in the Old World. Use of allozymic characters. Epidemiological and phyletic significance. *Annales de Parasitologie humaine et Comparée* **56**, 575–591.
- MANTEL, N. (1967). The detection of disease clustering and a generalised regression approach. *Cancer Research* **27**, 290–320.
- MARTY, P., LACOUR, J. P., PRATLONG, F., PERRIN, C., DEL GIUDICE, P. & LE FICHOUX, Y. (1998). Localized cutaneous leishmaniasis due to *Leishmania infantum* MON-1 contracted in northern Algeria. *Bulletin de la Société de Pathologie Exotique* **91**, 146–147.
- MARTY, P., LE FICHOUX, Y., PRATLONG, F. & GARITOUSSAINT, M. (1994). Human visceral leishmaniasis in Alpes-Maritimes, France: epidemiological characteristics for the period 1985–1992. *Transaction of the Royal Society of Tropical Medicine and Hygiene* **88**, 33–34.
- MAURICIO, I. L. & HOWARD, M. K. (1999). Genomic diversity in the *Leishmania donovani* complex. *Parasitology* **119**, 237–246.
- MAURICIO, I. L., STOTHARD, J. R. & MILES, M. A. (2000). The strange case of *Leishmania chagasi*. *Parasitology Today* **16**, 188–189.
- MINODIER, P., PIARROUX, R., GAMBARELLI, F., JOBLET, C., DUMON, H. & VICKERMAN, K. (1997). Rapid identification of causative species in patients with Old World leishmaniasis. The evolutionary expansion of the trypanosomatid flagellates. *Journal of Clinical Microbiology* **35**, 2551–2555.
- MORENO, G., RIOUX, J. A., LANOTTE, G., PRATLONG, F. & SERRES, E. (1986). Le complexe *Leishmania donovani* sl. Analyse enzymatique du complexe *Leishmania infantum*. Corrolaires biogéographiques et phylétiques. A propos de 146 souches originaires de l'Ancien et du Nouveau Monde. In *Leishmania. Taxonomie et Phylogénèse. Applications Eco-Epidémiologiques*. IMEEE, Montpellier, pp. 105–117.
- RIOUX, J. A., LANOTTE, G., PRATLONG, F., DEREURE, J., JARRY, D., MORENO, G., KILLICK-KENDRICK, R., PERIERES, J., GUILVARD, E., BELMONTE, A. & PORTUS, M. (1985). La leishmaniose cutanée autochtone dans le sud-est de la France. Résultats d'une enquête éco-épidémiologique dans les Pyrénées-Orientales. *Médecine et Maladies Infectieuses* **11**, 650–656.
- RIOUX, J. A., LANOTTE, G., SERRES, E., PRATLONG, F., BASTIEN, P. & PERIERES, J. (1990). Taxonomy of *Leishmania*. Use of isoenzymes. Suggestions for a new classification. *Annales de Parasitologie humaine et Comparée* **65**, 111–125.
- ROSSI, V., WINCKER, P., RAVEL, C., BLAINEAU, C., PAGES, M. & BASTIEN, P. (1994). Structural organisation of microsatellite families in the *Leishmania* genome and polymorphisms at two (CA)_n loci. *Molecular and Biochemical Parasitology* **65**, 271–282.
- SAITOU, N. & NEI, M. (1987). The neighbor-joining method: a new method for reconstructing phylogenetic trees. *Molecular Biology and Evolution* **4**, 406–425.
- SAMBROOK, J., FRITSCH, E. F. & MANIATIS, T. (1989). *Molecular Cloning: A Laboratory Manual*. Cold Spring Harbor Laboratory Press, NY.
- SELANDER, R. K. & LEVIN, B. R. (1980). Genetic diversity and structure in *Escherichia coli* populations. *Science* **210**, 545–547.
- TIBAYRENC, M. (1998). Genetic epidemiology of parasitic protozoa and other infectious agents: the need for an integrated approach. *International Journal for Parasitology* **28**, 85–104.
- TIBAYRENC, M. (1999). Toward an integrated genetic epidemiology of parasitic protozoa and other pathogens. *Annual Review of Genetics* **33**, 449–477.
- VICKERMAN, K. (1994). The evolutionary expansion of the trypanosomatid flagellates. *International Journal for Parasitology* **24**, 1317–1331.
- WILLIAMS, J. G., KUBELIK, A. R., LIVAK, K. J., RAFALSKI, J. A. & TINGEY, S. V. (1990). DNA polymorphisms amplified by arbitrary primers are useful as genetic markers. *Nucleic Acids Research* **18**, 6531–6535.