

Genetic diversity of mungbean (*Vigna radiata* L.) germplasm in Indonesia

Puji Lestari^{1,2}, Sue Kyung Kim¹, Reflinur^{1,2}, Yang Jae Kang¹,
Nurwita Dewi² and Suk-Ha Lee^{1,3*}

¹Department of Plant Science and Research Institute for Agriculture and Life Sciences, Seoul National University, San 56-1, Sillim-dong, Gwanak-gu, Seoul 151-921, Republic of Korea, ²Indonesian Center for Agricultural Biotechnology and Genetic Resources Research and Development, IAARD, Bogor 16111, Indonesia and ³Plant Genomics and Breeding Institute, Seoul National University, Seoul 151-921, Republic of Korea

Abstract

Despite widespread mungbean [*Vigna radiata* (L.) Wilczek] consumption in Indonesia, few molecular studies have been carried out on accessions and available data are minimal. In this study, we used 30 newly developed simple sequence repeat (SSR) markers designed from the mapped sequence scaffolds of the Korean Sunhwanokdu and Gyeonggiarae 5 mungbean genomes. These markers were used to examine loci in 83 mungbean accessions collected from diverse geographical areas in Indonesia. A total of 107 alleles were detected among the accessions with 29 polymorphic markers. However, the mean of polymorphic information content (0.33) value and diversity index (0.38) value was indicative of low genetic diversity in this germplasm. The mungbean population structure was not clearly differentiated and the number of subpopulations was unclear. Neighbour-joining tree analysis revealed that the genetic cluster did not reflect the geographical origin of the accessions. Interestingly, the most agriculturally improved varieties were genetically similar to some landraces from one of the main mungbean-producing regions. These newly developed SSR markers could be useful for detecting genetic variability as a basis for establishing a conservation strategy for mungbean germplasm with the aim of enhancing Indonesian breeding programmes.

Keywords: genetic diversity; Indonesia; mungbean; SSRs

Introduction

Mungbean [*Vigna radiata* (L.) Wilczek] can be used in a wide range of food applications and is also an ideal nutritional complement that can contribute to a healthy and balanced diet, particularly in poor and anaemic women and children (AVRDC, 2002). The analysis of key agronomical traits, such as seed quality and disease susceptibility, highlights the importance of conserving diversity within the mungbean germplasm (Lambrides and Godwin, 2007; Chankaew *et al.*, 2011). Mungbean

development and breeding have been carried out for several decades; despite this, molecular genomic datasets are limited. Recently, next-generation Illumina HiSeq technology has allowed genome-wide marker development for mungbean, enabling further genomic studies (Van *et al.*, 2013).

In Indonesia, mungbean ranks second only to soybean as an economically significant legume. Mungbean landraces (local cultivars) are widely dispersed in several regions of different main islands, of which many are still cultivated by traditional farming despite their superior agronomical traits. The landraces are usually endowed with tremendous genetic variability, which could be useful for mungbean improvement (Jambormias *et al.*, 2003; Tripathy *et al.*, 2010). Moreover, genetic

*Corresponding author. E-mail: sukhalee@snu.ac.kr

differences between landraces and commercial varieties and the level of heterogeneity still existing in landraces are not yet well characterized. In this study, we estimated the genetic diversity of the Indonesian mungbean germplasm using a newly developed simple sequence repeat (SSR) marker analysis, with the aim of obtaining information of future importance for the selection and breeding of mungbean varieties with desirable traits.

Materials and methods

In this study, we selected 83 accessions from diverse mungbean-producing areas of Indonesia (Table S1, available online). The samples comprised 16 cultivars and 67 landraces and were obtained from the germplasm collection of the Indonesian Center for Agricultural Biotechnology and Genetic Resources Research and Development (ICA-BIOGRAD). Genomic DNA was extracted from young leaves using a standard cetyltrimethylammonium bromide (CTAB) method (Gelvin and Schilperoort, 1995). DNA samples were equalized for concentration and then PCR-amplified using 30 SSR primer sets that were developed based on sequence variations between two mungbean genomes (Sunhwanokdu and Gyeongjjaerae 5; Table S2, available online). PCR products were separated using the fluorescence-based capillary electrophoresis Fragment Analyzer CE System (Advanced Analytical Technologies, Inc., Ames, Iowa, USA). Population structure was estimated using the model-based program STRUCTURE (Version 2.3; Falush *et al.*, 2003). Polymorphic information content (PIC)

values were calculated for the total population and for SSR markers (Liu, 2001). Genetic diversity index and phylogenetic tree were generated based on Nei's method, and support for clusters was evaluated by bootstrap analysis of 1000 permutations using PowerMarker V3.25 (Felsenstein, 1985; Liu and Muse, 2005).

Results and discussion

Genetic diversity within crop collections is beneficial for their effective conservation and management and for crop improvement strategies (Mondini *et al.*, 2009; Vetriventhan *et al.*, 2012). The molecular evaluation of landraces carried out in the present study could be useful as an integral part of pre-breeding process. Of the 30 SSR markers, 29 exhibited polymorphism, in which a total of 107 alleles were detected and average per locus was 3.7, which is relatively comparable to that in other reports in mungbean (Kumar *et al.*, 2002; Gwag *et al.*, 2010). The PIC value, reflecting allele diversity for a particular marker, ranged from 0.012 (SSR_34) to 0.819 (SSR_33), with an average of 0.33. Diversity index for the full 29-marker set was in range of 0.012–0.833 with an average of 0.38. The mean PIC value and diversity index were indicative of low levels of genetic diversity in the tested mungbean germplasm. Interestingly, although mungbean is a self-pollinating crop, a number of loci displayed heterozygosity, probably due to residual heterozygosity in the collection as well as outcrossing in some cultivars (Blair *et al.*, 2009).




Fig. 1. Geographical distribution of Indonesian mungbean germplasm used in this study. Black dots indicate the collection sites of genotypes as listed in Table S1 (available online). The primary mungbean producers, Java and Nusa Tenggara, are indicated with dotted-line boxes.

Admixture model-based simulations for the inferred population structure were performed by varying K from two to ten iterations using all the 83 accessions, which were mostly collected from Java and Nusa Tenggara (Fig. 1). However, no clear number of subpopulations was determined in this analysis. All the F_{st} estimates revealed non-significant differences in the overall genetic structure, and the low F_{st} values clearly revealed low levels of differentiation in the mungbean population. The fact that the mungbean landraces evaluated in this study could not be discretely distributed by their regions of origin is in good agreement with a previous study in mungbean and common bean (Burlé *et al.*, 2010; Gwag *et al.*, 2010), which is probably due to high homogeneity among the accessions (Evanno *et al.*, 2005). To provide a comprehensive picture of genetic structure, however, a geographical broadly representative collection of mungbean landraces across Indonesian islands is needed.

A neighbour-joining tree with bootstrap analysis demonstrated differentiation among the accessions,

which largely grouped according to the individual genetic characteristics rather than geographical origin (Fig. 2 and Table S1, available online). All clusters consisted of a mix of landraces of different origins and varieties. The majority of accessions from the regions of Java (West, Central and East) and Nusa Tenggara (West and East) were dispersed in all the four main clusters, which is consistent with the diversity index. By contrast, most of the landraces from West Nusa Tenggara and East Nusa Tenggara dominated clusters I and IV, respectively, and were relatively closely grouped therein. Of the 16 improved varieties, 13 were genetically similar to local cultivars, most of which were from Java and fewer originated from Nusa Tenggara (cluster II). Indonesian varieties may be developed mainly with a genetic background of Java landraces, as supported by their closest genetic relationship (0.042) in comparison with other landrace populations from other regions. Additionally, Nusa Tenggara has a number of dry regions and this may have induced adaptation to drought tolerance in




Fig. 2. Neighbour-joining tree based on the distance matrix of 29 simple sequence repeat markers genotyped in 83 mungbean accessions estimated with a bootstrap analysis using 1000 permutations. Landraces and varieties are edge labelled with a serial number and are depicted in black and blue colours, respectively.

mungbean varieties from these regions (IAARD, 2012) and would explain the clustering of these accessions. The landrace from the region furthest west (Sumatra; Local Adnan Lubis) was genetically distant to those from eastern parts of Indonesia (North Maluku; Plastik and Local Pasar Jailolo). The diverse nature of these landraces may be a reflection of the prevalent diverse agroclimatic conditions of each region (Burle *et al.*, 2010), but dissemination and extensive distribution with their origin lead to a low association of germplasm with geographical diversity (Poehlman, 1991).

In summary, the phylogenetic clusters derived from our mungbean genotyping provide a strong basis for ongoing diversity analysis of local cultivars. Further data are required to build upon the initial dataset and refine the definition of fine structure within the mungbean collection, for example, agromorphological data collection that is underway. In addition, genetic differentiation between the mungbean accessions would be enhanced by the use of additional molecular markers developed from the genomes of mungbean or its close relatives. The collection of diverse local cultivars and their subsequent genotyping would enhance germplasm diversity and provide information, both of which are beneficial for developing collection strategies and breeding purposes with desirable agromorphological characteristics. Overall, these newly developed SSR markers could be useful for detecting genetic diversity, which presents an opportunity for the identification and characterization of loci responsible for important and desirable traits.

Supplementary material

To view supplementary material for this article, please visit <http://dx.doi.org/10.1017/S1479262114000343>

Acknowledgements

This study was supported by a grant from the Next-Generation BioGreen 21 Program (no. PJ008117) of the Rural Development Administration, Republic of Korea.

References

- Asian Vegetable Research and Development Centre (AVRDC) (2002) High yielding, MYMV-resistant mungbean lines for South Asia. *AVRDC Progress Report, Shanbua, Taiwan*. AVRDC Publication No. 02-542. Shanbua: AVRDC, p. 25.
- Blair MW, Diaz LM, Buendia HF and Duque MC (2009) Genetic diversity, seed size associations and population structure of a core collection of common beans (*Phaseolus vulgaris* L.). *Theoretical and Applied Genetics* 119: 955–972.
- Burle ML, Fonseca JR, Kami JA and Gepts P (2010) Microsatellite diversity and genetic structure among common bean (*Phaseolus vulgaris* L.) landraces in Brazil, a secondary center of diversity. *Theoretical and Applied Genetics* 121: 801–813.
- Chankaew S, Somta P, Sorajjapinun W and Srinives P (2011) Quantitative trait loci mapping of Cercospora leaf spot resistance in mungbean, *Vigna radiata* (L.) Wilczek. *Molecular Breeding* 28: 255–264.
- Evanno G, Regnaut S and Goudet J (2005) Detecting the number of clusters of individuals using the software structure: a simulation study. *Molecular Ecology* 14: 2611–2620.
- Falush D, Stephens M and Pritchard JK (2003) Inference of population structure: extension to linked loci and correlated allele frequencies. *Genetics* 164: 1567–1587.
- Felsenstein J (1985) Confident limit on phylogenies: an approach using the bootstrap. *Evolution* 39: 783–791.
- Gelvin SB and Schilperoort RA (1995) *Plant Molecular Biology Manual*. Norwell, MA: Kluwer Academic Publisher.
- Gwag JG, Dixit A, Park YJ, Ma KH, Kwon SJ, Cho GT, Lee GA, Lee SK, Kang HK and Lee SH (2010) Assessment of genetic diversity and population structure in mungbean. *Genes & Genomics* 32: 299–308.
- Indonesian Agency for Agricultural Research and Development (IAARD) (2012) Kacang hijau di lahan kering (Mungbean in dry land). Sinar Tani edition, 25–31 January, no. 3441: 8–9.
- Jambormias E, Madubun EL and Hitijanhubessy FJD (2003) Yield potential, natural genetic variability and qualitative characteristics heritability of local variety Jamdena. *Jurnal Pertanian Kepulauan* 2: 100–105.
- Kumar SV, Tan SG, Quah SC and Yusoff K (2002) Isolation of microsatellite markers in mungbean, *Vigna radiata*. *Molecular Ecological Notes* 2: 96–98.
- Lambrides CJ and Godwin ID (2007) Mungbean. In: Kole C (ed.) *Genome Mapping and Molecular Breeding in Plants, Vol. 3: Pulses, Sugar and Tuber Crops*. Berlin/Heidelberg: Springer, pp. 69–90.
- Liu J (2001) PowerMarker V3.25 Manual. Available at: <http://www.powermarker.net>
- Liu K and Muse SV (2005) PowerMarker: integrated analysis environment for genetic diversity in core collection accessions of wild barley, *Hordeum vulgare* ssp. *Spontaneum*. *Hereditas* 136: 67–73.
- Mondini L, Noorani A and Pagnotta MA (2009) Assessing plant genetic diversity by molecular tools. *Diversity* 1: 19–35.
- Poehlman J (1991) *The Mungbean*. New Delhi: Oxford and IBH Publishing Co, pp. 195–222.
- Tripathy SK, Suchinnata S, Lenka D and Sahoo S (2010) Genetic diversity of mungbean [(*Vigna radiata* (L.) Wilczek)] genotypes based on SDS-PAGE albumin seed storage protein. *Legume Research – An International Journal* 33: 54–57.
- Van K, Kang YJ, Han K-S, Lee Y-H, Gwag J-G, Moon J-K and Lee S-H (2013) Genome-wide SNP discovery in mungbean by Illumina HiSeq. *Theoretical and Applied Genetics* 126: 2017–2027.
- Vetriventhan M, Upadhaya HD, Anandakumar CR, Senthilvel S, Parzies HK, Bharathi A, Varshney RK and Gowda CLL (2012) Assessing genetic diversity, allelic richness and genetic relationship among races in ICRISAT foxtail millet core collection. *Plant Genetic Resources: Characterization and Utilization* 10: 214–223.